

Animal Health Research Institute,  
Assiut, Regional Laboratory.

**OCCURRENCE OF *YERSINIA ENTEROCOLITICA*  
AND *AEROMONAS HYDROPHILA*  
IN PASTEURIZED MILK IN SOHAG CITY**  
(With 3 Tables)

By  
**SOHAIR, Z. HUSSEIN and EMAN K. AHMED**  
(Received at 25/12/2002)

تواجد ميكروب اليارسينيا أنتيروكوليتكا والأيرومونات هيدروفيليا  
في اللبن المبستر في مدينة سوهاج  
سهير زين العابدين ، إيمان قرشي أحمد

أجريت هذه الدراسة لمحاولة عزل ميكروبي اليارسينيا أنتيروكوليتكا والأيرومونات هيدروفيليا من خمسين عينة من اللبن المبستر جمعت عشوائياً من أكشاك الألبان والسوبر ماركت بمدينة سوهاج وقد أسفرت النتائج على أن ميكروب اليارسينيا أنتيروكوليتكا أمكن عزلها بنسبة ٤%. أما ميكروب الأيرومونات هيدروفيليا أمكن عزله بنسبة ١٦% [عترات الأيرومونات هيدروفيليا ٦%، عترات الأيرومونات كافي ٨%، وعترات الأيرومونات سوبريا ٢%]. وعند إجراء اختبار حساسية البكتريا المعزولة للمضادات الحيوية المختلفة، كانت عترات اليارسينيا أنتيروكوليتكا المعزولة حساسة للمركبات (تتراسيكلين، ستربتومايسين، كلورامفينيكول والسلفا ميزوكسازول) وكانت مقاومة للمركبات (الأمبيسلين والنورفلوكساسين). أما عترات الأيرومونات هيدروفيليا المعزولة فكانت أكثر حساسية للمركبات (كلورامفينيكول، سلفاميزوكسازول، وتتراسيكلين) ولكنها كانت مقاومة للمركبات (الأمبيسلين، النورفلوكساسين والنيوميسين).

**SUMMARY**

In Sohag governorate a trial was made for isolation of *Y. enterocolitica* and *A. hydrophila* from fifty random samples (400 ml each) of pasteurized milk collected from different supermarkets. *Y. enterocolitica* recovered from 4% of pasteurized milk samples. *Y. enterocolitica* isolates were susceptible to Tetracycline (TE<sub>30</sub>); Streptomycin (S<sub>10</sub>); Chloramphenicol (C<sub>30</sub>) and Sulphamethoxazole (SXT<sub>25</sub>) and resistant to Ampicillin (AM<sub>10</sub>) and Norfloxacin (NOR<sub>10</sub>). *A. hydrophila* group isolated from pasteurized milk were *A. caviae* (8%); *A. sobria* (2%) and *A. hydrophila* (6%). All isolates

were highly susceptible to chloramphenicol (C<sub>30</sub>); Sulphamethoxazole (SXT<sub>25</sub>) and Tetracycline (TE<sub>30</sub>) but resistant to Ampicillin (AM<sub>10</sub>); Norfloxacin (NoR<sub>10</sub>) and Neomycin (N<sub>30</sub>).

**Keywords:** *Yersinia enterocolitica*, *Aeromonas hydrophila*, pasteurized milk.

## INTRODUCTION

*Yersinia enterocolitica* and *Aeromonas hydrophila* are currently considered to be of great importance as a foodborn pathogen. Symptoms in human are characterized by diarrhea and abdominal pain in primary stage, but a number of complications or secondary symptoms may also occur (Vidon and Delmas, 1981; Abeyta *et al.*, 1986; Walker, 1987; Stelma, 1988; Jama, 1989; Varnam and Evans, 1991 and Palumbo *et al.*, 1992).

The organisms have psychrotrophic nature (able to grow and multiply to large population at normal refrigeration temperature) and they have the ability to produce toxins in the food kept at refrigerator temperature. (Boyce *et al.*, 1979; Stern *et al.*, 1980; Kapperud and Langeland, 1981; Turnbull *et al.*, 1984; Okren *et al.*, 1987; Palumbo *et al.*, 1989; Varnam and Evans, 1991 and Freitas *et al.*, 1993; and Kirov *et al.*, 1993. In 1982, Lovett *et al.*, mentioned that, pasteurization process of dairy products recommended by FDA are adequate to destroy large concentration of *Y. enterocolitica*. Moreover they found that the presence of *Y. enterocolitica* in pasteurized milk may probably result from substandard processing or recontamination after pasteurization. Several investigators could isolated *Y. enterocolitica* from pasteurized milk (Hughes, 1980; Bimet, 1983; Moustafa *et al.*, 1983; Jamshidian and Babakhani, 1999). Contamination of pasteurized milk even by one organism of *Y. species* could lead to high numbers of such organism during cold storage of milk (Bimet, 1983).

*Y. enterocolitica* could not survive after pasteurization temperature (62.8 for 30min) of brain heart infusion broth, skim milk and whole milk but could be recovered in few number in peptone sorbital bile broth after 8-10 days of incubation at 10°C (Kushal and Anand, 1999).

Motile *Aeromonas species* associated with human illness are *A. hydrophila*, *A. caviae* and *A. sobria*, (Stelma, 1989). Cousin in 1982, found that several strains of *Aeromonas species* have been isolated from pasteurized milk and the organism increased in

number during cold storage leading to deterioration of milk, while, Tibana *et al.* (1987) stated that, the presence of *Aeromonas species* in white cheese may be due to contaminated pasteurized milk used for production of such product or unsanitary conditions during manufacturing and storage. Several investigators could isolate *A. species* from the pasteurized milk (Drew and Greendway, 1990; Freitas *et al.*, 1993; Kirov *et al.*, 1993; Bahout, 1997 and El-Gamal, 1997) however Melas *et al.* (1999) mentioned that, the *Aeromonas species* could not be detected in the examined pasteurized milk.

The present study was carried out to investigate the incidence of *Yersinia enterocolitica* and *Aeromonas hydrophila* in pasteurized milk in Sohag city and to study their in vitro sensitivity to different antibacterial agents.

### **MATERIALS and METHODS**

Fifty random pasteurized milk samples (400ml each) were collected from different dairy shops and supermarkets in Sohag city, Egypt. The collected samples were transferred to the laboratory with a minimum of delay to be examined bacteriologically for the presence of *Y. enterocolitica* and *A. hydrophila*.

#### **Isolation:**

##### **(I) *Y. enterocolitica*:**

Trypticase soy broth tubes were prepared and each tube was inoculated with 1 ml of pasteurized milk samples. The inoculated enrichment broth tubes were incubated at 4°C for 14 days (Greenwood and Hooper, 1989).

Loopfuls from the incubated enrichment broth were streaked directly onto Cefsulodin Irgasan Novobiocin (CIN) agar. The plates were incubated at 35°C for 24h. The suspected organisms were identified morphologically and biochemically according to (Schiemann and Devenish, 1982) (urea hydrolysis, sugar fermentation, kligler iron agar and motility indol lysin sulphide).

##### **(II) *A. hydrophila*:**

One ml of each sample of pasteurized milk was added to 10 ml of sterile tryptone soya broth containing 10 µ/ml of Ampicillin in sterile tubes. The inoculated enrichment broth tubes incubated at 28°C for 24h.

A loopful from the incubated enrichment broth was streaked directly onto starch ampicillin agar and cefsulodin irgasan Novobiocin (CIN) agar, then the plates were incubated at 28°C for 24h.

The presumptive aeromonas colonies were subcultured onto nutrient agar slants and incubated at 28°C for 48h. Suspected organism was identified morphologically and biochemically according to Popoff and Veron (1976) (Oxidase test strips, motility test, esculin hydrolysis, gas production from glucose, gelatin liquefaction test and indol production test).

**In vitro sensitivity test:**

The antibacterial agents used in this study are described in Table 3.

**RESULTS**

The obtained results are recorded in Tables 1, 2, 3.

**Table 1:** Incidence of *Yersinia enterocolitica* and *Aeromonas hydrophila* in the examined pasteurized milk samples.

Number of Samples	<i>Yersinia enterocolitica</i>		<i>Aeromonas hydrophila</i>	
	Positive samples		positive samples	
	No.	%	No.	%
50	2	4%	8	16%

**Table 2:** Incidence of *Aeromonas hydrophila* group in examined pasteurized milk samples.

A.hydrophila group	No. of positive samples	%
<i>A. hydrophila</i>	3	6%
<i>A. caviae</i>	4	8%
<i>A. sobria</i>	1	2%
<b>Total</b>	<b>8</b>	<b>16%</b>

**Table 3:** In vitro sensitivity testing of *Y. enterocolitica* and *A. hydrophila*.

Antimicrobial discs	<i>Y. enterocolitica</i>	<i>A. hydrophila</i>
- Streptomycin (S <sub>10</sub> )	S	I
- Amicillin (AM <sub>10</sub> )	R	R
- Sulphamethoxazole (SXT <sub>25</sub> )	S	S
- Chloramphenicol (C <sub>30</sub> )	S	S
- Tetracycline (TE <sub>30</sub> )	S	S
- Cefotaxime (CTX <sub>30</sub> )	I	I
- Norfloxacin (NOR <sub>10</sub> )	R	R
- Neomycin (N <sub>30</sub> )	I	R
- Colistin sulphate (CT <sub>25</sub> )	I	I

S: Susceptible

I: Intermediate.

R: Resistant.

## DISCUSSION

The results recorded in Table 1 revealed that, 2(4%) of the examined pasteurized milk samples were positive for *Y. enterocolitica*, these finding simulate those reported by El-sherbini (1992) while they were lower than those reported by Delams and Vidon (1982) and higher than those recorded by Moustafa *et al.* (1983) Jamshidian and Babakhani (1999).

In vitro, all isolates were susceptible to Tetracycline (TE<sub>30</sub>); Streptomycin (S<sub>10</sub>); Chloramphenicol (C<sub>30</sub>) and Sulphamethoxazole (SXT<sub>25</sub>) and resistant to Ampicillin (AM<sub>10</sub>) and Norfloxacin (NOR<sub>10</sub>).

These results were similar to those recorded by Vidon, and Delmas (1981) Saad and Moustafa (1989) and quite different from those obtained by Adly and Yacoub (1995).

The results recorded in Table 1 revealed that, 8(16%) out of 50 pasteurized milk samples were positive for *Aeromonas species*. These result, were lower than that reported by Freitas *et al.* (1993) and Kirov, *et al.* (1993) and higher than those recorded by Chen *et al.* (1988); Drew and Greenaway (1990) and El Gamal (1997) but disagree with those recorded by Melas *et al.* (1999) who could not isolate *A. species* from pasteurized milk.

*Aeromonas hydrophila*; *A. Caviae* and *A. sobria* could be isolated in a percentages of 6%, 8%, and 2% of examined samples respectively. These results were nearly similar to that reported by Bahout *et al.* (1997) [*A. hydrophila* (9.4%), *A. caviae* (11.8%); *A. sobria* (3.5%)], and simulated that reported by Freitas *et al.* (1993) in case of *A. sobria* (2.5%) but lower in case of *A. hydrophila* (12.8%) and *A. caviae* (58.9%).

*Aeromonas* organisms are sensitive to temperatures above 48°C, therefore, pasteurized milk are expected to be free of *Aeromonas*. The presence of such organisms in pasteurized milk samples could be attributed to post pasteurization contamination. Palumbo *et al.* (1989). *A. sobria* was more virulent in the cytotoxic activity than the *A. caviae* and *A. hydrophila* (Janda *et al.*, 1984). In vitro, all isolates were susceptible to chloramphenicol (C<sub>30</sub>); sulphamethoxazole (SXT<sub>25</sub>) and tetracycline (TE<sub>30</sub>) and resistant to ampicillin (AM<sub>10</sub>); Norfloxacin (NOR<sub>10</sub>) and Neomycin (N<sub>30</sub>). These results were similar to those recorded by Molero *et al.* (1989)

but were quite different from those obtained by Dixon, and Voran (1992) and El- Gamal (1997).

## CONCLUSION

From this study it can be concluded that, some pasteurized milk samples were contaminated by *Y. enterocolitica* and *Aeromonas species* and this may reflect the lack of hygienic supervision and poorly sanitized processing equipment.

## REFERENCES

- Abeyta, C.Jr.; Kaysner, C.A.; Wekell, M.M; Sullivan, J.J and Stelma, G.N. (1986): Recovery of *Aeromonas hydrophila* from oysters implicated in an outbreak of foodborness. J. Food Prot., 49: 643-646.
- Adly, Y.H and Yacoub, Th.K. (1995): Prevalence and Characterization of *Yersinia enterocolitica* isolated from raw milk and ice cream in Minia city. J. Egypt. Vet. Med. Assoc. 55, 1,2: 59-66.
- Bahout, A.A. (1997): Incidence of *Aeromonas species* in pasteurized milk. Alex. J. Vet. Sci., 13 : 31-34.
- Bimet, F. (1983): Survival of *Y. enterocolitica* in milk. Technique laitiere. 977 : 43-45, 47-51. Dairy Sci. Abst. 46, 6, 409, 1984.
- Boyce, J.M.; Evans, D.J. Evans, D.G. and Dupont H.L. (1979): Production of heat stable, methanol soluble enterotoxin by *Y. enterocolitica*. Infect immun. 25 : 532-537 Environmental Bacteriology, Vol. 32, No. 2.
- Chen, I.H. Tsai, S.J.; Change, R.M.; Fang, S.W.; Wang, J.Y. and Liou, C.M. (1988): Microbial Survey of Raw and pasteurized milk. Food Sci., China, 15: 73-80.
- Cousin, M.A. (1982): Presence and activity of psychrotrophic microorganisms in milk and dairy products : A review J. Food Prot., 45 : 172-207.
- Delmas, C.L. and Vidon, D.J.M. (1982): Contamination of milk in Alsace by *Y. enterocolitica*. Lait. 62 (621/622): 288-704. Dairy Sci. Abst. 45, 2, 251-259.
- Dixon, B.A. and IVoran, G.S. (1992): The activity of ceftiofur sodium for *Aeromonas spp.* isolated from ornamental fish. Department of Biological Sciences, California State University, Hayward. J. Wildl Dis. 1992 Jul: 28 (3): 453-456.

- Drew, P.G and Greenaway, C. (1990):* Incidence of *Aeromonas* species in pasteurized milk in Victoria, Australia. XX111 International Dairy Congress, Montreal October 8-12, Vol. I. Brusseis, Belgium, International Dairy Federation, 122 : (219) Dairy Sci., Abst. (1991), Vol 53, No. 7.
- El- Gamal, A.M. (1997):* Incidence of motile *Aeromonas* species in raw milk and some dairy products: Vet. Med. J. Giza, 45 : 553-560.
- El- Sherbini, M. (1992):* Prevalence of *Y. enterocolitica* in pasteurized milk and cream. In proceeding of the 3<sup>rd</sup> world Congress on foodborne infections and intoxication, Berlin, Germany, 16-19 June, Vol. 1.
- Freitas, A.C.; Nunes, M.P.; Milhomem, A.M and Ricciardi, I.D. (1993):* Occurrence and characterization of *Aeromonas* species in pasteurized milk and white cheese in R10. De. Janeiro, Brazil. J. Food. Prot. 56: 62-65.
- Greenwood, M.H. and Hooper, W.L. (1989):* Improved methods for the isolation of *Yersinia* species from milk and foods. Food Microbiol. 41 (3): 99-104.
- Hughes, D. (1980):* Repeated isolation of *Y. enterocolitica* from pasteurized milk in a holding vat. at a dairy factory. J. of Appl. Bacteriol. 48 (3) : 383-385.
- Jama, J. (1989):* A multistate outbreak of infections caused by *Y. enterocolitica* transmitted by pasteurized milk. J. Am. Med. Ass. 27, Vol. 25 (4) 438-488.
- Jamshidian, M. and Babakhani, A. (1999):* Investigation of *Y. enterocolitica* in raw and pasteurized milk in khozestan area. J. of the Faculty of Vet. Med. University of Tehran 54 (1) 8-10.
- Janda, J.M.; Brenden, R. and Botton, E. J. (1984):* Differential susceptibility to human serum by *Aeromonas* spp. Curr. Microbiol. 11. 325-328.
- Kapperud, G. and Langeland, G. (1981):* Enterotoxin production at refrigeration temperature by *Y. enterocolitica* like bacteria. Curr. Microbiol. 5 : 119-122.
- Kirov, S.M.; Hui, D.S. and Hayward, L.J. (1993):* Milk as a potential source of *Aeromonas* gastrointestinal infection. J. Food. Prot. 56 : 306-312.
- Kushal, R. and Anand, S.K. (1999):* Repair and recovery of thermally injured cells of *Y. enterocolitica* in milk. Dairy microbiology Division, National Dairy Research Institute, Karnal. India. J. Food. Prot. 62 (10): 1203-1205.

- Lovett, J.; Bradshaw, J.G. and Peeler, J.T. (1982): Thermal inactivation of *Yersinia enterocolitica* in milk. App. Environ. Microbiol. 44 : 517-519.
- Melas, D.S.; Papageorgiou, D.K. and Mantls, A.L. (1999): Enumeration and confirmation of *Aeromonas hydrophila A.caviae* and *A.sobria* isolated from raw milk and other milk products in Northern Greece Department of F. hygiene and Technology, School of Vet. Med. Aristotle University of Thess a Ioniki, Greece. J. Food Prot, 62 : 5, 463-466.
- Molero, X; Bartolome, R.M; Vinuesa, T; Guarner, L.; Accarino, A; Casellas, F. and Garcia, R. (1989): Acute gastroenteritis due to vibrio parahaemolyticus in Spain. (presentation of 6 cases) Med. Clin. Barc. Jan 14; 92 (1): 1-4.
- Moustafa, M.K.; Ahmed, A.A. and Marth, E.H. (1983): Occurrence of *Y.enterocolitica* in raw and pasteurized milk. J. Food Prot. 46 (4) : 276-278.
- Okren, A.J.; Rose, B.E. and Bennet, B. (1987): Incidence and toxigenicity of *Aeromonas species* in retail poultry, beef and pork. J. Food, Prot. 50: 509-513.
- Palumbo, S.; Abeyta, C. and Stelma, G.N. (1992): *Aeromonas hydrophila* group Inc. Vanderzant and F.Splittstoesser (ed.). Compened methods for the microbiological examination of foods. 3rd American public health association, Washington. D.C.
- Palumbo, S.A.; Bencivengo, M.M.; Corral, F.D.; Williams, A.C. and Buchanan, R.L. (1989): Characterization of *Aeromonas hydrophila* group isolated from retail foods of animal origin. J. Clin. Microbiol., 27 : 854-859.
- Popoff, M. and Veron, M. (1976): A taxonomic study of the *Aeromonas hydrophila*. *Aeromonas panctata* group. J. Gen. Microbiol., 94: 11-22.
- Saad, N.M. and Moustafa. S. (1989): Prevalence of *Yersinia enterocolitica* in raw milk in Assiut city. Department of food hygiene, Faculty of Vet. Med. Assiut. Assiut. Vet. Med. J. Vol. 22 (43) : 95-99.
- Schiemann, D.A. and Devenish, J.A. (1982): Relationship of Helacell in activity to biochemical, Serological and virulence characteristics of *Y.enterocolitica* Infect. Immun. 35 : 497-506.
- Stelma, G.N., Jr. (1988): Virulence factors associated with pathogenicity of *Aeromonas* isolates. J. Food safety 9(1) 1-9.



- Stelma, J.R. (1989): Aeromonas hydrophila.* Doyle, MP (ed) foodborne bacterial pathogens. PP. 1-19. New York, Marcel Dekler.
- Stern, N.J.; Pierson, M.D. and Kotula, A.W. (1980):* Effect of PH and Sodium chloride on *Y. enterocolitica* growth at room and refrigeration temperature J. Food Sci. 45: 64-97.
- Tibana, A; Warnken, M.B.; Nunes, M.P. Ricclardi, I.D. and Noieto, A.L.S. (1987):* Occurrence of *Yersinia species* in raw and pasteurized milk in Rio de Janeiro, Brazil J. Food Prot., 50 : 580-583.
- Turnbull, P.C.B.; Lee, J.V.; Miliotis, M.D.; Van De walle, S.; Koornhof, H.J.; Jeffery, L. and Bryant, T.N. (1984):* Enterotoxin production in relation to taxonomic grouping and source of isolation of *Aeromonas species*. J. Clin. Microbiol., 19 : 175.
- Varnam, A.H. and Evans, M.G. (1991): Aeromonas.* In foodborne pathogens. Wolfe publishing Ltd. London.
- Vidon, D.J.M. and Delmas, C.L. (1981):* Incidence of *Yersinia enterocolitica* in raw milk in Eastern France. Applied and Environmental Microbiol. 41 (2): 355-359.
- Walker, S.J. (1987): Yersinia enterocolitica* A review of foodborn pathogen technical Bulletin No. 63 CFRRRA Chipping Campden.