

Dept. of Animal Hygiene & Zoonoses.
Faculty of Veterinary Medicine, Assiut University

**OCCURRENCE OF *E. COLI* O157:H7 IN
APPARENTLY HEALTHY DAIRY CATTLE AND
RETAIL MILK.**

(With 3 Tables and 1 Figures)

By

AMAL S.M. SAYED AND ASMAA A.A. HUSEIN

(Received at 29/3/2003)

**مدى تواجد ميكروب الايشريشيا كولاي O157:H7 في الأبقار الحلوب
السليمة ظاهريا و الألبان المتداولة في الأسواق**

أمل سيد ، أسماء حسين

إن المخاطر التي يتعرض لها الإنسان بعدوى الايشريشيا كولاي O157:H7 نتيجة لتناول اللبن الأبقار الخام لا تزال في ازدياد و تظهر عادة على هيئة حالات فردية أو في صورة اوبئة. و حيث أن لبن الأبقار يعتبر من المصادر الهامة لنقل هذا الميكروب، لذلك فقد أجريت هذه الدراسة لتحديد دور الأبقار الحلوب السليمة ظاهريا كمصدر للعدوى بهذا الميكروب. وقد تم جمع عدد ١١٠ عينة بواقع ٨٠ عينة من الأبقار السليمة ظاهريا (٤٠ عينة ألبان و ٤٠ عينة براز) هذا بالإضافة إلى عدد ٣٠ عينة من الألبان المعروضة للبيع في مناطق مختلفة من محافظة أسيوط و ذلك في الفترة من شهر يونيو إلى سبتمبر ٢٠٠٢. و قد أسفرت النتائج عن وجود هذا الميكروب في العينات المختبرة بنسبة ١٢,٧٣% . و قد تم عزله من 12 (30%) ، 1 (2.5%) و 1 (3.33%) من عينات البراز و الألبان و الألبان المعروضة للبيع على التوالي. و قد تم عزل الميكروب من عينات البراز فقط بنسبة ٢٧,٥% و عينات الألبان و البراز من نفس الحيوان بنسبة ٢,٥%. و قد تبين أن الأبقار السليمة ظاهريا تلعب دور هام كمصدر لميكروب الايشريشيا كولاي O157:H7 مما يزيد من احتمال تلوث الألبان و الذي يشكل خطورة على صحة الإنسان.

SUMMARY

Risk of acquiring verocytotoxin *E.coli* infections through ingestion of raw cow's milk continues to increase causing either sporadic cases or outbreaks. During June-September 2002, one hundred and ten samples were screened for the presence of verocytotoxin-producing *E.coli* (VTEC). Both milk samples (40) and fecal swabs (40) were collected from the same apparently healthy cows in dairy farms, moreover thirty

samples of retail milk marketed in Assiut province were collected. Verocytotoxin-producing *E.coli* was isolated from 14 (12.73%) of 110 samples after enrichment in mTSB (modified tryptic soya broth) supplemented with novobiocin, plating onto SMAC (sorbitol MacConkey agar) and then were identified using latex agglutination test. *E.coli* O157:H7 strains were confirmed by using polymerase chain reaction to detect hyl A gene (target of *E.coli* O157:H7). Verocytotoxin producing *E. coli* was recovered from 12 (30%), 1 (2.5%) and 1 (3.33%) of fecal swabs, milk samples and retail milk, respectively. *E. coli* O157:H7 was isolated from 27.5% fecal swabs only and 2.5% of both milk and feces of the same animal, however we could not isolate the organism from the milk samples only. It was concluded that apparently healthy dairy cows may act as reservoirs of verocytotoxin-producing *E.coli* with fecal shedding representing a risk factor of milk contamination which pose a public health hazard.

Key words: *E.coli* O157 : H₇ – Dairy cattle – Milk – PCR..

INTRODUCTION

E.coli O157:H7, is a new emerging, clonally distinct form of *Escherichia coli* was first identified as a significant food-borne zoonotic pathogen (Riley *et al.*, 1983). Verocytotoxin-producing *E.coli* O157:H7 (VTEC O157) has been emerged as a major food-borne pathogen which has the most sever effect on young children and the elderly (Griffin, 1995 and Feng *et al.*, 1998).

Although over 150 different OH serotypes of verocytotoxin producing *E.coli* (VETC) have been associated with human illness, the vast majority of reported outbreaks and sporadic cases of VTEC infections in humans have been associated with serotype O157:H7 causing a range of symptoms from mild non bloody diarrhoea to haemorrhagic colitis, life threatening haemolytic uremic syndrome and thrombocytopenic purpura through the production of the shiga-like toxins and other probable virulence factors. (Riley *et al.*, 1983; Slutsker *et al.*, 1997 and Nelson *et al.*, 1998).

Epidemiological investigations have revealed that dairy cattle, especially young animals are the principal reservoirs of *E.coli* O157:H7 which can be excreted by healthy carriers and diarrhoeic cattle (Hancock *et al.*, 1994; Zhao *et al.*, 1995; Porter *et al.*, 1997 and Clarke *et al.*, 1998). Most reported human cases are associated with contaminated

food of bovine origin especially raw milk and milk products (Griffin, 1995).

This study aimed to determine the existence of *E.coli* O157:H7 in feces and milk of apparently healthy cows as well as in the retail milk. At the meantime, we applied the polymerase chain reaction on the isolated strains to detect hly A target gene of *E.coli* O157:H7.

MATERIALS and METHODS

1. Samples collection:

One hundred and ten samples including individual cow's milk (40) and fecal swabs (40) of apparently healthy cows from dairy farms as well as retail milk (30) were collected from Assiut Province during the period from June to September 2002. Samples were transferred to the laboratory in an icebox for bacteriological examination.

2. Enrichment Technique:

Milk and fecal samples were enriched in modified Tryptic Soya Broth (mTSB) supplemented with novobiocin (20 mg/liter). The inoculated broth was incubated at 37 °C for 24 hours (De Boor and Heuvelink, 2000).

3- Isolation on Sorbitol MacConkey agar:

Loopful from the incubated broth was streaked onto Sorbitol MacConkey agar plates and incubated at 37 °C for 24 hours (De Boor and Heuvelink, 2000).

4- Identification of *E.coli* O157:H7:

4.1- Identification of *E.coli* :

Non sorbitol fermenter colonies were identified morphologically by Gram's stain and biochemically as *E.coli* according to Varnam and Evans, (1991) by the conventional IMViC(indole, methyl red, Voges Proskauer and Citrate utilization) and Triple sugar iron agar.

4.2- Latex agglutination test:

A latex agglutination test (*E.coli* O157, Oxoid diagnostic reagents 620 M) for identification of *E.coli* serogroup O157 had been done on 18 isolates of the *E.coli* culture. The Oxoid *E. coli* O157 latex was demonstrated by slide agglutination of *E.coli* strains possessing the O157 serogroup antigen (Borczyk *et al.*, 1987, Krishnan *et al.*, 1987 and Vernozy-Rozand, 1997). For each test sample, bacterial colonies were screened by a rapid latex test to ascertain their O157 infection status. If the test was positive, the suspected bacterial colony was subcultured

onto trypticase soya broth and incubated overnight at 37 °C for DNA extraction and amplification.

4.3- Detection of the E.coli O157:H7:

A. DNA extraction:

Nucleic acid was extracted from the trypticase soya broth culture as described by Pitcher *et al.* (1989), Boom *et al.* (1990) and Fathi *et al.* (2002) by using the chaotropic agent Guanidium thiocyanate (GuSCN).

B. PCR procedure:

Amplification was performed in a total volume of 25 µl of DNA extracts, 25 pmol of each primer (*E.coli* O157-3 & O157-4) according to Wang *et al.*, 1997), and PCR mixture (Ready To. Go™ PCR Beads, Amersham Pharmacia Biotech, Austria). The PCR was conducted in a Biometra thermal Cycler (Biometra-Germany). The amplification condition was one cycle of 94 °C for 15s, then 35 cycles of 94 °C for 3s, 50 °C for 10s and 74°C for 35s, and finally one cycle of 74 °C for 2 min. and 45 °C for 2s. The PCR products (10-15 µl of each) were separated by electrophoresis in 2% agarose gels containing ethidium bromide (1 µg/ml), and visualized under UV illumination.

RESULTS

Table 1: Existence of Non-sorbitol fermenters *E.coli* in Dairy cows

Source of samples	No. of samples	NSF <i>E.coli</i> Strains	
		No.	%
Feces	40	15	37.5
Milk	40	2	5
Retail milk	30	1	3.33
Total	110	18	16.36

*. NSF = Non Sorbitol fermenter

Table 2: Surveillance of verocytotoxin *E.coli* in Dairy cows.


Source of samples	No. of samples	<i>E.coli</i> O157:H7	
		No.	%
Feces	40	12	30
Milk	40	1	2.5
Retail milk	30	1	3.33
Total	110	14	12.73

Table 3: Occurrence of verocytotoxin *E.coli* in feces and milk of Dairy cows.

No. of isolated <i>E.coli</i> O157:H7	Feces only		Milk only		Milk & Feces	
	No.	%	No.	%	No.	%
		11	27.5	-	-	1

Fig 1: PCR examination of *E.coli* O157:H7

E.coli O157:H7 hly A gene O157-3, GTAGGGAAGCGAACAGAG and O157 4, AAGCTCCGTGTGCCTGAA (Amersham Pharmacia Biotech, Austria). The size of amplicon product (bp) ranged from 300-600 in this study.



PCR amplification of DNA extracted from *E.coli* O157 samples in a 2% agarose gel. Lane (M) is 100 bp DNA size marker (100 Base-Pair Ladder, Amersham Pharmacia Biotech); Lane (C) is PCR assay reagent control, Lane 1 to 10 are the positive isolates for *E.coli* O157:H7 of the 14 positive samples.

DISCUSSION

Dairy cattle have been implicated as a major reservoir of verocytotoxin *Escherichia coli* (Wells *et al.*, 1991). It is noteworthy that *E.coli* O157:H7 are shedded in feces of either healthy carriers or diarrhoeic cattle (Clarke *et al.*, 1998). Unlike typical *E.coli*, isolates of *E.coli* O157:H7 do not ferment sorbitol, therefore this criteria is commonly used for selective isolation of *E.coli* O157:H7 (Tarr *et al.*, 1997). It was recently documented that for better isolation of *E.coli* O157:H7 from milk and feces, enrichment in mTSB (modified tryptone soya broth) supplemented with novobiocin is recommended, then plating onto SMAC (Sorbitol MacConkey agar) (De Boor and Heuvelink, 2000). Furthermore, Johnson *et al.* (1995) suggested that mTSB plus novobiocin enhances the availability of O157 antigen.

Non sorbitol fermenter (NSF) *E.coli* strains were isolated from 15 (37.5%) of 40 cow's feces, 2 (5%) of cow's milk and 1 (3.33%) of 30 retail milk samples as illustrated in Table 1. *E.coli* strains were typed as *E.coli* O157 by using a latex agglutination test (*E.coli* O157, Oxoid diagnostic reagents 620 M). *E.coli* O157 were confirmed as *E.coli* O157:H7 by using polymerase chain reaction to detect hyl A gene (target of *E.coli* O157:H7). In this paper we used Guanidium thiocyanate (GuSCN) in the purification and detection of DNA because of its potency to lyse cells combined with its potency to inactivate nuclease (Thompson & Gillespie, 1987 and Boom *et al.*, 1990).

Polymerase Chain Reaction (PCR) techniques could be developed as a routine procedure for food borne pathogen detection (Wang *et al.*, 1997). In our study, fourteen isolates were positive by using PCR technique (Fig. 1). This result agreed with that mentioned by Wang *et al.* (1997) who recorded that multiple DNA bands might be observed in the PCR detection from pure culture. The obtained results were comparable to that reported by Holland *et al.* (2000) who used the multiplex PCR for detection of *E.coli* O157:H7. McDonough *et al.* (2000), justify the using PCR techniques as an important tool for the detection of shiga-like toxins producing *E.coli*.

Verocytotoxin *E.coli* (*E.coli* O157:H7) was recovered from 12 (30%) of the examined cow's feces (Table 2). Geue *et al* (2002) reported variable rates of *E.coli* O157:H7 recovery in feces of cows ranged from 29-82%. However lower rates of verocytotoxin *E.coli* prevalence in cow's feces were detected by Chapman *et al.* (1993b); Chapman *et al.* (1997); Mechie *et al.* (1997); Dutta *et al.*(2000); Abdel-Khalek *et al.*

(2001); Lahti *et al.* (2001) and Nielson *et al.* (2002) who estimated a prevalence rate of 9.4%, 16.1%, 14%, 1.7%, 3%, 1.31% and 21%, respectively. On the contrary, Wells *et al.* (1991) failed to isolate *E.coli* O157:H7 from feces of milking cows. These differences in the estimated prevalence rates could be attributed to epidemiological variation in the different geographical localization.

E.coli O157:H7 was detected in 1 (2.5%) of the examined farm cow's milk (Table 2). Higher prevalence rate (10%) was estimated in individual cow's milk by Chapman *et al.* (1993 a), while Abdel-Khalek *et al.* (2001) reported a slightly lower prevalence with a rate of 2%. The variation in the detected prevalence rates in milk may be due to the difference in sources of contamination. It was reported that the risk of acquiring *E.coli* O157:H7 infections through ingestion of raw milk continues to increase (Ryser, 1998) causing either sporadic cases or outbreaks of VTEC infection (Griffin and Tauxe, 1991 and CDC, 1993). We could able to recover *E.coli* O157:H7 from 1 (3.33%) of the examined retail milk samples (Table 2). Our results are nearly similar to those detected by Abdel-Khalek *et al.* (2001) who estimated a prevalence rate of 3%. Results in Table 3 declared that verocytotoxin *E.coli* could be recovered from 11 (27.5%) of fecal samples only. The high percent of recovery of verocytotoxin *E.coli* in the examined feces is of particular concern indicating a major risk of milk contamination probability that pose a health hazard. Although not currently recognized as a cause of mastitis, presence of verocytotoxin *E.coli* in milk usually indicates contamination from feces of the infected or carrier animals (D'Aoust, 1989). This criteria explains both the ability to isolate the organism from both milk and feces of the same animal (with a rate of 2.5%) as well as the inability to recover verocytotoxin *E.coli* from milk samples only, as illustrated in Table 3.

It is concluded that dairy cows may act as carriers of verocytotoxin *E.coli* with fecal shedding which act as a major source of milk contamination. Moreover, it has been isolated from both farm raw milk and retail milk which act as a major risk factor for consumers that can lead to life threatening sequelae especially in young children and the elderly (Feng *et al.*, 1998). Indeed, this situation needs to be reviewed regularly and addressed if circumstantial evidence indicates that animals and animal products may be a significant reservoir of human *E.coli* O157:H7 infections (Thorns, 2000).

REFERENCES

- Abdel Khalek, A.; El Gaml, A and El Sherbini, M. (2001):* Prevalence of *Escherichia coli* O157 in milk and feces of dairy farm animals in Dakahlia Province. 1st cong. Of Hygiene and Human Health. 6-8 February 2001. Dept. of Food Hygiene. Fac. Vet. Med., Assiut University, 271-279.
- Boom, R.; Sol, C.J.A.; Salimans, M.M.M.; Jansen, C.L.; Wertheim van Dillen, P.M.E. and Van der Noordaa, J. (1990):* Rapid and simple method for purification of nucleic acids. J. Clin. Microbiol., 28: 495-503.
- Borczyk, A.A.; Karmali, M.A.; Lior, H. and Duncan, L.M.C. (1987):* Bovine reservoir for Verotoxin producing *Escherichia coli*. Lancet, 98.
- CDC (Center for Disease Control) (1993):* Update: Multistate outbreak of *Escherichia coli* O157:H7 infection from Hamburgers-Western United-States/ 1992-1993. J. of the American Medical Association, 269, 2194-2196.
- Chapman, P.A.; Wright, D.J. and Higgins, R. (1993a):* Untreated milk as a source of verotoxingenic *E.coli* O157. Vet. Record 133, 171-172.
- Chapman, P.A.; Wright, D.J.; Norman, P.; Fox, J. and Crick, E. (1993b):* Cattle as a possible source of verocytotoxin-producing *Escherichia coli* O157 infections in man. Epidemiol. Infect. 111: 434-447.
- Chapman, P.A.; Siddons, C.A.; Cerdan Malo, A.T. and Harkin, M.A. (1997):* A one year study of *Escherichia coli* O157 in cattle, pigs, sheep and poultry. Epidemiol. Infect. 119, 245-250.
- Clarke, R.; McEwen, S.; Harnett, N.; Lior, H. and Gyles, C. (1998):* The prevalence of verotoxin-producing *E.coli* (VTEC) in bovines at slaughter. Abst. Of the annual meeting, American Society for Microbiology, p-48, p 282.
- D'Aoust J.Y. (1989):* Manufacture of dairy products from unpasteurized milk: a safety assessment. J. Food. Protec., 52: 906.
- De Boor, E. and Heuvelink, A.E. (2000):* Methods for the detection and isolation of shiga-toxin producing *E.coli*. J. Appl. Microbiol. Symposium supplement 88, 133-143.

- Dutta, Sh.; Deb, A.; Chattopadhyay, U.K. and Tsukamoto, T. (2000): Isolation of Shiga toxin-producing *Escherichia coli* including O157:H7 strains from dairy cattle and beef samples marketed in Calcutta, India. J. Med. Microbiol. (49): 765-767.
- Fathi, Sh. M.; Asmaa, A.A. Hussein; Essa, H.; Amal, A. and Thabet A. (2002): *Escherichia coli* O157:H7 in Fresh chicken cuts and offals collected from a chicken slaughter house in Assiut, Upper Egypt. 11th European Poultry Science, Bremen, 6-10 September 2002, Germany, p.185.
- Feng, P.; Lampel, K.A.; Karch, H. and Whittam, T.S. (1998): Genotyping and phenotyping changes in the emergence of *Escherichia coli* O157:H7. J. Infect. Dis. 177; 1750-1753.
- Geue, L.; Segura-Alvarez, M.; Conraths, F.J.; Kuczius, T.; Bockemuhl, J.; Karch, H.; Gallien, P.; (2002): A long-term study on the prevalence of shiga toxin-producing *Escherichia coli* (STEC) on four German cattle farms. Epidemiol. Infect. 129 (1): 173-185.
- Griffin, P.M. (1995): *Escherichia coli* O157:H7 and other enterohemorrhagic *Escherichia coli*. In Blaser, M.J.; Smith, P.D.; Rovdin, J.I.; Greenberg, H.B. and Guerrant, R. L. (ed) Infections of the gastrointestinal tract. Raven Press Ltd. New York. p 739-762.
- Griffin, P.M. and Tauxe, R.V. (1991): The epidemiology of infections caused by *Escherichia coli* O157:H7, other enterohemorrhagic *E.coli*, and the associated hemolytic uremic syndrome. Epidemiol. Rev. 13: 60-98.
- Hancock, D.D. Besser, T.E.; Kindsel, M.L.; Tarr, P.I.; Rice, D.H. and Paros, M.G. (1994): The prevalence of *Escherichia coli* O157:H7 in dairy and beef cattle in Washington State. Epidemiol. Infect. 113: 199-207.
- Holland, J.L.; Louie, L.; Simor, A.E. and Louie, M. (2000): PCR detection of *Escherichia coli* O157:H7 directly from stools.: Evaluation of commercial extraction methods for purifying fecal DNA. J. Clin. Microbiol. 38 (11) 4108-4113.
- Johnson, J. L.; Rose, B.E.; Sharar, A.K.; Ramson, G.M.; Lattuada, C. P. and McNamara, A. M. (1995): Methods used for detection and recovery of *Escherichia coli* O157:H7 associated with a food borne disease outbreak. J. Food Protec. 58: 597-603.
- Krishnan, C.; Fitzgerald V.; Dakin, S. and Behme. R. (1987): J. Clin. Microbio. 25, 1043-1047.

- Lahti, E.; Keskimaki, M.; Rantala, L.; Hyvonen, P.; Siitonen, A. and Honkanen-Buzalski, T. (2001): Occurrence of Escherichia coli O157 in Finnish cattle. Vet. Microbiol. 79 (3): 239-251.*
- McDonough, P.L.; Rossiter, C.A.; Rebhun, R.B.; Stehman, S. M.; Lein, D.H. and Shin, S.J. (2000): Prevalence of Escherichia coli O157:H7 from cull dairy cows in New York State and comparison of culture methods used during preharvest food safety Investigations. J. Clin. Microbiol. 38 (1) 318-322.*
- Mechie, S.C.; Chapman, P.A.; and Siddons, C.A. (1997): A fifteen month field study of E.coli O157 in a dairy herd. Epidemiol. Infection. 188, 17-25.*
- Nelson, S.; Clarke, R.C. and Karmali, M.A. (1998): Verocytotoxin-producing Escherichia coli (VTEC) infection. In Palmer, S.R.; Lord Soulsby and Simpson, D.I. H. Zoonoses. Biology, Clinical practice and public health control. Oxford University press. 89-104*
- Nielson, E. M.; Tegtmeier, C.; Andresen, H.J.; Gronbaek, C. and Andersen, J.S. (2002): Influence of age, sex and herd characteristics on the occurrence of verocytotoxin-producing Escherichia coli O157 in Danish dairy farms. Vet. Microbiol. 88 (3): 245-257.*
- Pitcher, D.G.; Saunders, N.A. and Owen, R.J. (1989): Rapid extraction of bacterial genomic DNA with guanidium thiocyanate. Lett. Appl. Microbiol. 8: 151-156.*
- Porter, J.; Mobbs, K.; Hart, C.A.; Sounders, J.R.; Pickup, R.W and Edwards, C. (1997): Detection, distribution and probable fate of Escherichia coli O157 from asymptomatic cattle on a dairy farm. J. Appl. Microbiol., 83, 297-306.*
- Riley, L.W.; Remis, R.S.; Helgerson, S.P; McGee, H.B.; Wells, J.G.; Davis, B.R.; Hebert, R.J.; Olcott, E.S.; Johnson, L.M.; Hargrett, N.T.; Blake, P.A. and Cohen, M.L. (1983): Hemorrhagic colitis associated with a rare Escherichia coli serotype. N. Engl. J. Med., 308: 681-685.*
- Ryser, E.T. (1998): Public health concerns. In: Applied Dairy Microbiology. Marth, E.H. and Steele, J.L.; Marcel Dekker. Inc. New York p.263-404.*
- Slutsker, H.; Ries, A.A.; Gveene, K.D.; Wells, J.G.; Hutwanger, L. and Griffin, P.M.; (1997): E.coli O157:H7 diarrhoea in the United States clinical and epidemiologic features. Annals of Internal. Medicine. 126, 503-513.*

- Tarr, P.I.; Besser, Th.E.; Hancock, D.D.; Keenc, W.E. and Goldoft, M. (1997): Verotoxigenic Escherichia coli infection: US overview. J. Food Protec. 60 (11) 1466-1471.*
- Thompson, J. and Gillespie, D. (1987): Molecular hybridization with RNA probes in concentrated solutions of guanidine thiocyanate. Anal. Biochem. 163: 281-291.*
- Thorns, C.J. (2000): Bacterial food-borne zoonoses. Rev. Sci. tech. Off. Int. Epiz., 19 (1), 226-239.*
- Varnam, A.H. and Evans, M.G. (1991): Food borne pathogens. An illustrated Textbook. Wolfe Publishing Ltd, New York, pp 101-118.*
- Vernozy-Rozand, C. (1997): A review: Detection of E. coli O157:H7 and other verotoxin producing E. coli (VTEC) in food. J. Appl. Bacteriol. 82, 537-551.*
- Wang, R.F.; Cao, W.W. and Cerniglia C.E. (1997): A universal protocol for PCR detection of 13 species of foodborne pathogens in foods. J. Appl. Microbiol. 83, 727-736.*
- Wells, J.G. ;Shipman, L.D.; Greene, K.D.; Sowers, E.G.; Green, J.H.; Cameron, D.N.; Downes, F.D.; Martin, M.L.; Griffin, P.M.; Ostroff, S.M.; Potter, M.E.; Tauxe, R.V. and Wachsmuth, I.K. (1991): Isolation of Escherichia coli serotype O157:H7 and other Shiga-like toxin-producing E.coli from dairy cattle. J. Clin. Microbiol. 29: 985-989.*
- Zhao, T.; Doyle, M.P.; Shere, J. and Garber, L. (1995): Prevalence of enterohemorrhagic Escherichia coli O157:H7 in a survey of dairy herds. Appl. Environ. Microbiol. 61: 1290-1293.*