

## **GENE EXPRESSION FOR BIOCHEMICAL MECHANISM AGAINST DOWNY MILDEW DISEASE IN MELON (*Cucumis melo* L.).**

**El-Shimi, I.Z.; L.A.A. Badr\* and M. I.Ghonim\*\***

**Veg. Res. Dept., Hort. Res. Inst., Agric. Res. Center**

**\*Dep. of Hort., Fac. of Agric., Moshtohor, Zag. Univ. "Benha Branch"**

**\*\*Plant Pathol. Inst., El-Kassasien Hort. Res. Station, Agric. Res. Center.**

### **ABSTRACT**

Three resistance introduced varieties melon were namely, PI 140471 as P<sub>1</sub>; PI 124111 as P<sub>2</sub> and PI 183227 as P<sub>3</sub> were used as (males) with three commercial varieties namely Hale's Best, Honey dew and Ismaelawi (females) as susceptible parents to achieve this investigation during 2001-2002 in the farm of El-Kassasien Horticulture Research Station.

All the six parental genotypes and their F<sub>1</sub>'s were evaluated under the artificial infection in open field in complete blocks randomized, with three replicates, to investigate the relationship between peroxidase activity, lignin content, infection severity and the nature of inheritance for this traits and the obtained results can be summarized as following:

1. Analysis of variance show that, there were highly significant differences for all studied traits.
2. Gene expression of infection severity, peroxidase activity and lignification of mean performance for parents and F<sub>1</sub> hybrids were high.
3. Additive and non-additive variances has important role in the expression of these traits.
4. Over dominance was governed the expression of these traits.
5. Results of general and specific combining ability show that, the best parents for low infection severity and high level of peroxidase activity were PI 140471; PI 183227 then PI 124111 while Ismaelawi was the best parent for lignification, meanwhile the best F<sub>1</sub> for the above mentioned characters were PI 140471 × Ismaelawi; PI 183227 × Honey dew while, the F<sub>1</sub> PI 140471 × Honey dew exhibited the highest values for lignin content.
6. These results could be used as markers for selection program for breeding program for resistance against downy mildew in melon.
7. The artificial infection in early stage of melon plants led to significant depression for average fruit weight, fruit yield per plant and total fruit yield per feddan, while the number of fruits per plants, flesh thickness and total soluble solids did not significantly affected.

### **INTRODUCTION**

Melon is considered one of important vegetable crop, for exportation or local consumption in Egypt. Downy mildew disease attack melon plants in the open field or under green house conditions, or under low tunnels in winter season.

The causal organism of downy mildew is *Pseudoperonospora cubensis*, it's obligate parasite, which encyst to the cells by natural openings like stomata. The pathogen produce a haustoria and intera cellular hypha which make an elicitation which transmitted from the surface of plasma membrane to heterokromatin by the receptores that exist on the surface of plasma membrane. This transmitted elicitation called "signaling"; when the signal reach to DNA, the biochemical events happen which mRNAs are encoding to produce pathogenesis related proteins (PR) Metraux *et al.* (1988) or enzymes like peroxidase, polyphenoloxidase.

Major dominant gene resistance in plants is often expressed as a hypersensitive response. That is, the vicinity of an infection, the cells of a resistant plant die, thereby isolating the pathogen. Such cell necrosis is commonly associated with the accumulation of phenolic compounds and increased activity of oxidative enzymes such as peroxidase Jennings *et al.* (1969) and Retig, (1974)

The positive associated between the rapidity of the response of peroxidase activity to fungal infection and disease resistance Bi and Zhang, (1993) has led to the hypothesis that peroxidase and other oxidative enzymes may be an integral part of the host plants defense mechanism.

Enhanced peroxidase activity is very often associated with resistance phenomena such as lignin production, or, lignification, Hammerschmidt and Kuc' (1982)

In this investigation, the resistant plants might be predicted on the basis of peroxidase values. So, the germplasm, which characterized with resistance to any pathogen or a high level of plant defense must be a great value in efficient program for improving any plant species. So, we try to discover the relationship between infection severity, peroxidase activity and lignification.

## **MATERIAL AND METHODS**

### **Material:**

Six parental genotypes of melon (*Cucumis melo* L.) represented by three plant introductions from the North Central Regional Plant Introduction Station, Ames, Iowa, USA; PI 140471 (Texas, USA) as P<sub>1</sub>, PI 12411 (India) as P<sub>2</sub> and PI 183227 (Egypt) as P<sub>3</sub> which are downy mildew resistant, (Cohen *et al.* 1987; Balass *et al.* 1992; Thomas and Jourdian 1992; Guirgis *et al.* 1998). These inbred lines were used as testers (males). The other three parental genotypes were two commercial foreign varieties, Hale's Best Jumbo (Sabetha seed) as P<sub>4</sub>, Honey dew (USA) as P<sub>5</sub> and one local variety, Ismaelawi as P<sub>6</sub>. These three genotypes are susceptible to downy mildew infection and were used as lines (females). This present investigation was carried out at the farm of EL-Kassasien Horticulture Research Station, Ismailia Governorate.

These materials were sown under green house conditions, to obtain purity for one season, (January 2000), then all the parental genotypes were

crossed, with out reciprocals, to obtain F<sub>1</sub> seeds (January 2001), under green house conditions.

#### Methods:

A- All seeds of six parental genotypes populations and their 9 F<sub>1</sub> hybrids were directly sown in Nily season of (15 August 2002), in randomized complete block design with three replicates. The plot area was 20 m<sup>2</sup> (10m-long × 2m-wide).

B- Fungal inoculation: The inoculum was prepared from infected cucumber leaves with the causal organism of downy mildew disease, *Pseudoperonospora cubensis*, which were collected from different locations around the experimental farm of EL-Kassasien. The detached leaves were then dipped in distilled water and gently shaken to wash off the exudates. The spore suspension was filtered through several layers of cheese cloth and its concentration was adjusted to  $7.5 \times 10^3$  sporangia/ml. using the haemocytometer slide, Fadl *et al.* (1996)

The inoculation procedures of melon populations were accomplished by spraying the plant materials according to Thomas *et al.* (1987). The first sample (control) was taken after 48 h of inoculation; Guirgis *et al.* (1999) to recorded these data: infection severity, which was estimated according to Horsfall and Barratt (1945).

Peroxidase activity was expressed as changes in absorbance at 470nm × 10<sup>2</sup> per minute per gram fresh weight and statistically analyzed with least significant difference test (P=0.05) according to Allan and Hollis (1972), as well as lignin content (gm/gm dry weight) was determined according to Soutar and Margared Bryden (1955). Also, correlation coefficient between the above mentioned characters was computed according to Kearsy and Pooni (1996).

To, estimate the effect of infection of downy mildew disease on yield components and quality of melon populations, five plants for each plot were randomly chosen and labeled to collect these data: fruit yield (Kg/plant), total fruit yield (ton/fed.) as well as flesh thickness (cm) and total soluble solids (%).

## RESULTS AND DISCUSSION

Analysis of variance of six melon genotypes and their F<sub>1</sub> hybrids for studied characteristics were presented in Table (1). Highly significant differences were detected among genotypes for all studied traits. This indicates that, there was a large amount of variability among parental genotypes concerning these traits. Both parents and their F<sub>1</sub> hybrids exhibited highly significant differences for all characters, indicating the presence of gene effects in the inheritance of these traits. Mean squares due to parents vs crosses were highly significant for all studied traits except the

healthy melon plants of infection severity indicating the presence of heterosis in  $F_1$  crosses for these traits.

Regarding the data presented in Table (2) showed that, there were a significant differences between resistant and susceptible plants genotype for all studied traits. Data of mean performance for infection severity percentage showed that, mean percentage of infected plants of resistant parents ranged from 8.14 % to 12.26%, meanwhile the percentage of susceptible infected

**Table (1) analysis of variance of six melon parental genotypes and their  $F_1$  crosses for infection severity, peroxidase activity, and lignin content.**

Source of variance	Mean squares					
	Infection severity (%)		Peroxidase activity ( $10 X^2$ )		Lignin content (gm/gm. d.w.)	
	Healthy	Infected	Healthy	Infected	Healthy	Infected
Reps.	1.376	0.245	3.078	2.918	0.022	0.0007
Genotypes	10.329**	114.118**	1230.92**	846.499**	0.157**	0.1803**
Parents(P)	15.79**	68.055**	7.390**	801.722**	0.318**	0.3887**
Crosses(C)	8.133**	76.494**	1747.96**	902.500**	0.055*	0.0285**
P vs C	0.545	645.42**	3212.29**	622.38**	0.1655**	0.3528**
Error	1.401	3.347	0.5930	1.707	0.0194	0.0053

\*,\*\* Indicate significance at 0.05 and 0.01 levels of probability, respectively.

plants ranged from 15.67% to 21.46%. This mean that resistance to downy mildew infection was exhibited by the parental genotypes of PI140471, PI124111 and PI 183227 showing low percentage of infected plants.

This finding was in agreement with those obtained by Thomas and Jouredian, (1992), which reported that PI 124111 was resistance to downy mildew, meanwhile Guirgis *et al.* (1998), reported that inbred lines of PI 140471 and PI 124111 had a low infection with *Pseudoperonospora cubensis*, (1.39and 2.79%) respectively while Haneydew and Ismaelawi varieties had high percentage of infected plants (5.62% and 9.62%) respectively.

For peroxidase activity, Table (2) revealed that, activities of peroxidase, in true leaves, on resistant parental genotypes were higher than that of susceptible parents, respectively. The average of peroxidase activity pre-infection was 14.19 rapidly increased up to 28.99 post-infection.

These results were in agreement with those obtained by Reuveni and Karachi, (1987); Yurina *et al.* (1990); and Yun-Xing Fu. *et al.*, (1995). which they reported that, activities of peroxidase in infected cucumber leaves with downy mildew was 28.76 and resistant genotypes had high peroxidase activity while those with low peroxidase activity were susceptible.

For lignin content, data Table (2), showed that, resistant parents, inbred lines of PI 140471; PI 124111 and PI 183227 had high content of lignin

compared with susceptible parents. Results indicated that, inbred line PI 140471 had the highest value of lignin content (1.075gm) followed by Ismaelawi (0.383 gm) in healthy case, but after inoculation with *Pseudoperonospora cubensis*, the expression of these genotypes changed, while the inbred line PI 140471 recorded the first rank followed by Honeydew; then PI 183227 and the inbred line Hale's Best Jumbo came the last. This mean that, after inoculation with *Pseudoperonospora cubensis*, the gene expression of all parental genotypes had changed which, rapid increase in lignin content were observed, these finding in agreement with those obtained by Luo-Gui Fen *et al.* (1997); which reported that, the sugar and lignin contents of leaves were increased by pre-inoculation treatments. They added that, sugar and lignin in cucumber leaves are involved in defense mechanisms. Matsumoto, (1994), reported that, the most resistant cultivar of Japanese radish had lignified cell walls prior to attack by *Peronospora parasitica* and the hypha could not penetrate cells. Also, Luo-GuiFen and Luo-FG, (1995), reported that, resistance to downy mildew was higher in plants with a greater lignin content.

A considerable difference was observed for  $F_1$  between the average of parents and that of  $F_1$ 's for infection severity, peroxidase activity and lignin content, the  $F_1$ 's showed an average 21.11% for infected plants superiority over their parents. Meanwhile, peroxidase activity for  $F_1$  hybrids, increased from 28.99 for parental genotypes, up to 33.04 for  $F_1$  crosses. Meanwhile, lignin content had significantly increased from (0.665 gm) to (0.770gm) for  $F_1$  hybrids showed higher values for biochemical characteristics than parents, suggesting the presence of dominant effects, which governed these traits. In this regard, Balass *et al.* (1992), reported that,  $F_1$  hybrid plants (Hemed  $\times$  PI 124111) that displayed only a partial resistance against *P. cubensis*. Back-cross progeny plants (Hemed  $\times$  PI 124111)  $\times$  Hemed) segregated 1:2:1 partly resistant: susceptible: highly susceptible to the disease.

It is worth to mention that, increasing of infection severity led to increase in peroxidase activity accompanied by increasing in lignin content. These findings were found to be in agreement with those reported by Nadonly and Sequeira, (1980); Jennings *et al.* (1969); and Svalheim and Robertsen, (1990). they reported that, in resistant plants, rapid significant increase in soluble peroxidase activity induced by fungal infection. Also, Guirgis *et al.* (2000); reported that, in resistant melon genotypes, rapid increase in peroxidase isozymes activity was observed for PI 140471, PI 124111 and PI 183227 while susceptible genotypes showed no rapid changes in the activity of peroxidase isozymes.

Moreover Hammerschmidt and Kuc<sup>2</sup>, 1982; Dean and Kuc<sup>2</sup>, (1987), reported that, lignification of the host cell wall may be a mechanism for induced systemic resistance in the cucumber. That is, lignification occurred more rapidly and to a greater extent in protected cucumbers with *Collectotrichum lagenarium* than in the control after challenge with *Cladosporium cucumerinum*. Further more, Balass *et al.* (1993); reported that, lignin like deposits was observed in infected PI 124111 leaves but not observed in cultivar Hemed muskmelon. They added that, peroxidase activity

and lignin deposits were higher in the resistant inbred lines PI 124111 and F<sub>1</sub> as compared with Hemed at 21° not at 12°.

The relative magnitude of additive ( $\sigma^2A$ ) and dominance ( $\sigma^2D$ ) variances was presented through the  $\sigma^2A/\sigma^2D$  ratio (Table 3). This ratio was equal unity for infection severity in healthy and infection stress conditions. Mean while, the ratio was less than unity for Pox activity and lignin content. This indicated that, additive and non-additive variance plays an important role in the inheritance of these characters. Values of the average of dominance "a" were greater than unity for all studied traits, indicating that over dominance plays an important role in the inheritance of all studied biochemical traits.

Broad sense heritability estimates for all studied traits were high. The heritability broad sense was ranged from 79.43 % for infection severity to 99.96 % for Pox activity (in healthy plants), respectively estimates of narrow sense heritability ( $h^2n$ ) were lower values than the broad sense. This indicate that dominant gene effects are governed these biochemical traits. These findings were found to be in harmony with reported by Shetty *et al.* 2001; where stated that, the large magnitude of dominance in the inheritance of peroxidase activity in pearl millet would tend to hinder progress in improving downy mildew resistance.

Estimates of general combining ability effects for the parental genotypes of melon are given in Table (4). The positive estimates of the gca effects were considered desirable for all studied traits, while the negative gca effects were considered desirable for percentage of infected plants.

For infection severity, the inbred line PI 183227 had the lowest negative gca estimate (-0.146), while Ismaelawi melon land race, had the highest positive and highly significant gca estimate. Meanwhile, the inbred line PI124111 had the highest negative and highly significant gca estimate for peroxidase activity, while Honeydew had the lowest positive gca estimate (healthy case). Under stress infection, the inbred line PI140471, had the highest positive and highly significant gca estimate followed by Ismaelawi, mean while, the only negative and highly significant gca estimate for lignin content was recorded by Ismaelawi.

Further more, the gca effects in Table (4) indicate that, the parental inbred lines, PI140471; PI124111 and PI18227 which showed resistance to downy mildew disease had the highest and highly significant gca effects for peroxidase activity (11.759, 4.615,-7.143), respectively under infection stress.

The best parent for low infection is PI140471; for gca estimates which exhibit high peroxidase activity and moderate lignin content. These findings were found in harmony with those obtained by Kim-ZhooHyeon and Kim-ZH, (1996), which reported that, GCA effect was very highly significant when they estimated the inheritance of resistance to downy mildew in *Cucumis melo*

Specific combining ability (sca) effects of the F<sub>1</sub> crosses for the studied traits are presented in Table (5). It is clear that, the number of crosses showing significant sca effects (either positive or negative) ranged from 4 crosses, for infection severity, to 8 crosses (for lignin content), mean

while, all the estimates of sca effects for the  $F_1$  crosses of peroxidase activity were highly significant. This indicate that, all the crosses of  $F_1$  hybrids exhibit a moderate resistance when inoculated by the pathogen of downy mildew disease and these  $F_1$ 's will be able to decrease the infection severity because of the resistant genes had activated by the fungal elicitation and encoded for transcript mRNAs for a high level of peroxidase enzymes. In this concern, Guirgis *et al.* (1998), reported that, a group of elicitors might be induced thereby affect the regulation mechanisms of peroxidase enzymes during the time course of infection.

Also, LaRosa *et al.*, (1992); Nelson *et al.*, (1992), and Liu *et al.* (1994), reported that, gene expression of some PR proteins which have antifungal activity Bol *et al.*, (1990) is regulated by a large number of hormonal and environmental signals which include fungal infection.

**Table (2): Mean performance of infection severity, peroxidase activity and lignin content in six parental melon genotypes and their  $F_1$  hybrids.**

Genotypes	Infection severity (%)		Peroxidase activity (10 $X^2$ )		Lignin content (gm/gm. d.w.)	
	Healthy	Infected	Healthy	Infected	Healthy	Infected
<b>Parents</b>						
PI140471 (1)	1.513	8.14	16.61	61.23	1.075	1.206
PI124111 (2)	1.843	12.26	13.97	19.63	0.380	0.593
PI183227 (3)	1.796	11.14	12.53	19.65	0.322	0.520
Hale's Best Jumbo (4)	4.550	16.30	15.53	19.46	0.231	0.385
Honey dew (5)	5.460	15.67	13.56	23.59	0.317	0.776
Ismaelawi (6)	6.963	21.46	12.98	30.39	0.383	0.410
Average	3.687	14.16	14.19	28.99	0.418	0.665
L.S.D. <sub>0.05</sub>	4.27	12.51	0.41	6.03	0.16	0.46
L.S.D. <sub>0.01</sub>	6.07	17.79	0.59	8.57	0.23	0.66
<b><math>F_1</math> hybrids</b>						
1 $\times$ 4 $F_1$	3.70	13.18	14.56	14.74	0.360	0.923
1 $\times$ 5 $F_1$	1.79	20.83	12.76	17.67	0.538	0.748
1 $\times$ 6 $F_1$	4.76	28.15	67.25	71.83	0.236	0.770
2 $\times$ 4 $F_1$	3.35	18.16	12.12	13.86	0.320	0.716
2 $\times$ 5 $F_1$	1.87	23.20	14.92	19.14	0.302	0.900
2 $\times$ 6 $F_1$	5.75	22.26	13.82	17.35	0.628	0.626
3 $\times$ 4 $F_1$	2.70	13.62	17.05	31.41	0.315	0.831
3 $\times$ 5 $F_1$	4.88	25.00	13.43	67.29	0.343	0.730
3 $\times$ 6 $F_1$	6.40	25.62	12.29	44.09	0.368	0.690
Average	3.91	21.11	19.80	33.04	0.367	0.770
L.S.D. <sub>0.05</sub>	9.41	9.31	6.35	10.29	1.24	0.44
L.S.D. <sub>0.01</sub>	12.96	12.82	8.75	14.17	1.71	0.61

**Table (3): Variance components for all studied traits in melon genotypes and their  $F_1$  hybrids.**

Characters		Parameters						
		$\sigma^2A$	$\sigma^2D$	$\sigma^2A/\sigma^2D$	"A"	$\sigma^2G$	$\sigma^2PH$	Heritability
								$h^2_b$ $h^2_n$
Infection severity (%)	Healthy	0.902	0.902	1.000	1.41	1.804	2.271	79.43 39.72
	Infected	9.63	9.52	1.012	1.40	19.15	20.26	94.52 47.53
Pox activity (10 $X^2$ )	Healthy	104.37	516.18	0.202	3.14	620.55	620.75	99.96 16.81
	Infected	47.07	308.61	0.152	3.62	355.68	356.25	99.84 13.21
Lignin content (gm/gm. D.w.)	Healthy	0.00	0.027	-0.024	-9.48	0.027	0.028	96.42 0.00
	Infected	0.001	0.009	0.147	3.67	0.010	0.010	98.32 9.83

\*Negative estimates are considered zero

**Table (4): Estimates of relative gca effects of parental melon genotypes for infection severity, peroxidase activity and lignin content.**

Parents	Mean squares					
	Infection severity (%)		Peroxidase activity (10 X <sup>2</sup> )		Lignin content (gm/gm. d.w.)	
	Healthy	Infected	Healthy	Infected	Healthy	Infected
Testers PI140471 (1)	-0.493	-0.170	1.669**	11.759**	0.010	0.035
PI124111 (2)	-0.254	0.316	-17.823**	-4.615**	0.015	-0.015
PI183227 (3)	0.748	-0.146	16.153**	-7.145**	-0.025	-0.020
Lines Hale's BestJumbo(4)	-0.661	-5.903**	-12.018**	-6.242**	-0.036	0.045
Honey dew (5)	-1.062**	2.117**	0.213	-4.653	-0.006	0.029
Ismaelawi (6)	1.724**	3.785**	11.804**	10.896**	0.042	-0.075**
Standard error	0.394	0.609	0.256	0.435	0.046	0.024

\*,\*\* : Significant the 0.05 and 0.01 probability levels respectively

The best sca effect for percentage of infected plants with (DM) disease was obtained from the cross (1 × 6 F<sub>1</sub>), (2 × 4F<sub>1</sub>), (2 × 6F<sub>1</sub>) and (3 × 5F<sub>1</sub>). These crosses could be considered promising genotypes for melon resistance to downy mildew disease. For Pox activity, the highest significant and positive sca effects were observed in the crosses, (1 × 6 F<sub>1</sub>), (2 × 4F<sub>1</sub>), (2 × 6F<sub>1</sub>) and (3 × 5F<sub>1</sub>). For lignin content, the highest significant and positive sca effects were shown in the crosses, (1 × 5F<sub>1</sub>) and (2 × 5F<sub>1</sub>). It is worthy to not that, the crosses (1 × 6 F<sub>1</sub>), (2 × 4F<sub>1</sub>), (2 × 5F<sub>1</sub>) and (2 × 6F<sub>1</sub>) showed high significant and positive sca effects for all the studied traits. These crosses are recommended for breeding programs to improve the melon resistance to downy mildew disease.

**Table (5): Estimates of sca effects of melon F1 hybrids for infection severity, peroxidase activity, and lignin content.**

Cross	Infection severity (%)		Peroxidase activity (10 X <sup>2</sup> )		Lignin content (gm/gm. d.w.)	
	Healthy	Infected	Healthy	Infected	Healthy	Infected
1 × 4 F <sub>1</sub>	0.942	-1.635	-6.350**	-12.357**	0.0181	0.0485
1 × 5 F <sub>1</sub>	-0.559	-2.01	-20.565**	-10.836**	0.1664*	-0.0875*
1 × 6 F <sub>1</sub>	-0.382	3.645**	26.916**	23.193**	-0.1846*	0.0390*
2 × 4 F <sub>1</sub>	0.357	2.857**	10.518**	3.321**	-0.0274	-0.0842*
2 × 5 F <sub>1</sub>	-0.725	-0.123	1.086**	7.009**	0.1740*	0.1379**
2 × 6 F <sub>1</sub>	0.368	-2.734**	-11.604**	-10.330**	0.2014*	-0.0537
3 × 4 F <sub>1</sub>	-1.299	-1.222	-4.168**	9.035**	0.0092	0.0357
3 × 5 F <sub>1</sub>	1.284	2.133**	19.479**	3.827**	0.0075	-0.0503
3 × 6 F <sub>1</sub>	0.014	-0.911	-15.311**	-12.863**	-0.0168	0.0146
Standard error	0.683	1.056	0.444	0.754	0.0809	0.0421

\*,\*\* : Significant the 0.05 and 0.01 probability levels respectively



Regarding the nature of associations among the studied traits at the genotypic, phenotypic and environmental levels the data represented in Table (6) showed that, there was a significant and positive genetic and environmental correlation between infection severity, peroxidase activity, and lignin content under healthy condition, meanwhile, significant and positive genetic and environmental correlation was found between peroxidase activity and lignin content under infection stress.

This means that, infected melon plants with *Pseudoperonospora cubensis* led to rapid increase in (pox) activity and (pox) catalyze the final step of polymerization of lignin synthesis. (Hammerschmidt and Kuc<sup>1</sup>, 1982 and Dean and Kuc<sup>1</sup>, 1987). In this regard, Reuveni *et al.* (1990) found that, high correlation was demonstrated between peroxidase activity of uninoculated melon plants and resistance to *P. cubensis* determined in vivo. Yun-XingFu *et al.* (1995), declare that, there was positive correlation between the activities of peroxidase and resistance to downy mildew of cucumber. Guirgis *et al.* (2000); reported that, a positive Spearman rank correlation coefficient (0.73) was observed between low infection severity and the highest changes in peroxidase isozyme activity scored at 27h after fungal infection of melon with downy mildew agents. Furthermore, data in Table (6) declare that, there were positive and insignificant genetic, phenotypic, and environmental correlation coefficient was observed between infection severity and lignin content in melon populations. Negative insignificant genetic and environmental correlation coefficients were observed between (pox) activity and lignin content under healthy conditions.

Table (6): Genotypic (Gr), phenotypic (Pr) and environment (Er) correlation coefficients for association between infection severity, peroxidase activity and lignin content.

		Infection severity (%)		Peroxidase activity (10 X <sup>2</sup> )		Lignin content (gm/gm. d.w.)	
		Healthy	Infected	Healthy	Infected	Healthy	Infected
Infection severity (%)	Gr			0.336*	0.034	-0.394**	0.043
	Pr			-0.168	-0.014	-0.062	0.081
	Er			0.311*	0.034	0.351*	0.045
Peroxidase activity (10X <sup>2</sup> )	Gr					-0.270	0.357*
	Pr					0.064	-0.275
	Er					-0.252	0.349*
Lignin content (gm/gm. d.w.)	Gr						
	Pr						
	Er						

\*,\*\* : Significant at the 0.05 and 0.01 probability levels, respectively

Data in table (7) declare that, there were significant differences between resistant and susceptible parents as well as between healthy and infected melon populations for all studied traits. For average fruit weight, the men performance of resistant melon parents did not affect by inoculation of *P. cubensis* while, the susceptible melon parents had decreased by the inoculation with *P. cubensis*. Meanwhile, the total yield, per plant and ton/fed., had affected by the inoculation of the pathogen, but number of fruit per plant did not affect.

Table (7): Mean performance of parental genotypes and their F<sub>1</sub> hybrids for yield and fruit quality under healthy and infection stress.

Genotypes	Average fruit weight (Kg)		Number of fruits/plants		Total fruit yield/ plant (Kg)		Total fruit yield (ton/fed.)		Flesh thickness (cm)		Total soluble solids (%)	
	Healthy	Infected	Healthy	Infected	Healthy	Infected	Healthy	Infected	Healthy	Infected	Healthy	Infected
<b>Parents</b>												
PI140471 (1)	0.069	0.064	4.93	4.89	0.387	0.349	2.326	2.153	0.583	0.583	6.00	6.00
PI124111 (2)	0.372	0.327	3.63	3.60	1.234	1.181	6.406	5.963	1.256	1.526	10.32	10.30
PI183227 (3)	0.665	0.586	2.15	2.13	1.432	1.251	7.592	6.692	2.566	2.566	10.65	10.65
Hale's Best Jumbo (4)	0.475	0.395	2.99	2.97	1.442	1.173	7.536	6.414	2.466	2.400	11.34	11.32
Honey dew (5)	0.462	0.391	3.22	3.21	1.483	1.257	7.900	6.673	2.233	2.200	11.76	11.75
Ismaelawi (6)	3.577	2.809	1.31	1.20	4.700	3.646	8.170	6.436	3.00	2.866	12.04	12.00
Average	0.936	0.762	3.04	3.00	1.776	1.476	6.655	5.721	2.062	2.023	10.35	10.33
L.S.D. <sub>0.05</sub>	0.419	0.407	2.62	2.68	1.95	1.27	1.33	1.85	0.513	0.631	2.96	2.95
L.S.D. <sub>0.01</sub>	0.596	0.578	3.73	3.81	2.77	1.81	1.90	2.63	0.729	0.897	4.21	4.20
<b>F<sub>1</sub> hybrids</b>												
1 × 4 F <sub>1</sub>	0.274	0.244	4.42	4.42	1.243	1.079	7.455	6.789	1.380	1.366	8.79	8.76
1 × 5 F <sub>1</sub>	0.245	0.203	4.19	4.19	1.061	0.850	6.368	5.070	1.433	1.366	9.21	9.20
1 × 6 F <sub>1</sub>	0.546	0.397	4.35	4.35	2.380	1.727	9.464	6.814	2.350	2.30	10.62	10.60
2 × 4 F <sub>1</sub>	0.496	0.459	3.54	3.54	1.752	1.620	8.760	8.043	2.380	2.35	9.77	9.76
2 × 5 F <sub>1</sub>	0.582	0.561	3.66	3.66	2.131	2.057	10.323	9.36	2.433	2.40	10.49	10.50
2 × 6 F <sub>1</sub>	0.512	0.402	3.21	3.20	1.643	1.287	9.524	7.615	2.866	2.70	10.76	10.75
3 × 4 F <sub>1</sub>	0.598	0.517	2.34	2.34	1.650	1.135	9.900	8.571	2.633	2.566	10.66	10.65
3 × 5 F <sub>1</sub>	0.538	0.410	2.44	2.43	1.326	0.995	3.956	5.993	2.6	2.533	10.82	10.82
3 × 6 F <sub>1</sub>	1.759	1.346	2.30	2.28	4.040	3.078	8.453	9.019	2.866	2.733	10.77	10.76
Average	0.616	0.504	3.38	3.37	1.913	1.536	8.689	7.527	2.326	2.257	10.21	10.20
L.S.D. <sub>0.05</sub>	0.282	0.257	1.55	1.44	1.41	1.33	3.37	4.06	1.02	0.78	1.79	1.777
L.S.D. <sub>0.01</sub>	0.389	0.355	2.14	2.98	1.94	1.83	8.41	5.59	1.41	1.08	2.46	2.43

As for fruit quality, i.e. flesh thickness and total soluble solids, data reveal that, there were no significant differences between healthy and infected melon plant as well as, the mean performance of resistant parents was less than the susceptible parents, also, the inoculation with *P. cubensis* did not effect in flesh thickness and total soluble solids.

In this regard, Tsabolov-P-Kh (1990), reported that, the highest early and total yields were obtained from resistance varieties of cucumber to *Pseudoperonospora cubensis*. Meanwhile, Medvedeva and Medvedev, (1983) reported that, less severity of symptoms and highest yields were obtained from calipso, when they evaluated 1644 genotypes of melon from 45 countries. Also, Thakur et al. (1996), revealed that, yield of 30 Momordica charautia genotypes was not significantly affected by *P. cubensis*.

Furthermore, ULLasa and Amin, (1988), reported that, infection of *Luffa acutangula* with *P. cubensis* decreased crop yields by 61%. They added

that, early infection caused significant yield reduction while late infection did not cause significant yield reduction.

On the level of F<sub>1</sub> hybrids, data on Table (7) show that there were no differences between healthy and infected melon populations for average fruit weight, fruit numbers per plant, flesh thickness and total soluble solids, but total fruit yield per plant and ton/feddan, were significantly affected by the pathogen of downy mildew diseases.

The best crosses for average fruit weight and total yield per plant as well as ton/feddan were the three crosses of the resistant parent PI 183227, for number of fruit per plant, were the three crosses of the resistant parent PI 140471, and for flesh thickness and soluble solids, the three crosses of susceptible parent, Ismaelawi.

In this concern, Sumner and Phatak, (1987), reported that, downy mildew reduced yields in Marketes, cucumber. Furthermore; Koh-Young Jin *et al.* (1996), reported that, large irregular lesions and small necrotic polygonal lesions appeared on the lower leaves, developing on to the upper leaves of downy mildew infected netted melon plants during the flowering season, resulting in death of whole leaves. They added, in a sever case, the disease devastated whole plantation, which made it impossible to harvest melons.

## **REFERENCES**

- Allan, A. L. and J.P. Hollis (1972). Sulfide inhibition of oxidases in rice root. *Phytopathology*, 62: 634-639.
- Balass, M.; Y.Cohen,; and M. Bar-Joseph (1992). Identification of a constitutive 45Kda soluble protein associated with resistance to downy mildew in muskmelon (*Cucumis melo* L.), line PI12411F. *Physiological and Molecular Plant Pathology*, 41:6, 387-396 17ref.
- Balass, M.; Y.Cohen,; M. Bar-Joseph- (1993). Temperature dependent resistance to downy mildew in muskmelon: Structural response. *Physiological and Molecular Plant Pathology*, 43: 1, 11-20; 17 ref.
- Bi, Y. and W. Y. Zhang, (1993). On changes respiratory rate, ethylene evolution and peroxidase activity of the infected melon. *Acta Phytopathol. Sinica*, 23: 1, 69-73.
- Bol, J. F., Linthorst, H. J. M. and B.J.C. Cornelissen (1990). Plant pathogenesis-related proteins induced by virus infection. *Annu. Rev. Phytopathol.* 28, 113-138.
- Cohen, Y. and Eyal, H. (1987). Downy mildew, powdery mildew and Fusarium wilt-resistant muskmelon breeding line PI-124111. *Phytoparasitica*, 15, (3): 187-195.
- Dean, R. A. and J. Kuc (1987). Rapid lignification in response to wounding and infection as a mechanism for induced systemic protection in cucumber. *Physiol. Mol. Plant Pathol.* 31, 69.
- Fadl, F. A.; Yehia; A. H.; Dawlat A. Abd El-Kader; M. M. Mahrous and M. I. Ghonim 1996. Occurrence and importance of *Cucurbits* downy mildew disease in Egypt. *Egypt. J. Agric. Res.*, 74 (4), 889-902.

- Guirgis, A. A.; Abd A. A El-Raheem .; S. E. Mohamedein, and I. Z. A. El-Shimi (1998). Molecular polymorphism of esterase isozymes in melon generations and their response to fungal inoculation and wounding. *Zagazig J. Agric. Res.* Vol. 25 No. (6), 991-1004.
- Guirgis, A. A.; A.A. Abd El-Raheem; A. I. Baz and I.Z.A. El-Shimi, (1998). Wound-induced molecular alterations and its relationship to peroxidase isozymes of fungal infected melon generations. *Egypt. J. Genet. Cytol.*,
- Guirgis, A. A; A. A. Abd El-Raheem and I. Z. A. El-Shimi (2000). Wound-induced molecular alterations and its relationship to peroxidase isozymes of fungal infected melon generations. *J. Agric. Sci. Mansoura Univ.*, 25 (4): 2075-2089.
- Hammerschmidt, R. and J. Kuc' (1982). Lignification as a mechanism for induced systemic resistance in cucumber. *Physiol. Plant Pathol.*, 20, 61.
- Horsfall, J. C. and R. W. Barratt (1945). An improved grading system for measuring plant diseases. *Phytopathol.*, 35: 655.
- Jennings, P. H., B. L. Brannaman and F. B. Zscheile, Jr. (1969). Peroxidase and polyphenoloxidase activity associated with *Helminthosporium* leaf spot of Maize. *Phytopathol.*, 59: 963-967.
- Kim-ZhooHyeon; Z. H. Kim., (1996). Inheritance of resistance to downy mildew (*Pseudoperonospora cubensis*) in *Cucumis melo*. *Journal of the Korean Society for Horticultural Science*, 37:2, 242-247; 8ref.
- Kearsy, M. J. and H. S. Pooni (1996). The genetical analysis of quantitative traits. *Champan. & Hall, London*.
- Koh-YoungJin; Ahn-MiYeon; Seo-JeongKyu; Koh-YJ; Ahn-MY; Seo-JK; (1996). Occurrence and control of downy mildew of netted melon in Chonnam province. *Korean-Journal of Plant Pathology*; 12: 4, 4 59-462, 7 ref.
- LaRosa, P.C.; P.M. Hasegawa and R. A. Bressan (1992). Osmotin gene expression is postranscriptionally regulated. *Plant Physiol.*, 100: 409-415.
- Liu, d.; K. G. Raghothama; P.M. Hasegawa and R.N. Bressan (1994). Resistance to the pathogen *Phytophthora infestans* in transgenic potato that over-express osmotin. *Proc. Natl. Acad. Sci. USA*, 91: 1888-1892.
- Luo-GuiFen and FG. Luo (1995).. Relationship between lignin content in cucumber leaves and resistance to *Pseudoperonospora cubensis*. *Journal of Jilin Agricultural University*, 17:2, 18-21, 4 ref.
- Luo-Gui Fen; Cui-Jun Tao; Gao-YuFang;Zhang-Li;Luo-GF;Cui-JT.; Gao-YF; and Zhang-L, (1997). Relationship between sugar, lignin content and resistance to downy mildew of cucumber. *Acta-Phytopathologica-Scinca*, 27: 1.65-69.
- Matsumoto, I. (1994). Lignin induction in the root tissues of Japanese radish infected with downy mildew fungus. *Memoirs of the College of Agriculture, Ehime- University*, 39: 1, 1-73.

- Medvedeva, N. I. and A. V. Medvedev, (1983). Agronomic and biological assessment of cucumber varieties with potential for breeding for resistance to downy mildew. *Trudy-po-Prikldoni-Botanike, Genetike-i-Selektsii, rec'd*, 77: 25-28, 5 ref.
- Metraux, J.P.; L. Streit and T. Staub (1988). A pathogenesis related protein in cucumber is a chitinase. *Physiological and Molecular Plant Pathology*, 33:1,1-9;22ref
- Nadonly, L. and L. Sequeira, (1980). Increase in peroxidase in peroxidase activities are not directly involved in induced resistance in tobacco. *Physiol. Plant. Pathol.*, 16:1-8.
- Nelson, D. E.; K. G. Raghothama; N.K Singh.; P.M. Hasegawa and R. A. Bressan (1992). Analysis of structure and transcription activation of an osmotin gene *Plant. Mol. Biol.*, 19:577-588.
- Retig, N. (1974). Changes in peroxidase and polyphenoloxidase associated with natural and induced resistance of tomato to *Fusarium* wilt. *Physiol. Plant. Pathol.*, 4: 145-150.
- Reuveni, R. and Karchi (1987). Peroxidase activity-a possible marker for resistance of melon against downy mildew. *Phytopathology*, 77: 12,1724.
- Reuveni, R. and Z. M ShimoniKarchi (1990). A rapid assay for monitoring peroxidase activity in melon as a marker for resistance to *Pseudoperonospora cubensis*. *Journal of Phytopathology*, 12'9: 4, 33-338; 14 ref.
- Shetty, H.S.; N.S.Vasanthi; B.R. Sarosh, and K.R. Kini (2001). Inheritance of downy mildew resistance, B-1,3-glucanase and peroxidase in pearl millet (*Pennisetum glaucum* L. R.Br.) crosses. *Thero. Appl. Genet.*, 102:1221-1226.
- Scutar, T. H. and Margaret Bryden (1955). Differentiation of jute and other vegetable fibers by acetyle content. *J. Text. Inst. Trans.*, Vol. 46: 528
- Sumenr-D. R. and S.C. Phatak (1987). Control of foliar disease of cucumber with resistant cultivars and fungicides. *Appl. Agric. Res.*, 2: 5, 324-329, 15 ref.
- Svaleheim, O. and B. Robertsen (1990). Induction of peroxidase in cucumber hypocotyle by wounding and fungal infection. *Physiol. Plantarum.*, 78: 2, 261-267.
- Thakur, J. C.; A. S. Khahra; and K. C. DhanJu (1996). Evaluation of bitter gourd genotypes against disease and their correlation with other quantitative characters. *Punjab-Vegetable Grower*, 31: 25-28, 5ref.
- Thomas, C.E.; Y. Cohen; E. L. Jourdain and H. Eyal (1987). Use of reaction types to identify downy mildew resistance in muskmelons. *Hort. Sci.*, 22(4): 635-640.
- Thomas, C.E.; E. L. Jourdain (1992). Evaluation of melon germplasm for resistance to downy mildew *HortScience*. 27:5.434-436; 8 rf.
- Tsabolov-P-Kh, (1990). We select varieties giving early yield. *Kartofel-i-Ovoschi*, No. 3.22-23.

- Ullase, B. A. and K. S. Amin, (1988). Ridge gourd downy mildew epidemics in relation to fungicidal sprays and yield loss. Mysore Journal of Agricultural Sciences. 22(1) 62-67. (C. F. review of plant pathology 1991, Vol. 70, No. 4, 4 ref.)
- Yun-Xing Fu; Cui-ShiMao; Huo-XiuWen, Yun-XF; Cui-SM; Huo-XW (1995). The relationships between the activity of several enzymes in cucumber tissues and their resistance to downy mildew of cucumber. Act-Agriculturae Boreali-Sinica, 10: 1, 92-98.
- Yurina, OV.; Anikina, I.I.; Yurina-T.P.,1990. Search for resistance forms. Kartofel-I-Ovoschchi, No.3, 25-26.

## التعبير الجيني لميكانيكية الدفاع الكيموحيوية ضد مرض البياض الزغبي في الشمام

إبراهيم زكي عبد الوهاب الشيمي-لطفى عبد الفتاح عبد الرحمن بدر\*-  
مجدي إبراهيم غنيم\*\*

أقسام بحوث الخضر-معهد بحوث البساتين-مركز البحوث الزراعية

\*قسم البساتين-كلية الزراعة-بمشتهر-جامعة الزقازيق-فرع بنها

\*\*معهد بحوث أمراض النبات-محطة بحوث البساتين-مركز البحوث الزراعية

تم التهجين بين ثلاثة آباء من القاوون تحمل صفة المقاومة لمرض البياض الزغبي المتسبب عن الفطر *Pseudoperonospora cubensis* وهي PI 140471 و PI 124111 و PI 183227 وثلاثة آباء أخرى حسنة للمرض وهي أصناف تجارية منتشرة مثل هالزست و هني ديو وإسماعيلوي وذلك في النوبة المقامة بمحطة بحوث البساتين بالقصاصين عام ٢٠٠١ بغرض الحصول على بذور الجيل الأول.

تم زراعة الآباء والهجن في قطاعات كاملة العشوائية-بحقل الأمراض-بغرض تقييم الآباء والهجن تحت ظروف العزى الصناعية بالمسبب المرضي بهدف البحث عن العلاقة بين نشاط إنزيم البيروكسيداز وعلاقته بعملية اللجننة الحادثة في الورقة نتيجة مهاجمة الكائن الممرض وكذلك معرفة طبيعية توريث بعض الصفات البيوكيميائية. وقد أشارت الدراسة إلى:

- ١- تشير نتائج تحليل التباين إلى وجود فرق معنوي جدا بين الآباء والهجن لكل الأصناف المدروسة.
- ٢- يشير معدل الأداء للآباء والهجن إلى أن التعبير الجيني لشدة الإصابة ونشاط إنزيم البيروكسيداز وكذلك عملية اللجننة كان مرتفعاً في الآباء المقاومة للمرض عنه في الآباء الحساسة وأن أفراد الجيل الأول كانت متوسطة بين الأبوين.
- ٣- تشير النتائج إلى أن كلا من التباين الوراثي المضيف وغير المضيف يلعب دوراً في التعبير الجيني لكل من شدة الإصابة ونشاط إنزيم البيروكسيداز وعلاقته بعملية اللجننة.
- ٤- كم تشير النتائج إلى أن السيادة الفائقة تحكم التعبير الجيني لكل الصفات المدروسة.
- ٥- تشير تقديرات القدرة العامة على الانتلاف أن أفضل الآباء التي حققت أقل نسبة إصابة مع محتوى عالي من نشاط إنزيم البيروكسيداز هو PI 140471 يليه PI 183227 ثم PI 124111 بينما كان الصنف الإسماعيلوي الأسرع في تكوين اللجنين.
- ٦- تشير بيانات القدرة الخاصة على الانتلاف إلى أن أفضل الهجن التي حققت أقل نسبة أصابه بالمسبب المرض وكذلك نشاط عالي من إنزيم البيروكسيداز هي:

PI 140471 × Ismaelawi و Hale's Best × PI 124111

PI 124111 × Ismaelawi و PI 183227 × Honey dew

PI 140471 ×

بينما كانت أفضل الهجن التي حققت نشاط عالي في تكوين اللجنين هي

Honey dew و PI 124111 × Honey dew

٧- يمكن استخدام هذه النتائج كأداة انتخابية عند التربية لمرض البياض الزغبي في الشمام والكتنولوب.

٨- أدت الإصابة بالعدوى الصناعية في الحقل ابتداء من بداية حياة النبات إلى حدوث انخفاض معنوي في متوسط وزن الثمرة وكمية المحصول للنبات الواحد وكمية المحصول الكلي للذئبان بينما لم تتأثر كل من عدد الثمار للنبات و سمك اللحم وكذلك المواد الصلبة الذاتية الكلية بالإصابة في الحقل معنوياً.