

## PRODUCTIVITY OF *Majorana hortensis* L. PLANTS AS INFLUENCED BY THE INTERACTIONS BETWEEN MINERAL AND BIOLOGICAL FERTILIZATION

Kandeel, A.M.<sup>1</sup> and M.S.Sharaf<sup>2</sup>

1- Hort. Dept., Fac. Agric., Ain Shams Univ., Cairo, Egypt .

2- Agric. Microbiol. Dept., Fac. Agric., Ain Shams Univ., Cairo, Egypt .

### ABSTRACT

A field experiment was carried out to investigate the effect of pre-sowing inoculation of marjoram with *Azotobacter chroococcum*, *Bacillus circulans*, vesicular arbuscular mycorrhizae(VAM) or their mixture in the presence of full or half of the recommended field rate of the inorganic NPK fertilization on the plant growth, microbial activities in the rhizosphere, volatile oil yield and chemical composition. Parameters related to the above-mentioned measures of performance were recorded. Data of this study showed that mixed inoculation with the three biofertilizers supplemented with half dose of NPK gave the highest and significant increases in the growth parameters, oil % and oil yield feddan<sup>-1</sup>, as well as, the N, P and K % in plant herb . The above-mentioned treatment increased the concentration of the constituents volatile oil, geranyl acetate, cineole, linalool, limonene and  $\alpha$ -terpenolene, while decreased the concentration of  $\alpha$ -terpineol compared with the control. Mixed inoculation also, raised total microbial densities and also those of azotobacters and *B.circulans* in marjoram rhizospheric soil and increased CO<sub>2</sub> evolution, N<sub>2</sub>-ase activity and mycorrhizal root infection, as well as, spore production .On the other hand, inoculation with either of *A.chroococcum* or *B.circulans* amended with half dose of NPK decreased the above-mentioned parameters compared with uninoculated plants supplemented with the full dose of NPK fertilization, but VAM applied alone or in conjugation with the bacterial partners significantly enhanced those parameters .

### INTRODUCTION

Recently, there has been an increasing awareness of the undesirable impact of mineral fertilizers on the environment, as well as, the potentially dangerous effects of chemical residues in plant tissues on the human health. As a result of this awareness, strict regulations have been imposed in several countries (especially in the European markets) prohibiting the import of "chemically-grown" products. These restrictions are also, applied for medicinal plants as the addition of heavy doses of chemical fertilizers adversely affects extraction of active constituents . This finding is faced by the high requirement of many medicinal plants for NPK mineral fertilizers . This has led growers of crop, medicinal and aromatic plants in many countries to adopt organic and biological agricultural practices . Biofertilization is known to compensate apart of the mineral fertilizers to minimize high rates of mineral fertilizers and consequently reduce agricultural costs, as well as, soil pollution. *Majorana hortensis* L. plant is considered as one of the most important medicinal and aromatic crop in Egypt for local and foreign markets with high production and great applications ( foods, medicine and perfumes ). Many trials have been conducted in this concern

for raising the productivity of this plant . In this respect, El-Gadban *et al.* (2002) reported that marjoram plants amended with the highest level of compost ( $15 \text{ m}^3 \text{ feddan}^{-1}$ ) either alone or in conjugation with a mixture of  $\text{N}_2$ -fixing bacteria showed considerable increments in growth characters and contents of N,P and K, as well as ,essential oil (%) and also components of oil ( $\alpha$ -terpineol,  $\alpha$ -terpenolene, cineole and linalool) compared with mineral fertilization treatment. Also, Abo El-Ala (2002) in his study on marjoram and basil showed that root colonization by  $\text{N}_2$ -fixers and P-dissolving bacteria was mostly stimulated in the rhizosphere of inoculated plants with a mixture of these microorganisms . This inoculation treatment, also significantly improved plant growth and oil yield. Similarly, Kandeel *et al.* (2002) reported that dual inoculation of sweet basil with *Azotobacter* + *Azospirillum* supplemented with half or full dose of the recommended mineral N-fertilizer, significantly increased plant growth, oil % and yearly oil yield  $\text{plot}^{-1}$  compared with uninoculated plant given full dose of nitrogen . Balabel (1997) showed highly significant yield responses of potato to K fertilization and inoculation with *B.circulans*. Concerning the effect of VAM, Gautam and Sharma (1996) found that VAM colonization varied in different medicinal plant roots and spores count ranged between 200 to 8900 spores  $100 \text{ g}^{-1}$  of soil . Higher spore count supported good growth . Abdul-Khaliq and Janardhanan (1998) studied the influence of 3 VA mycorrhizal fungi, *Glomus aggregatum*, *G. fasciculatum* and *G. mosseae* on 6 cultivated mints and found that inoculation with VAM fungi increased the shoot biomass of all species . Azcon and Barea (1998) demonstrated that mycorrhizal inoculation largely improved N,P and K uptake by lavender plant roots . EL-Sawy *et al.* (1998) stated that inoculation with a mixture of *Azotobacter*, *Azospirillum* and VAM amended with the full dose of P as rock-phosphate and inorganic N-fertilization, in combination with VAM inoculation, remarkably increased the growth of *Ammi visnaga* plant and its production of khellin . This increase, was accompanied by a high percentage of mycorrhizal root infection and rhizospheric  $\text{N}_2$ -ase activity .

This study aimed to evaluate the effect of inoculation with *A.chroococcum*, *B.circulans* and VAM or their mixture in the presence of half or full dose of the recommended field rate mineral NPK-fertilizers, on the growth, oil yield and chemical composition of marjoram plant. The positive responses of marjoram plants to increased levels of mineral fertilization was considered in this study. Therefore, inoculation with either of 3 biofertilizers, i.e. *A.chroococcum*, *B.circulans* and VAM or their mixture was conjugated with the full and the half of the recommended field of NPK-fertilization . The 2 above-mentioned rates were splitted into 3 doses to: a) minimize the adverse effect of the batch application of mineral nutrients on  $\text{N}_2$ -fixation or mycorrhizal colonization . b) stimulate a more prolonged effect of those biofertilizers . Moreover, the sustained effectiveness of biofertilizers under the above-mentioned conditions, will accommodate the high nutrient requirements of marjoram plants.

## MATERIALS AND METHODS

A field experiment was conducted at the Experimental Farm, Faculty of Agriculture, Ain Shams University, Shobra El-Kheima, for two successive seasons (2000/2001 and 2001/2002). The experiment aimed to study the effect of the interaction between NPK fertilization and single or combined inoculation with 3 biofertilizers (*A.chroococuum*, *B.circulans* and vesicular arbuscular mycorrhizae ) on the growth, oil yield and chemical composition of marjoram plant .

### Seeds and soil

Seeds of marjoram (*Majorana hortensis* L., Family:Lamiaceae) were obtained from Medicinal and Aromatic Research Department of Horticulture Institute, Dokky, Cairo, Egypt, to be used in this study. The soil of the Experimental Farm of Fac. Agric.,Ain Shams Univ., was used for experimentation. The chemical properties of the soil were determined before cultivation using a method described by Page (1982) . The data of the analysis are in Table (1) .

**Table (1): Physical and chemical analysis of the soil used for growing marjoram plant .**

Physical analysis	Sand %		24.5
	Silt %		22.7
	Clay %		52.6
	Soil texture		Clay
Chemical analysis	Electrical conductivity (dSm <sup>-1</sup> ) .		1.3
	PH		7.3
	Soluble anions ( meq/L )	CO <sub>3</sub> <sup>2-</sup>	---
		HCO <sub>3</sub> <sup>-</sup>	4.0
		Cl <sup>-</sup>	3.8
		SO <sub>4</sub> <sup>2-</sup>	2.7
	Soluble cations ( meq/L )	Ca <sup>+</sup>	3.8
		Mg <sup>+</sup>	2.6
		Na <sup>+</sup>	2.2
		K <sup>+</sup>	1.9
	CaCO <sub>3</sub> %		1.1
	Available micronutrients (ppm)	Fe	91.4
		Mn	35.6
		Zn	18.3
Cu		16.4	
Available phosphorus (mg g <sup>-1</sup> )		0.3	
Total nitrogen (mg 100g <sup>-1</sup> )		102	
Organic matter %		1.22	

### Organic manure ,mineral and bio-fertilizers

a)-A farm yard manure was provided from the Milk Replacer Center, Fac. Agric, Ain Shams Univ., Shobra El-Kheima, Cairo, Egypt. It has an organic carbon of 24.3 and 22.8%, total nitrogen of 1.3 and 1.1%, C/N ratio of

(18.69 :1) and (20.73:1), organic matter of 45.5 and 47.2%, moisture content of 5.6 and 7.4% ,P%= 0.4 and 0.3, K%= 1.7 and 1.5 and pH were 6.7and 6.6 for the first and second seasons, respectively, and chemically analyzed according to Page (1982) .

b)-Ammonium nitrate (33.5 % N) ,calcium superphosphate (15.5 % P<sub>2</sub>O<sub>5</sub>) and potassium sulphate (48 % K<sub>2</sub>O ) were used for NPK fertilization, respectively. The inorganic NPK were used at two rates:

1-Full of the recommended field rate ,i.e. 300,200 and 100 kg fed.<sup>-1</sup>of ammonium nitrate, calcium superphosphate and potassium sulphate equal (100, 31 and 48 kg fed.<sup>-1</sup>) of the N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O respectively .

2-Half of the recommended dose ,i.e. 150,100 and 50 kg fed.<sup>-1</sup> equal (50,15.5 and 24 kg fed.<sup>-1</sup>) of N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O respectively . Calcium superphosphate was added during soil preparation, while N and K were added into 3 equal doses as follows: a) 30 days after transplantation b) 2 weeks after the first harvest, and c) 2 weeks after the second harvest . Nine treatments were represented in complete randomized block design with 3 replicates as follows

1- Uninoculated(Control) + full dose of NPK .

2- *A.chroococcum* + half dose of NPK .

3- *A.chroococcum* + full dose of NPK .

4- *B.circulans* + half dose of NPK .

5- *B.circulans* + full dose of NPK .

6- VAM + half dose of NPK .

7- VAM + full dose of NPK.

8-A mixture of *A.chroococcum* +*B.circulans* +VAM+half dose of NPK.

9-A mixture of *A.chroococcum* +*B.circulans* +VAM+ full dose of NPK.

c)-*Azotobacter chroococcum* Az.M15 originated from the rhizosphere of *Majorana hortensis* L. and the silicate bacterium (*Bacillus circulans* ATCC 4513) which was provided from Cairo Mircen, Fac. Agric., Ain Shams Univ., Shoubra El-Kheima, Egypt, were used in this investigation . The two strains were grown separately on modified Ashby's medium (Abd El-Malek and Ishac,1968 ) and modified Alexandrov's medium (Zahra, 1969) for 7 and 4 days at 28 ± 2 °C, respectively. Vesicular arbuscular mycorrhizae (VAM) spores extracted from the rhizosphere of marjoram plants by wet-sieving and decanting technique (Gerdemann and Nicolson ,1963) .and were propagated on roots of maize plants under green house conditions for 12 weeks and then used in this study. The spores represented two main genera of VAM,i.e. *Glomus* and *Gigaspora* .

## **Experimental techniques**

### **Seeds and seedlings inoculation**

Seeds and developed seedlings of marjoram used for inoculation treatments were inoculated prior to sowing in nursery (mid November 2000 or 2001 ) and directly before transplantation in the field. Inoculation was carried out by soaking of seeds or immersing the seedling roots in cell suspension of either *A chroococcum* or *B.circulans* (contained about 10<sup>8</sup> cells ml<sup>-1</sup>) or their mixtures for 30 minutes. Cell culture prepared for bacterial inoculation was supplemented with arabic gum (16%) as an adhesive agent. Inoculated

seeds were dried at room temperature for 1 hour before sowing in nursery . VAM spores manipulated as a single biofertilization treatment or in combination with bacterial mixture, was applied by pipetting 3 ml of spore suspension (contained about 100 spores ml<sup>-1</sup>) in seedbed in nursery .

#### **Layout of experiment**

The farm yard manure was added to the whole soil area used for experimentation one month before cultivation (January) at a rate of 25 m<sup>3</sup> fed<sup>-1</sup> . At the beginning of February, the soil was prepared into 27 plots . The area of each plot was 4m<sup>2</sup>(2x2 m) which contained 3 rows. Ten weeks old seedlings were transplanted in the soil with a distance of 30 cm between transplants. The total number of transplants was 18 plot<sup>-1</sup> . Ammonium nitrate, calcium superphosphate and potassium sulphate were used for NPK fertilization .

At full blooming (during June, August and October), the plants were harvested 3 times and a range of parameters were recorded as follows:

#### **(1)Growth parameters**

Plant growth parameters of marjoram, i.e. plant height, number of branches plant<sup>-1</sup> and herb fresh and dry weights plant<sup>-1</sup> or fed.<sup>-1</sup> for each harvest during both seasons were recorded .

#### **(2)Microbiological parameters**

Microbiological parameters of marjoram rhizosphere were evaluated at the 1<sup>st</sup>, 2<sup>nd</sup> and 3<sup>rd</sup> cut by determining total microbial count and counts of *B.circulans* on modified Bunt and Rovira (Abd El-Hafez, 1966)and modified Alexandrov (Zahra ,1969) media, respectively, using the decimal plate count technique. Estimates of *Azotobacter* populations were determined by MPN technique using Cochran's Tables (Cochran, 1950) .The rates of CO<sub>2</sub> evolution were determined at each cut according to the method of Pramer and Schmidt (1964) modified by Shehata (1972). Nitrogenase activity in the rhizosphere of plants was estimated by acetylene reduction assay (ARA) according to the method of Schollhorn and Burris (1967) .

Mycorrhizal infection (%) in roots of marjoram plants was recorded in root samples collected from plants grown under different treatments according to the method described by Phillips and Hayman (1970) and mycorrhizal spore densities was also carried out according to Gerdemann and Nicolson (1963).

#### **(3)Quantification and analysis of volatile oil**

The volatile oil percentage of plant herb at each harvest was determined according to Guenther (1961) .Gas liquid chromatography (GLC) analysis was also performed on oil samples taken from the first harvest during 2000/2001 season in the Central Lab., Fac. Agric., Ain Shams Univ., Shobra El-Kheima, Cairo, Egypt . The volatile oil constituents of marjoram oil were identified according to Bunzen *et al.* (1969) .

#### **(4)Plant chemical analysis**

NPK contents in dry plant herb were determined during the first growing season after wet digestion . Nitrogen % was determined according to A.O.A.C.(1990) and P as well as K were determined according to the method described by Chapman and Pratt (1978) .

### Statistical analysis

Data of the 3 harvests of either of the 2 seasons were statistically analyzed according to the procedure described by Snedecor and Cochran (1981) and Duncan's Multiple Range Test was used to differentiate means of CO<sub>2</sub> evolution and N<sub>2</sub>-ase activity by using the method of Waller and Duncan (1969).

## RESULTS AND DISCUSSION

### 1-Influence of biological and NPK fertilizers on the microorganisms number and microbial activities in rhizosphere of marjoram plant:

#### 1.1.Densities of total microbial flora, silicate bacteria and azotobacters:

Data presented in Table (2) showed that densities of total microbial flora, silicate bacteria and azotobacters were considerably affected by inoculation, cutting time and NPK mineral fertilization. The highest densities of total microbial flora and azotobacters were  $157.8 \times 10^6$  and  $70.3 \times 10^4$  cfu g<sup>-1</sup> dry rhizospheric soil, respectively at the second cut of the plants inoculated with a mixture of the three biofertilizers in the presence of half dose of NPK fertilizers. Density of silicate bacteria was reached to level ( $108.2 \times 10^3$  cfu g<sup>-1</sup> dry rhizospheric soil) at the third cut in mixed inoculation treatments in the presence of full dose of NPK mineral fertilizers. This may be due to the maximal production of biologically fixed nitrogen during the maturity stage of plant growth (Nelson, 1983). Monib *et al.* (1979) found that inoculation of barley grains with *A.chroococcum*, significantly increased azotobacters population, especially in the rhizosphere.

**Table (2): Effect of inoculation with *A.chroococcum*, *B.circulans*, VAM or their mixture in the presence of full or half dose of NPK fertilizers on counts of total microbial flora, silicate bacteria and azotobacters population in rhizosphere soil of marjoram plant at 2000/2001 seasons.**

Treatments		TM <sup>5</sup>	SB <sup>6</sup>	AZ <sup>7</sup>	TM	SB	AZ	TM	SB	AZ
Inoculation	NPK rate	First cut			Second cut			Third cut		
Uninoculated	fd <sup>1</sup>	24.5	12.3	9.3	54.5	19.2	16.3	32.6	35.1	14.2
	hd <sup>2</sup>	45.6	16.4	23.4	86.6	25.5	67.3	73.4	45.2	49.5
<i>A.chroococcum</i>	fd	32.8	20.5	12.2	77.2	34.6	45.6	54.7	51.6	38.4
	hd	74.9	34.6	16.5	122.3	52.7	22.9	105.0	71.8	20.4
<i>B.circulans</i>	fd	63.8	53.8	14.3	111.2	65.8	18.6	81.3	92.3	16.0
	hd	46.6	21.5	17.4	72.4	38.6	27.9	63.2	61.6	23.1
VAM <sup>3</sup>	fd	33.2	23.8	15.9	68.3	44.8	20.7	52.1	68.3	18.6
	hd	98.3	57.6	47.2	157.8	71.4	70.3	124.5	97.5	58.2
Mixture <sup>4</sup>	fd	84.7	62.3	35.7	145.2	86.2	52.6	115.3	108.2	47.8

1 fd:full dose. 2 hd:half dose 3 VAM: Vesicular arbuscular mycorrhizae.

4 Mixture: *A.chroococcum* + *B.circulans* + VAM.

5 TM : Total microbial flora  $\times 10^8$  cfu g<sup>-1</sup> dry soil.

6 SB : Silicate bacteria  $\times 10^3$  cfu g<sup>-1</sup> dry soil.

7 AZ:Azotobacter population  $\times 10^4$  cfu g<sup>-1</sup> dry soil.

#### 1.2CO<sub>2</sub> evolution and nitrogenase activity

CO<sub>2</sub> evaluation and nitrogenase activity in marjoram rhizosphere were periodically estimated to evaluate microbial activities in the root zone as affected by application of bacterial and VAM inoculants. Data in Table (3)

**Table (3): Effect of inoculation with *A.chroococcum*, *B.circulans*, VAM or their mixture in the presence of full or half dose of NPK fertilizers on CO<sub>2</sub> evolution and nitrogenase activity in the rhizosphere of marjoram plant at 2000/2001 season**

Treatments		CO <sub>2</sub> emitted <sup>5</sup>	N <sub>2</sub> -ase activity <sup>6</sup>	CO <sub>2</sub> emitted	N <sub>2</sub> -ase activity	CO <sub>2</sub> emitted	N <sub>2</sub> -ase activity
Inoculation	NPK Rate	First cut		Second cut		Third cut	
Uninoculated	fd <sup>1</sup>	31.5	14.2	51.4	28.4	38.3	18.6
<i>A.chroococcum</i>	hd <sup>2</sup>	42.6	35.3	64.3	77.5	55.2	41.4
	Fd	36.3	29.5	58.5	61.2	51.6	35.2
<i>B.circulans</i>	Hd	57.2	26.8	71.3	55.6	67.5	32.8
	Fd	50.5	25.6	66.2	43.2	62.4	29.5
VAM <sup>3</sup>	Hd	34.4	24.4	60.8	64.6	54.8	34.6
	Fd	29.2	22.5	53.0	52.7	41.7	30.3
Mixture <sup>4</sup>	Hd	62.2	40.2	90.6	110.4	73.7	58.6
	Fd	51.5	37.3	78.2	84.3	64.1	49.8

1 fd: full dose.

2 hd:half dose

3 VAM: Vesicular arbuscular mycorrhizae 4 Mixture: *A.chroococcum* + *B.circulans* + VAM

5 in mg CO<sub>2</sub> 100g<sup>-1</sup> dry soil

6 in n moles C<sub>2</sub>H<sub>4</sub> h<sup>-1</sup>g<sup>-1</sup> dry soil

Main effects		CO <sub>2</sub> emitted	N <sub>2</sub> -ase activity
Inoculation			
	Uninoculated	47.06 d	15.07 d
	<i>A. chroococcum</i>	61.29 c	30.02 b
	<i>B. circulans</i>	72.35 b	20.41 c
	VAM	52.81 a	19.10 cd
	Mixture	81.38 a	40.09 a
NPK rate			
	Full dose	65.11 a	23.82 b
	Half dose	64.30 a	28.81 a
Cutting time			
	1 <sup>st</sup> cut	53.95 c	19.19 c
	2 <sup>nd</sup> cut	75.77 a	31.71 a
	3 <sup>rd</sup> cut	64.53 b	27.19 b
Significance			
	Inoculation	**	**
	NPK rate	**	*
	Cutting time	**	**
	Interaction	NS	NS

a,b,c,d : Means with the same letter are not significantly different .

\*\* = (P < 0.01)

\* = (P < 0.05)

NS = Not significant

showed that CO<sub>2</sub> evolution and N<sub>2</sub>-ase activity, significantly increased in the rhizosphere of marjoram plants to reach their maximal levels at the second cut then decreased at the third cut. Under that conditions the maximum level of CO<sub>2</sub> evolved and N<sub>2</sub>-ase activity were recorded with a mixed inoculation in the presence of half dose of NPK fertilizers, being, 90.6 mg CO<sub>2</sub> 100g<sup>-1</sup> dry rhizospheric soil and 110.4 n moles C<sub>2</sub>H<sub>4</sub> h<sup>-1</sup>g<sup>-1</sup> dry rhizospheric soil, respectively . Rhizosphere soil of uninoculated marjoram plants gave the lowest levels of CO<sub>2</sub> evolution. These results were in accordance with those of Abo El-Ala (2002) who reported that statistical main effects on the rate of CO<sub>2</sub> evolution due to application of multi-biofertilizer inoculant in the presence of half or full dose of mineral N and P was significantly higher, compared with that of uninoculated marjoram plants at the second cut . Heavy application of

inorganic N-fertilizers is known to inhibit N<sub>2</sub>-ase activity of N<sub>2</sub>-fixers (Pedersen et al., 1978). Low doses of N-fertilizers, on the other hand, promote the establishment of well functional N<sub>2</sub>-fixing association (Neyra and Dobreiner, 1977).

### 1. 3.Mycorrhizal root infection and spore production

Data in Table (4) revealed that mycorrhizal infection of marjoram roots, remarkably increased by inoculation with a mixture of azotobacters, silicate bacteria and VAM and supplementation with half dose of inorganic NPK fertilizers, being 48.83% at the 3<sup>rd</sup> cut. The corresponding spores density produced under this condition was  $7.62 \times 10^3$  spores g<sup>-1</sup> soil. Data, also show that inoculation with VAM appeared to strengthen the stimulatory effect of inoculation with diazotrophs. These findings confirm in part with those obtained by Bagyaraj and Menge (1978) who reported that tomato plants inoculated with both *Glomus fasciculatus* and *A.chroococcum* gave a highest mycorrhizal root infection and spore production. The positive effect noticed on VAM inoculated plants, may be due to that mycorrhizae promote absorption of nutrients from soil by the growing plants. This fact is demonstrated by Ross and Harper (1970) and Allen(1991)

**Table(4): Effect of inoculation with *A.chroococcum*, *B.circulans*, VAM or their mixture in the presence of full or half dose of NPK fertilizers on mycorrhizal root infection of marjoram plant at 2000/2001 season.**

Treatments		Root infection %	Spore number ( $\times 10^3$ g <sup>-1</sup> soil)	Root infection %	Spore number ( $\times 10^3$ g <sup>-1</sup> soil)	Root infection %	Spore number ( $\times 10^3$ g <sup>-1</sup> soil)
Inoculation	NPK rate	First cut		Second cut		Third cut	
Uninoculated	fd <sup>1</sup>	1.21	0.21	7.12	1.10	13.83	2.23
	hd <sup>2</sup>	3.92	0.77	15.71	1.93	19.82	3.43
<i>A.chroococcum</i>	fd	2.75	0.45		1.52	16.76	2.25
	hd	2.43	0.63	14.62	1.45	18.63	2.26
<i>B.circulans</i>	fd	1.82	0.34	12.32	1.32	17.34	2.23
	hd	7.60	1.23	22.27	2.43	37.55	6.25
VAM <sup>3</sup>	fd	5.20	1.14	19.35	2.21	35.00	5.87
	hd	11.22	1.71	29.71	4.60	48.83	7.62
Mixture <sup>4</sup>	fd	9.37	1.55	27.65	2.54	42.60	6.45

1 fd: full dose  
2.hd: half dose

3. VAM: Vesicular arbuscular mycorrhizae  
4. Mixture: *A.chroococcum* + *B.circulans* + VAM

## 2.Influence of biological and NPK fertilizers on plant growth, oil yield and chemical composition of marjoram plant :

### 2.1. Plant growth parameters :

Data of growth performance of marjoram plants are expressed in plant height (Table 5), number of branches plant<sup>-1</sup> (Table 6) and herb fresh and dry weights (Tables 7 and 8) during the two seasons. It is clear from these Tables that, inoculation with *A.chroococcum* or *B.circulans* in the presence of half dose of inorganic NPK-fertilization, remarkably decreased the different plant growth parameters compared with the uninoculated control



given the full dose of NPK. In the meantime, addition of full dose of NPK to *Azotobacter* inoculated plants, increased the recorded values over the control. However, increases were insignificant in the first season, and reached to significant levels during the second one compared with the control. Also, plants inoculated with *B.circulans* and amended with full dose of NPK showed insignificant stimulation in different plant growth characters against the control. VAM inoculation in the presence of half or full dose of inorganic NPK, significantly increased vegetative growth compared with the control. However, the most significant increments in different plant growth parameters were detected during both seasons with mixed inoculation treatment in the presence of half dose of NPK compared with the uninoculated plants given full dose of NPK (control). Application of the triple inocula with full dose of NPK reduced growth parameters compared with those obtained from plants amended with half dose of NPK and grown under the same condition. This finding may be due to the effect of higher dose of the mineral fertilizer that inhibited the activity of bacterial and fungal mixture. The adverse effect may also extend to the ability of the N<sub>2</sub>-fixing bacteria to release phytohormones which are known to stimulate plant growth, absorption of nutrients and photosynthesis (Fayez *et al.*, 1985). The silicate bacteria are also known to decompose aluminosilicate minerals, thus releasing potassium in an available form and enhance K uptake by the plants (Lauwers and Heinen, 1974). The VAM colonization also enhance nutrients uptake of nitrogen, phosphorus, potassium, calcium and water (Lampkin, 1994). The overall results of plant growth reported in this study, were in harmony with those obtained by El-Sawy *et al.* (1998) on *Ammi visnaga*, Abdul Khaliq and Janardhanan (1998) on mint, El-Gadban *et al.* (2002) on marjoram, Abo El-Ala (2002) on marjoram and sweet basil and Kandeel *et al.* (2002) on sweet basil plants.

**Table(5): Effect of inoculation with *A.chroococcum*, *B.circulans*, VAM or their mixture in the presence of full or half dose of NPK fertilizers on marjoram plant height.**

Treatments		Plant height (cm)					
		2001 season/First 2000			Second 2001/2002 season		
Inoculation	NPK rate	1 <sup>st</sup> cut	2 <sup>nd</sup> cut	3 <sup>rd</sup> cut	1 <sup>st</sup> cut	2 <sup>nd</sup> cut	3 <sup>rd</sup> cut
Uninoculated	fd <sup>1</sup>	45.67	50.17	43.17	48.17	49.67	40.67
<i>A.chroococcum</i>	hd <sup>2</sup>	48.50	49.33	41.83	50.50	47.33	40.17
	fd	49.17	56.17		53.83	52.83	48.33
<i>B.circulans</i>	hd	46.33	48.83	41.17	47.50	44.33	38.67
	fd	47.67	53.33	44.33	54.33	49.67	46.83
VAM <sup>3</sup>	hd	51.83	55.33	45.17	53.50	53.67	47.83
	fd	52.83	59.83	47.83	57.33	56.83	50.33
Mixture <sup>4</sup>	hd	55.17	58.67	48.50	56.67	58.33	51.83
	fd	47.17	53.33	42.33	57.50	48.33	43.33
L.S.D. at 5 %		5.85	2.72	3.59	4.47	6.04	6.97
L.S.D. at 1 %		8.05	3.75	4.95	6.16	8.32	9.60

1 fd: full dose  
2 hd: half dose

3. VAM: Vesicular arbuscular mycorrhizae  
4. Mixture: *A.chroococcum* + *B.circulans* + VAM

**Table (6): Effect of inoculation with *A.chroococcum*, *B.circulans*, VAM or their mixture in the presence of full or half dose of NPK fertilizers on number of branches marjoram plant .**

Treatments		Branches number plant <sup>-1</sup>					
		2001 season/First 2000			Second 2001/2002 season		
Inoculation	NPK rate	1 <sup>st</sup> cut	2 <sup>nd</sup> cut	3 <sup>rd</sup> cut	1 <sup>st</sup> cut	2 <sup>nd</sup> cut	3 <sup>rd</sup> cut
Uninoculated	fd <sup>1</sup>	7.67	11.33	17.67	8.33	12.67	16.67
<i>A.chroococcum</i>	hd <sup>2</sup>	8.67	10.67	17.67	7.33	11.00	15.67
	fd	8.33	14.33	19.33	8.67	14.67	19.33
<i>B.circulans</i>	hd	8.33	10.33	17.33	8.00	11.33	15.67
	fd	7.00	12.67	18.67	8.33	13.33	17.67
VAM <sup>3</sup>	hd	8.00	13.67	21.33	8.33	14.33	19.00
	fd	8.67	14.33	22.67	8.67	15.67	21.33
Mixture <sup>4</sup>	hd	9.33	15.67	23.33	8.33	15.33	22.33
	fd	7.67	13.67	20.33	8.00	12.67	19.00
L.S.D. at 5 %		1.38	2.09	1.73	NS	1.41	1.24
L.S.D. at 1 %		1.89	2.88	2.38	NS	1.95	1.70

1 fd: full dose  
2 hd: half dose

3 . VAM: Vesicular arbuscular mycorrhizae  
4. Mixture: *A.chroococcum* + *B.circulans* + VAM

**Table (7): Effect of inoculation with *A.chroococcum*, *B.circulans*, VAM or their mixture in the presence of full or half dose of NPK fertilizers on the herb fresh and dry weights of marjoram plant during 2000/2001 season**

Treatments		FW <sup>5</sup> of herb (g)plant <sup>-1</sup>			DW <sup>6</sup> of herb(g)plant <sup>-1</sup>			SY <sup>7</sup> of herb (ton)fed <sup>-1</sup>	
Inoculation	NPK rate	1 <sup>st</sup> cut	2 <sup>nd</sup> cut	3 <sup>rd</sup> cut	1 <sup>st</sup> cut	2 <sup>nd</sup> cut	3 <sup>rd</sup> cut	FW	DW
Uninoculated	fd <sup>1</sup>	110.33	183.50	153.50	27.50	47.50	43.17	8.052	2.127
<i>A.chroococcum</i>	hd <sup>2</sup>	116.17	169.83	149.17	28.17	44.83	42.17	7.833	2.073
	fd	120.17	194.83	155.83	29.50	49.83	44.67	8.475	2.232
<i>B.circulans</i>	hd	106.67	166.67	153.50	26.33	44.17	43.67	7.683	2.055
	fd	117.33	185.67	164.33	29.33	48.67	45.67	8.412	2.226
VAM <sup>3</sup>	hd	126.50	208.50	167.83	31.83	51.17	47.83	9.051	2.355
	fd	125.33	263.33	182.17	31.17	61.50	52.50	9.789	2.613
Mixture <sup>4</sup>	hd	139.33	257.17	192.33	35.83	66.83	55.50	10.608	2.847
	fd	121.83	210.33	162.50	30.67	54.33	54.67	8.904	2.352
L.S.D. at 5 %		15.22	24.24	8.83	3.90	5.76	2.42	0.434	0.106
L.S.D. at 1 %		20.97	33.38	12.16	5.37	7.94	3.34	0.597	0.145

1fd: full dose  
2 hd: halfdose  
3 VAM: Vesicular arbuscular mycorrhizae  
4 Mixture: *A.chroococcum* + *B.circulans* + VAM

5.FW: Fresh weight of herb plant<sup>-1</sup> (g) .  
6 DW: Dry weight of herb plant<sup>-1</sup> (g)  
7 SY: Stimated yield of herb fed<sup>-1</sup> (ton)

**Table (8): Effect of inoculation with *A.chroococcum*, *B.circulans*, VAM or their mixture in the presence of full or half dose of NPK fertilizers on the herb. fresh and dry weights of marjoram plant during 2001/2002 season.**

Treatments		FW <sup>6</sup> of herb (g)plant <sup>-1</sup>			DW <sup>6</sup> of herb(g)plant <sup>-1</sup>			SY <sup>7</sup> of herb (ton)fed <sup>-1</sup>	
Inoculation	NPK rate	1 <sup>st</sup> cut	2 <sup>nd</sup> cut	3 <sup>rd</sup> cut	1 <sup>st</sup> cut	2 <sup>nd</sup> cut	3 <sup>rd</sup> cut	FW	DW
Uninoculated	fd <sup>1</sup>	114.83	195.67	150.50	29.17	51.17	42.50	8.298	2.211
<i>A.chroococcum</i>	hd <sup>2</sup>	115.50	186.83	146.83	29.50	49.83	41.33	8.085	2.172
	fd	130.33			33.67	57.83	46.17	9.366	2.478
<i>B.circulans</i>	hd	102.50	172.67	144.67	26.50	45.00	40.83	7.557	2.022
	fd	118.33	203.83	156.50	30.33	52.33	43.67	8.616	2.274
VAM <sup>3</sup>	hd	123.67	223.33	168.67	32.17	57.17	46.67	9.282	2.445
	fd	131.17	240.50	184.83	34.17	61.33	52.83	10.017	2.670
Mixture <sup>4</sup>	hd	144.67	243.50	192.33	37.63	63.67	54.50	10.449	2.808
	fd	125.33	201.67	153.67	32.17	52.00	43.83	8.652	2.304
L.S.D. at 5 %		6.78	27.56	17.95	2.25	6.61	3.98	0.569	0.096
L.S.D. at 1 %		9.33	37.95	24.72	3.09	7.73	5.44	0.783	0.132

1fd: full dose

2 hd: halfdose

3 VAM: Vesicular arbuscular mycorrhizae

4 Mixture: *A.chroococcum* + *B.circulans* + VAM

5.FW: Fresh weight of herb plant<sup>-1</sup> (g) .

6 DW: Dry weight of herb plant<sup>-1</sup> (g)

7 SY: Stimated yield of herb fed<sup>-1</sup> (ton)

## 2.2.Herb contents of volatile oil and its constituents :

### 2.2.1.Volatile oil content

Data presented in Table (9) clearly showed that volatile oil percentage of marjoram herb was considerably influenced by different inoculation treatments combined with inorganic NPK–fertilization . The least records were generally observed in uninoculated control plants in all harvests during the two growing seasons . The highest oil percentage, on the other hand, was produced by plants treated with the triple inocula in the presence of half dose of NPK, in the second rank plants inoculated with VAM given full dose of NPK lied of magnitude .The increases in oil % obtained from these two above-mentioned treatments were significant when compared with the control. However, the differences between these two treatments were not significant . Inoculation with *Azotobacter* or *Bacillus* given half dose of NPK showed insignificant increases in oil percentages of marjoram compared with the control . On the other hand, inoculation with *A.chroococcum*, *B.circulans* ,and their mixture with VAM in the presence of full dose of NPK or inoculation with VAM alone with half dose of NPK, significantly increased the oil % over the control . Plants treated with the mixed inoculation treatment and amended with the half dose of NPK gave the highest significant increases compared with other treatments. This treatment yielded about 1.5 fold of oil yield in comparison with the control at both seasons . The second higher value was recorded for plants inoculated with VAM given full dose of NPK. However, plants inoculated with *A.chroococcum* or *B.circulans* amended with half dose of NPK gave lower level of the yearly oil yield fed<sup>-1</sup> compared with the control, but those inoculated with either *A.chroococcum*, *B.circulans*, or their mixture plus VAM given full dose of NPK or inoculation with VAM alone given half dose of NPK , significantly increased oil yield over the control . Similar findings were obtained by El Gadban *et al.* (2002) on marjoram, Kandeel *et al.* (2002) on basil and Abo El-Ala (2002) on marjoram and basil plants.

**Table (9): Effect of inoculation with *A.chroococcum*, *B.circulans*, VAM or their mixture in the presence of full or half dose of NPK fertilizers on percentages and yearly volatile oil yield (fed<sup>-1</sup>) of marjoram herb.**

Treatments		Volatile Oil %							
		First 2000/2001 season				Second 2000/2001 season			
Inoculation	NPK rate	1 <sup>st</sup> cut	2 <sup>nd</sup> cut	3 <sup>rd</sup> cut	EYO <sup>5</sup>	1 <sup>st</sup> cut	2 <sup>nd</sup> cut	3 <sup>rd</sup> cut	EYO
Uninoculated	fd <sup>1</sup>	1.51	1.43	1.37	30.24	1.48	1.41	1.39	31.38
<i>A.chroococcum</i>	hd <sup>2</sup>	1.57	1.43	1.40	29.82	1.56	1.40	1.39	30.60
	fd	1.60	1.51	1.44	33.60	1.59	1.49	1.45	37.20
<i>B.circulans</i>	hd	1.51	1.45	1.40	29.70	1.52	1.39	1.40	29.28
	fd	1.53	1.49	1.41	32.82	1.57	1.45	1.43	33.48
VAM <sup>3</sup>	hd	1.65	1.55	1.44	36.12	1.65	1.56	1.49	36.42
	fd	1.72	1.63	1.51	41.94	1.75	1.65	1.52	43.38
Mixture <sup>4</sup>	hd	1.75	1.63	1.53	46.14	1.80	1.60	1.55	46.32
	fd	1.79	1.63	1.51	38.22	1.67	1.56	1.55	36.90
L.S.D. at 5 %		0.07	0.03	0.06	1.73	0.05	0.04	0.03	1.65
L.S.D. at 1 %		0.09	0.04	0.09	2.38	0.07	0.05	0.05	2.28

1 fd: full dose

2.hd: half dose


3. VAM: Vesicular arbuscular mycorrhizae

4. Mixture: *A.chroococcum* + *B.circulans* + VAM

5.EYO: Estimated Yearly oil yield fed<sup>-1</sup> (liter)

### 2.2.2. Volatile oil constituents

GLC analysis of marjoram volatile oil mixture showed that oil samples obtained from uninoculated control plants showed the presence of 12 components (see chromatogram 1). Ten of them were identified, while the other two components (peaks No.9 and 12) were not identified due to the lack of the authentic samples. Data presented in Table (10) indicate that the major oil components in marjoram oil sample from plants of the control treatment were:  $\alpha$ -terpineol, geranyl acetate, cineole and citronellol, while the other components were found in small quantities. Mixed inoculation in the presence of half or full dose of NPK, resulted in increased concentrations of geranyl acetate, cineole, linalool, limonene and  $\alpha$ -terpenolene while decreased concentration of  $\alpha$ -terpineol was detected in comparison to the uninoculated control plants. Geraniol concentration tended to increase due to inoculation with VAM or *A.chroococcum* plus application of full or half dose of NPK compared with the control. Citronellol concentration on the other hand, was not affected due to different treatments. Similar results were recorded by El-Gadban *et al.* (2002) on marjoram plant.



**Chromatogram (1):** A typical sample GLC trace shows the separation of constituents of marjoram volatile oil for the control treatment (given full dose of inorganic NPK-fertilization)

**Table (10): The obtained constituents of marjoram volatile oil and their percentage as affected by single or mixed inoculation with *A.chroococcum*, *B.circulans* and VAM combined with inorganic NPK fertilization during 2000/2001 season.**

Peak No.	Components	Treatments									
		Inoculation									
		Uninoculated	<i>A.chroococcum</i>		<i>B.circulans</i>		VAM <sup>3</sup>		Mixture <sup>4</sup>		
		NPK rates									
	fd <sup>1</sup>	hd <sup>2</sup>	fd	hd	Fd	hd	Fd	hd	fd	hd	
1	$\alpha$ -Pinene	1.89	1.82	1.71	2.92	2.13	1.85	1.81	0.89	0.72	
2	$\beta$ -Pinene	6.81	7.63	6.53	7.85	5.91	6.93	6.64	4.31	4.81	
3	Limonene	5.80	5.77	5.31	5.38	5.43	6.05	6.21	7.79	7.73	
4	Cineole	10.72	10.11	10.22	10.29	10.44	10.89	11.43	12.40	14.25	
5	Linalool	3.23	3.11	3.83	3.42	3.62	3.32	3.00	5.51	5.70	
6	Citronellol	6.24	6.31	6.34	6.71	5.71	8.31	8.33	6.11	5.96	
7	Geraniol	4.53	6.87	6.90	5.09	4.11	7.12	7.65	5.30	4.63	
8	$\alpha$ -Terpineol	33.32	27.50	29.33	23.58	26.55	28.93	27.39	23.71	21.66	
9	Unidentified	1.95	2.59	1.79	3.37	2.31	1.51	2.05	0.97	1.12	
10	Geranyl acetate	18.61	20.74	21.32	22.91	26.01	19.87	20.72	26.21	26.44	
11	$\alpha$ -Terpendene	5.87	5.61	5.65	9.13	6.00	6.00	6.02	6.62	5.91	
12	Unidentified	1.03	1.94	1.06	2.34	1.78	1.24	0.75	0.17	1.07	

1 fd: full dose  
2.hd: half dose

3 . VAM: Vesicular arbuscular mycorrhizae  
4. Mixture: *A.chroococcum* + *B.circulans* + VAM

### 2.3. N , P and K percentages in plant herb

Data in Table (11) indicated that N, P and K percentages in marjoram plant herb were considerably influenced by the inoculation with *A.chroococcum*, *B.circulans*, VAM or their mixture combined with NPK fertilizers .The highest N,P and K percentages were recorded for mixed inoculation of marjoram supplementation of half dose of NPK . Plants inoculated with VAM given full dose of NPK, and half dose of NPK represented the second and third order of descending magnitude where N, P and K % were higher than other treatments . On the other hand, plants inoculated with *A.chroococcum* or *B.circulans* in the presence of half dose of NPK, showed decreased N, P and K % in their tissues compared with the uninoculated control plants. Treating plants with *A.chroococcum*, *B.circulans* or their mixture plus VAM with full dose of NPK amendment increased N,P and K % in marjoram plant herb over control plants . Similar results were obtained by Azcon and Barea (1998) on lavender, El-Gadban *et al.* (2002) on marjoram and Kandeel *et al.* (2002) on basil plants.

**Table (11): Effect of inoculation with *A.chroococcum*, *B.circulans*, VAM or their mixture in the presence of full or half dose of NPK fertilizers on N, P and K% in marjoram plant herb at 2000/2001 season.**

Treatments		N %			P %			K %		
Inoculation	NPK rate	1 <sup>st</sup> cut	2 <sup>nd</sup> cut	3 <sup>rd</sup> cut	1 <sup>st</sup> cut	2 <sup>nd</sup> cut	3 <sup>rd</sup> cut	1 <sup>st</sup> cut	2 <sup>nd</sup> cut	3 <sup>rd</sup> cut
Uninoculated	hd <sup>1</sup>	2.41	2.67	2.74	0.47	0.52	0.57	2.73	2.81	2.79
<i>A.chroococcum</i>	hd <sup>2</sup>	2.30	2.49	2.67	0.42	0.53	0.57	2.69	2.80	2.83
<i>A.chroococcum</i>	fd	2.49	2.72	2.75	0.48	0.54	0.59	2.72	2.93	2.97
<i>B.circulans</i>	hd	2.29	2.43	2.59	0.37	0.47	0.54	2.65	2.79	2.81
<i>B.circulans</i>	fd	2.35	2.57	2.75	0.47	0.54	0.59	2.84	2.94	2.95
VAM <sup>3</sup>	hd	2.68	2.83	2.98	0.57	0.61	0.73	2.79	2.89	2.99
VAM	fd	2.97	3.01	3.11	0.63	0.69	0.77	2.81	2.97	3.08
Mixture <sup>4</sup>	hd	3.12	3.21	3.25	0.65	0.69	0.86	2.87	2.98	3.12
Mixture	sfd	2.42	2.73	2.89	0.59	0.66	0.71	2.77	2.88	2.93

1 fd: full dose  
2.hd: half dose

3. VAM: Vesicular arbuscular mycorrhizae  
4. Mixture: *A.chroococcum* + *B.circulans* + VAM

## REFERENCES

- A.O.A.C.(1990). Official methods of Analysis of the Association of Official Analytical Chemist. Pub. A.O.A.C. INC. Suite 400,2200 Wilson Boulevard Arlington, Virginia 22201 USA 15<sup>th</sup> Ed. vol. I, pp. 17-22
- Abdel Hafez, A.M.(1966). Some studies on acid producing microorganisms in soil and rhizosphere with the specialreference to phosphate dissolvers Ph.D.Thesis, Fac.Agric., Ain Shams Univ., Cairo, Egypt.
- Abd El-Malek , Y. and Y. Z. Ishac (1968). Evaluation of methods used in counting azotobacters . J. Appl. Bact. , 31: 267-275 .
- Abdul-Khaliq, M. and K. K. Janardhanan (1998). Influence of vesicular-arbuscular mycorrhizal fungi on the productivity of cultivated mints. J. Medic. and Aromat. Plant Sci., 19 (1):7-10 ( c.f.Hort. Abst., 68: 2543).
- Abo El-Ala, H.K. (2002). Biofertilization techniques used for improving production of some medicinal plants in desert soil. Ph.D. Thesis, Inst. Environ. Studies and Res. Ain Shams Univ., Cairo Egypt.
- Allen, M. F. (1991). The ecology of mycorrhizae. Cambredge Univ. Press, pp.74-78.
- Azcon, R.and J.M. Barea (1998). Mycorrhizal dependency of a representative plant species in Mediterranean shrublands (*Lavandula spica* L.) as a key factor to its use for revegetation strategies in desertification threatened. Appl. Soil Ecol., 7 (1): 83-92.
- Bagyaraj, D. J. and J. A. Menge (1978). Interaction between a VA mycorrhizae and *Azotobacter* and their effects on rhizosphere microflora and plant growth . New Phytol., 80 : 567-573.
- Balabel, Naglaa M. A. (1997 ). Silicate bacteria as biofertilizers. M.Sc. Thesis, Fac. Agric., Ain Shams Univ., Cairo, Egypt.
- Bunzen, J. N. G.; J. Labbe; P. Prevot; J. Sperpint and J. Tranchant (1969). Practical manual of gas chromatography. J. Tranchant Ed. El-Seievier Publ. Comp. Amsterdam, London.

- Chapman, H.D. and P.E. Pratt (1978). Methods of analysis for soils, plants and waters. Univ. California, Div. Agric. Sci., pp. 1.
- Cochran, W. G. (1950). Estimation of bacterial densities by means of the "Most Probable Number" . *Biomtrics*, 6 :105-115.
- El-Gadban, E. A. E.; A.M. Ghallab and A. F. Abdel-Wahab (2002). Effect of organic fertilizer (Biogreen) and biofertilization on growth, yield and chemical composition of marjoram plants grown under newly reclaimed soil conditions. Proceedings of the 2<sup>nd</sup> Congress on Recent Technologies in Agriculture. Fac.Agric. Cairo Univ., 28-30 Oct. 2002 vol. II., pp. 344 -361.
- El-Sawy, M.; E. A. Saleh; I. Fendrik and M. S. Sharaf (1998). Effectiveness of dual inoculation with diazotrophs and vesicular arbuscular mycorrhizae on the growth and khellin content of *Ammi visnaga*. Arab Univ. J. Agric. Sci., Ain Shams Univ. , Cairo, 6 (2): 367 - 371.
- Fayez, M.; N. F. Emam and H. E. Makboul (1985). The possible use of nitrogen fixing *Azospirillum* as biofertilizer for wheat plants. Egypt. J. Microbiol., 20 (2): 199 - 206 .
- Gautam, S.P. and R.B. Sharma (1996). Improvement in the medicinal plants through VAM fungi. J. Phytol. Res., 9 (1): 35-37 .
- Gerdemann, J.W. and T.H. Nicolson (1963). Spore of mycorrhizal *Endogone* species extracted from soil by wet sieving and decanting. Trans. Brit. Mycol. Soc., 46: 235-244 .
- Guenther, E. (1961). The Essential Oils. D. Van Norstrand Com. Inc. Canada , Vol. III, 4<sup>th</sup>. Ed., pp. 399-433 .
- Kandeel, A.M.; N.S. Abo-Taleb and A.A. Sadek (2002). Effect of biofertilizers on the growth, volatile oil yield and chemical composition of *Ocimum basilicum* L. plant. Annals Agric. Sci., Ain Shams Univ., Cairo, 47 (1): 351-371 .
- Lampkin, N. (1994). Organic farming. Published by Farming Press Books and Videos Wharfedale Road, Ipswich I P I4 LG, UK., pp.13-85.
- Lauwers, A.M. and W. Heinen (1974 ). Biodegradation and utilization of silicate and quartz. Arch. Microbiol., 95: 67-78 .
- Monib, M.; Y. Abd El-Malek; I. Hosny and M. Fayez (1979). Effect of *Azotobacter* inoculation on plant growth and soil nitrogen .Zbl. Bakt. II Abt., 134 :140-148.
- Nelson, A.D. (1983). Nitrogen fixation associated with grasses in Oregon. Can. J. Microbiol., 22 :523-530 .
- Neyra, C. A. and J. Dobereiner (1977). Nitrogen fixation in grasses . Adv. Agron., 29 :1-38.
- Page, A.L. (1982). Methods of soil analysis. part 2. pp. 39-41. Chemical Microbiological Properties SSSP. INC., Madison, Wisconsin, USA.
- Pedersen, W.L.; K. Chakrabarty; R.V. Klucas and A. K. Vidaver (1978). Nitrogen fixation (acetylene reduction) associated with roots of winter wheat and sorghum in Nebraska. Appl. Environ. Microbiol., 35 (1):129-135.



- Phillips, J.M. and D.S. Hayman (1970). Improved procedures for clearing roots and staining parasitic and vesicular arbuscular mycorrhizal fungi for rapid assessment of infection. Trans. Brit. Mycol. Soc., 55: 158-161.
- Pramer, D. and E.L. Schmidt (1964). Experimental soil microbiology. Burgess Publishing Company, Minnesota, USA.
- Ross, J.P. and J.A. J. Harper (1970). Effect of *Endogone* mycorrhiza on soybean yield. Phyto., 60 :1552-1556 .
- Schollhorn, R. and R.H. Burris (1967). Acetylene as a competitive inhibitor on N<sub>2</sub>-fixation. Proc. Nat.Acad. Sci., USA, 85: 213-220.
- Shehata, S.M. (1972). Evolution of some biological tests as parameters for microbial activities related to soil fertility. Ph. D. Thesis, Fac. Agric., Cairo, Univ., Egypt.
- Snedecor, G.W. and W.G. Cochran (1981). Statistical Methods. 7<sup>th</sup> Ed., Iowa State Univ. Press, Ames. Iowa, USA, p.305 .
- Waller, R.A. and B. D. Duncan (1969). A way for the symometric multiple comparison problem. Amer. Stat. Assoc. J., 3.: 1485-1503.
- Zahra, M. K. (1969). Studies on silicate bacteria. M. Sc. Thesis, Fac. Agric., Cairo Univ., Egypt, pp. 71-76.

### إنتاجية نبات البردقوش تحت تأثير التداخل بين التسميد المعدني و الحيوي عواض محمد قنديل<sup>1</sup> ، محمد سعيد شرف<sup>2</sup>

١-قسم البساتين - كلية الزراعة - جامعه عين شمس -القاهرة- مصر.

٢-قسم الميكروبيولوجيا الزراعية-كلية الزراعة - جامعه عين شمس -القاهرة- مصر.

تم إجراء تجربته حقلية لدراسة تأثير تلقيح نبات البردقوش ببكتريا الأروتوباكتر كروكوسم ، والباسيلس سيركيولانس ، وفطريات الميكوريزا المكونة للحويصلات والتفرعات الشجرية أو خليط منهم في وجود المعدل الحقلية الموصى به كاملاً أو نصف هذا المعدل من التسميد المعدني من انتروجين والفسفور والبيوتاسيوم على نمو النبات ، والأنشطة الميكروبية في الريزوسفير (تمثلة في معدل انطلاق ك<sup>1</sup> واختزال الأستيلين وإصابة الجذر بفطريات الميكوريزا) ، ومحصوليه الزيت الطيار (مع استخدام الكروماتوجرافى لتحديد مكوناته) ، ومحتوى المجموع الخضري من ن، فو ، بو وقد تم تسجيل هذه القياسات في ثلاث حشاشات خلال موسم النمو ، وقد أظهرت نتائج هذه الدراسة أن التلقيح المختلط من مخصبات حيوية مجتمعه مع إضافة نصف المعدل السمادي المعدني (ن ، فو ، بو ) أعطى أعلى زيادات معنوية في القياسات الخاصة بالنمو ، والنسبة المئوية للزيت الطيار وكذا المحصول السنوي من الزيت للقدان وكذلك محتوى المجموع الخضري من ن ، فو ، بو كما أدى لزيادة نسبة بعض الزيوت الطيارة مثل geranyl acetate ، cineole, linalool, limonene and α-terpineol بينما انخفضت نسبة مادة α-terpineol مقارنة بالكنترول كما أدى التلقيح بالخليط إلى زيادة كتافه أعداد الميكروبات الكلية وكذلك الأروتوباكتر وبكتريا السليكات في ريزوسفير البردقوش ، كما زاد معدل انطلاق ك<sup>1</sup> ونشاط إنزيم النتروجينيز ومعدل الإصابة وإنتاج جراثيم الميكوريزا . من جهة أخرى أدى التلقيح بأي من الأروتوباكتر أو الباسيلس نبي وجود نصف معدل التسميد المعدني (ن ، فو ، بو) إلى انخفاض القياسات السابقة الذكر مقارنة بالنباتات غير الملحقة مع التسميد بالمعدل الكامل من التسميد المعدني ولكن أدى استخدام الميكوريزا بمفردها أو بالاشتراك مع خليط البكتريا إلى زيادة معدلات الصفات المختبرة