

## **PATHOGENICITY OF *Beauveria bassiana* TO *Bemisia tabaci* (Gnandius) (HOMOPTERA: ALEYRODIDAE).**

**Aly, Safaa H.**

**Agricultural Research Center, Plant Protection Research Institute,  
Pesticides Tests of cotton Pests Department, Egypt**

### **ABSTRACT**

Present test reported successful infection by *Beauveria. Bassiana* against the whitefly, *Bemisia. Tabaci*. The fungus tested for its pathogenicity to the third – instar nymphs, and tested also against one- and 4-day-old eggs. Also the tests proved a reduction in total protein, fat, and  $\alpha$ -amylase enzyme in the treated insects. Eggs were found to be immune to infection, but mortality of hatching nymphs reached 80-90%. The rate of hatching nymphs infection depended on the age of which the eggs were treated. Mortality of nymphs recorded on the third day after treatment and LC<sub>50</sub> was  $1.12 \times 10^3$  and LC<sub>90</sub> was  $3.8 \times 10^3$  spores/ml.

### **INTRODUCTION**

*Bemesia tabaci*, is a major pest of a wide range of wild and cultivated plants in warm climates worldwide (Osborne, and Landa, 1992). It affects crops grown in greenhouse and field, all year round with short development cycles and high fecundity. It is known as a principal pest of some crops such as vegetables, caused direct damage sucking plants sap and indirect damage by transference the viruses. Although mainly chemical control has been used against *B.tabaci*, it is not always successful, possibly due to the waxy nature of the cuticle (Byrne and Bellows, 1991). And rapid development of resistance (Smith, 1993). The reduced efficiency of insecticidal control and ecological awareness revived the interest in biological control. In this regard, *B. tabaci* is attacked by parasitoids, predators and entomopathogenic fungi. This fungi is among natural occurring antagonists to be considered in aspects of biological control. *Beauveria bassiana* provide good control of aphids and whiteflies (James and Elzen, 2000). The fungus work by attaching to the host surface, then penetrating into the body and killing it. The fungus is available commercially for greenhouse ornamentals and vegetables (Greer, 2000). The potential use of a fungus as a biological control agent is greatly influenced by the susceptibility of the various development stages of the insect. To the pathogene as well as its ability to initiate infection in early stages of the whitefly's development. *Beauveria bassiana* is an insect pathogen for good biological control of whitefly, aphids, thrips, psyllids, weevils and mealybugs in ornamental and vegetables in greenhouses and fields. Bastiaan (1997) used the *B. bassiana* against sweetpotato, silverleaf whitefly in the greenhouse. The present study provides information on the pathogenicity of *B.bassiana* against *B.tabaci*.

### **MATERIALS AND METHODS**

#### **Tested Insects:**

*Bemisia tabaci* was reared on tomato plants under controlled conditions in glasshouse at  $25 \pm 3$  °C,  $50 \pm 20\%$  RH and a photoperiod of 16:8 (light: dark). In order to obtain *B.tabaci* individuals of uniform age, 50-

## **Aly, Safaa H.**

100 adults were placed on small tomato plants for 24-36 h. Then all adults were removed and plants with eggs were transferred to environmental growth chambers for further development of homogeneous populations.

### **Tested fungicide:**

The product Bio-fly ( $30 \times 10^6$ ) cells / ml L is based on the entomogenous fungus *Beauveria bassiana*.

Was prepared by adding 1cm to 1L. of water, and then we prepared 4 concentrations of suspension; ( $7.5 \times 10^3$ ,  $3.75 \times 10^3$ ,  $1.875 \times 10^3$ ,  $0.937 \times 10^3$  spores /ml.) and control.

### **Bioassay procedure for nymphs and eggs of *B.tabaci*:**

Individual tomato leaves with uniformly insects were selected to treatments. Leaf sectors with approximately 50 to 100 insects were used. These leaf pieces bearing 3<sup>rd</sup> instar *B.tabaci* nymphs were immersed in a spore suspension and control for 10 sec. To prevent development of saprophytic fungi, treated leaves were placed for 20-30 min on filter paper to remove excess moisture. The leaves were then placed in petri dishes which were incubated in growth chamber at alternating temperatures of 25 °C (14 h in light and 10 h in the dark). Relative humidity close to 100% was reached by placing a moist filter paper in each petri dish. For aeration purposes, each petri dish was opened daily for 25-30 min. This procedure was necessary to avoid development of saprophytic fungi on whitefly honeydew. Larval mortality was determined daily by counting the number of infected or non-infected individuals per leaf. The test was repeated twice using 4 replicates. Eggs of uniform age (0-1 –day old age, and 4-day old age) were obtained as described earlier. Pathogenicity was determined by calculating the percentage of infected larvae among the total number of emerged larvae.

### **Biochemical analysis:-**

Sampling of individuals started 72hr. after they were immersed in the suspensions. Subsequently, samples were collected at random from each treatment as well as from control. Each sample consisted of about 150-200 alive insects that were weighed.

### **Determination of total protein:-**

The insect was immersed in 96% ethyl alcohol and left 24hr. in alcohol then removed and the extract was taken for soluble protein analysis. The extract was concentrated to 2 ml, and then transferred to tightly closed bottle and kept in the refrigerator until analysis. Total protein content was determined by the method of (Lowry *et al.* 1951).

### **Determination of fat content of treated insects:**

The rapid method of Bligh and Dyer (1959) was applied. Each sample was weighed and homogenized with a mixture of chloroform and methanol to produce a biphasic system, the chloroform layer contained the lipids. This layer was taken in a clean dry beaker (weight before) and chloroform was evaporated by air current. Thenafter, the remaining fat residues and beaker were re-weighed and the lipid content was calculated.

### **Determination of – Emylaze Enzyme of treated insects:**

The enzyme activity was assayed according to Rick and Stegbauer (1974).

## RESULTS AND DISCUSSION

### Pathogenicity of *B. Bassiana* on nymphs:-

Present data indicate that the nymphs of *B. tabaci* are susceptible to fungus *B. bassiana*. The successful infection by *B. bassiana* was also reported by Bastiaan, (1997) who reported that *B. bassiana* is a fungal pathogen of whiteflies and used it in the management of whiteflies in the greenhouse. Data in table (1) show that the nymphs of *B. tabaci* are susceptible to fungus and the high hazard appeared at the higher concentration than those at the lower concentration. The  $LC_{50}$  for the third-instar nymphs was  $1.12 \times 10^3$  and  $LC_{90}$  was  $3.8 \times 10^3$  spors/ml. [Fig (1)].

Table(1): Suseptibility of 3th instar larvae of *B. tabaci* to the entomopathogenic Fungus *B. bassiana* three days after treatment.

Concentrations	No. of treated Larvae (mean 3 Rep.)	Mort (Mean) (%)	Correct Mort. (%)
$7.5 \times 10^3$ spores/ml	100	98	96.04
$3.75 \times 10^3$ spores/ml	70	89	87.22
$1.875 \times 10^3$ spores/ml	100	72	70.56
$0.937 \times 10^3$ spores/ml	90	40	39.2
Control	100	2	0.0




Fig (1): Mortality response among 3<sup>rd</sup>-nymph instar of *B. tabaci* treated with *B. bassiana*

**Aly, Safaa H.**

These data are in agreement with wraight, *et. al.* (1998), who recorded the pathogenicity of *Beauveria bassiana* against the silverleaf whitefly, *Bemisia argentifolii*. The fungi appeared clearly on the treatment dead nymphs after putting in 100% moisture at 25 C as recommended by Butt & Goettel (2000), [see picture (1)].

Picture (1): nymph instar of *B.tabaci* infested by *B.bassiana*



**Pathogenicity of *B.bassiana* on eggs and hatched nymphs:**

Eggs of *B.tabaci* are immune to infection by *B. bassiana* however, in preliminary studies we have noticed that when there is a population consisting of different stages, and eggs are found in the vicinity of infected nymphs or adults, the eggs may become covered with fungal hyphae. Although the chorion of these eggs was not invaded by any of the fungi, the eggs covered with hyphae either did not hatch or hatched with a delay of 3-4 days. The hyphae present on the eggs were found to infect the nymphs immediately upon hatching. 0-1 and 4-day-old eggs were treated with suspension of *B.bassiana*  $10^7$  spors/ml (Gindin *et al.*,2000)and the mortality of hatching nymphs was recorded (Fig (2)).

Always, first-instar emergence began in 7-8 day old eggs, regardless of the time of treatment, and reached approximately 80-90%. The rate of infection of hatching nymphs was found to depend on the age at which the eggs were treated. The firstinfection of nymphs emerging from treated one-day old eggs were observed after 8 days, and the mortality mean for 3 Replication was 12.33%. And the mortality mean was 21.27% after 10 days of treatment. A significant increase in nymphs mortality was obtained when 4-

day-old eggs were treated. Mortality mean was 9.83% at 6 day, 36.5% at 8 day and 81.5% at 10 days after treatment of 4-day-old eggs. These date indicate that the egg treatment with *B.bassiana* has no effect on egg hatch rate, but does affect the mortality of hatching nymphs.




Fig.(2):Pathogenicity of *B. tabaci* eggs and hatched nymphs. Mean percent hatchability and mortality after treatment of 1-and 4-day-old eggs

**Aly, Safaa H.**

The rate of infection depends on the time required for nymph emergence after treatment. The treatment of one- 4-day-old eggs caused lower infection of hatching nymphs at the same time after nymph emergence. This fact reflects a decline in efficiency of inoculation with time on the one hand, but on the other, indicates the survival of the inoculum on leaves for at least 8 days. These data was in agreement with Gindin, *et al.*(2000) who studied the pathogenicity of *V.lecanii* to different stages of *B. argentifolii* and proved that the rate of infection depends on the time required for nymph emergence after treatment.

**Determination of total protein:**

Present data in table (2) indicate the effect of *B.bassiana* on the total soluble protein of *B.tabaci*. the data revealed that the fungus reduced the amount of soluble protein in the treated insects than the control. The mean of the total protein in the treated insects was 1.356 protein per weight (gr.) and control was 2.6858 per gr. The percentage of the decrease than control was 49.512. These data are in agreement with Eman & Sewify, (1991), who recorded a decrease in concentration of the total protein in Aphis insects treated with *V.lecanii*. These data also are in agreement with Gardner *et al.*(1979) and Cheung and Gula (1980), who recorded a decrease in certain haemolymph proteins, amino acids and carbohydrates in insects infected by the fungi, and they mentioned that, this reduction is due to the pathological action of the fungi particular those of higher virulence. Also are in agreement with Gabriel, 1968, Kucera, 1980, Ignofoo, 1981 and Brey and Latge 1986, who stated that the ability of fungi to produce extracellular enzymes lead to changes in haemolymph proteins and amino acids by breaking down proteins bound to chitin and to deterioration of the attached organs. Also this data are in agreement with Jackson *et al.* (1985) who stated that the highly significant quantitative differences in haemolymph protein and amino acids in Aphis due to infection by fungus, *V.lecanii*, and they referred to the ability of all isolates of *V.lecanii* to degrade lipid and protein by extracellular enzymes in the host. These data are in agreement also with Leger *et al.*(1986) who cleared the potentiality of fungal enzymes to degrade the protein and chitin in locust cuticle.

**Table (2): Effect of fungus infection on the total soluble protein of *B.tabaci*.**

Replicates	Amounts of total soluble protein mg per gm	Control	Decrease than control %
1	1.465	3.143	-
2	1.39	2.540	-
3	1.27	2.180	-
4	1.30	2.880	-
Mean	1.356	2.6858	49.512

**Determination of fat content of treated insects:**

Present data in table (3) indicate the effect of *B.bassiana* on the lipid contents of *B.tabaci* insects. The data revealed that the fungus reduced the lipid content in the treated insects than the control. The percentage of the lipids content from 4 replicates of sample treated with fungus was 13.353% but in control was 22.8570%. These data indicated that the fungus infected larvae decrease the lipid contents as effect on the metabolism in the treated insects. These data are in agreement with Smith and Gula, (1982), who stated that a wide variety of natural compounds such as glucose, several amino acids, chitin, starch and fatty acids can be used as carbon and energy source for germination of conidia of fungi, *B.bassiana*, and this fungus can colonize the haemolymph of Colorado beetle larvae, starting in the degradation process, Cermakova and Samsinakova (1960). These results also are in agreement with Jackson et.al,(1985) who referred to the ability of all isolates of *V.lacanii* to degrade lipid and protein by extracellular enzymes in the host. Also data are in agreement with Jagatap, (1973), who stated that the fungus spreads through the blood system, fat bodies, glandular tissues, digestive tract and nervous tissues of the host.

**Table (3):Effect of fungus infection on the lipid contents of *B.tabaci*.**

Replicates	Sample weight (gm)	Lipid content	Lipids content %
Treatment	1	0.500	0.0245
	2	0.5276	0.0772
	3	0.7144	0.1056
	4	0.530	0.1060
	Mean	0.568	0.0783
Control	1	0.565	0.110
	2	0.5578	0.174
	3	0.493	0.1004
	4	0.500	0.102
	Mean	0.52895	0.1216

**Determination of  $\alpha$  - amylase enzyme of treated insect:**

Data in table (4) indicate the effect of *B.bassiana* on the  $\alpha$ -amylase enzyme in the treated insect. The data showed a reduction in the amount of  $\alpha$ -amylase of *B.tabaci* insects treated with the fungus *B.bassiana* 0.11 than control w 82.1 gm. These results demonstrate that the fungi toxin is an inhibitor of insect digestive enzymes act as growth inhibitors of insects. The pathological action of entomopathogenic fungi on various insect species has been studied in relation to the qualitative and quantitative modifications of the haemolymph components (Gardner et al.,1979, Cheung and Gula, 1980). These data are in agreement with Samsinakova and Misikova, (1973) who examined the degradative enzymes as chitinase, protease, and lipase by fungal strains of diverse origins, in relation to their virulence against greater wax moth. Also these data are in agreement with Gardner et al.,(1979) and Cheung and Gula, (1980) who recorded a decrease in certain haemolymph

**Aly, Safaa H.**

proteins, aminoacids and carbohydrate in insects infected by the fungi. Data are in agreement also with Smith and Grula, (1982) who stated that a wide variety of natural compounds such as glucose, several aminoacids, chitin, starch and fatty acids can be used as carbon and energy source for germination of conidia of fungi, *B.bassiana*. Also the data are in agreement with Zacharuk, (1981) who stated that the degradative changes in insect tissues and organs occur before the fungus hyphal invasion due to certain metabolites of fungal origin that are mainly toxic substances.

**Table (4):Effect of fungus infection on the  $\alpha$ - amylase of *B.tabaci*.**

Replicates	Amounts of $\alpha$ -amylase per gm	Control
1	0.09	1.230
2	0.183	1.243
3	0.067	1.01
4	0.10	1.245
Mean	0.11	1.182

## REFERENCES

- Bacchus, S. (2000). Biopesticide technical document. Biopesticide Programs- Environmental Protection Agency, crystal mall 2 (7511C) 1921 Jefferson Davis Highway Arlington, VA 22202 (703): 308-8097.
- Bastiaan M. Dress (1997). Sweetpotato/Silverleaf whitefly management on Texas Greenhouse-grown poinsettia. Texas Agricultural Extension Service. The Texas A & M University System.
- Bligh, E.G. and W.Y. Dyer (1959): A rapid method of total lipid extraction and purification. Can. J. Biochem. Physiol, 37(8):911.
- Butt, T.M. and M.S. Goettel (2000). Bioassays of entomogenous fungi. In: Navon, A and K.R.S. Ascher (eds.): Bioassays of entomopathogenic microbes and nematodes. CAB International, Wallingford, UK, 141-195.
- Brey, P.T. and J.P. Latge. (1986). Integumental penetration of the pea aphid, *Acyrtosiphon pisum* by *Conidiobolus obscurus* (Entomophthoraceae) J. Invertber. Patholl., 38: 335-344.
- Byrne, D.N. and T.S. Jr. Bellows (1991). Whitefly biology. Annu. Rev. Entomol, 36: 431-437.
- Cermakova, A. and A. Samsinakova. (1960). Uber den Mechanismus des Durchdringens des Pilzes *Beauveria bassiana* Say. Ceskoslov. Parasitol., 7:231-236. (C.F. Comprehensive insect physiology, biochemistry and pharmacology, Pergamon Press, Oxford, Chapter 129, Fungal control, 1983).
- Cheung, P.Y.K. and E.A. Grula. (1980). Alteration of haemolymph in corn earworm larvae by the entomopathogen *Beauveria bassiana*. Abstr. Annu. Meet., Am. Soc. Microbiol., 80.

- Eman B.Moursy and G.H. Sewify. (1991). Effect of the entomopathogenic fungus, *Verticillium Lacanii* on haemolymph protein and amino acids of *Aphis Craccivora* and *Brevicoryne brassicae*. Egypt. J.Biol.P.Cont.7(2), 1991,121: 127.
- Gabriel, B.P.(1968). Enzymic activities of some entomophthorous fungi.J.inverteber. Pathol., 11: 70-81.
- Gardner, W.A.; R.M.Sutton and R.Noblet. (1979). Effect of infection by *Beauveria bassiana* on haemolymph proteins of noctuid larvae. Ann. Ent.Soc.Amer., 72: 224-228.
- Gindin,G; N.u.Geschtovt; B.Racah and I. Barah (2000). Pathogenicity of *Verticillium Lacanii* to different developmental stages of the silverleaf whitefly, *Bemisia argentifolii*. Phytoparasitica 28:3, 2000.
- Greer, L.(2000), Greenhouse IMP: Sustainable Aphid control. Pest management Technical Note. Appropriate Technology Transfer for Rural Areas (ATTRA) NCAT Agriculture specialist.
- Ignoffo, C.M.(1981). The fungus *Nomuraea rileyi* as a microbial insecticide. IN microbial control of pests and diseases 1970-1980, PP. 513-538, Ed.H.D.Burges, Academic Press, London.
- Jackson, C.W.; J.B.Heale and R.A.Hall. (1985). Traits associated with virulence to the aphid *Macrosiphoniclla sanborni* in eighteen isolates of *Verticillium Lecanii*. Ann. Appl. Biol., 106: 39-48.
- Jagatap, A.P., (1973). Studies of the entomogenous fungus *M. anisopliae* affecting *Pyrilla* sp. On sugar cane. 3. Pathologic Histology. Maharashtra vinyan Mandir Patrika, 8: 25-30.
- James, R.R.and G.W. Elzen (2000). Antagonism between *Beauveria bassiana* and Imidacloprid when combined for silverleaf whitefly control. United States Department of Agriculture. Agricultural Research Service. Taktran.
- Kucera, M.(1980). Pteases from the fungus *Metarhizium anisopliae* toxic for *Galleria mellonella* larvae. J.Inverteber. Pathol., 35: 304-310.
- Leger, R.J.ST.; R.M. Cooper and A.K. charnley. (1986). Cuticle degrading enzymes of entomopathogenic fungi: cuticle degradation in vitro by enzymes from entomopathogens. J.Invertber. Pathol., 47: 167-177.
- Lowry, O.H., N.J. Rosebrough, A.L. Farriand R.J. Randall. (1951): Protein measurement with the folin. J.Biol. chem. 193: 265-275.
- Obsorne, L.S. and Z. Landa (1992). Biological control of whiteflies with entomopathogenic fungi. Fla. Entomol. 75: 456-471.
- Rick, W. and H.P.Stegbauer. (1974). Amylase measurement of reducing groups. In H.V. Bergmeyer (ed.), Methods of Enzymatic Analysis, 2nd edn., vol. 2,Academic Press, New York.
- Samsinakova, A., and S. Misikova (1973): Enzyme activities in certain entomophagous representative of Deuteromycetes in relationship to their virulence. Ceska mykol., 27: 55-60.
- Smith, P.(1993). Control of *Bemisia tabaci* and the potential of *Paecilomyces fumosoroseus* as a biopesticide. Biocontrol News Gnf. 14:71 N-78N.

**Aly, Safaa H.**

- Smith, R.J., and E.A. Grula (1982). Toxic components of the larval surface of the corn earworm and their effects on germination of *B. bassiana*. J. Invertebr. Pathol., 39: 15-22.
- Wraight, S.P.; R.I. Carrulthers; C.A. Bradley; S.T. Jaronski; L.A. Lacey; P.Wood and S. Galaini-Wraight (1998). Pathogenicity of the entomopathogenic fungi paecilomyces spp and *Beauveria bassiana* against the silverleaf whitefly, *Bemisia argentifolii*. Journal of Invertebrate pathology 71(1), 217-226.
- Zacharuk, R.Y. (1981). Fungal diseases of terrestrial insects. In Pathogenesis of Invertebrate Microbial Diseases. Edited by E.W. Davidson, P. 367-402, Allanheld Osmun Publish.

التأثير الممرض للفطر *Beauveria bassiana* على الذبابة البيضاء  
*Bemisia tabaci*  
صفاء حسنين علي

معهد بحوث وقاية النبات - مركز البحوث الزراعية - مصر

هذه الدراسات أثبتت نجاح الفطر *B.bassiana* في إصابة الذبابة البيضاء *B.tabaci* تم دراسة تأثير الفطر على حوريات عمر ثالث وعلى بيض عمر يوم واحد و أربعة ايام من الوضع. أثبتت الدراسات انخفاض نسبة البروتين والدهون وأيضاً أنزيم الأميليز داخل الحشرات المعاملة. وقد وجد مناعة البيض للإصابة بالفطر *B.bassiana* ولكن نسبة موت الحوريات الفاقسة وصل ٨٠-٩٠%. وقد وجد أن نسبة إصابة الحوريات الفاقسة تعتمد على عمر البيض المعامل. وقد سجل للحوريات عمر ثالث  $LC_{50}$   $1.1 \times 10^3$ ،  $LC_{90}$   $3.8 \times 10^3$  كونيديا/ملل. أظهرت النتائج أنه من الممكن استخدام الفطر *B.bassiana* في مكافحة البيولوجية لحشرة *B.tabaci*.