

EFFECT OF PHOSPHORUS FERTILIZATION AND GRAINS INOCULATION WITH PHOSPHATE DISSOLVING BACTERIA ON MICROBIOLOGY OF RHIZOSPHERE, WHEAT YIELD AND YIELD COMPONENTS

El-Agrodi, M.W.¹; H.A. El-Fadaly¹; H.A. Shams El-Din² and A.M. El-Shehawy³

¹Soil Sci. and Microbiology Depts., Fac. of Agric., Mansoura University, Mansoura, Egypt.

²Soil, Water and Environ. Res. Inst., ARC, Cairo, Egypt.

ABSTRACT

Two field experiments were carried out at Sakha Agricultural Research Station. In this investigation inoculated wheat grain cultivars Sakha 63, *Triticum aestivum* L., were cultivated in clayey soils during the two successive seasons of 1998/1999 and 1999/2000. The inoculation was done using phosphate dissolving bacterial strains namely *Bacillus megatherium* var. *phosphaticum* to study the effect of inoculation with different levels of superphosphate on microbiology of wheat rhizosphere, NPK uptake, and the plant growth as well as yield and yield components.

Obtained data showed increase in five bacterial groups examined in wheat rhizosphere. These groups are P-dissolving bacteria, total viable bacteria, *Rhizobium* spp., *Azotobacter* spp. and *Azospirillum* spp.

Regarding the growth characters, the values of plant height and number of tillers were significantly increased as results of bacterial inoculation as well as P fertilization.

For yield and yield components, the obtained values of the spike weight, 1000-grain weight, grain yield and straw yield also increased as a result of bacterial inoculation with P-dissolving bacteria. After the harvesting, the plant analysis showed increases in the content of nitrogen, phosphorus and potassium either in grains or in wheat straw as well.

This can also take part in reducing the pollution of the soil from the chemicals added every year. Furthermore, this also increased the yield of wheat and the nutritive value as well.

INTRODUCTION

Wheat is the most important food crop for many of the world population especially in the developing countries, such as Egypt. There are several factors that affect the production of wheat plant and the most important one of these factors is the fertilization. The intensive use of expensive mineral fertilizers in recent years, which results in environmental pollution problems has focused the attention of research on the possibility of using biofertilizers as an alternative or complementary for mineral fertilization (Raju *et al.*, 1997). Besides the other plant essential nutrients, phosphorus comes next to nitrogen as a vital nutrient for plants and microorganisms. The inorganic forms of the element in soil are compounds of calcium, iron, aluminum and fluorine. The organic forms are compounds of phytins, phospholipids and nucleic acids, which come mainly by way of decaying vegetation (Subba-Rao, 1988). Besides the other important functions of P in plants, it plays as an

agent of energy transfer and deficiency of available P is more likely to limit crop production than any other material except water. Recently, emphasis has been paid to the possibility of great utilization of unavailable P-forms such as rock-phosphate by the action of P-solubilizing microorganisms. Published data demonstrated that the significant increases in the yield of different crops were possible when inoculated with P-dissolvers (El-Faramawi, 1994). Biofertilizers denote preparations containing living microorganisms, such as bacteria; *Rhizobia*, *Azotobacter*, *Azospirillum* and phosphate dissolvers, such as; *Bacillus megatherium*. These microorganisms can improve the soil fertility by changing unavailable sources of atmospheric nitrogen and soil phosphorus into available form for growing crops. Biofertilizers are considered to be cheap way to recycle the elements to conserve natural resources and to act as protection factor against increasing pollution due to the extensive use of mineral fertilizers (Zaghloul *et al.*, 1996 and Neeru *et al.*, 2000). In the developed countries, chemically produced water-soluble P fertilizers are routinely applied to crops. The P in these fertilizers is initially available for plant use, then rapidly reacts with soil components and becomes progressively less available for plant uptake. In the developing countries, where chemical fertilizers are not available or are too expensive, ground rock P offers a less costly option (Nahas, 1996). This form of P is much less available to plants than standard fertilizers, except in acidic soils. Many soil microorganisms are able to solubilize otherwise unavailable forms of bound P. These organisms are found in most soils but the numbers and types vary from soil to soil. The ability of microorganisms to solubilize P in soil, and make it available for plant use was demonstrated. In the 50 years since this discovery, researchers world-wide have worked to isolate P-solubilizing organisms, study their characteristics, and use them in inoculants to make P more readily available in commercial agriculture (Krishan *et al.*, 1996 and Singh and Kapoor, 1999).

This work was carried out to examine the inoculation of wheat grains with *Bacillus megatherium* var. *phosphaticum* together with different level of superphosphate on microbiology of wheat rhizosphere, The yield and yield components, wheat growth and NPK uptake were also undertaken in this investigation.

MATERIALS AND METHODS

This investigation was carried out at Sakha Agricultural Research Station. In this investigation, wheat grains cultivars Sakha 69 were used. Field experiments were conducted during the two successive seasons of 1998/1999 and 1999/2000.

Soil samples:

Soil samples of experimental fields were collected from Sakha Agricultural Research Station before setting up the trial. Twenty surface samples were taken at ten different locations at 0-20 cm depth. Mixed sample was obtained and saved in plastic bags. Mechanical analysis was carried out using the method described by Piper (1950). The organic carbon content of

soil samples was determined according to Jackson (1958). The soluble cations and anions were determined according the method described by Richards (1954). The electrical conductivity was determined according to the method described by Richards (1954). Their analysis either physical or chemical are presented in Table (1).

Table (1): Some physical and chemical properties of the experimental clayey soil during 1998/1999 and 1999/2000 seasons

Soil properties	Season of cultivation	
	1 st (1998/1999)	2 nd (1999/2000)
Clay (%)	49.19	49.95
Silt (%)	25.74	24.95
Sand (%)	24.28	23.92
CaCO ₃ (%)	2.10	2.15
Organic matter (%)	0.96	1.05
pH (1:2.5 soil water suspension)	8.10	8.00
EC dSm ⁻¹ (Soil paste extract at 25°C)	2.10	2.30
Total phosphorus (ppm)	1200	1250
Available macronutrients (ppm):		
N	22.00	35.00
P	8.30	8.50
K	825.5	914.50
Available micronutrients (ppm):		
Zn	0.98	1.10
Fe	7.85	8.30
Mn	7.50	8.20
Soluble cations meq/liter:		
Ca ⁺⁺	5.55	5.35
Mg ⁺⁺	4.63	4.00
Na ⁺	10.55	11.11
K ⁺	0.40	0.36
Soluble anions meq/liter:		
CO ₃ ⁻	—	—
HCO ₃ ⁻	2.50	2.00
Cl ⁻	10.86	11.17
SO ₄ ⁻	7.77	7.65

Bacterial strain:

An active phosphate dissolving bacterial strain namely *Bacillus megatherium* var. *phosphaticum* was provided from Microbiology Department, Agric. Res. Center, Ministry of Agriculture, Egypt. This strain was maintained on nutrient agar slant at 5°C till used. Strain of *Bacillus megatherium* var. *phosphaticum* was grown on a liquid medium of modified Bunt and Rovira (Abdel-Hafez, 1966) using 250 ml Erlenmeyer flasks containing 100 ml of the medium. The flasks were incubated at 28°C for 7 days. The bacterial growth was washed and suspended in a physiological mineral solution. Grains of wheat plants were washed and soaked for 30

El-Agrodi, M.W.

minutes in the physiological mineral solution contained about 10^6 cells/ml. An adhesive agent Arabic gum was used to obtain coated grains. The coated grains were then air dried in the shade for 30 minutes and sown immediately.

The two field experiments were designed in a split split design. The arrangement of the field experiment involves two factors. The first is the microbial inoculation with phosphate dissolving bacteria as main plots, two treatments were conducted for wheat included uninoculated and inoculated one. The second factor is P-fertilization using superphosphate (15.5% P_2O_5) included five treatments, i.e., 0, 5, 10, 15 and 20 kg P_2O_5 /fed. The treatments were replicated 4 times, and the plot area was $3 \times 3 = 9 \text{ m}^2$.

Wheat grains were sown on 18th November, 1998 in the 1st season and on 21st November, 1999 in the 2nd season. All treatments received nitrogen fertilizer (Ammonium nitrate 33% N) at the rate of 75 kg N/fed in 3 doses, the 1st dose represented 20% of nitrogen applied rate and added during sowing as activator dose, the 2nd dose represented 40% of the nitrogen applied rate and applied before the 1st irrigation (3 weeks from planting) and 3rd dose was 40% of the nitrogen applied rate which broadcasted before the 2nd irrigation (3 weeks from the 1st irrigation). Phosphorus fertilizer was added in one dose before the 1st irrigation.

Harvesting was in 30th May, 1999 and 5th June, 2000 in the 1st and 2nd season, respectively.

Before harvesting, 10 plants from wheat were randomly collected, put in paper bags, then immediately carried to the laboratory. These plants were then used to study the behaviour of plant growth as affected by addition of P_2O_5 and bacterial inoculation with phosphate dissolving bacteria as well. The subjected examination were plant height (cm), and number of tillers.

Yield and yield components:

The grain yield of wheat was determined by threshing the harvested plants and collecting the grains of the area of 9 m^2 (plot), separated from straw, weighed and determined per plot and the obtained results were expressed as ton/fed. Straw yield was obtained by subtracting the values of grain yield from the values of total yield of accumulated weight (grain + straw). The obtained results were expressed as ton/fed.

Plant analysis:

For determination of nitrogen content, samples of shoots after 60 days from planting, grain, as well as straw at maturity from treatments were ground in a mill, portions of each grain and straw (0.2 g) were digested according to Chapman and Pratt (1963). The digested samples were distilled by micro-kjeldahl procedure. The nitrogen content (%) of distillate was determined by titration according to Black *et al.* (1965). Phosphorus content in the digested samples was colorimetrically measured according to the method described by Snell and Snell (1967). Potassium content was measured by the method described in AOAC (1980) using the Perkin Elmer flame photometer.

Microbiological measurements:

The count of total viable content was determined in rhizosphere soil samples according to Vincent (1970) using the soil extract agar medium

(Allen, 1959). By using Bunt and Rovira medium modified by Abdel-Hafez (1966), phosphate dissolving bacteria were counted by the decimal plate count technique. Symbiotic N-fixers expressed as *Rhizobium spp.* were determined by yeast extract-mannitol agar (YMA) medium (Allen, 1959). Non-symbiotic N-fixers experiment as both *Azotobacter spp.* and *Azospirillum spp.* by using the technique of most probable number (MPN) as described by Vincent (1970), the total number of *Azotobacter spp.* were counted in rhizosphere samples. The medium of Ashby modified by Abdel-Malek and Ishac (1968) was used. The positive tubes were distinguished by the presence of the pellicle and by examining stained preparation. By using the semi-solid malate medium of D bereiner (1978), the total number of viable *Azospirillum spp.* were counted in rhizosphere samples. The technique of most probable number (MPN) described by Vincent (1970) was used. The number of viable bacteria were calculated using Cochran's tables (Cochran, 1950) and related to oven dry weight sample.

Statistical analysis:

The obtained results were subjected for statistical analysis according to the procedure outlined by Gomes and Gomes (1984).

RESULTS AND DISCUSSION

Microbiological value of wheat rhizosphere:

Data in Table (2) clearly show that the counts of P-dissolving bacteria in rhizosphere of wheat plant increased with time to reach the maximum values after the 2nd month of sowing under all phosphatic and inoculation treatments.

Table (2): Changes in counts of phosphate dissolving bacteria in rhizosphere of wheat as affected by different rates of P₂O₅ and P-dissolving bacterial inoculation.

P ₂ O ₅ (kg/fed)	Uninoculated					Inoculated				
	Months				Mean	Months				Mean
	1	2	3	4		1	2	3	4	
	P-dissolving bacteria x 10 ³ /cells/g soil					P-dissolving bacteria x 10 ⁴ /cells/gv				
0	30	55	35	30	37.50	30	65	45	30	42.5
5	30	60	40	31	40.25	30	70	45	31	44.0
10	30	62	43	31	41.50	30	75	50	32	46.8
15	31	65	45	32	43.25	30	80	52	33	48.8
20	32	70	50	33	46.25	33	85	55	35	52.0
Mean	30.6	62.4	42.6	31.4	41.75	30.6	75.0	49.4	32.2	46.8
	P	T	I							
F test	**	**	**							
LSD 0.05	22	7.57	9.85							
0.01	30	10.3	10.63							

The inoculation with P-dissolving bacteria high significantly increased the count of bacteria, under all P fertilization treatment compared to uninoculated ones. The highest mean count value is 75.0 x 10⁴ cfu/g soil

El-Agrodi, M.W.

under the inoculated treatment compared to 62.4×10^2 cfu/g soil for uninoculated treatment.

P-fertilization high significantly affected the count of P-dissolving bacteria where the count increased with increasing the level of added P. Under the inoculated treatment the mean count of P-dissolving bacterial increased from 37.5×10^2 cfu/g soil (control) to 40.25, 41.50, 43.25 and 46.25 $\times 10^2$ cfu/g soil as a result of applying 5, 10, 15 and 20 kg P_2O_5 /fed, respectively. In the case of inoculated treatment the mean count of P-dissolving bacteria increased from 42.5×10^4 cfu/g soil (control) to 44.0, 46.8, 48.8 and 52.0×10^4 cfu/g soil for the above mentioned P level, respectively.

Data in Table (2) show also that the inoculation gave several fold increases in the count of P-dissolving bacteria under P-applied levels compared to uninoculation treatments. For example after 2 months from sowing, the count of P-dissolving bacteria was 85.0×10^4 cfu/g soil for the treatment of inoculation and 20 kg P_2O_5 /fed compared to 70.0×10^5 cfu/g soil for uninoculated + 20 kg P_2O_5 /fed treatment, the increase represents 121.4 fold.

The count of P-dissolving bacteria dropped sharply after the 3rd and 4th month from planting. The count after the 4th month is nearly equal the count after the 1st month which means that the increase in the count of P-dissolving bacteria is related to the activity of plant growth.

Data in Table (3) showed that the count of total viable bacteria in the rhizosphere of wheat plant which took the trend of P-dissolving bacteria. Raising the rate of P_2O_5 /fed accompanied with an increase in the count of total viable bacteria but the increase under the inoculation is higher than the increase under the uninoculation state. The mean count increased from 37.5×10^4 cfu/g soil to 43.5×10^4 cfu/g soil under uninoculated treatment, while the mean increase was from 38.8×10^5 cfu/soil to 45.0×10^5 cfu/g soil under inoculated treatment due to raising the rate of P_2O_5 from zero to 20 kg/fed. Maximum increase was attained after 2 months from sowing. The count after 2 months and with 20 kg P_2O_5 /fed was to 65.0×10^4 and to 70.0×10^5 cfu/g soil for uninoculated and inoculated treatment.

As shown in Table (4), data proved the proliferation of the count of *Rhizobium spp.* in wheat rhizosphere with the inoculation with P-dissolving bacteria. In case of uninoculation, the count reach its maximum value after the 2nd month of planting with 20 kg of P_2O_5 /fed. This value was found to be 65×10^2 cfu/g soil. The counter part number in case of inoculation experiments was 80×10^4 under the same conditions. The mean value of *Rhizobium spp.* changed from 26.5×10^2 cfu/g without P fertilization to 35.8×10^2 /g soil fertilized with 20 kg of P_2O_5 /fed in case of uninoculation. These figures became 41.3×10^4 cfu/g and of unfertilized soil which changed to 50.0×10^4 cfu/g of 20 kg P_2O_5 /fed of fertilized soil as can be seen in the same table. The increasing of such bacteria in wheat rhizosphere means more N-fixation and increasing the amount of growth promoting substances. Therefore, all parameters of the plant growth and yield would be increased as found by Ahmed (1995).

Table (3): Changes in counts of the total viable bacteria in rhizosphere of wheat plant as affected by different rates of P₂O₅ and P-dissolving bacterial inoculation.

P ₂ O ₅ (kg/fed)	Uninoculated					Inoculated				
	Months				Mean	Months				Mean
	1	2	3	4		1	2	3	4	
	Total viable bacteria x 10 ⁴ /cells/g soil					Total viable bacteria x 10 ⁵ /cells/g soil				
0	30	55	35	30	37.5	30	60	35	30	38.8
5	30	60	40	30	40.0	30	65	38	32	41.3
10	30	62	43	31	41.5	30	65	40	32	41.8
15	31	63	45	31	42.5	30	68	42	31	42.8
20	32	65	45	32	43.5	33	70	45	32	45.0
Mean	30.6	61.0	41.6	30.8	41.0	30.6	65.6	40.0	31.4	41.9
	P	T	I							
F test	**	**	**							
LSD 0.05	0.1	0.5	0.07							
0.01	0.2	0.7	0.08							

Table (4): Changes in counts of *Rhizobium spp.* in rhizosphere of wheat plant as affected by different rates of P₂O₅ and P-dissolving bacterial inoculation.

P ₂ O ₅ (kg/fed)	Uninoculated					Inoculated				
	Months				Mean	Months				Mean
	1	2	3	4		1	2	3	4	
	<i>Rhizobium spp.</i> x 10 ² /cell/g soil					<i>Rhizobium spp.</i> x 10 ⁴ /cell/g soil				
0	15	50	30	11	26.5	30	65	45	25	41.3
5	15	52	33	12	28.0	30	70	50	30	45.0
10	17	55	35	12	29.8	32	75	50	30	46.8
15	16	60	40	13	32.3	31	78	55	30	48.5
20	18	65	45	15	35.8	31	80	60	30	50.3
Mean	16.2	56.4	36.6	12.6	30.5	30.8	73.6	52.0	29.0	46.4
	P	T	I							
F test	**	**	**							
LSD 0.05	2.23	1.94	7.97							
0.01	3.08	2.64	13.78							

Data recorded in Table (5) reveal that the counts of *Azotobacter spp.* in rhizosphere of wheat increased to reach the maximum value of 83 x 10³ cfu/g soil fertilized with 15 kg P₂O₅/fed under inoculation condition after the 2nd month. The lowest counts was found to be 10 x 10² cfu/g of not fertilized soil under uninoculated treatments after the 1st month. The mean value of *Azotobacter spp.* count was increased from 20.3 x 10² cfu/g soil without P fertilization to reach 22.3 x 10² cfu/g soil with 20 kg P₂O₅/fed and from 37.0 x 10³ cfu/g soil without P to 47.0 x 10³ cfu/g soil with 20 kg P₂O₅/fed in case of uninoculated and inoculated treatments, respectively. This may enhance the N-fixation and consequently encouraged plant growth and nutrient uptake by wheat plants.

Table (5): Changes in counts of *Azotobacter spp.* in rhizosphere of wheat plant as affected by different rates of P₂O₅ and P-dissolving bacterial inoculation.

P ₂ O ₅ (kg/fed)	Uninoculated					Inoculated				
	Months				Mean	Months				Mean
	1	2	3	4		1	2	3	4	
Azotobacter spp. x 10 ² /cells/g soil					Azotobacter spp. x 10 ³ /cells/g soil					
0	10	35	24	12	20.3	21	74	38	15	37.0
5	12	37	25	10	21.0	35	80	50	30	48.8
10	14	37	25	12	22.0	42	81	52	32	51.8
15	12	40	24	10	21.5	40	83	54	33	52.5
20	14	40	25	10	22.3	38	81	51	18	47.0
Mean	12.4	37.8	24.6	10.8	21.4	35.2	79.8	49.0	25.6	47.4
	P	T	I							
F test	**	**	**							
LSD 0.05	2.31	0.21	1.07							
0.01	3.29	0.29	1.48							

The recorded data in Table (6) reveal that the inoculation of wheat grains with *Bacillus megatherium* resulted in an increase in the densities of *Azospirillum spp. var. phosphaticum* colonized the rhizosphere region. The mean increases under zero P₂O₅ were found to be from 17.5 x 10² cfu/g of uninoculated soil to 34.3 x 10³ cfu/g of inoculated soil. The maximum value of *Azospirillum spp.* densities was 78 x 10³ cfu/g soil, which was observed after the 2nd month of inoculation and 10 kg P₂O₅/fed. At the next months, the bacterial counts were sharply dropped and reached their lowest values being 10 x 10² cfu/g soil after the 4th month without addition P and uninoculated treatments as shown in Table (10). This results clearly show an interesting relation between phosphate dissolving bacteria and the number of *Azospirillum spp.* in rhizosphere. These results came in harmony with those obtained by Zaghloul *et al.* (1996). They found that the highest count of *Azospirillum* was activated in soil inoculated with *Bacillus megatherium*.

Wheat growth:

Data in Table (7) show the effect of phosphate fertilization level and grain inoculation of wheat plant on some growth parameters in 1998/99 and 1999/2000 seasons in the field experiments. The data reveal that increasing phosphorous (P₂O₅/fed) resulted in high significant increases in plant height and number of tillers / plant under inoculated and uninoculated treatments. Increasing the rate of P₂O₅/fed increased plant height in case of uninoculation to reach the maximum values to be 101.3 and 101.25 cm in 1st and 2nd seasons, respectively. On the other hand, in case of inoculation the increase of P₂O₅ leads to increase the height of wheat plant to the maximum value to be 104.75 cm for both cultivated seasons. This could be attributed to the role of microorganisms by supplying wheat roots with their fixed nitrogen and improving the vegetative growth. The mean values of plant height under phosphate fertilization were 98.50 and 103.60 cm in the 1st season, while they were 99.00 and 103.30 cm in the 2nd season under uninoculated and inoculated treatments, respectively.

Table (6): Changes in counts of *Azospirillum spp.* in rhizosphere of wheat plant as affected by different rates of P₂O₅ and P-dissolving bacterial inoculation.

P ₂ O ₅ (kg/fed)	Uninoculated					Inoculated				
	Months				Mean	Months				Mean
	1	2	3	4		1	2	3	4	
	Azospirillum spp. x 10 ² /cell/g soil					Azospirillum spp. x 10 ³ /cell/g soil				
0	10	30	20	10	17.5	19	71	33	14	34.3
5	11	32	22	10	19.0	34	75	42	16	41.8
10	12	35	22	11	19.8	38	78	45	20	45.3
15	12	40	25	10	21.8	35	74	44	20	43.3
20	14	45	30	10	24.8	21	70	30	12	33.3
Mean	11.8	36.4	23.8	10.2	20.6	29.4	73.6	38.8	16.4	39.6
	P	T	I							
F test	NS	**	**							
LSD 0.05	--	1.05	1.18							
0.01	--	1.42	1.45							

Table (7): Effect of phosphorus fertilization and bacterial inoculation with *Bacillus megatherium var phosphaticum* on wheat growth.

P ₂ O ₅ (kg/fed)	First season		Second season	
	Plant Height (cm)	No. of tillers / plant	Plant Height(cm)	No. of tillers / plant
	Uninoculated			
0	95.60	3.95	96.180	3.98
5	97.20	3.90	97.75	4.07
10	98.90	4.05	99.25	4.13
15	99.50	4.43	100.63	4.53
20	101.30	4.30	101.25	4.82
Mean	98.50	4.10	99.00	4.30
F test	**	NS	**	**
LSD 0.05	1.36	--	1.73	0.30
0.01	1.55	--	2.93	0.41
	Inoculated			
0	102.40	4.88	102.50	4.50
5	103.00	4.75	102.25	4.95
10	103.75	4.85	103.25	5.05
15	104.00	4.95	103.50	5.08
20	104.75	5.15	104.75	5.60
Mean	103.60	4.90	103.30	5.00
F test	**	**	**	**
LSD 0.05	1.88	0.24	1.50	0.20
0.01	2.18	0.33	2.01	0.27

Concerning the number of tillers / plant, it could be noticed that the same trend with plant height was realized. Under the uninoculated treatments, the number of tillers per plant was significantly increased in 1st season and high significantly increased in the 2nd season, the highest value

El-Agrodi, M.W.

was achieved with applying 20 kg P₂O₅/fed (4.82 tillers/plant). In case of bacterial inoculation, raising the rate of P₂O₅/fed induced high significantly increases in the number of tillers in both seasons. Maximum values were attained due to applying 20 kg P₂O₅/fed (5.15 and 5.60 tillers/plant, in the 1st and 2nd season, respectively). The increase in plant height may be due to the role of fertilization in stimulation of plant cell division and internode elongation (Peng and Li, 1991). The positive effect of phosphorous may be due to the role of phosphorous in photosynthesis and respiration in addition to its role in cell division and development of meristematic tissues as reported by Ashour (1998).

Inoculation of wheat grains by phosphate solubilizing bacteria increased the mean values of tillers per plant in both growing seasons. The mean values of tillers / plant were 4.10 and 4.30 in case of uninoculation, while they were 4.90 and 5.00 in case of bacterial inoculation for the first and the second seasons, respectively. This increase may be due to the amount of biological nitrogen fixed by the three different bacterial groups found in plant rhizosphere. This in addition to vitamins and auxin-like growth substances and the considerable amount of ascorbic acid, which usually liberated by rhizosphere microbes. Wheat plant can assimilate these materials and reflected their beneficial effect on all parts of the plant (Hegazy *et al.*, 1995).

Yield and yield components:

Data in Table (8) show the values of spike length (cm), spike weight (g), 1000-grain weight (g), grain and straw yield (ton/fed) as affected by phosphate fertilization levels and grain inoculation by phosphate dissolving bacteria in 1998/99 and 1999/2000 cultivation seasons. In case of the uninoculated treatment, in the first season the data clearly show that increasing phosphate levels significantly increase spike length to reach the maximum length of 9.7 cm when using 20 kg P₂O₅/fed. Spike weight also increased to reach 1.94g when using 20 kg P₂O₅/fed. For the 1000-grain weight, the same trend was observed, so the weight of 1000 grain was 42.35 g under the highest P₂O₅ level (20 kg/fed)..

Similar effect of P₂O₅ was found with the grain yield as well as straw yield in ton/fed. The values were 2.51 and 5.14 ton/fed for grain and straw yield, respectively. Similar trend was found in the 2nd season, where all studied parameters were high significantly increased as the rate of P₂O₅/fed increased giving the highest values of 10.1 cm, 2.02 g, 42.56 g, 2.52 ton/fed and 5.13 ton/fed for spike length, spike weight, 1000-grain weight, grain yield and straw yield, respectively at the highest P₂O₅ rate (20 kg P₂O₅/fed)..

The bacterial inoculation of wheat grain with phosphate dissolving bacteria high significantly increased the spike length, giving the maximum value in the 1st season of 9.88 cm when using 15 kg P₂O₅/fed. The obtained value of the spike weight is 2.34g when using 20 kg of P₂O₅/fed. The other values obtained in the 1st season are 43.41, 5.53 and 5.29 for 1000-grain weight (g), grain yield (ton/fed) and straw yield (ton/fed) by using 20 kg of P₂O₅/fed, respectively. The increase in spike length due to adding P₂O₅ together with *Bacillus megatherium* var. *phosphaticum* leads to increase in cell elongation and cell division. This is due to increase in photosynthesis and

nitrogen fixed by bacteria present in wheat rhizosphere. These explanation is in accordance with that of Ishac (1988), Sharief *et al.* (1998) and El-Naggar (1999). Also Singaram and Katharaman (1991) revealed that using *Bacillus megatherium* increased grain and straw yield of maize.

Regarding the values obtained in the 2nd season, it could be seen that using of 20 kg P₂O₅/fed achieved the highest value for each examined parameters as shown in Table (8). These values are 10.30 cm, 2.29g, 43.75g, 2.55 ton/fed and 5.56 ton/fed for spike length, spike weight, 1000-grain weight, grain yield and straw yield, respectively. The increase in straw yield may be mainly due to the increase in plant height and number of tillers per plant.

Table (8): Effect of phosphorus fertilization and bacterial inoculation with *Bacillus megatherium* var. *Phosphaticum* on wheat yield and yield components in field experiments.

P ₂ O ₅ (kg/fed)	First season					Second season				
	Spike Length (cm)	Spike Weight (g)	1000-grain weight (g)	Grain yield (ton/ fed)	Straw yield (ton/ fed)	Spike Length(cm)	Spike Weight (g)	1000-grain weight (g)	Grain yield (ton/ fed)	Straw yield (ton/ fed)
Uninoculated										
0	8.85	1.83	39.86	2.24	4.18	8.97	1.83	40.01	2.24	4.14
5	9.00	1.87	40.61	2.31	4.36	9.10	1.90	40.68	2.31	4.37
10	9.03	1.88	41.11	2.38	4.90	9.15	1.94	41.18	2.38	4.88
15	9.00	1.92	41.94	2.49	5.04	9.80	1.95	42.21	2.49	5.21
20	9.70	1.94	42.35	2.51	5.14	10.10	2.02	42.56	2.52	5.13
Mean	9.12	1.89	41.18	2.38	4.72	9.42	1.93	41.33	2.39	4.74
F test	**	**	**	**	**	**	*	**	**	**
LSD	0.46	0.09	0.22	0.02	0.26	0.45	0.03	0.20	0.03	0.13
0.05	0.63	0.15	0.29	0.03	0.35	0.64	—	0.28	—	0.18
0.01										
Inoculated										
0	8.97	1.90	41.03	2.30	4.32	9.05	1.91	41.01	2.30	4.35
5	9.25	1.94	41.61	2.35	4.49	9.30	1.93	41.69	2.36	4.51
10	9.63	2.06	42.10	2.40	4.95	9.50	2.09	42.28	2.41	5.24
15	9.88	2.16	42.83	2.52	5.28	9.90	2.08	43.15	2.54	5.74
20	9.73	2.34	43.41	2.53	5.29	10.30	2.29	43.75	2.55	5.56
Mean	9.45	2.08	42.19	2.42	4.87	10.35	2.06	42.37	2.43	5.08
F test	**	**	**	**	**	*	*	**	*	**
LSD	0.18	0.13	0.24	0.01	0.11	0.20	0.17	0.24	0.02	0.10
0.05	0.24	0.21	0.33	0.01	0.14	—	—	0.33	—	0.14
0.01										
P x I	NS	S	NS	NS	NS	NS	NS	NS	NS	NS

N, P and K content:

Data presented in Table (9) show that the N content of wheat organs increased due to raising the P₂O₅ so that the maximum values in case of uninoculation were obtained when using 20 kg P₂O₅ / fed. These values are 1.93, 2.38; 2.04, 2.29 and 0.380, 0.343% in the 1st and 2nd season for shoots at boating stage, grain and straw at maturity, respectively.

El-Agrodi, M.W.

Table (9): Effect of phosphorus fertilization and bacterial inoculation with *Bacillus megatherium* var. *Phosphaticum* on N content (%) of wheat plant.

P ₂ O ₅ (kg/fed)	First season			Second season		
	Boating stage	Grain at maturity	Straw at maturity	Boating stage	Grain at maturity	Straw at maturity
Uninoculated						
0	1.81	1.86	0.285	1.93	1.99	0.293
5	1.84	1.90	0.298	2.08	2.09	0.305
10	1.87	1.93	0.315	2.24	2.19	0.323
15	1.89	1.94	0.340	2.29	2.25	0.335
20	1.93	2.04	0.380	2.38	2.29	0.343
Mean	1.87	1.93	0.324	2.18	2.16	0.320
F test	**	**	**	**	**	**
LSD 0.05	0.02	0.07	0.010	0.04	0.037	0.01
0.01	0.03	0.10	0.014	0.06	0.051	0.02
Inoculated						
0	1.86	1.93	0.298	2.04	2.04	0.305
5	1.92	2.01	0.313	2.18	2.12	0.333
10	1.93	2.11	0.325	2.30	2.25	0.343
15	1.94	2.14	0.345	2.32	2.29	0.368
20	1.97	2.16	0.400	2.42	2.35	0.388
Mean	1.92	2.07	0.336	2.25	2.21	0.347
F test	**	**	**	**	**	**
LSD 0.05	0.02	0.06	0.018	0.04	0.03	0.019
0.01	0.03	0.09	0.017	0.06	0.05	0.021
P x I	NS	NS	NS	S	NS	NS

On the other hand, in case of inoculated treatment, tabulated data prove also that the use of 20 kg P₂O₅ / fed was the best for all tested parts of plants. The obtained values were 1.97, 2.42; 2.16, 2.35 and 0.400, 0.388% for shoots at boating stage, grain and straw at maturity in the 1st and 2nd season, respectively. Furthermore, data in Table (9) show clearly that the inoculation with P-dissolving bacteria magnified the nitrogen content of wheat plant organs as compared with uninoculated treatment under the phosphorus application treatments. For example, the mean values of N% of grain, under uninoculated treatment are 1.93 and 2.16% for the 1st and 2nd season. This values are 2.07 and 2.21% under inoculation treatments.

Data presented in Table (10) clearly show that the increasing of P-fertilization (P₂O₅) levels resulted in significant increases in P-content (%) in wheat shoot at boating stage after 60 days of planting. The contents of phosphorus were also increased in wheat grains and straw at maturity either in case of uninoculated or inoculated treatments. The maximum values of P-contents were noticed under the treatments of 20 kg P₂O₅ / fed either in case of uninoculation or inoculation treatment.

Regarding the data of non inoculated treatments, the values of P% were 0.365, 0.428; 0.350, 0.400 and 0.118, 1.43 in the 1st and 2nd season for wheat shoots at boating stage, grain and straw at maturity, respectively. These values, on the other hand, become 0.398, 0.435; 0.355, 0.418 and

0.125, 0.153 in the 1st and 2nd season for shoots at booting stage, grain and straw at maturity, respectively, when wheat grains were inoculated before planting by P-dissolving bacteria.

The increase of P content in different wheat plant organs due to phosphorus fertilization may be attributed to the increase in soil available P and consequently the high efficiency of the roots in absorbing various nutrients including phosphorus. Neeru *et al.* (2000) reported that the higher levels of phosphorus in soil were associated with increase in N and P content of plant. The observed increase was definitely due to the synergistic effect between the examined bacterial groups in plant rhizosphere. These findings could be attributed to that phosphorus dissolving bacteria increased the soluble phosphorus in root zone, so the plant absorbs more P under the inoculation treatments. These results are similar to those obtained by Saber *et al.* (1981) and Singh and Kopper (1999).

Table (10): Effect of phosphorus fertilization and bacterial inoculation with *Bacillus megatherium* var. *Phosphaticum* on P content (%) of wheat plant.

P ₂ O ₅ (kg/fed)	First season			Second season		
	Booting stage	Grain at maturity	Straw at maturity	Booting stage	Grain at maturity	Straw at maturity
	Uninoculated					
0	0.300	0.295	0.095	0.308	0.320	0.105
5	0.310	0.305	0.103	0.325	0.343	0.118
10	0.323	0.320	0.108	0.340	0.350	0.123
15	0.345	0.338	0.115	0.367	0.378	0.128
20	0.365	0.350	0.118	0.428	0.400	0.143
Mean	0.329	0.372	0.108	0.356	0.358	0.123
F test	**	**	**	**	**	**
LSD 0.05	0.012	0.011	0.007	0.015	0.015	0.010
0.01	0.017	0.015	0.010	0.021	0.020	0.014
	Inoculated					
0	0.310	0.305	0.105	0.315	0.338	0.113
5	0.345	0.318	0.108	0.340	0.343	0.125
10	0.353	0.328	0.115	0.345	0.358	0.135
15	0.385	0.343	0.123	0.405	0.395	0.140
20	0.398	0.355	0.125	0.435	0.418	0.153
Mean	0.358	0.329	0.115	0.368	0.370	0.133
F test	**	**	**	**	**	**
LSD 0.05	0.016	0.012	0.008	0.015	0.014	0.009
0.01	0.022	0.017	0.012	0.021	0.020	0.013
P x I	S	NS	NS	NS	NS	NS

Data presented in Table (11) reveal that the phosphorus fertilization used for wheat plant resulted in significant increases of K% content in wheat shoots at booting stage, grain and straw at maturity under uninoculated and inoculated treatments. So the highest K% was achieved when P₂O₅ added at the rate of 20 kg P₂O₅ / fed however the wheat grains before planting inoculated or not but the values of K% was higher under inoculated

El-Agrodi, M.W.

treatments. The obtained values in case of uninoculated treatment are 2.01, 2.25; 0.755, 0.875 and 1.16, 1.42 in the 1st and 2nd season for shoots at boating stage, grain and straw at maturity, respectively. These values were increased due to inoculation with P-dissolving bacteria to be 2.18, 2.33; 0.823, 0.893 and 1.23, 1.46 in the 1st and 2nd season for shoots at boating stage, grain and straw at maturity, respectively. It is known that increasing soluble P in the root zone well increase the root system, so absorb more nutrients. These results are in agreement with those obtained by Neeru *et al.* (2000).

Table (11): Effect of phosphorus fertilization and bacterial inoculation with *Bacillus megatherium* var. *Phosphaticum* on K content (%) of wheat plant.

P ₂ O ₅ (kg/fed)	First season			Second season		
	Boating stage	Grain at maturity	Straw at maturity	Boating Stage	Grain at maturity	Straw at maturity
Uninoculated						
0	1.82	0.680	0.900	1.99	0.715	1.23
5	1.88	0.698	0.953	2.09	0.752	1.33
10	1.89	0.728	1.01	2.14	0.798	1.35
15	1.92	0.737	1.11	2.19	0.850	1.39
20	2.01	0.755	1.16	2.25	0.875	1.42
Mean	1.90	0.719	1.03	2.13	0.798	1.34
F test	**	**	**	**	**	**
LSD 0.05	0.08	0.027	0.050.07	0.06	0.032	0.020.03
0.01	0.11	0.037		0.09	0.044	
Inoculated						
0	1.92	0.665	0.982	2.11	0.750	1.28
5	2.06	0.702	1.08	2.18	0.808	1.34
10	2.12	0.750	1.14	2.22	0.838	1.35
15	2.12	0.792	1.16	2.28	0.863	1.42
20	2.18	0.823	1.23	2.33	0.893	1.46
Mean	2.08	0.746	1.12	2.22	0.830	1.37
F test	**	**	**	**	**	**
LSD 0.05	0.08	0.025	0.070.09	0.07	0.027	0.03
0.01	0.11	0.033		0.09	0.038	0.04
P x I	NS	S	NS	NS	NS	NS

Concerning the combination of the three elements, nitrogen, phosphorus and potassium, many researchers in different countries revealed that NPK have active role to increase grain yield and yield components. NPK have also supreme effect on plant growth and hence affect grain yield, yield components and grain quality as well. Nitrogen, phosphorus and potassium are essential for wheat plants as well as they are basic for nucleic acid, NAD, NADP, co-enzymes, ATP, meristemic cells of plants. nucleic proteins and they are share through ATP in active amino acids to protein synthesis.

On the light of the obtained results. it can be, generally, concluded that the inoculating wheat plants *Triticum aestivum* L variety Sakha 69 cultivated in clayey soil with an active bacterial strain as phosphate dissolver namely *Bacillus megatherium* var. *phosphaticum* is of great importance. This lead to significant increases in the biofertility of soil as well as the yield of the plant growth. The biofertility of soil expressed in increasing the number of different

bacterial groups in rhizosphere area. These groups are the phosphate dissolving bacteria, the total viable bacteria, *Rhizobium spp.*, *Azotobacter spp.* and *Azospirillum spp.* These groups have active role in releasing phosphorus in addition to the N₂-fixation process and the degradation of the organic materials by the enzymatic systems they have.

This can also take part in reducing the pollution of the soil from the chemicals added every year. Furthermore, this also increased the yield of wheat and the nutritive value as well.

REFERENCES

- Abdel-Hafez, A.M. (1966). Some studies on acid producing microorganisms in soil and rhizosphere with special reference to phosphate dissolvers. Ph.D. Thesis, Fac. of Agric., Ain Shams Univ., Cairo, Egypt.
- Abd El-Malek, Y. and Y. Ishac (1968). Evaluation of methods used in counting *Azotobacter*. J. Appl. Bacteriol., 31:261-270.
- Ahmed, A.A. (1995). Response of wheat plants to nitrogen and biological fertilization under condition of north west Coast of Egypt. M.Sc. Thesis, Fac. of Agric., Ain Shams Univ., Cairo, Egypt.
- Allen, T.N. (1959). Experiments in Soil Bacteriology. Burgess Publishing Co, Minneapolis, Minnesota, USA.
- A.O.A.C. (1980). Official Methods of Analysis. 13th ed., Association of Official Analytical chemists. Washington, DC, USA.
- Ashour, S.A. (1998). Influence of biofertilization and phosphorus application on growth and yield of potato (*Solanum tuberosum*, L.). Vegetable Res. Dept., Hort. Res. Inst., Agric. Res. Center, Cairo, Egypt.
- Black, C.A.; D.D. Evans; L.E. Ensminger; J.L. White and F.E. Clark (1965). Methods of Soil Analysis. Part I, Chap. 18, Water Capacity by Klute A., ASA Monograph Series No. 9, Amer. Soc. Agro., Madison, Wisconsin, USA.
- Chapman, H.D. and P.F. Pratt (1963). Method of Analysis for Soils, Plant and Waters. California Univ., Div. of Agric. Sci.
- Cochran, W.G. (1950). Estimation of bacterial densities by mean of the "Most probable number". Biometrics, 6:105-115.
- D bereiner, G. (1978). Influence of environmental factors on the occurrence of *Azospirillum lipoferum* in soil and root. In environmental role of N-fixing blue green and a symbiotic bacteria *E. coli*. Bull. Stockholm, 26:343-352.
- El-Faramawi, Fatma, K. (1994). Role of mycorrhizal fungi in the growth of tomato plant and in the biological control of fusarium wilt. M.Sc. Thesis, Fac. of Sci., Ain Shams Univ., Egypt.
- El-Naggar, E.M.A. (1999). Efficiency use of bio and chemical fertilizers on wheat. Ph.D. Thesis, Fac. of Agric., Mansoura Univ., Egypt.
- Gomes, K.A. and A.A. Gomes (1984). Statistical Procedures for Agricultural Research. John Wiley and Sons, Inc., New York.
- Hegazy, M.H.; F.K. Abd El-Fattah and D. Abadi (1995). Effect of algalization, nitrogen fertilization and plant density on rice yield and its components. Annals Agric. Sci., Ain Shams Univ., Cairo, 40(2):547-557.

El-Agrodi, M.W.

- Ishac, Y.Z. (1988). Inoculation With Associative N₂-Fixers in Egypt. Plant and Soil. Kluwer Academic Publishers Dordrecht, Boston, London, P 114.
- Jackson, M.L. (1958). "Soil Chemical Analysis". Prentice-Hall, Inc., England, Cliffs, 219-221.
- Krishan, C.; K.J.Singh and K. Chandra (1996). Effect of Azotobacter and phosphate solubilizing microorganisms on yield of rice crop. Environ. Ecol., 14(1):39-41.
- Nahas, E. (1996). Factors determining rock phosphate solubilization by microorganisms isolated from soil. World J. Microbiol. and Biotechnol., 12:567-572.
- Neeru, N.; K. Vivek; K. Rishi; D. Annette; G. Andreas and M.Wolfgana (2000). Effect of P-solubilizing *Azotobacter chroococcum* on N, P, K uptake in P responsive wheat genotypes grown under greenhouse conditions. J. Plant Nut. and Soil Sci., 163(4):393-398.
- Peng, J.H. and Y.C.Li (1991). Relationship of genetic variances of quantitative characters in Indica rice to nitrogen level. Inter. Rice Res. Newletters, 16(3):6-7.
- Piper, C.S. (1950). Soil and Plant Analysis. 1st ed. Int. Publ. New York, USA, 47-79.
- Raju, R.A.; K.M.N. Reddy and K.A. Reddy (1997). Performance of phosphorus carriers in wetland rice (*Oryza sativa*) on iverina alluvials. Indian J. Agronomy, 42(2):272-274.
- Richards, L.A. (1954). Diagnosis and improvement of saline and alkaline soils. US salinity Laboratory. California, Handbook No., 60:94-106.
- Saber, M.S.M.; M. Yousry and M. O. Kabesh (1981). Effect of inoculation with phosphate dissolving bacteria on K-uptake by pea plants cultivated in a calcereous soil. Egypt. J. Soil Sci., 21(2):143-152.
- Sharief, A.E.; S.E. El-Kaila; A.A.Leilah and H.E.M. Mostafa (1998). Response of some wheat cultivars to nitrogen fertilizer levels and biological fertilization. J. Agric. Sci. Mansoura Univ., 23(12):5807-5816.
- Singaram, P. and G.V. Kathadaraman (1991). Direct, residual and cumulative effects of phosphatic fertilizers on the yield attributes and field of fingermillet, maize and blackgram grown in a cropping system. Fertilizer News, 36(8):21-27.
- Singh, S. and K.K. Kapoor (1999). Inoculation with phosphate-solubilizing microorganisms and a vesicular-arbuscular myco-rrhizal fungus improves dry matter, yield and nutrient uptake by wheat grown in a sandy soil. Biol. Fertil. Soils, 28:139-144.
- Snell, F.D. and C.T. Snell (1967). Coloimetric Methods of Analysis. Van Nostrand Company Inc., 551-552.
- Subba-Rao, N.S. (1988). Biofertilizers in Agriculture. Oxford and TBH Pub. Co. Ltd., New Delhi, Bombay & Calcutta, 134-141.
- Vincent, I.N. (1970). A Manual for the Practical Study of the Root Nodule Bacteria. Blackwell Scient. Publications, Oxford & Ediburgh, UK.
- Zaghloul, R.A.; M.H. Mostafa and A.A. Amer (1996). Influence of wheat inoculation with mycorrhizal fungi, phosphate solubilizing bacteria and *Azospirillum* on its growth and soil fertility. Annals Agric. Sci. Moshtohor, 34(2):611-626.

تأثير التسميد الفوسفاتي وتلقيح حبوب القمح بالبكتيريا المذيبة للفوسفات على
ميكروبيولوجيا الريزوسفير ومحصول القمح الناتج ومكوناته
محمد وجدى العجرودى^١ - حسين الفضالى^٢ - حسن إبراهيم شمس الدين^٣ -
عادل محمد بهجت الشهاوى^٣
^١ قسم الأراضى - قسم الميكروبيولوجى - كلية الزراعة - جامعة المنصورة
^٢ قسم تحسين الأراضى - معهد بحوث الأراضى والمياه والبيئة - مركز البحوث الزراعية -
مصر .

تم دراسة تأثير تلقيح حبوب القمح (سحا ٦٩) بالبكتيريا المذيبة للفوسفات وذلك على
ميكروبيولوجيا ريزوسفير النبات وكذلك على محصول القمح الناتج ومكوناته وذلك خلال التجارب
الحقلية .

أجريت تجربتان حقليتان بمحطة بحوث سحا بكفر الشيخ فى أراضى طميية خلال موسم
الزراعة ١٩٩٨ / ١٩٩٩ ، ١٩٩٩ / ٢٠٠٠ حيث تم زراعة حبوب القمح صنف سحا ٦٩ والتي
تم تلقيحها بالبكتيريا المذيبة للفوسفات متمثلة فى ميكروب *Bacillus megatherium*
var.phosphaticum

بينت النتائج المتحصل عليها زيادة تعداد البكتيريا المذيبة للفوسفات فى منطقة الريزوسفير
مقارنة بالكنترول هذا علاوة على زيادة بعض المجاميع البكتيرية الأخرى مثل التعداد الكلى
للبكتيريا الحية وتعداد ميكروبات الريزوبيم وكذلك بكتيريا الأزوتوباكتر والأزوسبيريلم .
وبالنسبة لقياسات النبات فقد إزدادت أطوال النباتات مع التلقيح البكتيرى جنبا إلى جنب مع
زيادة للنبات تحت الدراسة .

دلت القياسات الخاصة بمحصول القمح ومكوناته زيادة ملحوظة فى وزن السنبله ، وزن
١٠٠٠ حبة علاوة على محصول الحبوب والقش الناتج .

ولقد تم تحليل النبات الناتج كيمائيا والتي بينت نتائجها زيادة فى النسبة المئوية لكل من
النيتروجين والفسفور والبوتاسيوم لكل من الحبوب والقش وذلك علاوة على زيادة محصول القمح
وصلاحية العناصر الغذائية بالتربة . وقد دلت نتائج هذا البحث على أهمية تعظيم استخدام الأسمدة
الحيوية والتي أدت إلى تقليل نسبة استخدام السماد الكيماوى مما يقلل من تلوث التربة الزراعية
علاوة على حل مشكلة الفوسفات المترسبة فى الأراضى المصرية نتيجة إضافة الأسمدة الفوسفاتية
سنويا نظرا لطبيعة الأراضى المصرية لكونها أراضى تميل للقلوية ومحتوية على كربونات
كالسيوم .