

IMMUNOSUPPRESSIVE ROLE OF FLORFENICOL IN BUFFALO CALVES

H.M. HASAAN and M.H, NASSER*

Animal Reproduction Research Institute, P.o. 12556 El-haram, Giza, Egypt.

* Dept. of Medicine; Fac. of Vet. Med.; Kafr El-Sheikh, Tanta Univ.

Received: 24.10.2002

Accepted: 2 . 12.2002

SUMMARY

A study was designed to investigate the effect of florfenicol therapeutic dose, 20mg/kg body weight intramuscularly ; 2 injections 48 hours apart , on the immune competence of Egyptian female buffalo calves (3- 6 months) vaccinated with brucella abortus strain 19 . Florfenicol administration elucidated a significant decline in lymphocyte blastogenesis , chemotaxis and phagocytic activity.

Throughout the estimation of antibody titre , the results disclosed remarkable decrease of antibody levels by standard tube agglutination (SAT) and rivanol tests . It was concluded that the drug significantly suppress both cellular and humoral immune performance in Egyptian buffalo calves.

INTRODUCTION

Florfenicol roots trace back to chloramphenicol, whose P-nitro group caused human cancers. The florfenicol molecule differs structurally from chloramphenicol in two therapeutically important ways . First , The P- nitro group on chloramphenicol has been replaced by methyl sulphonyl group because p-nitro group is implicated in chloramphenicol induced idiosyncratic aplastic anemia . Second , the hydroxyl group on the carbon 3 of chloramphenicol has been replaced with fluorine atom .The replacement of this group with a non acetylatable fluorine atom protect the molecule from inactivation by acetyl transferase , the common mechanism of bacterial resistance to chloramphenicol (Sams ,1994) .

During studding the immunopharmacological effect of florfenicol in non lactating dairy cows ,

fect of florfenicol in non lactating dairy cows , Bretzalff et al. (1987) reported that florfenicol significantly inhibited phagocytosis at all concentrations.

On the same line , Khodary and Rizk (2000) indicated that florfenicol at the recommended dose reduced the leucocytic count in calves. Recently, Khalil (2002) concluded that florfenicol treatment induced immunosuppressive effects in calves , fish and poultry .

Florfenicol seem to have a promising forthcoming immunotherapeutic application in bovine .The present work was designed to investigate its possible effects on both cellular and humoral immune response of buffalo calves.

MATERIAL AND METHODS

1-Drug: Florfenicol (Nuflor[®] Produced by Schering -Plough Animal Health , USA.

2-Animals : Ten , clinically healthy, Egyptian female buffalo calves (3- 6 months) were used in this work. They were divided into two equal groups. The first received the vaccine (Brucella abortus strain 19, Coopers Animal Health Inc., Kansas , USA) alone and served as a control. The second group was injected with florfenicol therapeutic dose, 20mg/kg body weight intramuscularly ; 2 injections 48 hours apart simultaneous with vaccination process .

3-Blood sampling :From each animal ,two samples (serum and heparinized blood) were collected 2 hours prior to drug injection and weekly during 8 weeks post treatment.

A -Cellular immune response : -

1. Lymphocyte transformation:

Blast transformation of peripheral blood lymphocytes was measured after De Cock et al. (1980) ; Ishikawa and Shirahata (1985) and Shimakaira et al. (1985). Lymphocytes were separated according to Brulles and Wells (1977). The cells were washed three times then suspended in RPMI - 1640 medium with 10% fetal calf serum . The lymphocytes were cultured with mitogen, phytohaemagglutinin (PHA), in flat bottom microtitre plate. The plates were covered and incubated at 37°C in a humidified Co₂ atmosphere (5-10%) for 48 - 72 hours. The concentration of residual glucose in RPMI - 1640 medium used in this technique was estimated in control and stimulated cells, according to by Charles et al.(1978).

2. Phagocytic activity:

Isolation and cultivation of phagocytic cells was done according to Antley and Hazen (1988). The mononuclear cells were prepared for phagocyte activity with freshly prepared culture of Candida albicans (Richardson and Ssmith, 1981).The total number of phagocytic cells (the number of phagocytes with ingested yeast cells) and the total number of ingested yeast cells in individual phagocytes were determined to calculate the percent-

age of phagocytosis.

3. Chemotaxis:

Chemotaxis under agarose was performed according to Nelson et al. (1975).

B-Humoral immune response :-

1- Brucella antibody titer :

The titer of antibodies was measured in serum using serum tube agglutination test according to the method adopted by Alton et al.,1988 . Simultaneously, rivanol test was carried out according to the procedures of the National Veterinary Services Laboratories, Ames Iowa, USA (1984).

Statistical analysis :

The results are represented as the mean \pm S.E. .Statistical significance was determined by student's (t) test for paired observations according to Snedecor (1971).

RESULTS

A- Effect of florfenicol on the cellular immune response:

Florfenicol administration displayed a significant reduction of chemotaxis. This reduction continued through the 4th week post treatment. After the 5th week, it increased to reach the control lev-

Table (1) : Effect of florfenicol on the chemotaxis index in calves.
(Mean \pm S.E.) (n=5)

Week (post vaccination)	Vaccinated Group	Treated vaccinated group
Prevaccination	2.45 \pm 0.31	2.56 \pm 0.20
1	2.51 \pm 0.15	1.72 \pm 0.07*
2	3.26 \pm 0.24	1.90 \pm 0.16*
3	3.67 \pm 0.11	1.57 \pm 0.25*
4	3.80 \pm 0.17	1.83 \pm 0.17*
5	2.05 \pm 0.28	2.11 \pm 0.22
6	2.25 \pm 0.35	2.31 \pm 0.29
7	2.58 \pm 0.36	2.08 \pm 0.40
8	1.72 \pm 0.22	1.93 \pm 0.24

*: Significant at P< 0.05

el (Table, 1).

The group treated with florfenicol depicted a significant reduction in lymphocyte stimulation indices after 1st. week till the end of experiment

(in the presence of PHA). Phagocytic cell percentage showed a significant decrease for 6 weeks (Table, 2).

Table (2) : Effect of florfenicol on lymphocyte transformation (LYT) indices (in the presence of PHA), phagocytic cell % (Phag %) in calves. (Mean \pm S.E.) (n=5).

Week (post vaccination)	Vaccinated group		Treated vaccinated group	
	LYT	Phag%	LYT	Phag%
Prevaccination	1.80 \pm 0.13	75 \pm 4.7	1.83 \pm 0.11	77.8 \pm 4.7
1	2.23 \pm 0.27	77.3 \pm 6.7	0.93 \pm 0.17*	61.7 \pm 3.6*
2	2.97 \pm 0.05	81.5 \pm 5.3	1.07 \pm 0.17*	52.3 \pm 4.1*
3	2.43 \pm 0.21	82.7 \pm 4.5	0.95 \pm 0.11*	50.7 \pm 4.2*
4	2.51 \pm 0.34	77.5 \pm 4.7	1.06 \pm 0.05*	56.2 \pm 5.7*
6	2.80 \pm 0.31	85 \pm 4.7	0.98 \pm 0.06*	55 \pm 3.7*
7	2.71 \pm 0.10	80.3 \pm 6.7	0.85 \pm 0.18*	70.3 \pm 6.7
8	2.60 \pm 0.11	77.6 \pm 4.8	1.36 \pm 0.13*	73.6 \pm 3.7

*: Significant at P < 0.05

B- Effect of florfenicol on humoral immune response.

Table (3) displayed a significant lowering in brucella antibody titres using either rivanol or stan-

durd tube agglutination titre (SAT) due to administration of florfenicol with strain 19 brucella vaccine . This reduction observed 1-4 weeks post vaccination.

Table (3): Effect of florfenicol on the antibody titre using Rivanol and Standard agglutination test (SAT) in vaccinated buffalo calves. (Mean \pm S.E.)

Week (post vaccination)	Vaccinated group		Treated vaccinated group	
	SAT	Rivanol	SAT	Rivanol
Prevaccination	0.0 \pm 0.0	0.0 \pm 0.0	0.0 \pm 0.0	0.0 \pm 0.0
1	66 \pm 10.3	25 \pm 0.0	0.0 \pm 0.0*	0.0 \pm 0.0*
2	299.3 \pm 15.2	90 \pm 10	90.82 \pm 10.5*	25.0 \pm 0.0*
3	897 \pm 91.6	160 \pm 15	305.0 \pm 20.64*	65.81 \pm 5.64*
4	417 \pm 42.2	400 \pm 37	211.6 \pm 3.55*	190.87 \pm 13.28*
5	382.6 \pm 35.1	200 \pm 20	351.48 \pm 55.8	188.57 \pm 21.81
6	245 \pm 25.7	150 \pm 15	219.98 \pm 5.42	159.37 \pm 10.28
7	90.81 \pm 9.52	30.0 \pm 5	82.80 \pm 10.4	55.64 \pm 10.87
	90.0 \pm 5.82	30.0 \pm 5	86.28 \pm 8.40	35.8 \pm 5.5

*: Significant at P < 0.05

DISCUSSION

Florfenicol is a new molecule belonging to the family of agents that includes thiamphenicol and chloramphenicol compounds. Because of their structural similarities, all compounds in this family have the same mechanism of antibacterial activity (Smas, 1994).

The present investigation was planned to gain more information on the effect of florfenicol on the immune competence of Egyptian buffalo calves.

Results of this study indicated that intramuscular administration of florfenicol restrain the immune system of calves. The inhibitory effect was pronounced at the 20mg/kg body weight intramuscularly ; 2 injections ,48 hours apart ,which have been recommended as a therapeutic dosage in many animal species . The immune suppressive effect was associated with decline in lymphocyte transformation , phagocytic ,chemotaxis activity , and antibody titre.

The previous data substantially reinforced the findings reported by Bretzlaff et al. (1987) and Paape et al. (1990) in cows . They documented that florfenicol induced immune suppression via inhibition of phagocytosis and alteration of neutrophil morphology .

Immunosuppressive effect of florfenicol may be attributed to the ability of the drug to penetrate lymphocyte and thereby inhibits its protein and DNA synthesis and eventually suppress the cellular function (Zhakov et al.,1980). Florfenicol was found to be highly effective against number of gram positive and negative bacteria (Graham et al., 1988) .Given this background , it could be predicted that vaccination with Br. abortus strain 19 would be subjected to static effect of florfenicol consequently reducing the reducing the body immune responsiveness to vaccine . Furthermore, the decrease of antibody titre may be referred to the suppressive effect of florfenicol on the haematopoietic activity resulting in acute depletion of circulating B and T lymphocytes , consequently declining the antibody titre (Zhakov et al. ,1980).

Unfortunately, our data do not provide a satisfactory clue for the hitherto documented immunosuppressive activity of florfenicol. Nevertheless, comparable effects to ours have been observed using other antibiotics such as chloramphenicol, that inhibited phytohaemagglutinin-induced lymphocyte transformation (Becker et al. , 1974). The immunosuppressive effect of chloramphenicol was tested in detailed study by Voiculescu et al (1983). They demonstrated that chloramphenicol strongly inhibited the antigen - dependent B cell blastogenesis in vitro . In addition chloramphenicol suppressed phagocytosis and completely blocked neutrophils activity

(Paape et al. , 1990).

Pieced together , our findings did clearly demonstrated a substantial immunosuppressive activity of florfenicol. In this context, the present study bias the possible future contraindicative use of this drug at least as antibacterial therapy in vaccinated animals.

REFERENCES

- Alton,G.G.; Jones,L.M.; Angus,S.D. and Verger , J.M. (1988) : Technique for the brucellosis laboratory " INRA. Paris. ISEN
- Antley, P.P. and Hazen , K.C. (1988) : Role of yeast cell growth temperature on *Candida albicans* virulence in mice .*Infect. Immunity* ,(56) :2884- 3890.
- Becker, D. ; Katz, J. and Metz, J. (1974): Comparative effect of thiamphenicol and chloramphenicol on haemopoiesis and lymphocyte transformation in vitro in man. *Postgard Med. J.*, 50: (supp.1) : 88-94
- Bretzlaff,K.N. ; Neff-Davis,C.A.; Ott.R.S. ; Koritz.G.D. ; Gustafsson. B.K. and Davis.L.E.(1987): Florfenicol in non- lactating dairy cows : pharmacokinetics , binding to plasma proteins ad effects on phagocytosis by blood neutrophils. *J. Vet. Pharmacol. Ther. Sep*; 10(3):233-240.
- Brulles,S. and Wells,P.W. (1977) : In vitro Stimulation of avian lymphocyte by various mitogens . *Res.Vet.Sci.*, (23) : 84-86.
- Charles , R. ; Carpenter , A .B. ; Henry , R . and Bose , J. R. (1978): Suppression of mitogen - stimulated blastogenic response during reticuloendotheliosis virus Induced tumorigenesis.*Immunol.* 120 . (4) : 1313 - 1320.
- De Dock, W.; De Cree,J. ; Van Wauwe,J. and Verhaegen , H. (1980): Measurment of mitogen stimulation of lymphocytes with a glucose consumptin test. *J . of Immun. Methods*,33:127-131.
- Graham, R. ; Palmer,D. ; Paratt,B.C. and Hart,C.. (1988) : invetro activity of florfenicol . *Eur. J. Clin. Microbiol. Dis.* ;7(5):691-694.
- Ishikawa , H. and Shirahata , T.(1985): Application of glucose consumption test for evaluating blastogengensis in bovine lymphocytes. *Jpn. J. Vet. Sci.* 48(1):111-115.
- Khalil ,H. M. (2002) : Immunosuppressive effect of florfenicol ; "A comparative study " PhD Thesis , Pharmacology Dept. Fac. Vet. Med. , Zag. Univ.
- Khodary , R.M. and Rizk, H.I. (2000) : Effect of florfenicol on some blood constituents of buffalo calves . *J.Vet. Med.Res. Vol II No.*, 1: 93-102
- Nelson , R. D . , Quic , P. G . and Simmons, R.L . (1975) : Chemotaxis under agrose:A new and simple method for measuring chemotaxis and spontaneous migration of human polymorph-nuclear leukocytes. *J. of Immunog.* ,(6): 1650-1660.
- Paape ,M.J. ;Miller, R.H. and Ziv.G. (1990) : Effects of florfenicol, chlorfenicol and thiamphenicol on phagocytosis , chemoiluminescence and morphology of bovine polymoirphnuclear neutrophil leukocytes. *J. Dairy Sci.* 73 (7):1734-1744.
- Richardson, M.D. and Smith, H. (1981): Resistance of virulent and attenuated strain of *C. albicans* to intracellular killing by human and mouse phagocytes. *J. Infect.Dis.*, (144): 554-565.

- Shimakuria, Y.; Kudo, T.; Honio ,H. and Khazawa , K. (1985) : Glucose consumption test for peripheral lymphocyte transformation in Shipa goat. Research Bulletin of faculty of agriculture , Gifu Univ. No 50 ,329-334.
- Snedecor, G.W. (1971): Statistical methods. 14 th. Ed. The Iowa State College Press. Ames. Iowa.
- Smas, R.A. (1994) : Florfenicol : chemistry and metabolism of a noval broad spectrum antibiotic. International Symposium on bovine respiratory disease, new therapeutic advances . 30 Aug . 1994, in association with the XVIII World Buiatrics congress.
- Voiculescu. C.; Stanciu, L. ; Voiculescu, M.; Rogos,S. and Dumirtriu,I. (1983): Experimental study of antibiotic induced immunosuppression in mice I. Humoral and cell mediated immune responsiveness related in- vivo antibiotic treatment . Comp. Immun. Microbiol. Infect. Dis., 6:291-299.
- Zhakove,M.S.; Karput,I.M. and Prudnikov, V.S.(1980): Influence of oxytetracycline in young piglet and rabbits in response to in activated salmonellosis vaccine . Moscow , USSR; " Kolos" 132-137