

*Annals Of Agric. Sc., Moshtohor,*  
*Vol. 42(4): 1791-1807, (2004).*

**BEEF SAUSAGE, AND BEEF BURGER PRODUCTION BY ADDING  
TREATED MUNG BEAN  
BY**

**Bahlol, H.El.M.\* and Abd El-Aleem, I.M.\*\***

\* Food Science Dept. and \*\* Agric. Chemistry Dept,  
Fac. of Agric. Moshtohor, Zagazig Univ. (Benha Branch).

**ABSTRACT**

This study was carried out to remove the antinutritional matters from mung bean seeds used to produce beef sausage and beef burger products. Crude protein content in mung bean seeds was 26.8%, while, the antinutritional factors: trypsin inhibitor, tannins,  $\alpha$ -amylase inhibitor, hemagglutinating, phytic acid, total vicine, raffinose, stachyose and verbascose were: 4.6 mg/g, 1.36%, 299 U/g, 1902 U/g, 1.93%, 4.7 mg/g, 0.28%, 0.63% and 1.46%, respectively.

Soaking mung bean seeds in sodium bicarbonate solution (pH 7.85) was effective in removing all antinutritional matters except phytic acid, however, germination was most effective for its reduction. But soaking in citric acid solution was more effective when compared in lowering the content of phytic acid, raffinose, stachyose and verbascose.

In order to reduce the cost of beef products, meat was replaced by rehydrated mung bean flour at the levels of 0, 5, 10, 15, 20, 25, 30 and 35% besides improving their physico-chemical properties.

**INTRODUCTION**

Mung bean is considered an important source of high quality plant protein for human consumption, however, it contains various antinutritional matters. It is of great importance to find a formula for beef sausage and beef burger which lowers the cost of these products, especially under local conditions of meat shortage and high price.

Several conventional processing methods such as soaking and heat treatment were required to remove the undesirable components from dry mung bean seeds for improving their nutritional quality.

Germination has been suggested as an inexpensive and effective method for improving the quality of legumes by enhancing their digestibility (Reddy *et al.*, 1985), increasing the level of amino acids (Chang and Harrold, 1988) and reducing the content of antinutritive matters (Vidal-Valverde *et al.*, 1994; Urbano *et al.*, 1995; Zaki, 1996; Abuel-Fetouh *et al.*, 1998 and El-Bagouty *et al.*, 1999).

The objective of the present work was to decrease antinutritive matters in mung bean seeds by using soaking, germination and autoclaving. After that using treated mung bean seeds in preparing beef sausage and beef burger.

## MATERIALS AND METHODS

### Materials:

Seeds of mung bean (*Vigna radiate* L.) variety 2010 were obtained from Agricultural Research Center, Giza, Egypt.

Raw beef meat and mutton fat purchased from supermarkets of Kalyobia. Spices ingredients (black pepper, cardamom, cloves, cubeb, cumin, garlic, nutmeg, fennel, coriander, laurel and cardamom) were purchased from local market.

### Treatments:

**Soaking:** Dry mature mung bean seeds were soaked in three treatments distilled water, 0.1% citric acid (pH 4.94) and 0.07% sodium bicarbonate (pH 7.85). The seeds were soaked for 3, 6, 9 and 12 h then drained.

**Germination:** Dry mature mung bean seeds were soaked for 2 h in distilled water then germinated in sterile beakers lined with filter paper, and placed in dark incubator at 25°C. Distilled water was sprinkled on seeds twice a day during germination. Seeds germinated for 24, 48 and 72 h.

**Autoclaving treatment:** Soaked, germinated and unsoaked seeds were autoclaved at 121°C for 10, 20, 30, and 40 min and dried in oven at 50°C then ground to flour.

### Preparation of beef sausage and burger samples:

Visible fat tissues were trimmed from lean meat, then minced by electric chopper. Mutton fat tissues also, were minced. Mung bean flour was rehydrated by mixing with distilled water at ratio 1 : 2 (w:w) and added by levels, 0 (control), 5, 10, 15, 20, 25, 30 and 35% replace with beef meat to prepare sausage and burger. The formula of beef sausage and burger is shown in Table (1) as mentioned by Moghazy and El-Shaarawy (2001) and Moghazy *et al.* (2004):

### Methods:

**Assay of trypsin inhibitors (TI):** The trypsin inhibitors activity (TIA) was measured as described by Stauffer (1993).

**Determination of Hemagglutinating (HA):** Lectin activity was determined by measuring its hemagglutinating action according to the method described by Lis and Sharon (1972).

**Determination of phytic acid:** Phytic acid content was estimated colorimetrically using Wade reagent (Latta and Eskin, 1980).

**Determination of total vicine:** Total vicine was extracted and determined according to the method of Collier (1976).

**Table (1): The formula of beef sausage and burger prepared in laboratory**

Component	Sausage	Burger	Spices mixture
Beef meat	70.60%	62.00%	For sausage: Fennel 59.76%, coriander 27.09%, cubeb 3.19%, black pepper 3.19%, clove 3.19%, laurel 1.99% and cardamom 1.59%.  For burger: Black pepper 5.61%, cardamom 2.24%, cloves 2.24%, cubeb 22.42%, cumin 11.21%, garlic 56.05%, and nutmeg 0.22%.
Mutton fat	14.00%	-	
Water (as ice flakes)	7.00%	10.00%	
Starch	4.65%	-	
Sodium pyrophosphate	0.30%	-	
Salt (NaCl)	2.00%	1.50%	
Garlic	0.24%	-	
Skim milk powder	0.40%	-	
Glucose	0.1%	-	
Ascorbic acid	0.04%	-	
Sodium nitrite	0.01%	-	
Ground onion	-	7.00%	
Egg	-	7.00%	
Semolina	-	12.00%	
Spices mixture	0.66%	0.50%	

Beef sausage and beef burger products were fried according to Modi *et al.* (2003).

**Determination of raffinose oligosaccharides:** Oligosaccharides were extracted from powdered samples with 80% (v/v) ethanol (Akpapunam and Markakis, 1979). Unidirectional descending paper (Whatman No. 1, 20x45 cm) chromatography using a solvent of n-butanol, ethanol and water (5:3:2 by volume) was conducted for 48 h to separate oligosaccharides. The sugars were identified on the basis of their  $R_f$  and  $R_G$  values according to Akpapunam and Markakis (1979). The concentration of the identified sugars was determined using the phenol sulphuric acid method of Dubois *et al.* (1956).

**Alpha-Amylase inhibitor assay:** The extraction of  $\alpha$ -amylase inhibitor was performed as described by Bernfeld (1955).

**In vitro protein digestibility:** The digestibility of protein *in vitro* was carried out as described by Santosh and Chauhan (1986).

**Chemical analysis:** Moisture, crude protein, ether extract, ash and crude fiber contents were determined according to A.O.A.C. (1995). Carbohydrates content was calculated by difference.

**Freshness tests:** Total volatile nitrogen (TVN) was determined according to the methods mentioned by Winton and Winton (1958). Thiobarbituric acid (TBA) value was determined according to Harold *et al.* (1981). The pH value of meat product was measured using digital pH-meter model SA 210 according to the method of Woye Woda *et al.* (1986).

**Water holding capacity (WHC) and plasticity** were measured according to the method described by Soloviev (1966).

**Cooking loss and shrinkage** were determined according to Darweash and Moghazy (1998).

**Sensory evaluation** was evaluated according to Watts *et al.* (1989).

Statistical analysis was applied on the results of organoleptic evaluation of different samples of beef sausage and beef burger which were treated as data for complete randomization design. Least significant difference (L.S.D.) was calculated at 0.05 level of significance according to Snedecor and Cochran (1980).

## RESULTS AND DISCUSSION

### Chemical composition of raw mung bean seeds:

Data in Table (2) show that crude protein content was 26.8% which was higher than 23.4% in ten mung bean genotypes studied by Ismail (1995), but, lower than 28.2% found by Abu Arab and Helmy (2001) for unspecified cultivar. This difference in protein content could be attributed to genetical and environmental factors. The antinutritive matters, trypsin inhibitor (TI), tannins,  $\alpha$ -amylase inhibitor ( $\alpha$ -AI), hemagglutinating (HA), phytic acid, total vicine, raffinose, stachyose and verbascose in the dry mature mung bean seeds were determined to be 4.60 mg/g, 1.36 g/100g, 299 U/g, 1902 U/g, 1.93%, 4.70 mg/g, 0.28%, 0.63% and 1.46%, respectively. Our data were different from those reported by Zaki (1996), but, agreed with the results of Abuel-Fetouh *et al.* (1998).

Table (2): Chemical composition of mung bean seeds.

Components		
Moisture	(%)	10.53
Crude protein*	(%)	26.80
Ether extract*	(%)	1.24
Ash*	(%)	4.35
Crude fiber*	(%)	3.10
Available carbohydrate**	(%)	64.51
Trypsin inhibitor *	(mg/g)	4.60
Tannins*	(g/100 g)	1.36
$\alpha$ -amylase inhibitor*	(U/g)	299.00
Hemagglutinating*	(U/g)	1902.00
Phytic acid*	(g/100 g)	1.93
Total vicine*	(mg/g)	4.70
Raffinose*	(%)	0.28
Stachyose*	(%)	0.63
Verbascose*	(%)	1.46


\* Calculated on dry weight basis

### Effect of soaking on antinutritional matters:

Soaking mung bean seeds in the three treatments resulted in gradual decline in all antinutritional matters as shown in Table (3) and illustrated in Fig. (1). These treatments were effective in removing about 24 to 29% of TI, 31 to 33% of  $\alpha$ -AI, 25 to 28% of HA, 11 to 13 of tannins, 13 to 36% phytic acid, 29 to 38% of vicine, 13 to 28% of raffinose, 25 to 36% of stachyose and 31 to 38% of verbascose contents. Fig. (1) included two lines only in the case of verbascose

**Table (3): Effect of soaking in distilled water, sodium bicarbonate and citric acid solutions on antinutritional Matters (on dry weight basis).**

Time (h)	Distilled water		NaHCO <sub>3</sub>		Citric acid		Distilled water		NaHCO <sub>3</sub>		Citric acid	
	mg/g	Residue (%)	mg/g	Residue (%)	mg/g	Residue (%)	U/g	Residue (%)	U/g	Residue (%)	U/g	Residue (%)
	<b>Trypsin inhibitor (TI)</b>						<b>Alpha-amylase inhibitor (A)</b>					
Raw	4.6	100	4.6	100	4.6	100	299	100	299	100	299	100
3	3.864	84	3.818	83	4.416	96	242.19	81	245.18	82	245.18	82
6	3.726	81	3.588	78	4.321	94	239.2	80	236.21	79	230.23	77
9	3.588	78	3.45	75	4.186	91	233.22	78	224.25	75	215.28	72
12	3.496	76	3.266	71	4.048	88	218.27	73	206.31	69	200.33	67
	<b>Hamagglutinating (HA)</b>						<b>Phytic acid</b>					
	U/g	Residue (%)	U/g	Residue (%)	U/g	Residue (%)	%	Residue (%)	%	Residue (%)	%	Residue (%)
Raw	1902	100	1902	100	1902	100	1.93	100	1.93	100	1.93	100
3	1616.7	85	1559.64	82	1997.68	84	1.7949	93	1.8142	94	1.7177	89
6	1578.66	83	1502.58	79	1599.64	82	1.737	90	1.7949	93	1.6019	83
9	1540.62	81	1445.52	76	1464.54	77	1.6598	86	1.7177	89	1.5247	79
12	1521.6	80	1369.44	72	1423.5	75	1.5826	82	1.6791	87	1.351	70
	<b>Total vicine</b>						<b>Tannins</b>					
	mg/g	Residue (%)	mg/g	Residue (%)	mg/g	Residue (%)	%	Residue (%)	%	Residue (%)	%	Residue (%)
Raw	4.7	100	4.7	100	4.7	100	1.36	100	1.36	100	1.36	100
3	3.948	84	3.819	77	3.854	82	1.292	95	1.3192	97	1.36	100
6	3.76	80	3.525	75	3.65	78	1.2648	93	1.3056	96	1.328	98
9	3.619	77	3.243	69	3.525	75	1.224	90	1.2512	92	1.3056	96
12	3.384	72	2.914	62	3.337	71	1.1832	87	1.2104	89	1.2648	93
	<b>Raffinose</b>						<b>Stachyose</b>					
	%	Residue (%)	%	Residue (%)	%	Residue (%)	%	Residue (%)	%	Residue (%)	%	Residue (%)
Raw	0.28	100	0.28	100	0.28	100	0.63	100	0.63	100	0.63	100
3	0.28	100	0.2716	97	0.2436	87	0.5607	89	0.5166	82	0.504	80
6	0.28	100	0.2548	91	0.2352	84	0.5292	84	0.4977	79	0.4599	73
9	0.2772	99	0.2492	89	0.2156	77	0.5103	81	0.4851	77	0.4347	69
12	0.2744	98	0.2436	87	0.2016	72	0.4977	79	0.4725	75	0.4032	64
	<b>Vaerbasose</b>											
	%	Residue (%)	%	Residue (%)	%	Residue (%)						
0	1.46	100	1.46	100	1.46	100						
3	1.2118	83	1.2118	83	1.118	80						
6	1.1524	79	1.1388	78	1.058	73						
9	1.095	75	1.095	75	0.9344	64						
12	1.022	70	1.0074	69	0.9052	62						



**Fig. (1): Effect of soaking on antinutritional matters.**

that distilled water and NaHCO<sub>3</sub> have the same line. The results clearly show that soaking in 0.07% sodium bicarbonate solution was more effective in removing all antinutritional matters except phytic acid. But soaking in citric acid solution was more effective when compared in lowering the content of phytic acid, raffinose, stachyose and verbascose.

These results are in agreement with those of Fernandez *et al.* (1993) who observed that after soaking faba beans in H<sub>2</sub>O, citric acid and sodium bicarbonate solutions, a decrease in TI took place except in citric acid soaking, due probably to the stability of the inhibitor in acidic pH.

**Effect of germination on antinutritional matters:**

Germination increases the activity of anabolic and catabolic reactions which has been applied to mung bean seeds to study its effect on the levels of various antinutritional matters. As shown in Table (4) and Fig (2) TIA gradually decreased to 27% of its original level in ungerminated seeds. These results are in agreement with those of Bau *et al.* (1997), Abdel-Galil (1998) and Zaki *et al.* (1999).

Germination for 72 h removed 81% of HA in mung bean seeds. Bau *et al.* (1997) observed disappearance of most HA activity in soybean after 4 days of germination.

The germination process led to sharp decrease in  $\alpha$ -AI levels to reach 22%. Decrease in  $\alpha$ -AI in germinated seeds could be attributed to proteolytic degradation of inhibitor during germination (Gupla and Wagle, 1980).

Germination is the most effective process for reduction of phytic acid content in mung bean seeds. These losses may be attributed to the activity of the phytase enzyme. Vidal-Valverde *et al.* (1994) noted that phytic acid was hydrolyzed during germination.

A sharp reduction was observed in verbascose sugars during the first 24 h of germination to 53% but, slight decrease was for stachyose (24%) and verbascose (8%) during the same period. At the end of germination (72 h) the reduction of sugars was 76%, 71% and 56% for verbascose, stachyose and raffinose, respectively.

**Table (4): Effect of germination time on antinutritional matters**

Germination time (h)	Trypsin inhibitor (%)	Alpha-amylase inhibitor (%)	Hemagglutinating (%)	Phytic acid (%)	Total vicin (%)	Tannins (%)	Raffinose (%)	Stachyose (%)	Verbasose (%)
Raw	100	100	100	100	100	100	100	100	100
24	76	56	48	72	89	100	92	76	47
48	42	34	26	44	75	70	70	73	38
72	27	22	19	15	45	40	44	29	24




Fig. (2): Effect of germination time on antinutritional matters.


The effect of autoclaving on the antinutritional matters in mung bean seeds:

Data in Table (5) and Fig. (3) indicated that autoclaving was more effective than soaking for inactivating antinutritional matters in mung bean seeds. However high reduction was observed in soaked-autoclaved seeds. Moreover, autoclaved germinated seeds revealed enormous reduction. The results of the present work suggest that a combination of two or more simple processing methods may be used to improve the nutritional value of mung bean seeds. Our data agreed with those of Khalil and Mansour (1995), on faba beans.

Table (5): Residual percentage from antinutritional matters after autoclaving

Treatments	Trypsin inhibitor (%)	Tannins (%)	Alpha-amylase inhibitor (%)	hemagglutinin (%)	Phytic acid (%)	Total vicine (%)	Raffinose (%)	Stachyose (%)	Verbascose (%)
Control without autoclaving	100	100	100	100	100	100	100	100	100
Dry seed	44	46	45	35	65	51	42	39	40
Soaking in water	18	17	12	0	29	38	13	11	9
Soaking in sodium bicarbonate	16	21	14	0	25	35	11	9	8
Soaking in citric acid	34	23	17	0	19	37	8	6	5
Germinated for 72 h	4	10	10	0	13	33	13	11	9





**Fig. (3): Effect of autoclaving on antinutritional matters.**

**Effect of processing on in-vitro protein digestibility:**

In-vitro protein digestibility of raw and treated mung bean seeds was performed and results are shown in Table (6). Soaking resulted a slight improvement in protein digestibility which was calculated to be 74.6, 73.9 and 75.2% after soaking for 12 h in water, citric acid and sodium bicarbonate solution, respectively compared to 72.4% of raw seeds.

**Table (6): Effect of treatments on protein digestibility index**

Treatments	Digestibility (%)
Raw seeds	72.4
Soaking in water (12 h)	74.6
Soaking in bicarbonate (12 h)	75.2
Soaking in citric acid (12 h)	73.9
Germination seeds (12 h)	77.7
Raw seeds + Autocl.	76.3
Soaking in water + Autocl.	79.1
Soaking in bicarbonate + Atocl.	79.8
Soaking in citric + Autocl.	76.8
Germination seeds + Autocl.	83.4

Germination improved the protein digestibility of mung bean seeds than autoclaved raw seeds however, germinated-autoclaved seeds showed the highest in-vitro protein digestibility of 83.4%. The improvement in digestibility could be induced by the combined effect of decrease in TI content and/or a greater susceptibility to enzyme attack of the degraded proteins formed during germination as described by Liener (1994).

### Chemical and sensory evaluation of prepared beef sausage and beef burger by adding rehydrated mung bean flour:

#### a) Beef sausage:

Data in Table (7) show the moisture, crude protein, ether extract, ash and total carbohydrates contents in beef sausage prepared in laboratory. Moisture content of fresh sausage ranged from 58.92 to 62.61%, while, it was from 49.27 to 52.57 % after frying. Moisture, crude protein and ether extract contents decreased by increasing the level of replacement with mung bean while, ash and total carbohydrates took the opposite direction. This is mainly due to the lower content of protein and fat content in the replace ingredients. On the contrary, ash and total carbohydrates increased by increasing the supplementation levels. These results are in agreement with those reported by Faheid *et al.* (1998).

Also, results in the same table indicated that moisture and ether extract decreased after frying, while crude protein, ash and total carbohydrates increased. Crude protein decreased in all treatments by increasing the levels of replacement for meat by prepared mung bean seeds. The percentage of decrease reached 10.22% for crude protein at level 35% replacement. However, increase of crude protein after frying may be due to decrease in ether extract content due to escape of some fats in cooking process and/or lowering of meat with increasing the level of replacement as mentioned by Nuzhat *et al.* (2002).

Table (7): Chemical composition of beef sausage with rehydrated mung bean.

Replacement with mung bean (%)	Moisture (%)		Crude protein* (%)		Ether extract* (%)		Ash* (%)		Carbohydrate* (%)	
	Fresh	After frying	Fresh	After frying	Fresh	After frying	Fresh	After frying	Fresh	After frying
0	62.61	52.57	38.94	42.16	44.16	38.99	7.59	7.98	9.31	10.87
5	62.08	51.64	38.65	41.32	43.88	38.43	7.61	7.63	9.86	12.62
10	61.55	51.70	37.77	39.27	43.53	38.67	7.65	7.71	11.05	14.35
15	60.89	51.98	35.79	38.88	43.20	37.48	7.70	7.78	13.31	15.86
20	60.12	51.67	33.94	37.90	42.86	37.51	7.74	7.82	15.46	16.77
25	59.93	51.50	32.20	36.78	42.60	37.27	7.77	7.83	17.43	18.12
30	59.39	49.99	30.48	35.27	42.35	37.00	7.82	7.90	19.35	19.83
35	58.92	49.27	28.72	34.77	42.08	36.44	7.88	7.95	21.32	20.84

\* On dry weight basis.

Data in Table (8) indicate the physico-chemical properties of beef sausage with series levels of mung bean replacement besides changes in TVN, TBA, pH value, WHC, plasticity and cooking loss in prepared beef sausage. TVN amounted to 9.60 mg/100 g in fresh beef sausage and increased after frying to 9.80 mg/100 g. Adding treated mung bean seeds at levels from 5 to 35% in fresh or fried beef sausage decreased TVN to 5.95 and 6.60 mg/100 g respectively. TBA took the same trend and revealed 0.64 and 0.65 mg/kg in fresh and fried beef sausage and decreased to 0.33 and 0.36 mg/kg respectively. On the contrary pH value was 5.87 and 5.95 in fresh and fried beef sausage and increased gradually to 6.06 and 6.08, respectively. Concerning W.H.C. it was 1.25 cm<sup>2</sup>/0.3

g in fresh beef sausage and decreased gradually to 0.50 cm<sup>2</sup>/0.3 g, while plasticity decreased from 3.20 to 1.70 cm<sup>2</sup>/0.3 g. Cooking loss % was 8.51 in fresh beef sausage and decreased gradually to 2.06. These results are in agreement with Faheid *et al.* (1998) and Modi *et al.* (2003).

**Table (8): Physico-chemical properties of beef sausage with rehydrated mung bean.**

Replacement with mung bean (%)	TVN (mg/100 g)		TBA (mg/kg)		pH value		WHC (cm <sup>2</sup> /0.3 g)		Plasticity (cm <sup>2</sup> /0.3 g)		Cooking loss (%)	
	Fresh	After frying	Fresh	After frying	Fresh	After frying	Fresh	After frying	Fresh	After frying	Fresh	After frying
	0	9.60	9.80	0.64	0.65	5.87	5.95	1.25	-	3.20	-	8.51
5	9.45	9.70	0.59	0.62	5.89	5.99	1.15	-	3.05	-	7.72	-
10	8.40	9.10	0.53	0.59	5.93	6.01	0.95	-	2.80	-	7.54	-
15	7.80	8.90	0.50	0.52	5.95	6.03	0.95	-	2.65	-	6.94	-
20	7.50	8.10	0.45	0.46	6.02	6.05	0.80	-	2.45	-	5.89	-
25	6.80	7.50	0.38	0.42	6.03	6.06	0.75	-	1.95	-	4.68	-
30	6.30	7.00	0.35	0.38	6.04	6.07	0.60	-	1.75	-	2.66	-
35	5.95	6.60	0.33	0.36	6.06	6.08	0.50	-	1.70	-	2.06	-

TVN: Total volatile nitrogen  
 WHC: Water holding capacity

TBA: Thiobarbituric acid

Data in Table (9) indicate the sensory evaluation (color, aroma, taste, texture, palatability and total scores) in beef sausage prepared in laboratory with replacement by mung bean (0 to 35%). Results show that there are significant differences (P<0.05) for color, taste, texture and palatability between control and all treatments, except there was no significant differences (P>0.05) in aroma between control and treatments with replacement level of 5, 10 and 15% rehydrated mung bean.

**Table (9): Sensory evaluation of beef sausage with rehydrated mung bean.**

Replacement with mung bean (%)	Color	Aroma	Taste	Texture	Palatability	Total score
Control	9.8±0.13 <sup>a</sup>	9.6±0.16 <sup>b</sup>	9.8±0.13 <sup>a</sup>	9.9±0.10 <sup>a</sup>	9.8±0.13 <sup>a</sup>	48.9±0.38 <sup>a</sup>
5	9.2±0.30 <sup>b</sup>	9.3±0.16 <sup>ab</sup>	9.2±0.13 <sup>b</sup>	9.3±0.17 <sup>b</sup>	9.2±0.15 <sup>b</sup>	46.2±0.59 <sup>ab</sup>
10	9.2±0.39 <sup>b</sup>	9.3±0.21 <sup>ab</sup>	9.1±0.10 <sup>b</sup>	9.2±0.16 <sup>bc</sup>	9.1±0.00 <sup>b</sup>	45.9±0.53 <sup>ab</sup>
15	9.0±0.46 <sup>bc</sup>	9.1±0.13 <sup>abc</sup>	9.0±0.16 <sup>bc</sup>	9.2±0.18 <sup>bc</sup>	9.0±0.15 <sup>b</sup>	45.3±0.59 <sup>b</sup>
20	8.8±0.30 <sup>bc</sup>	8.7±0.26 <sup>bc</sup>	8.8±0.10 <sup>bc</sup>	9.0±0.16 <sup>bc</sup>	8.9±0.21 <sup>bc</sup>	44.2±0.63 <sup>b</sup>
25	8.6±0.38 <sup>c</sup>	8.4±0.28 <sup>c</sup>	8.5±0.22 <sup>c</sup>	8.7±0.15 <sup>c</sup>	8.4±0.16 <sup>c</sup>	42.6±0.52 <sup>b</sup>
30	7.4±0.40 <sup>d</sup>	7.6±0.31 <sup>d</sup>	6.6±0.22 <sup>d</sup>	6.9±0.18 <sup>d</sup>	6.7±0.21 <sup>d</sup>	35.2±0.68 <sup>c</sup>
35	6.4±0.40 <sup>e</sup>	6.9±0.41 <sup>d</sup>	5.7±0.26 <sup>e</sup>	5.9±0.28 <sup>e</sup>	5.5±0.27 <sup>e</sup>	30.4±0.92 <sup>d</sup>
LSD	0.57	0.72	0.50	0.51	0.50	3.76

a, b, c, d & e: There is no significant difference between any two means, with the same attribute, have the same letter (P > 0.05).

Anyhow, the mung bean added to sausage samples could be separated into two groups, hence there is no significant differences ( $P>0.05$ ) between any two samples with the same group. The first group includes sausage treatments replacement levels 5, 10, 15 and 20% of mung bean.

The second group includes sausage treatments replacement with 25 to 35% mung bean. In the same time there is significant difference ( $P<0.05$ ) between the two groups.

#### b) Beef burger:

Data in Table (10) show the moisture, crude protein, ether extract, ash and total carbohydrate contents in beef burger prepared in laboratory with replacement levels of mung bean. Moisture content of fresh beef burger was 67.12%, while it ranged from 64.34 to 66.01% in all treatments. Moisture content decreased after frying in all treatments, where it ranged from 53.57 to 56.44%.

Crude protein and ether extract contents decreased with increasing the level of replacement by mung bean seeds, while, ash and total carbohydrates content increased with increasing replacement levels (0 to 35%).

Also, results in the same table indicate that moisture, crude protein, ash and total carbohydrate contents decreased after frying, wherever ether extract content increased.

The percentage of decrease reached to 9.25 and 7.05% for crude protein at level 35% compared to fresh and fried one, respectively. Decrease of crude protein after frying may be due to increase in ether extract. These results are in agreement with those reported by Abd El-Salam and Hassanin (1987), Mansour and Khalil (1999) and El-Mansy *et al.* (2002).

Table (10): Chemical composition of beef burger with rehydrated mung bean.

Replacement with mung bean (%)	Moisture (%)		Crude protein* (%)		Ether extract* (%)		Ash* (%)		Carbohydrate* (%)	
	Fresh	After frying	Fresh	After frying	Fresh	After frying	Fresh	After frying	Fresh	After frying
0	67.12	54.81	51.81	44.52	14.97	22.59	6.52	6.31	26.70	26.58
5	64.34	55.28	49.06	43.06	13.84	22.52	6.59	6.48	30.51	27.94
10	65.89	56.44	48.53	42.71	13.61	22.44	6.70	6.53	31.15	28.32
15	65.19	54.57	47.29	41.46	13.20	21.31	6.77	6.60	32.75	30.64
20	66.01	54.74	46.11	40.72	12.71	20.74	6.96	6.66	34.23	31.89
25	65.60	56.40	44.72	39.46	11.91	20.83	6.96	6.70	36.41	33.01
30	65.33	56.16	43.54	38.42	11.77	21.09	6.98	6.73	37.71	33.77
35	65.65	53.57	42.56	37.47	11.24	20.51	6.96	6.84	39.24	35.09

\* On dry weight basis.

Data in Table (11) indicate the physico-chemical properties of beef burger with series levels of mung bean replacement besides changes in TVN, TBA, pH value, WHC, plasticity, cooking loss and shrinkage in prepared beef burger. TVN amounted to 10.40 mg/100 g in fresh beef burger and increased after frying to 11.70 mg/100 g. Adding treated mung bean seeds at levels from 5 to 35% in fresh or fried beef burger decreased TVN to 5.80 and 7.40 mg/100 g respectively. TBA took the same trend and revealed 0.56 and 0.59 mg/kg in fresh and fried beef burger and decreased to 0.33 and 0.35 mg/kg respectively. On the contrary pH value was 5.40 and 5.64 in fresh and fried beef burger and increased gradually to 5.78 and 5.95 respectively. Concerning W.H.C. it was 4.90 cm<sup>2</sup>/0.3 g in fresh beef burger and decreased gradually to 3.30 cm<sup>2</sup>/0.3 g, while plasticity increased from 1.70 to 2.00 cm<sup>2</sup>/0.3 g. Cooking loss % and shrinkage after frying reached 13.26 and 13.42 fresh beef burger and decreased gradually to 3.54 and 6.75, respectively. Similar findings were reported by Lecomte *et al.* (1993) and Modi *et al.* (2003).

**Table (11): Physico-chemical properties of beef burger with rehydrated mung bean.**

Replacement with mung bean (%)	TVN (mg/100 g)		TBA (mg/kg)		pH value		WHC (cm <sup>2</sup> /0.3 g)		Plasticity (cm <sup>2</sup> /0.3 g)		Cooking loss (%)		Shrinkage after frying (%)
	Fresh	After frying	Fresh	After frying	Fresh	After frying	Fresh	After frying	Fresh	After frying	Fresh	After frying	
0	10.40	11.70	0.56	0.59	5.40	5.64	4.90	-	1.70	-	13.26	-	13.42
5	9.90	11.00	0.53	0.56	5.42	5.54	4.50	-	1.75	-	12.49	-	12.95
10	9.30	10.60	0.50	0.53	5.48	5.62	4.45	-	1.75	-	9.49	-	12.18
15	8.50	9.90	0.48	0.51	5.55	5.66	4.25	-	1.95	-	7.36	-	10.86
20	7.80	9.20	0.45	0.49	5.59	5.75	4.08	-	1.85	-	6.59	-	9.24
25	6.90	8.50	0.42	0.44	5.65	5.82	3.75	-	1.90	-	4.84	-	8.87
30	6.20	8.00	0.38	0.40	5.71	5.90	3.50	-	1.95	-	3.72	-	7.38
35	5.80	7.40	0.33	0.35	5.78	5.95	3.30	-	2.00	-	3.54	-	6.75

TVN: Total volatile nitrogen  
Water holding capacity

TBA: Thiobarbituric acid WHC:

Data in Table (12) indicated the sensory evaluation (color, aroma, taste, texture, palatability and total scores) in prepared beef burger with replacement by mung bean (0 to 35%). Results indicated that there were no significant differences ( $P>0.05$ ) for all properties tested except total score between control sample and all treatments for 10% replacement levels.

Anyhow, the treatment of beef burger samples could be separated into two groups, hence there is no significant differences ( $P>0.05$ ) between any two samples with the same group. The first group includes beef burger treatments replacement levels 5, 10, 15 and 20% of mung bean.

The second group includes beef burger treatments replacement with 25 to 35% mung bean. In the same time there is significant difference ( $P < 0.05$ ) between the two groups.

So, it could be recommended to apply replacement level with 20% mung bean from meat used in prepared sausage and beef burger products.

Table (12): Sensory evaluation of beefburger with rehydrated mung bean.

Replacement with mung bean (%)	Color	Aroma	Taste	Texture	Palatability	Total Score
Control	9.5±0.21 <sup>a</sup>	7.2±0.19 <sup>a</sup>	9.6±0.24 <sup>a</sup>	9.7±0.25 <sup>a</sup>	7.3±0.17 <sup>a</sup>	43.3±1.06 <sup>a</sup>
5	9.2±0.19 <sup>ab</sup>	6.8±0.13 <sup>ab</sup>	9.2±0.24 <sup>abc</sup>	9.3±0.24 <sup>ab</sup>	6.9±0.21 <sup>abc</sup>	41.4±0.93 <sup>ab</sup>
10	9.1±0.25 <sup>ab</sup>	6.8±0.14 <sup>ab</sup>	9.2±0.19 <sup>abc</sup>	9.1±0.33 <sup>ab</sup>	6.8±0.23 <sup>abc</sup>	41.0±1.02 <sup>ab</sup>
15	8.9±0.21 <sup>bc</sup>	6.6±0.08 <sup>bc</sup>	9.0±0.13 <sup>bc</sup>	8.8±0.30 <sup>bc</sup>	6.8±0.21 <sup>abc</sup>	40.1±0.72 <sup>b</sup>
20	8.8±0.13 <sup>bcd</sup>	6.4±0.07 <sup>bc</sup>	9.0±0.13 <sup>bc</sup>	8.7±0.23 <sup>bc</sup>	6.7±0.14 <sup>abc</sup>	39.6±0.40 <sup>bc</sup>
25	8.7±0.13 <sup>bcd</sup>	6.3±0.14 <sup>c</sup>	8.8±0.15 <sup>c</sup>	8.6±0.25 <sup>bc</sup>	6.6±0.18 <sup>bc</sup>	39.0±0.72 <sup>bc</sup>
30	8.5±0.17 <sup>cd</sup>	5.8±0.23 <sup>d</sup>	8.1±0.20 <sup>d</sup>	8.3±0.30 <sup>c</sup>	6.4±0.20 <sup>bc</sup>	37.1±0.90 <sup>cd</sup>
35	8.3±0.23 <sup>d</sup>	5.6±0.24 <sup>d</sup>	7.9±0.25 <sup>d</sup>	8.2±0.29 <sup>c</sup>	6.3±0.35 <sup>c</sup>	36.3±1.14 <sup>d</sup>
LSD	0.55	0.46	0.56	0.78	0.60	2.51

a, b, c, & d: There is no significant difference between any two means, with the same attribute, have the same letter ( $P > 0.05$ ).

## REFERENCES

- Abdel-Galil, M.L. (1998): Effect of maturity and processing on antinutritional factors and raffinose family sugars in seeds of two pea (*Pisum sativum*) varieties. *J. Agric. Sci. Mansoura Univ.*, 23: 1347-1369.
- Abd El-Salam, M.A. and Hassanin, T.M. (1987): Utilization of defatted soy flour and chickpea for preparing minced fish patties. *Agric. Res. Review*, 65(3): 531-540.
- Abou Arab, A.A. and Helmy, M.F. (2001): Removal of Antinutritional factors and improvement of protein digestibility of Mung bean seed by different soaking treatments. *J. Agric. Sci. Mansoura Univ.*, 26(7): 4343-4360.
- Abuel-Fetouh, S.A.; El-Morsi, E.A.; Ghazy, M.A.; Abdel-Salam, H.S. and Fawzy, A.M. (1998): Changes in the levels of lectin, trypsin inhibitor and  $\alpha$ -amylase inhibitor in some legume seeds during germination and processing. *J. Agric. Sci. Mansoura Univ.*, 23: 2919-2939.
- Alpapunam, M.A. and Markakis, P. (1979): Oligosaccharides of 13 American cultivars of cowpea (*Vigna sinensis*). *J. Food Sci.*, 44: 1317-1318.
- A.O.A.C. (1995): Official Methods of Analysis, Association of Official Analytical Chemists, 16<sup>th</sup> Ed., Virginia, U.S.A.
- Bau, H.M.; Villaume, C.; Nicolas, J.P. and Mejean, L. (1997): Effect of germination on chemical composition, biochemical constituents and antinutritional factors of soybean (*Glycine max*) seeds. *J. Sci. Food Agric.*, 73: 1-9.
- Bernfeld, P. (1955): Amylases  $\alpha$  and  $\beta$  in *Methods Enzymology*, 1 149-158.

- Chang, K.C. and Harrold, R.L. (1988): Changes in selected biochemical components *in vitro* protein digestibility and amino acids in two bean cultivars during germination. *J. Food Sci.*, 53: 782-787
- Collier, H.P. (1976): The estimation of vicine in faba beans by an ultraviolet spectrophotometric method. *Can. Inst. Fd. Sci. Technol.*, J. 9: 155.
- Darweash, B.M. and Moghazy, E.A. (1998). Evaluation of chemical and nutritional properties of beef burgers extended with beef ribs bone protein isolate. *Egypt. J. Appl. Sci.*, 13(2): 109-125.
- Dubois, M.; Gilles, K.A.; Hamilton, J.K.; Rebers, P.A. and Smith, F. (1956): Colorimetric method for determination of sugars and related substances. *Annal. Chem.*, 28: 350-356.
- El-Bagoury, A.A., Salem, M.A. and Hanan E.M. (1999): Effect of soaking and germination of legumes on chemical and nutritional values. *J. Agric. Sci. Mansoura Univ.*, 24(10): 5731-5745.
- El Mansy, H.A.; Mahmoud, M.H.M. and Abe El-Azm, H.A. (2002): Utilization of tempeh in some meat products. *J. Annals of Agric. Sc. Moshtohor*, 40(1): 971-985
- Faheid, S.M.M.; El-Akel, A.T. and Mohamed, S.E. (1998): Quality attributes of beef burger supplemented with different sources of plant proteins and the relationship between different cooking methods. *Egypt. J. Appl. Sci.*, 13(7): 217-237
- Fernandez, N.M.; Avanda, P.; Lopez-Jurado, M. and Urbano, G. (1993): Effect of processing on some antinutritive factors of faba beans. *Food Chem.*, 45: 161-166.
- Gupla, K. and Wagle, D.S. (1980): Changes in antinutritional factors during germination in (*Phaseolus mungoreous*) a cross between phaseolus mungo and phaseolus aureus. *Food Sci.*, 45: 394-397
- Harold, E., Ronald, S.K. and Ronald, S. (1981): Person's chemical analysis of foods. 8<sup>th</sup> Ed. Phototypeset by input Typesetting Ltd., London Sw198 Dr printed in 8 Great Britain by Butler and Tammer Ltd.
- Ismail, H.A. (1995): Biochemical studies on the albumins and globulins of promising ten genotypes of mung bean (*Vigna radiate*) seed. *Minia J. Agric. Res. & Dev.*, 17: 139-158.
- Khalil, A.H. and Mansour, E.H. (1995): Effect of cooking, autoclaving and germination on the nutritional quality of faba beans. *Food Chemistry* 54: 177-182.
- Latta, M. and Eskin, N.A.M. (1980): A simple and rapid colorimetric method for phytate determination. *J. Agric. Food Chem.*, 28: 1313-1315
- Lecomte, N.B.; Zayas, J.E. and Kastner, C.L. (1993): Functional and Sensory Characteristics improved in coumited meats. *J. Food Sci.*, 58: 464.
- Liener, I.E. (1994): Implications of antinutritional components in soybean foods. *Crit. Rev. Food Sci Nutr* 34: 31-67
- Liu, H. and Sharon, N. (1972). Soybean (*Glycyne max*) agglutinin. *Methods Enzymol.* 28: 360-366
- Mansour, E.H. and Khalil, A.H. (1999): Characteristics of low fat beef burgers or influenced by various types of wheat fibers. *J. Sci. Food and Agric.* 79(4): 493-498.

- Modi, V.K.; Mahendrakar, N.S.; Rao, D.N. and Sachindra, N.M. (2003): Quality of buffalo meat burger containing legume flours as binders. *Meat Sci.*, 66(1): 143-149.
- Moghazy, E.A. and El-Shaarawy, M.O.A. (2001): Quality attributes of beef burger as affected by using propyls and frozen storage. *Egypt. J. Agric. Res.*, 79(4): 1499-1512.
- Moghazy, E.A.; Sharaf, S.M. and El-Seesy, T.A. (2004): Effect of substitutines mutton tailfat with vegetable oil on quality attributes of helthy beef sausage. *Egypt. J. Appl. Sci.*, 19(5): 191-205.
- Nuzhat, H.; Imran, P.; Sultan, S.; Zobia, J. and Asif, I. (2002): Incorporation of legumes as non-meat protein extenders in fresh beef sausages. *Pakistan J. Agric. Sci.*, 39(3): 236-239.
- Reddy, N.R.; Pierson, M.D.; Sathe, S.K. and Salunkhe, D.K. (1985): Dry bean tannins: a review of nutritional implications. *J. Amer. Oil. Chem. Soc.*, 62: 541-549.
- Santosh, K. and Chauhan, B.M. (1986) Effect of domestic processing and cooking and *in vitro* protein digestibility of moth bean. *J. Food Sci.*, 51 (4): 1083-1084.
- Snedecor, G.W. and Cochran, W.G. (1980): *Statistical Methods*. 7<sup>th</sup> Ed. Oxford and IBIT Pub. Co.
- Soloviev, V.E. (1966). *Aging of meat*. Food industry Pub., Moscow.
- Stauffer, C.E. (1993): Trypsin inhibitor measurement. Effect of order of reagent addition. *Cereal Chem.*, 7: 111-112.
- Urbano, G.; Lupez-Jurado, M.; Hernandez, J., Fernandez, J., Fernandez, M., Moreo, M.C.; Frias, J.; Diaz-Pollan, C.; Prodanov, M. and Vidal-Valverde, C. (1995): Nutritional assessment of raw, heated and germinated lentils. *J. Agric.Food Chem.*, 43: 1871-1877
- Vidal-Valverde, C.; Erias, J.; Estrella, I.; Gorospe, M.J., Ruiz, R. and Bacon, J. (1994): Effect of processing on some antinutritional factors of lentils. *J. Agric. Food Chem.*, 42: 2291-2295.
- Watts, B.M.; Yamaki, G.L., Jeffery, L.E. and Elias, L.G. (1989). *Basic sensory methods for food evaluation*. 1<sup>st</sup> Ed., The International Development Res. Center Pub., Ottawa, Canada.
- Winton, A.L. and Winton, R.B. (1958): Okoloff magnesium oxide distillation volumetric method. "The analysis of Food P.848, John Willey, New York and Chapman, London.
- Woye Woda, A.D.; Shaw, S.J.; Ke, P.J. and Burns, B.G. (1986): *Recommended Laboratory Methods for Assessment of Fish Quality*. Canadian Technical Report of Fisheries and Aquatic Science, No. 1448.
- Zaki, A.M. (1996): Raffinose oligosaccharides, proteins and some antinutritional factors in the seeds of two mung bean genotypes as affected by germination and processing. *Annals Agric. Sci. Moshtohor*, 34 971-982.
- Zaki, A.M.; Ismail, H.A., Abdel Naem and El-Morsi, E.A. (1999): Antinutritional factors and *in vitro* digestibility in guar seed as affected by germination and autoclaving. *J Agric Sci. Mansoura Univ.* 24(1) 7037



## إنتاج السجق والبرجر البقرى باستخدام بذور فول المانج المعالجة

همام الطوخي محمد بهلول\* ، إبراهيم محمد عبدالعظيم\*\*  
\* قسم علوم الأغذية - \*\* قسم الكيمياء الحيوية الزراعية .  
كلية الزراعة بمشتهر - جامعة الزقازيق/البحر

أجريت هذه الدراسة بهدف استخدام بذور فول المانج بعد التخلص من المواد الضارة في إنتاج السجق والبرجر البقرى. وجد أن محتوى البروتين لبذور فول المانج ٢٦,٨%. المواد الضارة: مثبط التربسين، التينينات، مثبط أنزيم ألفا أميليز، اللكتين، حمض الفيتيك، الفيسين، سكريات الراهينوز، الامتكيوز والفريمكوز في البذور الجافة الغير معاملة كانت ٤,٦ مجم/جم، ١,٣٦%، ٢٩٩ وحدة/جم، ١٩٠,٢ وحدة/جم، ١,٩٣%، ٤,٧ مجم/جم، ٠,٢٨%، ٠,٦٣%، ١,٤٦% على الترتيب.

وجد أن النقع في محلول بيكربونات الصوديوم (رقم pH ٧,٨٥) كان أكثر تأثير في إزالة معظم مضادات التغذية فيما عدا حمض الفيتيك. ومن جهة أخرى وجد أن الإنبات (لمدة ٧٢ ساعة) أكثر تأثيرا في خفض نسبة حمض الفيتيك، بينما كانت المعاملة بالأتوكلاف بعد الإنبات (٧٢ ساعة) أفضل في تحسين القيمة الغذائية.

تم استخدام دقيق بذور فول المانج الناتج بعد الإنبات والمعاملة بالأتوكلاف في تصنيع السجق والبرجر البقرى وذلك بعد استرجاعها بالماء بنسبة (١ : ٢) (وزن : وزن) وتم الاستبدال بنسب صفر، ٥، ١٠، ١٥، ٢٠، ٢٥، ٣٠، و ٣٥% من كمية اللحم المستخدم في التصنيع.

وأظهرت نتائج التحليل الكيماوى والتقييم الحسى للمنتجات المصنعة أنه يمكن استخدام نسبة استبدال حتى ٢٠% لإنتاج منتج جيد الصفات الحسية.