

Animal Health Research Institute,  
Mansoura Lab., Egypt

## **ESCHERICHIA COLI ASSOCIATED WITH SWOLLEN HEAD SYNDROME IN BROILER CHICKENS**

(With 4 Tables)

By

**M. M. ABD ELATIF**

(Received at 23/3/2004)

**إيشيريشيا كولاي المصاحبة لظاهرة تورم الرأس في دجاج التسمين**

**محمود محمد محمود عبد اللطيف**

أجريت هذه الدراسة علي ١٥٠ عينة ممثلة لعدد خمسة قطعان دواجن تسمين ما بين الأسبوع الثاني والأسبوع السادس من العمر مصابة بحالات تورم الرأس في نطاق محافظة الدقهلية، حيث أنه تم أخذ العينات من دجاج سليم ظاهريا وآخر مريض إكلينيكيًا. وبالفحص البكتريولوجي للعينات تم عزل الميكروب القولوني من كل العينات التي تم فحصها بنسبة ٧٨,٧% ، حيث كانت النسبة في كل من السليم ظاهريا والمريض إكلينيكيًا ٧٢% و ٨٥,٣% على التوالي وبالتصنيف السيرولوجي للمعزولات كانت كالتالي O<sub>2</sub> بنسبة ٣٥,٦% ، O<sub>78</sub> بنسبة ٣٠,٥% ، O<sub>5</sub> بنسبة ٢٠,٣% وعترات كانت غير مصنفة بنسبة ١٣,٦%. ويعمل إختبار الحساسية للمعزولات وجد أنها أكثر حساسية لمادة الأنروفلوكساسين ، الجنتاميسين ، الإرثرومايسين ، سيفوتاكسين والامستربتوميسين.

### **SUMMARY**

A total of 150 samples were taken from five broiler chicken flocks showing swollen head syndrome between 2<sup>nd</sup> and 6<sup>th</sup> weeks of age in Dakhlia governorate, where the samples were taken from apparently healthy as well as clinically diseased chickens. Bacteriological examination of the total number of collected samples revealed the isolation of *E. coli* with percentage of 78.7% where the isolation from apparently healthy and clinically diseased with percentage of 72.0% and 85.3% respectively. Serological identification of the isolates revealed that the serotype O<sub>2</sub> (35.6%), O<sub>78</sub> (30.5%), O<sub>5</sub> (20.3%) and untyped strains (13.6%). By using antimicrobial agents, *E. coli* isolates were more sensitive to enrofloxacin, gentamycin, erythromycin, cefotaxin and streptomycin.

**Key words:** *Escherichia coli*, swollen head syndrome, broiler chickens.

## INTRODUCTION

Swollen head syndrome (S.H.S.) is an acute upper respiratory disease of chickens with multiple causes, adversely affecting broiler livability and growth rate in many areas of the world (Alexander, 1991 and Hafez and Lohren, 1990). The disease was initially observed in chicken in South Africa and was confirmed by (Morley and Thomson, 1984). SHS has been reported in Egypt by Ahmed, (1991); Hebat Allah, (1997) and Abd-Rabu (1999), it characterized by swelling of peri- and infra-orbital sinuses, torticollis, nasal discharges and variable mortality (O'Brien, 1985). Pneumovirus (PV) was reported to be the primary causative agent of SHS in broiler (Morley and Thomson, 1984 and Naylor and Jones, 1993 ).

*Escherichia coli* has been a consistent feature of SHS, and it was isolated from internal organs, middle ear, brain and subcutaneous exudate of the head in outbreaks of SHS in broilers (Pattison *et al.*, 1989; Droual and Woolock, 1994 and Nakamura *et al.*, 1997).

Bacteriological examination of SHS cases in broilers revealed the isolation and identification of pathogenic *E.coli* which invades the conjunctiva and subcutaneous regions of the head as a subsequent secondary infection and responsible for the clinical appearance of SHS (Shane, 1991, Jons *et al.*, 1991 and Stehling *et al.*, 2003 ).

Therefore, *E. coli* is thought to be the main secondary complicating agent of SHS in broiler chickens (Lu *et al.*, 1994; Georgiades *et al.*, 2001 and Wafaa and Tanios, 2002).

This study was carried out for isolation and identification of *E. coli* associated with SHS from broiler farms in Dakahlia governorate, and testing the activities of some antimicrobial agents on the *E. coli* isolates.

## MATERIALS and METHODS

### (1) Samples:

A total of 150 samples collected from five broiler flocks from apparently healthy and clinically diseased chickens between 2<sup>nd</sup> and 6<sup>th</sup> week of age in Dakahlia governorate.

### (2) Media:

Media used for bacteriological examination included nutrient broth, MacConkey agar, Triple sugar iron agar, peptone water 1% for indole test, glucose phosphate broth for methyl red and voges proskauer tests, Simmon's citrate media and 1% peptone, glucose, sucrose,

mannitol and lactose were used .The different media and reagents were prepared according to the techniques described by Cruickshank *et al.*, (1975).

**(3) Bacteriological examination:**

Samples were collected from caseous materials and exudate of heads, nasal secretions, sinus contents and tracheal scraping of apparently healthy and diseased or freshly dead chickens, inoculated in nutrient broth and incubated at 37°C for 24 hours.

Subcultures were made onto MacConkey agar plates and incubated at 37°C for 24 hours. Suspected colonies were picked up and identified morphologically and biochemically. (Cruickshank *et al.*, 1975 and Brenner, 1992 ).

**Table 1:** Biochemical activities of *E. coli*

<i>Isolated bacteria</i>	<i>Indole</i>	<i>M.R.</i>	<i>V.P.</i>	<i>Urease</i>	<i>H<sub>2</sub>S</i>	<i>Citrate</i>	<i>Lactose with gas</i>	<i>Glucose with gas</i>	<i>Mannitol</i>	<i>Sucrose</i>
<i>E. coli</i>	+ve	+ve	-ve	-ve	-ve	-ve	+ve	+ve	+ve	+ve

**(4) Serological identification:**

*E. coli* "O" antisera from Behring Werk Ag. Marburg, Lahn, Germany were used for serotyping of the isolated strains. The procedure outlined by using polyvalent and monovalent *E. coli* antisera (Edward and Ewing, 1972).

**(5) Antimicrobial sensitivity test :**

The obtained *E. coli* isolates were tested for their sensitivity to the antimicrobial agents according to the method of Cruickshank *et al.*, (1975) by measuring the inhibition zone which was interpreted according to Manual of Bio-Merieux, (1986). The antimicrobial sensitivity discs were kindly obtained from " Oxoid " .

**RESULTS**

The results of bacteriological examination , identification and antibiotic sensitivity test were recorded in table 2, 3 and 4

**Table 2:** Prevalence of *E. coli* among apparently healthy and clinically diseased broiler

Flock	Apparently healthy broiler			Clinically diseased broiler		
	No. of collected samples	No. of positive samples	%	No. of collected samples	No. of positive samples	%
1	15	12	80.0	15	15	100
2	15	11	73.3	15	12	80.0
3	15	12	80.0	15	14	93.3
4	15	10	66.7	15	12	80.0
5	15	9	60.0	15	11	73.3
<b>Total</b>	<b>75</b>	<b>54</b>	<b>72.0</b>	<b>75</b>	<b>64</b>	<b>85.3</b>

**Table 3:** Serological identification of isolated *E. coli* strains

Serogroup	No. of positive/total*	Percentage
O <sub>2</sub>	42	35.6
O <sub>78</sub>	36	30.5
O <sub>5</sub>	24	20.3
Untyped	16	13.6

\* Total = 118 isolates

**Table 4:** The antibiogram of isolated *E. coli* recovered from examined samples .

Type of antibiotics	Disc	Sensitive		Moderate		Resistant	
	Conc.	No.	%	No.	%	No.	%
Erythromycin	15µg	68	57.6	29	24.6	21	17.8
Flumoquine	10µg	12	10.2	31	26.3	75	63.6
Chlorphenicol	10µg	50	42.4	31	26.3	37	31.4
Gentamycin	10µg	70	59.3	23	19.5	25	21.2
Streptomycin	10µg	60	50.8	28	23.7	30	25.4
Ampicillin	20µg	35	29.7	33	27.9	50	42.4
Amoxycillin	30µg	36	30.5	36	30.5	46	38.9
Enrofloxacin	10µg	89	71.2	17	14.4	17	14.4
Tetracyclin	30µg	52	44.1	26	22.0	40	33.9
Cefotaxin	30µg	65	55.1	26	22.0	27	22.9

The percentage were calculated in relation to the total number of examined isolates .

## DISCUSSION

Swollen head syndrome (SHS) is an emerging condition with multiple causes, *E.coli* is the most secondary bacterial cause of SHS in broiler, (Shane, 1991).

This work showed that the prevalence of *E.coli* from most examined samples in a total percentage of (78.7%), which isolated from apparently healthy (72.0%) and clinically diseased (85.3%), Table (2). These results were similar to those recorded by Parreire and Yano, (1998) isolated *E.coli* (72.0%) from chickens with SHS. Georgiades *et al.*, (2001), who isolated *E.coli* (87.5%) from the infraorbital sinuses of broiler chickens with SHS. Also Wafaa and Tanios, (2002) recovered *E.coli* strains (84.6%) from natural cases of SHS in broiler chickens. A lower percentage was reported by Rizk and Bkhiet, (2001) who isolated *E. coli* with an incidence of (15.5%) from broiler chickens with SHS. High incidence of *E.coli* in this study may be attributed to environmental conditions which play a significant role in interacting with infectious agents in the productions of respiratory diseases in poultry. (Kleven and Glisson, 1997).

Serological serotyping in this study revealed that *E. coli* serotypes were O<sub>2</sub> (35.6%); O<sub>78</sub> (30.5%); O<sub>5</sub> (20.3%) and untyped strains (13.6%) with available antisera (Table 3). These serotypes were also isolated by Zallen, (1986); Al-Ankari *et al.*, (2001) and Wafaa and Tanios, (2002). Other serotype O<sub>1</sub> was indentified by White *et al.*, (1993).

As shown in Table (4) the results of antimicrobial sensitivity test for the *E.coli* isolates revealed that the *E.coli* isolates were more sensitive to enrofloxacin (71.2%), gentamycin (59.3%), erythromycin (57.6%), cefotaxin (55.1%), streptomycin (50.8%), tetracycline (44.1%), chloramphenicol (42.4%) and moderate sensitive to amoxycillin (30.5%), ampicillin (27.9%), flumoquine (26.3%) .

In this respect our results agree to some extent to those reported by Hamouda and Amer, (2000) who showed that the *E.coli* strains which isolated from SHS were sensitive to erythromycin, enrofloxacin and gentamycin .

From the results of our study, we can concluded that the *E. coli* infection is the main secondary bacterial cause of swollen head syndrome in broiler chickens and most serotypes of isolated *E. coli* were sensitive for enrofloxacin, gentamycin, erythromycin, cefotaxin and streptomycin. Also this study suggested that *E.coli* infection in broiler can not be eliminated by antibiotics treatment only but required hygienic measures to be taken to lower the infection risk.

## REFERENCES

- Abd Rabu, K.M.B (1999)*: Studies on swollen head syndrome in chickens Mvsc. Dis. of poultry and Rabbits Facult. of Vet. Med., Cairo University .
- Ahmed, A.A.S. (1991)*: Newly disease of chickens in Egypt. Aerosols, 4., sept .
- Al Ankari, A.R., Bard bury, J.M., Naylor, C.J., Worthington, K.J., Johnson, C.P. and Jones, R.C. (2001)*: Avian pneumovirus inoculated with Escherichia coli at different time intervals. Avian pathology 30, 257-267 .
- Alexander, D.J. (1991)*: Pneumovirus infections (Turkey rhinotracheitis and swollen head syndrome of chickens). In disease of poultry Ed. by Calenk, B.W., Barnes, H.J., Beared, C.W.; Reid, W.M. and Yoder, Jr, H.W. 10<sup>th</sup> Ed. (1997). Iowa state University press, Ames, USA.
- Bio-Marieux (1986)*: Laboratory reagents and products. Bacteriology, Morey L'Etoile 69266, charbonnieres, Les Bains, Frances.
- Brenner , D.G. (1992)*: Introduction to the family Enterobacteriaceae in :I Balows, A.; Truper, H.G.; Dworkin, M; Herder, W. and schleifer, K.H. (Eds), the prokaryotes. Springer-Verlag Newyork, pp, 2673-2695.
- Cruickshank, K.R.; Duguid, J.P.; Marmion, B.p. and Swain, R.H.A. (1975)*: Medical. Microbiology. 12 th ed. Vol. II, Churchill Livingstone Limited Edinberg, London and New York .
- Droual, R. and Woolcock, P.R. (1994)*: Swollen head syndrome associated with E. Coli and Infectious Bronchitis virus in control valley of California AV. Path, 23 : 733- 742 .
- Edwards, P.R. and Ewing, MW.M. (1972)*: Identification of enterobacteriaceae. 3rd Ed. Burgers publishing Co. Minneoplos.
- Georgiades, G.; Iodanidis, P. and Maria Konmbari (2001)*: Cases of swollen head syndrome in broiler chicken in Greece. Avian Dis, 45 : 745-750.
- Hafez, H.M and Lohren, U. (1990)*: Swollen head syndrome: clinical observations and serological examinations in west Germeny, DTW 97 : 322-324.
- Hamouda, A.S. and Amer. M.M. (2000)*: Diagnosis and control of swollen head syndrome in a broiler chickens farm. J. Egypt Vet. Med .Ass. 60,(5) 63 – 72.

- Hebat Allah, A.M. (1997)* : Some studies on swelling head syndrome in chickens Ph.D. thesis, Fac, of Vet .Med .Assiut university.
- Jones, R.C.; Naylor, C.J.; Bradbury, J.M.; Savage, C.E.; Worthington, K. and William, R.A. (1991)*: Isolation of turkey rhinotracheitis like virus from broiler breeder chickens in England.Vet. Res, 129 :509-510.
- Kleven, S.H. and Glisson, J.R. (1997)*: Multicausal respiratory diseases. In Diseases of poultry. 10<sup>th</sup> ed. B.W. Calnek, H.Jhon Brnes, C.W Beard, L.R.Mc Dougald and Y.M.Saif. Iowa atate University Press, Ames, Iowa. PP. 1008-1009.
- Lu, Y.S; Shein, Y.S.; Tasi, H.J.; Tseng, c.s.; Lee, S.H. and Lin D.F. (1994)*: Swollen head syndrome in Taiwan–isolation of an avian pneumovirus and serological survey Avian Pathol .,23:169–174.
- Morley, A.J. and Thomson. D.K. (1984)*: Swollen head syndrome in broiler chickens. Avian Dis., 28 : 238 – 243.
- Nakamura, K.; Mase, M.; Tanimura, N.; Yamaguchi, S. Nakazawa, M. and Yuasa, N. (1997)*: Swollen head syndrome in broiler chickens in Japan: its pathology, microbiology and biochemistry . Avian pathol.,26 : 139 – 154.
- Naylor, C.J. and Jopnes R.C. (1993)*: Turkey rhinotracheitis: a review. Vet. Bull.,63 (5) 439 – 449.
- O'Brien, J.D.P. (1985)*: Swollen head syndrome in broiler breeders. Vet. Rec ., 117 619 –620.
- Parreira VR anf Yano T. (1998)*: Cytotoxin produced by Escherichia coli isolated from chickens with swollen head syndrome (SHS). Vet. Microbiol. May; 62(2): 111-9.
- Pattison, M., Chettie, N.; Randall, C.J.and Wyeth, P.J. (1989)*: Observations on swollen head syndrome in broiler and broiler breeder chickens. Vet.Rec., 125 : 229 – 231.
- Rizk, M. Sh. and Bkhiat, A.A. (2001)*: Bacteriological aspects of facial odema in native and foreign chicken at Bohaira province. Assiut Vet. Med. J.Vol 45 No 90.
- Shane, S.M. (1991)*: The swollen head syndrome an emerging condition in broiler. Vineland updqte (38).
- Stehling, E. G, Yano, T, Brocchi. M. and da Silveira WD. (2003)*: Characterization of a plasmid-encoded adhesion of an avian pathogenic Escherichia coli ( APEC ) strain isolated from a case of swollen head syndrome (SHS). Vetmicrobiol. 2003 Aug 29; 95 (1-2) : 111-20.

- Wafaa, M.M. Hassan and Tanios, N.I. (2002):* Natural and experimental infection of E.coli in swollen head syndrome in Broiler chickens .J.Egypt Vet .Med. Ass. 62(6)207 – 216.
- White D.G, Dho-Moulin M, Wilson R.A, and Whittam T.S (1993):* Clonal relationships and variation in virulence among Escherichia coli strains of avian origin. Microb pathog. May; 14(5) : 399-409.
- Zallen GH.K. (1988):* Case report: swollen head syndrome in chickens. Proc. 37<sup>th</sup> , West. Poult. conf. Californilor, 139.