

Animal Health Research Institute, Assiut, Egypt

OCCURRENCE OF KLEBSIELLA IN SOME MEAT PRODUCTS AND THE EFFECT OF COLD STORAGE AND SOME MEAT ADDITIVES ON ITS GROWTH RATE IN ASSIUT.

(With 3 Tables and 4 Figures)

By

AMAL A. MOHAMED and A.M. NASSAR*

*Dept of Food Hygiene, Faculty of Vet. Med, Assiut Univ., Egypt

(Received at 26/9/2004)

مدى تواجد الكليبيسيلا في بعض منتجات اللحوم وتأثير التخزين بالتبريد وبعض اضافات اللحوم على معدل نموها في أسيوط

أمال احمد محمد ، احمد محمد نصار

تم تجميع مائة عينة عشوائية من منتجات اللحوم البيفبرجر واللاتشون والسجق والبسطرمة (٢٥ من كل نوع) من محلات السوبر ماركت المختلفة في أسيوط بهدف التعرف على العدد الكلى وعزل وتصنيف ميكروب الكليبيسيلا. أظهرت النتائج أن كل العزلات صنفت *Klebsiella pneumoniae*. تم عزل الميكروب بنسبة ٥٢، ٢٨ و ٨% من عينات البيفبرجر واللاتشون والسجق بمتوسطات ٣، ٢، ١٠ × ± ١، ٦، ١٠، ٥، ٩١ × ١٠ ± ٤، ١١ و ١، ٤ × ١٠ ± ٠، ٧، خلية/جم بالترتيب. ثبت من التحليل الاحصائي وجود فروق معنوية بين متوسطات البيفبرجر والسجق وكذلك بين اللاتشون والسجق بينما لا توجد فروق معنوية بين البيفبرجر واللاتشون ولم يتم عزل الميكروب من عينات البسطرمة المفحوصة. تم دراسة تأثير التخزين بالبرودة عند درجتى حرارة مختلفة وهي ٤ درجة مئوية لمدة ٤ أيام و-٤ درجة لمدة ٢١ يوم على نمو الميكروب فى اللحم المفرومة. حيث أظهرت النتائج أن تأثير الميكروب بدرجة -٤ أكبر من تأثيره بدرجة +٤ مما أدى الى تأخر نموه. وكذلك تمت دراسة تأثير بعض اضافات اللحوم مثل كلوريد الصوديوم بنسبة ١، ٢ و ٣% وكذلك نيتريت. الصوديوم بنسبة ١٠٠ و ٢٠٠ جزء فى المليون فى ثلاثة درجات حرارة مختلفة وهي ٤، ٣٧ و ٤٢ درجة مئوية على التوالي. أظهرت النتائج تأخر نمو الميكروب بزيادة درجة الحرارة وزيادة تركيز المادة المستخدمة كما ظهرت فروق معنوية فى التعداد اللوغاريتمى بين المجموعة الضابطة (control) والعينات المعاملة محل الدراسة. وقد تم مناقشة الأهمية الصحية للميكروب المعزول والاحتياطات الواجب اتخاذها لتقليل تواجدها والتخلص منه لحماية المستهلك.

SUMMARY

One hundred random samples of meat products; beef-burger, lunchon, sausage, and basterma (25 each) were collected from different

supermarkets in Assiut and subjected to enumeration, isolation and identification of *Klebsiella* organisms. The results showed that all the strains were identified as *Klebsiella pneumoniae*. *Klebsiella* organisms could be detected with percentage of 52, 28, and 8 from beef –burger, luncheon and sausage with mean values of $3.2 \times 10^4 \pm 1.6$, $5.9 \times 10^3 \pm 4.11$ and $1.4 \times 10^2 \pm 0.07$ cfu/g respectively. No significant difference could be detected among beef –burger and luncheon, while a significant difference appeared between beef burger and sausage and also between sausage and luncheon at $P < 0.05$. *Klebsiella* could not be detected in basterma. The effect of cold storage (chilling and freezing) on the growth rate of *Klebsiella* strains was determined at temperatures of 4°C and -4°C in minced meat. The results indicated that freezing showed reduction in the growth rate and inactivation more than chilling. The effect of different concentrations of sodium chloride (1,2 and 3%) and sodium nitrite (100, and 200 ppm) were tested against the isolated organism in minced meat at three different temperatures 4°C , 37°C and 42°C respectively. Highly significant differences were recorded between the control and the treated samples by increase temperature and concentration of sodium chloride and sodium nitrite. Public health significance and suggested measures for improving the quality of meat products to protect the consumer were given.

Key words: *Klebsiella*, cold storage, meat additives.

INTRODUCTION

Meat and meat products are liable to be contaminated during processing and handling by various microorganisms thus causing spoilage and economic losses.

The genus *Klebsiella* was named after Edwin klebs, a late 19th. century German microbiologist. All *Klebsiella* species are non motile, many strains hydrolyze urea producing a light pink color in the slant urea agar, *Klebsiella pneumoniae* is indole negative and *Klebsiella oxytoca* is indole positive (Koneman *et al*, 1992).

Klebsiella species are widespread through the environment and also carried by human. Both genera are well- recognized community and nasopathogens and cause significant infections. They are a common cause of respiratory and non respiratory infections. It is responsible for 1% to 5% of all case community –acquired pneumonia, that generally occur in patients with debilitating conditions such as alcoholism,

diabetes mellitus and chronic obstructive pulmonary disease with extensive necrosis and hemorrhage. Lung abscess occur after pneumonic process or secondarily to *Klebsiella* species infection and have high rates morbidity and mortality (Bouza and Cercenado, 2002). *Klebsiella pneumoniae* is an opportunistic pathogen responsible for nosocomial infection and trend toward multiple antibiotic resistance (De champs *et al.*, 2004). It is also responsible for acute renal failure to acute pyelonephritis with extensive necrosis of the kidney (Creighton *et al* 2001).

Sixty adult *klebsiella meningitis* patients have been identified, most cases were associated with debilitating disease and devastating metastitic septic abscesses (Lu CH *et al.*, 2002). Also a histamine producing strain of *Klebsiella pneumoniae* was isolated and could be detected by (Taylor *et al* 1979).

The most popular meat products are hamburger, luncheon, sausage and basterma; For the first time Sabota *et al.*, (1998) reported *Klebsiella pneumoniae* as enteroinvasive food-borne pathogen transmitted from hamburger and symptoms of gastroenteritis rapidly deteriorated to multiorgans failure. The public health significant of *Klebsiella* organisms in food materials particularly meat products and their role as source of *Klebsiella* infection have been studied by many investigators (Hechelmann *et al.*, 1974 and Okolo 1985).

this study was under taken to :

- 1-Investigate the occurrence of *Klebsiella* organisms in some meat products (beef-burger, luncheon ,sausage and basterma) in Assiut.
- 2-Count and identification (biotyping) of the isolated of *Klebsiella*.
- 3-Study the effect of cold storage chilling (+4C⁰) and freezing (-4C⁰) on the growth rate of the organism in minced meat.
- 4-Study the effect of different concentrations of sodium chloride and sodium nitrite on *klebsiella* growth rate at three different temperatures, 4, 37, 42 C⁰.

MATERIALS and METHODS

Collection of samples :

One hundred random samples of meat products; 25 each of beef-burger, luncheon, sausage and basterma were collected from different supermarkets in Assiut and transferred to the laboratory. All the samples were examined for enumeration, isolation and identification of *Klebsiella* organisms.

Preparation of samples:

The collected samples were prepared according to the technique recommended by ICMSF (1978) for enumeration of *Klebsiella* organisms. The surface plate technique described by Bagley and Seidler, (1978) was applied using double MacConkey- inositol- carbenicillin agar medium., The number of large mucoid and red colony with red pigment usually diffusing into the surrounding agar indicating fermentation of lactose and acid production was then calculated and recorded. The isolated *Klebsiella* organisms were identified according to morphological characters (Cruickshank *et. al*, 1975); biochemical characters (Edward's and Ewing, 1972) and biotyping according to Niazi *et. al.*, (1977).

Test organisms:

One strain of *Klebsiella pneumoniae* previously isolated and identified from meat products was grown in 10 ml of nutrient broth at 37C⁰/24 h. The culture was decimally diluted and plated to enumerate the organisms present. The culture was diluted to achieve an inoculum level of 10⁶/g for *Klebsiella pneumoniae*. 1kg of semitendinosus muscle of cattle carcass was passed twice through meat mincer then divided and inoculated with the organisms suspension and thoroughly mixed to have 3x10⁶ organisms /g.

Experimental part:

- 1- The effect of chilling and freezing on the growth rate of *Klebsiella* organism in minced meat for 4day for chilling and 21day for freezing.
- 2-Effects of different concentrations of sodium chloride (1,2,and 3%) and sodium nitrite (100, and 200 ppm) during storage at different temperatures on the growth rate of *Klebsiella* organism.

1- The effect of cold storage on the growth rate of *Klebsiella* organisms:

a- The effect of chilling:

50 gm of the inoculated sample were stored at chilling temperature (4⁰C) and examined daily for 4 days for detecting the number of *Klebsiella* per gram which was done by surface plate count technique.

b- The effect of freezing:

100 gm of minced meat which were inoculated with 3x10⁶/gm were stored at -4C⁰ and examined for the aimed organisms every two days till 21 days

2- The effect of some additives and storage temperature on growth rate of klebsiella organisms:

a- sodium chloride:

600 gm minced meat previously inoculated with *Klebsiella* 3×10^6 were divided into 12 parts, 50gm each as the following: 3 parts as control, 3 parts with 1% added NaCl, 3 parts with 2% added and 3 parts with 3% added. Then one part of the control and one part of each concentration were stored at 4°C, 37°C and 42°C. The samples were examined daily for 4 days at temperature 4°C and for 2 days at temperature 37°C and for one day at temperature 42°C. The total colony count was recorded in each plate.

b- Sodium nitrite:

The used sodium nitrite concentrations were 100 and 200 ppm. Three hundred grams of minced meat samples were divided into 2 parts; part was inoculated with 100 ppm and divided into 3 parts (50gm each) then stored at 4°C, 37°C and 42°C. The other part was inoculated with 200 ppm and divided into 3 parts according to the storage temperature 4°C, 37°C and 42°C. Then the samples were examined as mentioned above.

Statistical analysis:

The data were statistically analyzed by Mann-whitney Rank sum test (non-parametric test) according to Michigan State University (1988).

RESULTS

The results were explained in 3 tables and 4 figures.

DISCUSSION

It is evident from achieved results in table (1) and according to the biochemical identification that all strains were identified and biotyped as *Klebsiella pneumoniae*. The incidence percentages of *Klebsiella pneumoniae* recovered from different meat products were 52, 28, and 8 from beef-burger, luncheon and sausage respectively. The count ranged from 7×10^2 to 2×10^5 , 4×10^2 to 3×10^4 and 7×10 to 2×10^2 in beef-burger, luncheon and sausage respectively. The mean values were $3.2 \times 10^4 \pm 1.6 \times 10^4$, $5.9 \times 10^3 \pm 4.11$ and $1.4 \times 10^2 \pm .07$ cfu / g for the above mentioned products respectively. *Klebsiella* organisms could not be detected in basterma samples. The results show significant difference between the mean *Klebsiella* count value in case of beef-burger and

sausage and also between luncheon and sausage but no significant difference was found between mean values of beef- burger and luncheon. Sausage is liable to contamination from different sources during preparation, processing, handling storage (Chesnut *et al*, 1977). Our results for sausage is nearly similar to the results obtained by Morshdy (1991) and lower than Sedik *et al* (1988), Eldaly *et al* (1989), Khalafalla and El-Sherif (1993), and Tolba (1994), our results is higher than (Ahmed *et al* 1988 and Soriano *et al* 2000). Our results in hamburger are lower than Sedik *et al* (1988) and higher than El Mossalami *et al* (1982), Ahmed *et al* (1988) and Tolba (1994). In luncheon our findings are lower than Sedik *et al* (1988) and Refaie and Nashed (1989) and higher than Tolba (1994). *Klebsiella* microorganisms could not be detected in basterma and this agrees with Sedik *et al* (1988) and Tolba (1994) but Refaie and Nashed (1989) could detect *klebsiella* in basterma in a percentage of 11%. Basterma samples were contaminated to lesser extent than the other three products and this may attributed to the addition of nitrite which has a bacteriostatic effect (Libby, 1975) and the curing processing of the product which play a great effect in inhibiting the multiplication of the microorganisms (Fehlhaber 1981).

Klebsiella pneumoniae was reported as enteroinvasive food-borne pathogen transmitted from hamburger and has been documented to harbor heat- labile and heat stable enterotoxins (Sabota *et al*, 1998). The presence of *Klebsiella pneumoniae* has a short survival time in meat product and could indicate recent contamination which turn indicated unsanitary and unhygienic handling and suspected contamination of food with feces as modes of transmission causing sever vomiting (Newton *et al*, 1977). Kleeberger *et al* (1980) reported that the higher Enterobacteriace count might be attributed to contamination of flesh used for manufacture of such products. Moreover, mincing machines, grinder, equipment and knives are considered sources of infection and contamination of meat during processing (Frazier 1967 Chesnut *et al* 1977, and Eldaly, 1983).

It is evident from the results achieved in table (2) when minced meat were inoculated with *K.pneumoniae* (3×10^6 /g) and stored at $4C^0$ and $-4 C^0$ that chilling had no effect on growth till 4 days where no significant multiplication was noticed. Similar findings were reported by Hechelmann *et al*, (1974) and ICMSF (1978). While freezing showed a remarkable and gradual decrease in count during storage. Our results for freezing agree with Calcott *et al* (1976) who mentioned that cooling and warming rates affect bacterial survival like *Klebsiella*. Takahashi *et al*,

(1982) showed that these organisms were sensitive against freezing and thawing. Karim and Yu (1980) reported a gradual decline in number of the organisms in frozen and frozen thawed meat samples 2-4 weeks after their storage at -17°C . In the same site ICMSF (1978) reported that freezing may result in death of microorganisms.

Most bacteria were sensitive to salt on freezing and thawing. Psychrotrophic bacteria can develop and multiply under refrigeration temperature and constitute the most common cause of refrigerated food spoilage Frazier and Westhoff (1978).

The effect of meat additives and storage at various temperatures on the growth rate of *K.pneumoniae* experimentally inoculated into minced meat (count in Log_{10}) is reported in table 3. Storage temperature at 4°C showed market reduction in the count of *K.pneumoniae* in control sample and with the use of different concentrations of 1, 2 and 3% sodium chloride the count was reduced from 6.48 to 3.9 Log_{10}/g and the use of sodium nitrite 100 and 200 ppm. reduced the count to Zero when incubated for 1 day. While *K.pneumoniae* at 37°C storage temperature the control samples showed gradual rising of the count which reach to 8.1 Log_{10} , and by addition of sodium chloride concentration the count reduced to 5.6 Log_{10} , while addition of sodium nitrite reduced the count to 5.5 Log_{10} . after 2 days. Storage temperature at 4°C showed no significant change until 3 days and slight increase in the 4th day and by addition of sodium chloride 1, 2 and 3% the count was reduced to 5.70 Log_{10} and addition of sodium nitrite 100 and 200 ppm reduced the count to 4.85 Log_{10} after 4 days of storage.

These findings showed that Klebsiella organisms are one of psychrotrophic bacteria. It survives in meat products at temperature of refrigeration, the results agree with Hechelmann *et al*, (1974) and Patterson and Gibbs (1977), that the organisms of genus Klebsiella could contribute substantially to spoilage of refrigerated meat. Our results showed that increasing sodium chloride decreased the count of *Klebsiella pneumoniae* and this may be due to the inhibitory effect of sodium chloride on microorganisms, this agree with Muralidhararao and Nandy, (1977). Also Lechowich *et al*, (1978) stated that the salt decreases water content in microorganisms and subsequently reduces the microbial growth. The effect of temperature shows that the organism count was increased to its maximum growth at 37°C and this is similar to that reported by Johnson *et al* (1975) and Tsujia, *et al* (1982).

It is concluded that strict hygienic conditions should be applied and best quality of ingredients is to be used for meat processing. Food

establishment should be hygienically constructed and supplemented with good equipment's easily cleaned and disinfected. Moreover, concerned authorities should be taken an active part in controlling such products by imposing bacteriological standard to ensure a maximum safety to the consumers.

REFERENCES

- Ahmed, A.H.; Ahmed, H.A.; AbdEl-Rahman, H.A. and Moustafa, M.K. (1988): Incidence of enterobacteriaceae in some selected food stuffs. *Assiut Vet. Med. J. Vol. 20, No.40: 104-109.*
- Bagley, S.T. and Seidler, R.J. (1978): primary *Kelebsiella* identification with MacConkey – inositol – carbenicillin agar. *Appl. Environ. Microbiol.* 36, 536-580.
- Bouza, E. and Cercenada, E. (2002): *Klebsiella* and *Enterobacter* :antibiotic resistance and treatment implications. *Semin Respir Infect. Sep; 17(3): 215-30.*
- Calcott, P.H.; Lee, S.K. and Macleod, R.A. (1976): The effect of cooling and warming rate on the survival of variety of bacteria. *Can. J. Microbiol. Jan, 22(1) 106-9.*
- Chesunt, C.M.; Emswiler, B.S. and Kotula, A.W. (1977): Bacteriological quality of ingredients used in ground beef manufacture. *J. Ansci, 44; 213.*
- Creyghton, W.M.; Lobatto, S. and Weening, J.J. (2001): Acute renal failure caused by *Klebsiella pneumoniae* pyelonephritis. *Clin Nephrol. Nov; 56 (3): 391-3.*
- Cruickshank, R.; Duguid, J.; Marmion, B. and Swain, R. (1978): *Medical Microbiology* 12th Ed. Churchill Livingston, Ebinbrugh, London. New York.
- Dechamps, C.; Rich, C.; Chandezon, P.; Chanal, C. and Siro forestier (2004): Factors associated with antimicrobial resistance among clinical isolates of *Klebsiella pneumoniae* 1- year survey in a french university hospital. *Eur.J. clin. Microbiol.*
- Edward's, P.R. and Ewing, W.H. (1972): *Identification of Enterobacteriaceae.*, 3rd Ed. Burgess Publ. Co. Minneapolis.
- Eldaly, E. (1983): Sanitary status of meat serving establishments Ph.D.V.SC. Thesis Cairo University.
- Eldaly, E.; Mansour, N. and Saleh, E. (1989): Occurrence of Enterobacteriaceae in some Egyptian meat products. *Alex. J. Vet. Sci, Vol. 5, No.1, 345-357.*

- El-Mossalami, E.; Roushdy, S. and Ibrahim, A.M. (1982):* Sanitary condition of locally produced hamburger, J Egypt.Vet. Med. Assoc .Vol. 42. No. 4, 71-84.
- Fehlhaber, K. (1981):* Properties of Shigella with referance to food hugiene. Archi Experimentelle Veterinr Medizin 35: 955-964.
- Frazier, W.C. (1967):* Food Microbiologie 2nd Ed.,N.X..Mc.Graw Hill, New York Inc
- Frazier, W.C. and Westhoff, D.C. (1978):* Food Microbiology 3rd Ed. Tata Mc-Graw Hill Publ.Co. Lrd., New Delhi.
- Hechelmann, H.; Bom, Z.; Uchidd, K. and Lesstner, L. (1974):* Vorkommendes tribus Klebsiellae Kuhlge lagertem.flesh und Fleischwaren. Fleischwirtschaft, 54, 1515-1517.
- ICMSF (1978):* International Commission on Microbiological Specification for Food: Microorganisms in Foods, Their Significance and Methods of Enumeration, Vol. 1 2nd Ed. Univ. of Toronto Press, Toronto, Buffaloe, Canada.
- Johnson, R.; Colwell, R. and Tamura, K. (1975):* Numerical totonomy study of enterbacteriaceae Int. J. syst. bact. 25(1), 12-37.
- Karim, M.I. and Yu, S.Y. (1980):* Microbiological quality of fresh frozen and frozen thawed meat. Malazian. Appl. Bact., 9(2)75.
- Khalafalla, F.A. and El-Sherif, A. (1993):* Psychrotrophic bacteria in sausage. Die Nuhung 37, 5, 428-432.
- Konemen, E.W.; Allen, S.D.; Janda, W.M. Schrechenberer, and Winn, W.C. (1992):* Color Atlas and Texbook of Diagnostic Microbiology. 4 th Ed., winters, R. (ed.) J.B. Lippincot Company, Philadelphia.
- Kleeberger, A.; Schafer, K. and Busse, M. (1980):* Ecology of enterobacteria on slaughter house meat. Fleischwirtschaft, 60(8), 1529-1531.
- Lechowich, R.; Brown, W.; Deibel, R. and Somers, I. (1978):* The role of nitrite in production of canned cured meat products. Food Technol. 5, 45-58.
- Libby, J.A. (1975):* Meat Hygiene 4th Ed. LEA and Febi GER Philadelphia.
- Lu, CH., Chang, W.N, Chang, H.W. (2002):* Klebsiella meningitis in adults: clinical feature prognostic factors and therapeutic outcomes. J. Clin Neurosci Sep; 9(5): 533-8.
- Michign State University (1988):* MSTAT-C, a software program for the design, management and analysis of agronomic research experiments. Michigan, U.S.A.

- Morshdy A., Ahmed Ammar, Rabeisleh, El-Sayed, El-Naenaeey and El-Tahir, M. Haraun, E.M. (1991):* Sanitary studies of sausage marketed in El-Gassi, M., Zagazing Vet. J. Vol. 14 No. 1 115-122.
- Muralidhararao, N. and Nady, S. (1977):* Organisms of Enterobacteriaceae family associated with animal by products. Ind. J. Anim. Sci, 47(6) 344-384.
- Newton, K.C.; Harrison, J.C. and Smith, K.M. (1977):* Coliforms from hides and meats. Appl. Environ. Microbiol. Jan, 199-200.
- Niazi, Z.; Kirpal, G.; Amtsberg, G. and Refait, M. (1977):* Biochemistry, serology, pathogenicity to mice and the resistance to antibiotics of Klebsiella strains of various kind of animals. Berl. Munch. Tierarzt. Wschr. 90(22), 435-440.
- Okolo, M.I. (1985):* Observation on the bacteriological quality of market goat meat and public health significance. Bull. Anim. Hlth. Prod. Afr. 33. 307-310.
- Patterson, J.T. and Gibbs, P.A. (1977):* Incidence and spoilage potential of isolates from vacuum-packaged meat of high pH value. J. Appl. Bact. 43, 25-38.
- Refaie, R.S. and Nashed, S.M. (1990):* Bacteriological studies on enterobacteriaceae in some meat products. Assiut Vet. Med. J. Vol. 23 No. 45, April 110-115.
- Sabota, J.M.; Hoppes, W.L.; Ziegler, J.R.; Dupont, H.; Matherwson, J. and Rutecki, G.W. (1998):* A new variant of food poisoning :enteroinvasive Klebsiella pneumoniae and Escherichia coli from a contaminat hamburger. Am. J. Gastroenterol. Jan, 93(1): 118-9.
- Sedik, M.F.; Nouman, T.M; Zienab, M.; Niazi, and Gergis, A.F. (1988):* Klebsiella in meat and meat products. Vet. Med. J. 36(1), 37-50.
- Soriano, J.M.; Rico, H.; Molto, J.C. and Manes, J. (2000):* Microbial evaluation of Spanish potato omelette and cooked meat samples in University restaurants. J. Food Prot. Sep.,
- Takahashi, M.; Ito, S. and Yoshida, K. (1982):* Effects of freezing and thawing and storage by freezing on the biological and morphological properties of strain k-9 of *Klebsiella pneumoniae* and its variants. Z Allg Mikrobiol, 22(9): 649-60.
- Taylor, S.L.; Guthertz, L.S.; Leatherwood, M. and Liebre, E.F. (1979):* Histamine production by *Klebsiella pneumoniae* and an incident of scombroid fish poisoning, Appl Environ. Microbiol. Feb, 37(2): 274-8.

- Tolba, K. (1994): Microflora in locally processed frozen meat. Med.J.,Giza.Vol. 42, No. 2. 99-105.
- Tsuji, A.; Kaneko, Y.; Takahashi, K. ; Ogawa, M. and Gotos (1982): The effect of temperature and pH on the growth of eighteen and nine glucose non-fermenting of gram-negative organism. Microbiol Immuno. 1.; 26 (1): 15-24.

Table 1: Statistical values of *klebsiella* count/g of the examined meat products.

(n=25 of each)

Samples	No. of positive <i>Klebsiella pneumoniae</i>	%	+ve other k. species	%	Klebsiella count/g **			
					Min.	Max.	Mean *	S.E
Beef-burger	13	52	0	0	7x10 ²	2x10 ⁵	3.2x10 ⁴ E	±1.6x10 ⁴
Lunchon	7	28	0	0	4x10 ²	3x10 ⁴	5.9x10 ³ E	± 4.11
Sasusage	2	8	0	0	7x10	2x10 ²	1.4x10 ² F	±.07
Basterma	--	--	0	0	--	--	--	--

*Means have the same letter are not significant (p<0.05)

**Average of 3 trials.

Table 2: Effect of chilling and freezing on *klebsiella* in minced meat. (Log₁₀ cfu/g.)

Storage period/day	Chilling	Storage period/day	Freezing
0	6.48	0	6.48
1	6.49	2	6.45
2	6.5	4	6.39
3	6.54	6	6.08
4	7.30	8	5.90
5	Spoilage	10	5.89
		12	5.83
		14	4.69
		16	4.60
		18	4.18
		21	4

Table 3: The Effect of sodium chloride and sodium nitrite on *Klebsiella pneumoniae* organisms in minced meat at different temperature.

Temp.	Storage Period/day	Control	Sodium chloride			Sodium nitrite	
			1%	2%	3%	100 ppm	200ppm
42C ^o	0	6.48 B	6.48 B	6.48 B	6.48 B	6.48 B	6.48 B
	1	5.68 H	5.64 D	5.64 D	3.95 C	0.0 N	0.0 N
37C ^o	0	6.48 B	6.48 B	6.48 B	6.48 B	6.48 B	6.48 B
	1	6.69 I	5 J	5.94 A	5.70 H	6.73 I	6.15 K
	2	8.15 L	5.89 A	5.73 H	5.63 D	6.46 M	5.5 O
4C ^o	0	6.48 B	6.48 B	6.48 B	6.48 B	6.48 B	6.48 B
	1	6.49 B	6.46 B	6.48 P	6.30 F	6.11 Q	5.96 E
	2	6.52 B	6.32 F	6.40 E	6.15 R	6 E	5.64 A
	3	6.54 B	6.28 F	5.91 A	5.78 S	6 E	5.93 A
	4	7.30 T	6.04 E	5.90 A	5.61 D	5.70 H	4.85 U
	5	spoilage					

Log₁₀ cfu/g have the same letters are not significant (p<0.05).




Fig. 1: The effect of chilling and freezing on *Klebsiella Pneumoniae* in minced meat.




Fig. 2: The effect of sodium chloride and sodium nitrite on *Klebsiella Pneumoniae* in minced meat at 42°C.




Fig. 3: The effect of sodium chloride and sodium nitrite on *Klebsiella Pneumoniae* in minced meat at 37°C.




Fig. 4: The effect of sodium chloride and sodium nitrite on *Klebsiella Pneumoniae* in minced meat at 5°C.