

EFFECT OF TAFLA (EGYPTIAN CLAY) AS FEED SUPPLEMENT ON THE PERFORMANCE OF BROILER CHICKS.

Abd El Meged, A. H*; A. A Hemid** and F. Abdel Azeem**

* Poultry Nutrition Department, Animal Production Research Institute, Agricultural Research Center, Ministry of Agriculture, Dokki, Giza, Egypt.

**Department of Poultry Production, Faculty of Agriculture, Ain Shams University, Cairo, Egypt.

ABSTRACT

A total number of 150 unsexed 7day old Hubbard chicks were distributed into five experimental treatments. Each of 30 chicks in 3 replicates (10 chicks / replicate) was used in this study. The first experimental group was fed on the control diet, whereas, groups 2, 3, 4 and 5 were fed on the control diet supplemented with Tafla (local clay) at levels of 1.5, 3, 4.5, and 6% .A starter diet was used during the starter period (7-28 days of age) contained 22.03 % CP and 3105 kcal ME /kg diet, while the finisher diet containing 19.00 % CP and 3196 kcal ME /kg diet was used during the finisher period (29-49 days of age) all the nutrients were as recommended by NRC (1994). Diets were nutritionally evaluated through growth performance feed intake and feed conversion as well as percentage of dressing, digestibility trials and blood parameters.

Results obtained could be summarized as follows:

1. Average live body weight, body weight gain, survival rate and economical efficiency % were significantly ($p < 0.01$) increased in broiler groups fed on the diets supplemented with 3% Tafla compared with other groups.
2. Values of feed conversion were significantly improved, while average feed consumption was decreased significantly with Tafla supplementation 1.5% level.
3. Digestibility coefficients of DM, OM CP, EE, CF and NFE were significantly ($p < 0.01$) increased with Tafla supplementation at levels 1.5 and 3%.
4. Relative weights of carcass traits studied were significantly affected by 1.5 and 3% Tafla supplemented diets

In general, these results indicate that using (1.5 and 3% Tafla) as feed additives enhance the productive performance and economical efficiency with no adverse effects on blood constituents of broiler chicks.

Keywords: Clay, Tafla, broiler, diets, productive performance.

INTRODUCTION

Numerous mineral materials (natural clay) are receiving greater attention in poultry production. In this respect, the research and development efforts aimed to find out the possibility of employing of natural clay such as (Tafla, zeolite, clinoptilolite, and bentonite) in enhancement of poultry performance.

Dobeic (1994) found that, supplementation of poultry diet with clinoptilolite (zeolite) at 1, 2 or 4% levels increased feed conversion efficiency. The highest feed conversion was with 2% clinoptilolite. Ziobina (1990), Elliot and Edwards (1991) and Ward *et al.* (1993) also found that natural zeolite significantly improved feed efficiency. Fisinin *et al.* (1985) and

Karelina (1985) found that zeolite increased the efficiency of feed utilization for growth and increased the financial returns in broilers. Berrios et al (1983) clarified that, feed conversion was significantly more efficient, consumption of protein and ME were low and cost of diet was reduced with zeolite dietary supplementation. Kasumov *et al.* (1986) indicated that, addition of clinoptilolite to the diet of laying hens was economically advantageous. Ziobina (1990) also found that, daily weight gain and meat quality of broiler chickens were improved with 3% zeolite added in the diet given for 49 days. Dobeic (1994) postulated that, clinoptilolite (zeolite) at levels of 1, 2 or 4% in the diet increased live weight gain and final body weight. The highest growth rate was with 2% clinoptilolite. On the other hand, Ward *et al.* (1993) found that, growth rate decreased with addition of 0.75% sodium zeolite in chicks diet. Lemser *et al.* (1992) reported that, 1.5, 3 or 5% bentonite decreased body weight by 11.6, 14.5 and 12%, respectively, while zeolite supplements in broilers diet showed no effect. Kalyuzhnov *et al.* (1988) also reported that, with zeolite dietary supplement, growth rate was intensified in broiler chickens. Sodium bentonite increases chilling shrink and reduces perirenal fat in rabbits (Lambertini *et al.*, 1989). Bentonite, like other clays, is able to adsorb much water and cations, appears to improve the physical nature of pelleted feeds (Martin *et al.* 1969) and may in some ways influence absorption from the gut (Mendel, 1971). Andronikashvili *et al.* (1994) found that, clinoptilolite (zeolite) added to poultry diet delayed transit time of digesta through the digestive tract by 2 to 2.5 h and promoted absorption of nutrients. Digestibility of organic matter, fat and nitrogen-free extract and also N-utilization were increased by supplementary zeolite. The objectives of this experiment were to study the effect of Tafla on the productive performance of broiler chicks.

MATERIALS AND METHODS

The present study was carried out at the Poultry Nutrition Research Section, Department of Poultry Production, Faculty of Agriculture, Ain Shams University, Cairo, Egypt. A total number of one hundred and fifty unsexed Hubbard broiler chicks at 7-days of age and 145 g average initial body weight assigned to five experimental groups. Chicks were randomly placed in wire cages for each treatment, 30 chicks each. The Tafla (natural clay) was brought from Egypt Company for Refractories, then crushed, ground until become dust and added to the diets. Egypt Company for Refractories mentioned that, the chemical composition of Tafla were Cao (2%), MgO (1%), Na₂O (2%), K₂O (1.2%), Fe₂O₃ (4%), SiO₂ (50%) and Al₂O₃ (22%). Chicks were fed starter diet for (0- 7days) without supplemented Tafla, then fed on the experimental diet. A starter diet was used during the starter period (7-28 days of age) which contained 22.03 % CP and 3105 kcal ME /kg diet, while the finisher diet contained 19.00 % CP and 3196 kcal ME /kg diet was used during the finisher period (29-49 days of age) (Table 1) and supplemented with 5 levels of Tafla (0,1,5,3,4.5 and 6%). Feed and water were available ad libitum. Chicks were grown in brooders with raised wire floors and exposed to 24 hours of constant light. All chicks were kept under the same managerial

hygienic and environmental conditions. Individual body weight was recorded at 7, 28 and 49 days of age. Live body weight, Body weight gain (BWG), feed intake (FI) (and feed conversion (FC) were also recorded during these periods. At 49 days of age, 4 birds from each treatment were used to determine the digestibility coefficients of nutrients. The proximate analysis of the experimental diet and excreta were carried out according to A.O.A.C (1990). At the end of the experiment, eight chicks (4 male and 4 female birds) from each experimental treatment group were slaughtered and carcasses. Liver, heart, gizzard and abdominal fat were weighed as a percentage of live body weight. Blood samples were taken for determination of, albumin, total protein, total lipids, cholesterol, glutamic oxaloacetic transaminase (GOT) and pyruvic transaminase activity (GPT) by kits from Bio Mereux. Finally, the economical efficiency (%) was calculated from the input-output analysis based upon the differences in body gain. The Economical efficiency (EE) was calculated according to the equation $EE = [(A-B)/B] \times 100$ where A is the selling cost of the obtained gain and B is the feeding cost of this gain. performance index was estimated according to North (1981): performance index = (live body weight/feed conversion) x100. The obtained data were statistically analyzed using the general linear model procedure described in SAS User's Guide (SAS Institute, 1995). Differences among means were tested using Duncan's multiple range test (Duncan, 1955).

Table(1). Composition of basal diets and Chemical analysis:

Ingredients	Starter	Finisher
Yellow corn	49.25	57.88
Soybean meal (44%)	40.85	32.52
Soy Oil	6.10	6.00
Limestone	1.40	1.30
Di-calcium Phosphate	1.60	1.60
NaCl	0.30	0.30
DL- Methionine	0.20	0.10
Vit & Min Premix*	0.30	0.30
Total	100	100
Price LE/ton	1137	1081
Determined analysis:		
Crude protein %	22.03	19.00
Crude Fiber %	4.07	3.65
Ether Extract %	8.00	8.16
Calculated analysis:		
ME KCal/Kg diet	3105	3196
Calcium %	1.01	0.91
Available phosphorous %	0.45	0.37
Lysine %	1.32	1.09
Methionine %	0.56	0.43
Meth. + Cys. %	0.92	0.74

* Each 3 Kg of vitamin and mineral mixture contain : vit. A, 12000,000 IU; vit. D3, 2000,000 IU ; Vit. E, 10,000 mg; Vit. K3 2000 mg; Vit B1 1000 mg; vit B2 5000 mg; Vit B6, 1500 mg; Vit. B12 10 Mg; Pantothenic acid, 10,000 mg; Nicotinic acid, 30,000 mg; Folic acid, 1000 mg; Bioten, 50 mg; Choline , 50,000 mg; Copper, 10,000-mg ; iodine, 1000 mg; Iron, 30,000, Mg; Manganese, 60,000 Mg; Zinc, 55,000 mg; Selenium, 100 mg and Cobalt, 100 mg.

RESULTS AND DISCUSSION

Effect of feeding different levels of Tafla on broiler performance:

Live body weight (LBW).

The effect of Tafla on growth performance of experimental birds was presented in (Table2). During 28 and 49 days, analysis of variance showed highly significant differences ($P<0.01$) between birds fed on the diet containing 1.5 and 3% Tafla and all other treatments. Analysis of variance showed highly significant differences ($P<0.01$) in the final LBW. Results indicated that birds fed 1.5 and 3% Tafla recorded the highest final LBW followed by those fed the 4.5% Tafla level and control group. Average BWG during the experimental period were significantly higher ($P<0.01$) for birds fed 3% Tafla than any of the other groups, followed by those fed 1.5, 4.5 and 6% Tafla levels. It seems that, 3% Tafla was the most efficient for broiler among the experimental levels. The reduction in the growth for birds given the diets containing high levels of Tafla (4.5 or 6%) may be due to the effect of high levels Tafla which like other clays, is able to adsorb much water and cations, appears to reduce the growth and may in some ways influence absorption from the gut.

Table(2).Growth performance of broiler chicks as affected by dietary Tafla.

Items	Treatments				
	Control	1.5%Tafla	3% Tafla	4.5% Tafla	6%Tafla
Live body weight (g)					
7days	145.17 ^a ±0.54	143.05 ^a ±0.55	145.45 ^a ±0.43	144.14 ^a ±0.49	145.14 ^a ±0.49
28days	1010.07 ^b ±28.32	1054.76 ^a ±17.78	1056.15 ^a ±17.35	1000.10 ^b ±24.01	940.10 ^c ±19.71
49days	1866.87 ^b ±20.47	1948.59 ^a ±21.43	2033.84 ^a ±16.63	1882.77 ^b ±18.72	1754 ^c ±16.72
Body weight gain(g):					
7-28days	864.90 ^b ±17.25	911.71 ^a ±9.90	910.70 ^a ±13.33	855.96 ^c ±13.90	794.96 ^c ±13.90
29-49days	856.80 ^b ±17.39	893.83 ^a ±18.70	977.69 ^a ±17.83	882.67 ^b ±19.76	814.60 ^c ±19.76
7-49days	1721.7c±20.47	1805.54 ^b ±21.45	1888.39 ^a ±16.69	1738.63 ^c ±16.75	1609.56 ^d ±18.74
Feed Intake (g / bird):-					
7-28 days	1158.33 ^b ±23.15	1117.82 ^b ±43.50	1156.59 ^b ±53.56	1241.14 ^a ±33.71	1157.67 ^b ±33.7
29-49days	2225.67 ^a ±93.78	2023.74 ^b ±92.33	2177.00 ^a ±92.39	2095.14 ^{ab} ±97.86	1970.00 ^c ±95.86
7-49days	3384.00 ^a ±76.80	3141.56 ^b ±93.40	3333.59 ^a ±93.74	3336.28 ^a ±98.85	3327.60 ^a ±98.85
Feed conversion ratio:					
7-28 days	1.34 ^b ±0.03	1.23 ^c ±0.03	1.27 ^c ±0.06	1.45 ^a ±0.03	1.46 ^a ±0.02
29-49days	2.57 ^a ±0.07	2.20 ^c ±0.10	2.23 ^c ±0.10	2.37 ^b ±0.13	2.40 ^b ±0.14
7-49days	1.97 ^a ±0.03	1.74 ^c ±0.06	1.76 ^c ±0.07	1.92 ^b ±0.08	2.07 ^a ±0.07
Performance Index:					
7-49days	87.36	103.74	107.30	90.47	77.75
Mortality No.-					
7-49days	3	2	2	2	3

Means within the same row with different superscripts are significantly different, ($P<0.01$). NS=Not Significant

Results obtained are in agreement with those reported by Ziobina (1990) who found that, daily weight gain of broiler chickens were improved with 3% zeolite added in the diet given for 49 days. Dobeic (1994) postulated that, clinoptilolite (zeolite) at levels 1,2 or 4% in the diet increased live weight gain and final body weight. The highest growth rate was with 2% clinoptilolite.

On the other hand, Ward *et al* (1993) found that growth rate decreased with addition of 0.75% sodium zeolite in chicks diet. Lemser *et al.* (1992) reported that, 1.5, 3 or 5% bentonite decreased body weight by 11.6, 14.5 and 12%, respectively, while zeolite supplements in broilers diet showed no effect.

Feed intake (FI).

During the starter period (7-28 days), total feed intake (FI) Table (2) recorded the highest value for the birds given 4.5% Tafla followed by those fed control diet, 6, 3 and 1.5 % Tafla levels. The differences in FI due to Tafla level effect were highly significant ($P<0.01$). For the overall period (7-49 days), analysis of variance showed highly significant differences ($P<0.01$) in FI among birds given different Tafla levels 1.5, 3, 4.5 and 6 % respectively. These results indicate that, the inclusion of Tafla to broiler diets as a feed additive decreased FI at level of 1.5% Tafla than control group and increased with more higher level (3, 4.5 and 6%).

Feed conversion ratio (FCR):

During the first experimental period (7-28days), It is clear that the improvement in the FCR was associated with increasing the dietary Tafla level up to 3%. These results were, statistically, highly significant ($P<0.01$). Similarly, the best FCR was corresponding to the 1.5 and 3% Tafla levels in the finisher period. Birds fed 1.5 or 3% Tafla converted their ration into BWG more efficiently than the other groups during the overall experimental period. Birds fed on diets containing high level of Tafla (4.5% or 6%) and control group recorded the poorest feed conversion ratios. These results were statistically, significant ($P<0.05$). Diets used in the present experiment were formulated to contain equal. Levels of nutrients, therefore the improvement in growth and feed efficiency for broilers given the diets containing low levels (1.5 or 3%) of Tafla would suggest an improvement in digestive efficiency as a result of Tafla addition. Tafla may be increased each of ion exchange capacity (Grim, 1968) digestible nutrients (EL-Hakim *et al* 1994).

These results are in agreement with those reported by Dobeic (1994) who found that, supplementation of the diet with clinoptilolite (zeolite) at 1, 2 or 4% levels improved feed conversion efficiency. The best feed conversion was with 2% clinoptilolite. Ziobina (1990), Elliot and Edwards (1991) and Ward *et al.* (1993) also found that, natural zeolite significantly improved feed efficiency. The diets used in this study did not contain a growth promoter. This was done intentionally because of the potential beneficial effect of Tafla on gastrointestinal trace macro-organisms. It was hypothesized that a more positive effect would be seen without growth promoters. A positive effect was seen and the performance of the flock was excellent at 3.0% of Tafla, However it is not possible to determine if Tafla was acting as a growth promoter or could replace a growth promoter in the diet.

From the previous results, it could be concluded that dietary Tafla 1.5 up to 3% of the diet was shown to improve the growth, feed efficiency and performance index of chickens while numerically reduced mortality.

Nutrients Digestibility Coefficients:

The effect of Tafla on the digestion coefficient of nutrients are

presented in Table (3).It is worthy to note that digestibility of CP, EE and NFE were improved and surpassed that of the control by 1.5 or 3% Tafla. Next are the data obtained by control group while the lowest values nutrient digestibility were obtained by chicks having 4.5 or 6% Tafla .The overall digestibility results of the three nutrient were reflected as a results with organic matter digestibility The highest value being 79.7and 79.9 was for 1.5 or 3% Tafla supplemented groups. While the lowest value was recorded for the chick received diets supplemented with 4.5 or 6% Tafla .The remarkable increase in nutrient digestibility by low levels (1.5 to 3%) of Tafla may be due to its prevention effect of the mold growth so can reduce the bioavailability of mycotoxines (Resanovic et al, 1999) Khosravi and Modirsanei (1999) and accordingly led to higher utilization efficiency of nutrients in the feed. Andronikashvili et al. (1994) found that, (zeolite) increased the digestibility of organic matter, fat and nitrogen-free extract and also N utilization.

Table(3). Digestion coefficients as affected by dietary Tafla supplementation.

Items	Treatments				
	Control	1.5%Tafla	3% Tafla	4.5% Tafla	6%Tafla
Digestibility Coefficients (%):-					
DM	78.40 ^b ±0.06	80.70 ^a ±1.22	81.30 ^a ±1.20	78.70 ^{bc} ±0.70	77.30 ^c ±0.24
OM	77.80 ^b ±0.81	79.70 ^a ±1.16	79.9 ^a ±1.03	77.10 ^{bc} ±0.92	74.90 ^c ±0.01
EE	79.50 ^b ±0.61	84.00 ^a ±3.28	84.20 ^a ±0.95	77.60 ^c ±0.45	73.10 ^c ±1.28
CF	29.90 ^b ±5.28	25.10 ^a ±1.06	27.80 ^a ±1.00	27.60 ^a ±5.75	25.10 ^a ±1.00
CP	92.50 ^b ±0.04	94.20 ^a ±0.02	95.4 ^a ±1.10	92.90 ^b ±0.20	92.30 ^b ±0.02
NFE	94.90 ^b ±0.27	95.00 ^a ±0.06	95.40 ^a ±0.05	94.00 ^b ±1.17	94.60 ^b ±0.04

Means on the same row differently superscripted are significantly different. * (P<0.05)
NS=Not Significant

Carcass Traits :-

The data in Table (4) show that, dressing percentages as well as total edible parts were affected by dietary treatments., Dressing percentages and total edible parts significantly (P<0.01) higher in broiler chicks fed on 1.5,3 and 4.5% Tafla diet when compared with those fed on the 6%tafla and control group.

Table(4). Carcass traits of broiler chicks as affected by dietary Tafla supplementation.

Traits	Treatments				
	Control	1.5%Tafla	3% Tafla	4.5% Tafla	6%Tafla
Dressing %	70.11 ^b ±0.80	73.54 ^a ±2.06	73.51 ^a ±0.36	73.67 ^a ±0.82	71.22 ^b ±104
Liver %	2.70 ^a ±0.16	2.75 ^a ±0.61	2.71 ^a ±0.11	2.68 ^a ±0.08	2.65 ^a ±0.05
Heart %	0.50 ^a ±0.03	0.53 ^a ±0.01	0.50 ^a ±0.03	0.48 ^a ±0.01	0.49 ^a ±0.02
Gizzard %	1.63 ^a ±0.12	1.60 ^a ±0.06	1.60 ^a ±0.06	1.59 ^a ±0.07	1.62 ^a ±0.12
Total giblets %	4.03 ^a ±0.19	4.37 ^a ±0.11	4.22 ^a ±0.18	4.27 ^a ±0.08	4.25 ^a ±0.16
Total edible parts %	74.14 ^b ±0.82	77.91 ^a ±1.81	77.73 ^a ±0.51	77.94 ^a ±1.05	75.80 ^a ±1.16
Total inedible parts %	25.86 ^a ±0.39	22.09 ^b ±0.69	22.27 ^b ±0.34	22.60 ^b ±0.82	24.20 ^a ±1.16
Abdominal fat %	0.92 ^a ±0.26	0.84±0.28	0.82 ^a ±0.22	0.81 ^a ±0.20	0.80 ^a ±0.10
Gizzard fat %	0.37 ^a ±0.09	0.36 ^a ±0.06	0.35 ^a ±0.08	0.32 ^a ±0.07	0.30 ^a ±0.03
Heart Fat %	0.09 ^a ±0.01	0.08 ^a ±0.06	0.08 ^a ±0.01	0.06 ^a ±0.02	0.06 ^a ±0.01
Non carcass fat %	1.38 ^a ±0.30	1.28 ^a ±0.38	1.25 ^a ±0.26	1.19 ^a ±0.26	1.16 ^a ±0.17

Means on the same row differently superscripted are significantly different.

* (P<0.05)

NS=Nct Significant

Internal organs were not affected by dietary treatments. Although there were no significant differences among dietary treatments for abdominal fat and non-carcass fat, there were the numeric variations in treatments and reduced values by increasing Tafla levels. The improvements in carcass traits are in agreement with those obtained by Yamani *et al*, (1994) reported that, the improved the percentages of carcass traits over the control group by using Tafla in broiler diets. The percentage of liver and abdominal fat did not show any significant differences among the experimental group by the addition of Tafla .Lon *et al* (1993) reported that, Natural Cuban zeolites in broilers diets increased carcass yield from 62.9 to 64.8%, with lower abdominal fat content (2.13 and 1.69%).

In general, dressing % and total edible parts % significantly increased in experimental groups fed diets supplemented with Tafla up to 4.5%.

Blood Parameters: -

Effect of dietary treatments on blood parameters is shown in Table(5). Total protein, albumin, A/G ratio and globulin levels in all treatment groups (up to 4.5% Tafla) were significantly higher than those of control group. However, broiler chicks fed diet supplemented with 6% Tafla had the lowest ($P < 0.05$) level of these parameters than those fed anther levels. These results are in agreement with those obtained by Yang *et al* (1999) who reported that, Zeolite significantly increased contents of serum total protein.

Table(5). Blood constituents of broiler chicks as affected by dietary Tafla supplementation.

Traits	Treatments				
	Control	1.5% Tafla	3% Tafla	4.5% Tafla	6% Tafla
Total protein (g/dL)	3.60 ^b ±0.09	4.58 ^a ±0.19	4.95 ^a ±0.26	4.98 ^a ±0.18	3.55 ^b ±0.10
Albumin (g/dL)	1.41 ^b ±0.06	1.60 ^a ±0.08	1.58 ^a ±0.07	1.53 ^a ±0.07	1.39 ^b ±0.06
Globulin (g/dL)	2.19 ^b ±0.07	2.98 ^a ±0.07	3.37 ^a ±0.20	3.45 ^a ±0.18	2.16 ^b ±0.08
A/G ratio	0.64 ^a ±0.04	0.59 ^b ±0.03	0.47 ^b ±0.03	0.44 ^b ±0.03	0.64 ^a ±0.05
GOT (U/L)	20.40 ^a ±1.06	18.02 ^a ±1.98	20.65 ^a ±3.27	19.27 ^a ±2.47	19.25 ^a ±3.45
GPT (U/L)	14.22 ^a ±2.30	14.86 ^a ±2.05	15.40 ^a ±2.20	15.55 ^a ±2.83	14.96 ^a ±2.07
Total lipids (mg/dL)	220.80 ^b ±12.90	175.14 ^b ±14.77	170.33 ^b ±13.70	165.12 ^b ±12.30	160.00 ^b ±2.50
Cholesterol (mg/dL)	125.00 ^a ±9.30	95.33 ^b ±9.26	92.13 ^b ±8.36	91.12 ^b ±6.30	90.25 ^b ±5.46

Means within same row with different superscripts are significantly different.($P < 0.05$).

With respect to GPT and GOT, chicks fed diets 1.5, 3 and 4.5 recorded significantly higher ($P < 0.05$) levels of GPT and GOT than the control group or those fed diet contained 6% Tafla. This slightly changes in the blood transaminases level may depend on the rate of protein metabolism, which may be a function of birds age rather than any other factor. It is well known that, by the simple process of transamination, an amino radical is transferred to alfa- keto acid, while the keto oxygen is transferred to the donor of the amino radical which is promoted by transaminases. Analysis of variance showed non significant differences among treatment applied in GOT and GPT which may be an indicator for an optimum hepatic function (Guyton,1981). Cholesterol and total lipids were significantly reduced by Tafla supplementation. The control group had significantly ($P < 0.05$) higher levels of Cholesterol and total lipids as compared to all other experimental groups.

In general, data shown in Table (5) indicated that, Tafla supplementation is more efficient as hypo-cholesterolemic and hypo-lipemic agent when compared with control group. may be due to that addition of Tafla reduced body fat, which may be reflected as a decreased level in the blood. Moreover, this observed decrease in blood total lipids and cholesterol levels with Tafla applied as feed additives may be attributed to the decrease that may occur in the rate of total lipids and cholesterol absorption through the intestinal villi that may be reflected as a decrease in their level in the blood EL-Gendi, 1996). From data obtained on the previous blood parameters it could be concluded that addition of Tafla to broiler' chicks diet had no adverse effects on blood components, as well as it had no deleterious effects on liver (as measured by GPT and GOT levels).

Economic efficiency (EE):

The higher economic efficiency (EE) observed in the groups fed 1.5 and 3% (Tafla) may be attributed mainly to the heavier weight gain of these groups Table(6). Diets containing 4.5%, 6% and 0% (control group) Tafla gave lower (EE) compared with 1.5 and 3% (Tafla). These results are in agreement with those reported by Fisinin *et al.* (1985) and Karelina (1985) who found that, zeolite increased the financial returns in broilers. Berrios *et al.* (1983) clarified that, consumption of protein and ME were low and cost of diet was reduced with zeolite dietary supplementation. Kasumov *et al.* (1986) indicated that addition of clinoptilolite to the diet of laying hens was economically advantageous. Generally the addition of (Tafla) improved (EE) up to 3% level. It seems that the addition of Tafla up to 3% as a cheap feed additive to commercial growing broiler diets improved economical efficiency. In conclusion, the obtained results indicate that clay feeding at a rate of 3% could serve as an effective, for increasing performance in the same time without any detrimental effects on birds health.

Table (6). Economic efficiency of the experimental dietary Tafla treatments.

Items	Treatments				
	Control	1.5% Tafla	3% Tafla	4.5% Tafla	6% Tafla
Total gain(g)	1721.09	1805.54	1888.39	1738.63	1609.63
Price/kg feed(P.T)	111	112	114	115	117
Total cost/ Feed intake(L.E)	3.76	3.30	3.48	3.83	3.89
Total revenue/bird (L.E)	9.47	9.93	10.38	9.56	8.85
Net revenue/bird (L.E)	5.71	6.63	6.9	5.73	4.96
Economical efficiency (EE%)	152	201	198	150	128
Relative economic efficiency (%)	100	132.20	130.30	98.70	84.20

1-The selling price of each Kg. Live body weight (5.5 LE).

2-Assuming that the relative economic efficiency of the control equals 100.

REFERENCES

Andronikashvili T.G., B. S. Tsereteli; V. K. Dolidze and N. G. Iremashvili (1994). Zeolite supplements in diets for birds. *Zootekhnnya*, 5: 17-18.
 A.O.A.C. (1990). Association of Official Analytical Chemists. *Official methods of Analysis*. 15th Edition, Washington, D.C.

- Bogdanov, G. A. and Abd El-Rahim (1997). Milk production performance and blood constituents in lactating cows fed rations supplemented with clinoptilolite or humolite during summer and winter seasons. *International Conference on Animal Production and Health*. Cairo, Egypt. Pp, 237-246.
- Berrios I., M. Castro and M. Cardenas (1983). Zeolite inclusion in feeds for laying hens fed *ad libitum*. *Cuban Journal of Agricultural Science*, 17(2): 169-174.
- Dobeic M. (1994). Influence of clinoptilolites on chicken growth. *Veterinarska Stanica*, 25(6): 323-332.
- Duncan, D. B. (1955) Multiple range and multiple test F test *Biometrics* 11-42.
- EL-Gendi, G. M. (1996). Effect of feeding dietary herbal feed additives on productive and metabolic responses broiler chicks. *Egypt. Poult. Sci.*, 16(11) 395-412.
- El-Hakim, E. M.; H. M. EL-Gendi ; Abdel- Raouf.; A. M. Allim and M. K Mohasan (1994). Effect of various levels bentonite on availability in sheep diets containing high levels of urea. *Egypt. Journal of Applied Sci.*, 3 (2) : 188-195.
- Elliot, M. A. and H. M. Edwards (1991) .Comparison of the effects of synthetic and natural zeolite on laying hen. *Journal of Poultry Science*, 10 (10): 2115-2130.
- Fisinin V., Ageev V. Sintserova O., Lenkova T. and Kvashali N. (1985). Zeolites in diets for poultry. *Ptitsvodstvo*, 9: 25-26.
- Fitzpatrick, E. A. (1983). Soils, their formation. Classification and distribution. Published by Longman Inc, New York, USA.
- Grim, R. E. (1968). Clay mineralogy - Mc -Grow, Hill Book Co., New York.
- Guyton, A. C. (1981) .Text Book 1-Medical of Physiology W.B.Saunders Company (Sixth edition), London.
- Kalyuzhnov, V. T.; I. E. Ziobina and L. G. Nikulina (1988). physiological basis of including zeolite in diets of poultry. *Spolzovanie Tseolitov Sibiri i Dalnego Vostokav Seiskom khozyaistve*, Pp 15-20. Novosibirsk.
- Karelina, O. (1985). Zeolites for feeding broilers. *Ptitsvodstvo*, 9 :26.
- Kasumov, S.N.; A. G. Seidov and Mugalinskii Sh.. N. (1986). Clinoptilolite tuff - a mineral supplement in diets for laying hens. *Aktualnye Voprosy Razvitiya Krupnogo Rogatogo Skota V Azerbajjan*, Pp 59-62. Baku, Azerbaijan .
- Khosravi-AR; Modirsanei-M (1999). Comparison between several procedures for reducing the effects of aflatoxin on broiler chick performance. *Journal-of-the-Faculty-of-Veterinary-Medicine -University-of-Tehran*. 1999, 54: 2, 59-66.
- Lambertini L.; G. Zaghini; S. Galassi and M. C. Benassi (1989). Sodium bentonite in growing rabbit feed: animal performance and effects on the digestive tract. *Rivista di Coniglicoltura.*, 26 (3): 51-55.

- Lemser A.; WI. Ochrimenko and G. Richter (1992). Ion exchangers in rye-based broiler diets and N losses in excrement's of goats and poultry mixed with bentonite. Kongressband 1992 Gottingen. Vortrage zum Generalthema des 104, Gottingen; Okologische Aspekte Extensiver Landbewirtschaftung, 801-804
- Lon-Wo, E; V. Zaldivar; E. Margolles (1993). Effect of natural zeolites on poultry feeding with different nutritional levels or high mycotoxin contamination. Cuban-Journal-of-Agricultural-Science. 1993, 27: 2, 199-204.
- Martin L.C.; A.J. Clifford and A. D. Tillman (1969). Studies on sodium bentonite in ruminant diets containing urea. Journal of Animal Science, 29 (5): 777-782.
- Mendel V. E.. (1971). Montmorillonite clay in feed lot rations. Journal of Animal Science, 33(4): 891-894.
- North, M.O. (1981). Commercial chicken production Manual. Avi, Publishing Company. I. N. C. Westport Connecticut, USA.
- SAS (1995). SAS User's Guide: Statistical Analysis System Institute, Inc., Cary, N.C.
- Resanovic, R; T; Z. Palic; Nikolovski-Stefanovic; Z. Veljic; J-Rodic (1999). In vivo detoxication of aflatoxin B1 using modified clinoptilolite. Eighth Yugoslav Symposium on Poultry Production. Proceedings II. Sokobanja, Yugoslavia, 5-9 October 1999. Zivinarstvo. 1999, 34: 8-9, 21-24.
- Ward T.L.; K. L. Watkins and L. L. Southern (1993). Interactive effect of sodium zeolite A and Eimeria acervulina infection on growth and tissue minerals in chicks. Journal of Poultry Science, 12(11) 2172-2175.
- Yang-CaiMei; Chen-AnGuo; Yang-CM; Chen-AG (1999). Effects of dietary zeolite on growth performance and the mechanism in yellow-feather type broilers. Journal-of-Zhejiang-University-Agriculture-and-Life-Sciences., 1999, 25: 6, 619-622.
- Yamani, K.A.O. Rashwan A.A.and Magady, M.M.(1997). Effect of copper, zinc and tafia dietary supplementation on broiler performance International conference on Animal production and health, zagazig, Egypt.457-483.
- Ziobina, E. (1990) Retention of trace elements from zeolites in birds and the effect of Pegasin on the meat quality of broiler chickens. Sibirskoe Otdelenie, Sibirskii Nauchno Issledovatel'skii Proektno tekhnologicheskii institute Zhivotnovodstva 2:28-30

تأثير استخدام الطفلة المصرية كإضافات غذائية علي الأداء الإنتاجي لدجاج اللحم
أحمد حسين عبد المجيد- علاء الدين عبد السلام حميد* - فتحى عبد العظيم محمد*
معهد بحوث الإنتاج الحيواني- مركز البحوث الزراعية - وزارة الزراعة - الدقي - الجيزة
*قسم إنتاج الدواجن - كلية الزراعة - جامعة عين شمس- القاهرة- مصر

تهدف هذه التجربة إلي دراسة تأثير استخدام مستويات مختلفة من الطفلة المحلية على أداء دجاج التسمين واستخدم في هذه التجربة ١٥٠ ككتوت هابرد وقسمت إلى خمسة مجاميع تجريبية في كل مجموعة ٣٠ ككتوت (٣ مكررات- ١٠ ككتوت في كل مكرر) وتم تغذية الطيور على عليقه كتنترول في الأسبوع الأول وبدأت التغذية على المعاملات في عمر ٧ أيام وكانت المعاملات صفر و ١,٥ و ٣ و ٤,٥ و ٦% طفلة مضافة إلى العليقة الأساسية والتي كانت تحتوى على ٢٢% بروتين و ٣١,٥ كيلو كالورى خلال فترة البادي من (٧-٢٨ يوم) وكانت عليقه الناهي تحتوى على ١٩% بروتين و ٣١٩٦ كيلو كالورى من عمر (٢٩-٤٩ يوم) وتم تقدير المستهلك من العلف والزيادة في الوزن والكفاءة الغذائية كما تم إجراء تجربة ذبح لوزن أجزاء الذبيحة ووزن دهن الجسم والأجزاء المأكولة و أخذت عينات للدهن لتقدير البروتين الكلى واللييدات الكلية والكولستيرول والبيومين و GOT GPT كما تم إجراء تجربة هضم في نهاية التجربة لتقدير معاملات هضم البروتين والدهن والمادة الجافة والألياف وتم حساب التكلفة الاقتصادية وكانت نتائج التجربة كالتالي :-

- ١- يزيد وزن الجسم وتحسن الكفاءة الاقتصادية معنويا بإضافة الطفلة بمستوى ٣% بالمقارنة مع باقى المجموعات التجريبية.
- ٢- تتحسن الكفاءة الغذائية بإضافة الطفلة بمستوى ٣% بالمقارنة بالمجموعات الأخرى.
- ٣- كانت هناك زيادة معنوية في معاملات الهضم للبروتين والألياف والدهن والمادة الجافة والعضوية والمستخلص الخالى من الأزوت بإضافة الطفلة بمستوى ١,٥ و ٣%.
- ٤- ينقص مستوي الكولستيرول واللييدات الكلية الدم وكان ذلك مر تبطاً بانخفاض نسبة الدهن في الذبيحة بوجه عام.
- ٤- يزيد وزن الذبيحة ومجموع الأجزاء المأكولة منها معنويا بإضافة الطفلة بمستوى ١,٥ و ٣% وبوجه عام توضح نتائج التجربة انه يمكن استخدام الطفلة كإضافات غذائية في علائق التسمين بمستوى ١,٥ و ٣% مما يزيد الأداء الإنتاجي والكفاءة الاقتصادية وتقليل النسبة المئوية للدهن في الذبيحة ولييدات الدم دون حدوث تأثيرات سيئة على مكونات الدم.