

UTILIZATION OF OLIVE PULP BY-PRODUCTS AS AN UNCONVENTIONAL FEEDSTUFF IN DIETS OF NILE TILAPIA, *Oreochromis niloticus*, FINGERLINGS

Abd Elmonem, Asmaa I.

Department of Animal Production, Faculty of Agriculture, Alexandria University, El-Shatby, Alexandria, Egypt.

ABSTRACT

A feeding experiment was conducted to evaluate the effect of olive pulp at different levels on growth performance, feed and nutrients utilization, body composition, plasma glucose, total plasma lipids (TPL) and total plasma proteins (TPP), liver enzymes and cost-benefit analysis of Nile tilapia fingerlings. Ten glass aquaria with the dimensions of 70x30x40 cm were used to stock 10 fish/aquarium averaging 9 g/fish. five experimental diets were formulated to contain 0, 5, 10, 15 and 20% of Olive pulp instead of wheat milling by product. All the diets were isonitrogenous and isoenergetics. Fish fed 3-5% of body weight daily. The feed amount was given three times daily, six days a week for 77 days. Fish were weighed biweekly and feed amounts were adjusted on the basis of the new fish weight.

Results indicated that no significant differences ($P>0.05$) were observed in growth performance among all fish groups fed varied levels of olive pulp. Yet, there was a tendency of Nile tilapia fed 5% olive pulp to increase their body weight, gain and SGR. Lower amounts of feed intake were obtained when fish fed diets had 15 and 20% olive pulp. No significant differences ($P>0.05$) were observed in FCR, PER, FFEV% and ER% among all fish groups fed diets containing varied levels of olive pulp. However, carcass fat content and plasma glucose and plasma total lipids were increased by increasing levels of olive pulp. In contrast and total plasma protein values were decreased. Economic analysis showed lower incidence cost and higher profit index by increasing olive pulp levels in Nile tilapia diets.

Keywords: Nile tilapia, Olive pulp, Growth, Cost-benefit analysis

INTRODUCTION

The importance of developing cheap nutritionally balanced artificial diets for fish has been increased dramatically (Osman *et al.*, 2003). Incorporation of cheap untraditional feedstuffs, particularly energy sources in animal and fish diets may participate in solving the problem of feed shortage and alleviate the pollution problems. Olive pulp is one of the most important agro-industrial by-products that may be serve as unconventional energy sources. Olive pulp as energy source (Khamis *et al.*, 1989; Mohamed, *et al.*, 1997 and El-Aiatt, 1999; Mousa, 2000 and 2001 and Mousa and Abdel Samee, 2002). More than 95% of the estimated world's olive crops are grown in the Mediterranean area. Annual world's olive production amounts to 804 million tones, which gave rise to three million tones crude olive cake after industrial processing (Abdel-Naby, 1998). Khamis *et al.* (1989) reported that the annual available olive pulp and date seeds in Egypt were estimated to be about 7600 tones giving about 5935 tons of dry matter and about 3433 tons of total digestible nutrients.

Publications on olive pulp used in fish feeding are very little, except El-Aiatt (1999) who found that weight gain of hybrid tilapia fed 30% olive

waste diet was lower than those fed the control diet. The present study, therefore, was carried out to investigate the effect of different levels of olive pulp on growth performance, feed conversion, protein and energy efficiencies, body composition, blood analysis and cost-benefit analysis of Nile tilapia, *O. niloticus*, fingerlings.

MATERIALS AND METHODS

Experimental facilities:-

This work was carried out at the Wet Lab., Animal Production Department, Faculty of Agriculture, Alexandria University, Alexandria, Egypt. Ten glass aquaria with dimensions of 70x 30x 40 cm, 70-L total net volume, were used in this study. All aquaria were filled with dechlorinated tap water stored 24 hours before use. Fingerlings of Nile tilapia, *O. niloticus* were obtained from the Maryot Fish Farming Co, Alexandria Governorate, Egypt. Fish were transported in fiberglass tanks (300 L). After arrival, all fish were kept for one week to alleviate stresses and to be adapted to the new conditions. Ten fish in the same initial weight (9 g/fish) were selected and randomly allotted into each experimental aquarium. Fish were fed the control diet for two weeks, during this period healthy fish of the same weight replaced dead one.

Experimental diets:-

Five experimental diets were formulated to contain 0, 5, 10, 15 and 20% of olive pulp instead of wheat milling by-product. Each diet was fed to two randomly assigned (duplicate) aquariums. Ingredients composition of the experimental diets is presented in Table (1). Fishmeal was home made by collecting small fish and non-saled fish named locally "Wazafa" from fish market to dry in oven at 60 °C to dryness. Then dry fish were grounded and sieved prior to keep at -20 °C until use as an animal protein source. Soybean meal 48% was used as a plant protein source. While yellow corn, wheat bran and wheat milling by-product were served as energy sources. The experimental diets were isocaloric by using sunflower oil in different levels depended on the energy content in the rest ingredients.

The experimental diets were prepared by mixing dry ingredients with water and were pelleted using a meat mincer with a 1-mm holes diameter. The pellets were air dried and stored at -20° C until use. Fish fed 5, 4 and 3 % of the body weight daily for the 1-4, 5-8 and 9-11th week, respectively. The feed amount was given at three times a day (9 00, 1200 and 1500 hrs) in equal proportions. Feeding was performed for six consecutive days a week, for 77days. Fish were weighed biweekly and feed amounts were adjusted on the basis of the new weight of fish.

Environmental conditions:-

Fish were reared at 28±2 °C and 8.6±0.1 pH. Third water volume of all aquaria was exchanged daily, except the weighing day where about two thirds of water volume were changed.

Analytical methods:-

Diets and fish samples were analyzed according to AOAC (1990). The gross energy contents of the experimental diets and fish samples were calculated by using factors of 5.65, 9.45 and 4.2 kcal/g of protein, lipid and carbohydrate, respectively (NRC, 1993). Digestible energy content was calculated from standard physiological fuel values as 4 and 9 kcal/g of protein and carbohydrate and lipid, respectively (Garling and Wilson, 1976).

Table 1: Ingredients composition of the experimental diets

| Ingredients | Diets No. | | | | |
|---------------------------|-----------|-------|-------|-------|-------|
| | 1 | 2 | 3 | 4 | 5 |
| Fish meal | 15 | 15 | 15 | 15 | 15 |
| Soybean meal | 34 | 33 | 32 | 31 | 30 |
| Yellow corn | 5 | 5 | 5 | 5 | 5 |
| Olive pulp meal | 0 | 5 | 10 | 15 | 20 |
| Wheat milling by-product | 34.5 | 30.5 | 26.5 | 22.5 | 18.5 |
| Wheat bran | 5 | 5 | 5 | 5 | 5 |
| Sunflower oil | 4.5 | 3.5 | 2.5 | 1.5 | 0.5 |
| Vitamin mix. ¹ | 0.875 | 0.875 | 0.875 | 0.875 | 0.875 |
| Ascorbic acid | 0.125 | 0.125 | 0.125 | 0.125 | 0.125 |
| Mineral mix. ² | 1.0 | 1.0 | 1.0 | 1.0 | 1.0 |
| Total | 100 | 100 | 100 | 100 | 100 |

1. Vitamin mixture (g/100 g) was 960000 IU, 160000 IU, 0.8 g, 80 mg, 0.32 g, 0.12 g, 0.8 g., 0.8 mg 1.6 g, 80 mg, 4 mg, 40 g. of vitamin A, D3, E, K, B1, B2, B6, pantothenic acid, B12, niacin, biotin, choline chloride, respectively.

2. Mineral mixture (g/100g) was 12.75, 72.85, 0.55, 0.25, 0.02, 5, 2.5, 0.08, 0.05, 0.01 and 6 mg of $MgSO_4 \cdot 7H_2O$, $CaHPO_4 \cdot 2H_2O$, $ZnSO_4 \cdot 7H_2O$, $MnSO_4 \cdot 4H_2O$, $CaI_2 \cdot O_4 \cdot 6H_2O$, KCl, $FeSO_4 \cdot 7H_2O$, $CuSO_4 \cdot 5H_2O$, $CoSO_4 \cdot 7H_2O$, $CrCl_3 \cdot 6H_2O$ and NaCl, respectively.

Blood analysis:-

Blood was collected using heparinized syringes from the caudal vein of all the experimental fish at the termination of the experiment. Blood samples were centrifuged at 3000 rpm for 15 minutes to allow separation of plasma which was subjected to determine plasma total protein (Armstrong and Carr, 1969), total lipid (Frings *et al.*, 1972) and glucose (Trinder, 1969). The activities of aspartate aminotransferase (AST) and alanine aminotransferase (ALT) were determined colorimetrically by using commercial kits (Quimica Clinica Aplicada, S.A., Aspain) according to the method of Reitman and Frankel (1957).

Cost-benefit analysis:-

To evaluate cost-benefit analysis of olive pulp inclusion in the diets of Nile tilapia fingerlings, incidence cost (IC) and profit index (PI) were calculated according to New (1985).

Statistical analysis:-

Analysis of variance (ANOVA) was carried out according to Snedecor and Cochran (1982) using a completely randomized design (CRD). Differences were subjected to Duncan's Multiple Range-Test (Duncan, 1955) at a significance level of 0.05. Program of MSTAT-C was used to compute the statistical analysis (MSTAT-C, 1988).

RESULTS AND DISCUSSION

Nutritive value of olive pulp:-

Data presented in Table (2) show that olive pulp used in the present study had a moderate percentages of crude protein (11.56%) and crude fiber (16.53%) with higher percentages of ether extract (21.29%) and nitrogen free extract (38.02%) while ash was 2.6% on dry matter basis. The present results are in agreement with (Abasa *et al.* 1987 and Al-Shanti, 2003) except the ether extract that was higher than those obtained by the previous studies. This may be due to differences of oil extraction methods. The percentage of olive pulp differs according to the variety and degree of maturity of the fruit (Abdel Gabber, 1979), oil extraction procedures (Al-Shanti, 2003), the presence or absence of pis and elapsed time and treatment of pulp between pressing and processing (Belidasakis; 1984). In addition some varieties of olive fruit had higher oil content ranged from 43.5 to 50.82% (Kirtsakis, 1998 and Azza, 2002) such these varieties wasted fatty pulp in case of bad oil extraction.

Table 2: Chemical analysis of the experimental diets.

| Item | Olive pulp | Diet No. | | | | |
|-------------------------------------|------------|----------|-------|-------|-------|-------|
| | | 1 | 2 | 3 | 4 | 5 |
| Dry matter % | 94.68 | 94.41 | 94.1 | 93.99 | 94.12 | 94.23 |
| % on the DM basis:- | | | | | | |
| Crude protein , CP | 11.56 | 30.48 | 30.56 | 30.64 | 30.72 | 30.81 |
| Ether extract | 21.29 | 9.20 | 9.72 | 10.23 | 10.75 | 11.26 |
| Crude fiber | 16.53 | 2.90 | 3.69 | 4.49 | 5.29 | 6.09 |
| NFE | 48.02 | 46.6 | 45.34 | 44.07 | 42.79 | 41.52 |
| Ash | 2.60 | 10.82 | 10.69 | 10.57 | 10.45 | 10.32 |
| Calculated energy (kcal/100 g diet) | | | | | | |
| Gross energy , GE | - | 456 | 456 | 456 | 456 | 456 |
| Digestible energy, DE | - | 391 | 391 | 391 | 391 | 391 |
| P/E ratio mg CP/kcal DE | -- | 77.95 | 78.16 | 78.36 | 78.57 | 78.8 |

Growth performance and surviving:-

Survival rate of all fish groups fed varied levels of olive pulp was 100%. Data presented in Table (3) show the growth performance and feed and nutrient utilization of fish fed different levels of olive pulp. Results confirm a great tendency of Nile tilapia fed 5% olive pulp to increase body weight, gain and SGR. However, statistical analysis showed that no significant differences ($P>0.05$) among all fish groups fed varied levels of olive pulp. Results in the present study agree to a satisfactory extent with those of Tortuero *et al.* (1989); Morani (1990); El-Kerdawy (1997); Mousa and Abdel-Samee (2002) and Mousa and Shetawi (2002). In poultry feeding, Al-Shanti (2003) found that chicks fed diet contained 5% of either extruded full-fat soybean or olive cake had significantly higher weight gain values and economic efficiency. While in fish feeding, El-Aiat (1999) used higher levels of olive pulp in tilapia diets. He found that percentage of gain in weight decreased by the replacement of wheat flour in diets by 30% olive waste to about 62% than the control diet. The present work confirmed the above findings.

Table 3: Growth, survival rate, feed and nutrient efficiencies utilization of Nile tilapia fingerlings fed diets containing different levels of olive pulp.

| Item | | Diet No. | | | | | MSE ¹ |
|--------------------|--------|--------------------|--------------------|---------------------|--------------------|--------------------|------------------|
| | | 1 | 2 | 3 | 4 | 5 | |
| Initial weight | g/fish | 9.74 | 9.77 | 9.72 | 9.74 | 9.74 | 0.01 |
| Final weight | g/fish | 39.94 | 41.05 | 38.5 | 37.9 | 37.85 | 0.51 |
| Gain ² | g/fish | 30.2 | 31.29 | 28.78 | 28.16 | 28.12 | 0.51 |
| SGR ³ | % /day | 1.83 | 1.87 | 1.79 | 1.77 | 1.77 | 0.02 |
| Survival rate | % | 100 | 100 | 100 | 100 | 100 | 0.00 |
| Feed intake | g/fish | 58.9 ^{ab} | 59.82 ^a | 56.75 ^{bc} | 55.65 ^c | 54.25 ^c | 0.77 |
| FCR ⁴ | | 1.85 | 1.8 | 1.86 | 1.87 | 1.82 | 0.01 |
| PER ⁵ | | 1.79 | 1.82 | 1.76 | 1.75 | 1.79 | 0.01 |
| PPV ⁶ % | | 29.38 | 30.85 | 28.54 | 28.71 | 26.72 | 0.67 |
| ER ⁷ % | | 18.87 | 20.35 | 17.76 | 17.96 | 17.6 | 0.47 |

¹Values in the row having a common superscript letter are not significantly different (P>0.05).

1. Standard error of the means derived from the analysis of variance
2. Gain = (final weight - initial weight).
3. Specific growth rate = $100(\ln \text{ final weight} - \ln \text{ initial weight}) / \text{day}$.
4. Feed conversion ratio = DM intake/weight gain.
5. Protein efficiency ratio = weight gain/protein intake.
6. Productive protein value = $100(\text{protein gain} / \text{protein intake})$.
7. Energy retention % = $100 \times (\text{gross energy gain} / \text{gross energy intake})$.

Feed intake and feed utilization:-

The results of feed intake under the present study (Table, 3) indicated that fish fed diets containing 5% of olive pulp exhibited the highest feed intake. However, lower amounts of feed intake were obtained when fish fed diets had 15 and 20% olive pulp. These results are in agreement with those obtained by many researchers (Razzaque and El-Sheikh, 1982 and Youssef *et al.*, 2001). It was clearly shown that, no significant differences (P>0.05) were observed in feed conversion ratio, protein efficiency ratio, productive protein value and energy retention among all fish groups fed diets containing varied levels of olive pulp. Razzaque and El-Sheikh (1982) found that the use of olive cake did not show any significant effects on feed intake and FCR. This means that the animal fed the ration containing olive pulp consume normally due to a good palatability of olive pulp (Khamis *et al.*, 1989). However, Omar *et al.* (1995) observed that increasing olive pulp levels (0, 5, 10, 15 and 20%) impaired FCR, especially of lambs at the high levels of 15 and 20%.

Results in the present study indicated slightly improvement of feed and protein and energy utilization when fish fed 5-10 % olive pulp. It may be due to the nature of olive pulp oil which had an adequate amount of unsaturated fatty acid (HUFA), it consists mainly of about 96% HUFA (Zaza *et al.*, 2001). The mixture of unsaturated fatty acids has been determined as essential for growth (Murray *et al.*, 1991). Furthermore, Klatt (1986) stated that dietary unsaturated fatty acids are the subjects of current interest because they have been credited with number of beneficial effects.

Body composition:-

Data in Table (4) show the body composition of fish fed varied levels of olive pulp. No significant differences ($P>0.05$) were observed in crude protein and ash content. However, ether extract content of fish fed 5% olive pulp had higher values followed by fish fed the control diet than the other treatments. Similar trend was noticed with energy content. Also, fish at the start of the experiment had lower crude protein, ether extract, ash and energy contents, but they had higher moisture than at the end of the experiment. A significant of fat content of fish fed olive pulp may be due to its higher oil content, which affected deposition of lipid in fish tissue. Increasing dietary oil in fish diets resulted in a higher fat content in many species of fish (El-Dakar *et al.*, 1988 and Shalaby, 1998).

Table 4: Body composition (% fresh weight) of Nile tilapia fed different levels of olive pulp^{ns}.

| Diets | Initial fish | Olive pulp levels (%) | | | | | SEM |
|--------------------|--------------|-----------------------|-------|-------|-------|-------|------|
| | | 0 | 5 | 10 | 15 | 20 | |
| Moisture | 76.47 | 73.36 | 72.42 | 74.11 | 73.35 | 73.63 | 0.33 |
| Crude protein | 13.45 | 15.76 | 16.13 | 15.5 | 15.64 | 15.2 | 0.26 |
| Ether extract | 5.9 | 6.67 | 7.16 | 6.11 | 6.2 | 6.16 | 0.19 |
| Ash | 3.85 | 4.13 | 4.29 | 4.26 | 4.32 | 4.31 | 0.04 |
| Energy, Kcal/100 g | 99.67 | 152 | 159 | 145 | 147 | 144 | 3.64 |

ns means not significant

* Standard error of the means derived from the analysis of variance.

Blood parameters:-

Table (5) shows that plasma glucose, total plasma protein and total plasma lipid were affected by different levels of olive pulp. It was clear that increasing of olive pulp levels in tilapia diets resulted in increases in plasma glucose and total lipids. On the other hand, olive pulp gave a decrease in total plasma protein. Similar results were obtained by Mousa (2000) and Al-Shanti (2003). Aspartate aminotransferase (AST) showed significantly ($P>0.05$) higher activity in fish fed the control diet. However, it decreased dramatically as olive pulp increased in the experimental diets. At the same time, alanine aminotransferase (ALT) did not show significance in all fish groups. These results suggested that olive oil may influence through its essential oil to improve liver function. Mousa (2000) demonstrated that no significant differences were observed in adrenal, liver and kidney functions as well as serum total protein, albumin and globulin when lambs fed different levels of olive pulp. In addition, total lipid and cholesterol increased significantly ($P<0.05$) by 15 and 23% respectively in growing lambs fed 25 and 30% olive pulp.

Table 5: Blood parameters of Nile tilapia fed different levels of olive pulp.

| Item | Experimental Diets | | | | |
|-------------------------------|--------------------|-----------------------------|-------|-------|-------|
| | 1 | 2 | 3 | 4 | 5 |
| Plasma glucose mg/dl | 53.12 | 52.72 | 55.48 | 63.58 | 79.47 |
| Plasma total protein g/dl | 5.31 | 3.90 | 3.89 | 3.37 | 4.52 |
| Plasma total lipid g/dl | 3.79 | 5.53 | 6.30 | 6.17 | 6.76 |
| AST ¹ U/dL | 154 | 140 | 133 | 121 | 114 |
| ALT ² U/dL | 62 | 60 | 57 | 50 | 47 |
| 1- Aspartate aminotransferase | | 2- Alanine aminotransferase | | | |

Cost-benefit analysis:-

Cost-benefit analysis of fish fed olive pulp is presented in Table (6). Results indicated that feed cost per kg weight gain of fish (incidence cost) decreased by 96, 97, 96 and 91% of the control diet for fish fed diets containing 5, 10, 15, and 20% olive pulp, respectively. In contrary, change in profit index ranged between 3 and 10% more than those fed the control diet. Lower incidence cost and higher profit index was obtained at 20% olive pulp. Similar results were obtained by Mousa and Abdel-Samee (2002) and Abdel-Ghaffar (2002) who demonstrated that olive pulp replacement up to 20% of rabbit diets decreased feed cost per kg body weight gain by 18%, it increased considerably the economical efficiency.

Generally, it could be recommended to use safely olive pulp at 5-20% in fish feeding.

Table 6: Cost-benefit analysis of Nile tilapia fed different levels of olive pulp.

| Item | Diet No. | | | | |
|-----------------------------|----------|------|------|------|------|
| | 1 | 2 | 3 | 4 | 5 |
| Cost / kg feed | | | | | |
| Incidence cost ¹ | 4.60 | 4.43 | 4.48 | 4.41 | 4.21 |
| Change % | 100 | 96 | 97 | 96 | 91 |
| Profit index ² | 1.30 | 1.36 | 1.34 | 1.36 | 1.43 |
| Change % | 100 | 105 | 103 | 105 | 110 |

1- Incidence cost = feed cost to produce 1 kg fish

2- Profit index = value of fish /cost of feed consumed, 1 kg fresh fish equals 6 LE

Cost of 1 kg olive pulp was 0.10 LE.

Costs were as common commercial feeds in local markets (Prices in LE).

REFERENCES

- Abasa, M.N.: Nabawiya Hafiz, El-Torky, M.; Nour, A. M.; Borhami, B. E.; El-Shazly and Naga, M.A. (1987): Effect of sodium hydroxide or steam treatments on nutritive value of poor quality roughages in vitro. *Alex. J. of Agric. Res.*, 29 (3): 1178-1188.
- Abd El-Naby, M.R. (1998): Evaluation of olive cake as waste product of food industry in rabbits. M.Sc. Thesis, Fac. of Agric. Zagazig Univ., Egypt.
- Abdel-Gabbar, H. (1979) Chemical studies on olive oil, its cake and analogous oil M.Sc. Thesis, Fac. Of Agric. Cairo Univ., Egypt.

- Abdel-Ghaffar, A. (2002). Improving productivity of heat stressed babbits using genetic and nutritional techniques under Sinai conditions. M.Sc. Thesis, Fac. Environ. Agric. Sci., Suez Canal Uni.
- Al-Shanti, H. A. (2003): Effect of using olive cake or extruded full-fat soybean in broiler chicks diets. Egypt. Poult. Sci. Vol 23 (1): 1- 13.
- AOAC (1990). Official Methods Analysis of Association of Official Analytical Chemists. 15 th Ed .Published by the Association of Analytical Chemists. Virginia 2220/USA.
- Armstrong W.D. and Carr, C.W. (1969). Physiological Chemistry Laboratory Directions (3rd ed.). Purges publishing Co., Minn Capolis.
- Azza, A.A.A. (2002). Effect of extraction systems and processing technology on the quality of virgin oil oil. Ph.D. Thesis, Alexandria University, Alexandria, Egypt.
- Belidasakis, N. G. (1984): Olive cake pulp on the fattening of lambs. 2. Diets with a low proportion of olive cake pulp. Ellenike Kteninatrike Hellenic Veterinary Medicine, 28 (4) 222-230.
- Duncan, D.B. (1955). Multiple range and multiple F-test. Biometrics, 11:1-42.
- El-Aiatt, A.A.O. (1999). The use of non-conventional sources in diets of fish farms. M.Sc. Thesis, Fac. Environ. Agric. Sci., Suez Canal Uni.
- El-Dakar, A.Y.; Nour, A.M.; Omar, Eglal A. and El-Shazly, K.A. (1988). Fat utilization by common carp (*Cyprinus carpio* L.). The Proceedings of First Scientific Conf. on the Role of Scientific Research in the Development of Fish Resources. Alexandria University, Alexandria, 6-8 August 1988.
- El-Kerdawy, D. M.A. (1997): Olive cake as a new energy source for growing rabbits. Egyptian Journal of Rabbit Science, 7 (1): 1-12.
- Frings C.S.; Fendly W.; Dunn R.D. and Quen C.A. (1972). Improved determination of total serum lipid by the sulpho-phosph vaniline reaction. Clinical Chemistry, 18: 673-674.
- Garling, D.L. and Wilson R.P., (1976). Optimum dietary protein to energy ratio for channel catfish fingerlings, *Ictalurus punctatus*. J. Nutrition, 106: 1368- 1375.
- Khamis, H.S.; El-Shaer, H.M., Farid, M.F.A., Shalaby, A.S. and Salem, O.A. (1989): Utilization of date seeds and olive pulp as supplementary feed for lactating ewes in Sinai. Proc. of 3rd Egyptian-British Conf. on Animal, Fish and Poultry Production. Alexandria, 7-10 October:
- Kiritskis, A.K. (1998). Flavour components of olive oil, A review. J. Am. Oil Chem. Soc., 75 (7): 673-681.
- Klatt, L. (1986). The lure of omega-3- polyunsaturated fatty acids. Food Sci. News L. 16 -1-4 (Poultry Sci., 1991, 70: 1403-1411).
- Mohamed, M. I.; Awadalla, I. M.; Gihad, E. A. and El-Kady, R.I. (1997): Growing camel performance on different dietary regime. J. Agric. Mansoura Univ., 22(12): 4315- 4323.
- Morani, S. (1990): Full-fat Soya Handbook American Soybean Association, Brussels, Belgium.
- Mousa, M.R.M. (2000). Effect of feeding olive pulp on productive performance of Awassi ewes and their lambs under subtropical conditions. J. Agri. Sci. Monsoura University, 25 (8): 4967-4975.

- Mousa, M.R.M. (2001). Effect of fattening system of lambs under North Sinai conditions on some performance and metabolic parameters. J. Agri. Sci. Monsoura University, 26 (8): 715-726.
- Mousa, M. R. M. and Abdel-Samee, A.M. (2002): Effect of olive pulp on feeding on the growth performance and some related blood biochemical of growing rabbits under semi-arid conditions. Egyptian Journal of Rabbit Science, 1: 59- 68.
- Mousa, M.R.M. and Shetaewi, M..M. (2002): Milk yield and growth rate as influenced by feeding olive pulp and acacia shrubs for Awassi sheep under sime-arid conditions. Annals of Agric. Moshtohor ,40 (2): 843-856.
- M-STAT-C (1988). Rusell, D. Freed, m-stat director crop and soil sciences department, Michigan state university, USA
- Murray, R.; Granner, K.; Mayes, P.A. and Rodwell, V.W. (1991). Harper's Biochemistry twenty second edition, Appleton & Lange, Norwalk, Connecticut/Los Altos, California.
- National Research Council, NRC (1993). Nutrition Requirements of Fish. National Academy Press. Washington, D.C. 114pp.
- New, M.B. (1985). Feed and Feeding of Fish and Shrimp. A manual on the preparation and presentation of compound feeds for shrimp and fish in aquaculture, United Nation Development Program, FAO, Rome, pp. 14-35.
- Omar, J.M. A.; Govoret, L. and Abo-Omar J. M. (1995): Utilization of olive cake in fattening ratio of Awassi lambs. Review de Medicine verterinaire, 146 (4): 273- 276.
- Osman, M.F.; Kamal, S.M.; Hayam D. Tonsy and Nour A.A. (2003). The utilization of two salt tolerant plants un Nile tilapia feeding. Egyptian J. Nutrition and Feeds 6. 459-466.
- Razzaque, M.A. and El-Sheikh O. F. (1982): Olive oil cake in the rations of growing heifers. Libyan J. of Agric., 10: 25-35.
- Reitman, A. and Frankel, S. (1957). Determination of aspartate amino transferase and alanin aminotransferase. Am. J. Clin. Path., 28: 56.
- Shalaby, S.M.H. (1998). Studies on fish production. Nutritional requirements of rabbitfish, *Siganus rivulatus*, fingerlings. Ph.D. Thesis, Alexandria University, Alex., Egypt
- Snedecor, G.W. and Cochran, W.G. (1982). Statistical Methods, 6th ed. Iowa State University. Press, Ames. Iowa
- Tortuero, F.; Rioperez, J. and Rodriguez, M. L. (1989): Nutritional value for rabbits fed olive cake and the effects on their visceral organs. Animal Feed Science and Technology, 25: 79-86.
- Trinder B. (1969). Ann. Clin. Biochem. ,6:24-32.
- Yousef, K.M.; Fayed, Afaf M. and Khamis, H.S. (2001): Productive and reproductive performances of ewes and does fed non-conventional diets based on olive pulp in Sinai. Egyptian J. Nutrition and Feeds (2001) 4 (special issue): 591-604.

Zaza, G.; Fatma, G.F.; Abo-Donia; and Hammand, M.R. (2001): Effect of incorporating olive cake as a non-conventional feedstuff in diets of growing commercial rabbit. Egyptian Journal of Nutrition and feeds (Special Issue): 861 –868.

الاستفادة من تفل الزيتون كمصدر علفي غير تقليدي في تغذية اصبعيات البلطى النيلية

أسماء إبراهيم عبد المنعم
قسم الإنتاج الحيواني - كلية الزراعة-جامعة الإسكندرية

في تجربة غذائية أجريت لدراسة تأثير إضافة مستويات مختلفة من تفل الزيتون على النمو والاستفادة من الغذاء والعناصر الغذائية، والجلوكوز والدهون الكلية والبروتينات الكلية للبلازما، والتركيب الكيماوى لجسم الأسماك وتحليل التكلفة- الربح لأسماك البلطى النيلية. استخدمت عشرة أحواض زجاجية بأبعاد ٧٠ X ٣٠ X ٤٠ سم لتخزين عشرة أسماك لكل حوض بمتوسط وزن ٩ جم/سمكة. تم تركيب أربعة علائق تجريبية متزنة في البروتين و الطاقة لتحتوى على صفر، ٥، ١٠، ١٥، ٢٠ % من تفل الزيتون. قدمت العلائق ثلاثة مرات يوميا في ستة أيام كل أسبوع لمدة ٧٧ يوما. وكانت الأسماك توزن كل أسبوعين ويعدل على أساسها كميات الغذاء.

أشارت النتائج أنه على الرغم من عدم وجود اختلافات معنوية في أداء النمو للأسماك المغذاه على مستويات مختلفة من تفل الزيتون، إلا أن الأسماك التي تغذت على ٥% تفل زيتون كانت تميل الى إحداث زيادة خفيفة في وزن الجسم والزيادة الوزنية ومعدل النمو النوعي. كما أن كميات الغذاء المأكول الأكثر انخفاضا قد لوحظت في حالة الأسماك التي تغذت على ١٥، ٢٠% تفل زيتون. وأنه لا يوجد اختلافات معنوية في معدل التحويل الغذائى والكفاءة النسبية للبروتين وكفاءة احتجاز الطاقة والبروتين بين جميع الأسماك التي تغذت على المستويات المختبرة من تفل الزيتون. في حين سجلت زيادة كل من دهون الجسم وجلوكوز البلازما ودهونها الكلية بزيادة مستويات تفل الزيتون في علائق المختبرة وعلى العكس انخفضت قيم البروتينات الكلية في البلازما.

كما أظهر التحليل الاقتصادي الى انخفاض التكلفة الغذائية لانتاج اكجم سمك وزيادة دليل الربحية بزيادة مستويات تفل الزيتون في علائق البلطى النيلية.