

EFFECT OF REPLACING NON-FAT DRY MILK WITH GELATIN IN MAKING YOGHURT FROM COW'S MILK

Badran, I. I.

Dept. of Dairy Sci. and Technol., Fac. of Agric., Minufiya University, Shibin El-Kom, Egypt.

ABSTRACT

Ten batches of yoghurt were made from cow's milk. Control yoghurt (C₁) was made from cow's milk fortified with 3.0% non-fat dry milk (the total equal eleven batches). Another five batches were made from cow's milk fortified with non-fat dry milk at the rate of 1.5% and 0.0, 0.3, 0.35, 0.40 and 0.45% gelatin (C₂, T₁, T₂, T₃ and T₄), respectively. The other five batches were made from cow's milk fortified with 0.0, 0.45, 0.50, 0.55 and 0.60% gelatin (C₃, T₅, T₆, T₇ and T₈), respectively. All yoghurt treatments were sampled at 1, 3, 6, 9 and 12 days of storage period for chemical, rheological and sensory evaluation. The results showed that reduction of non-fat dry milk caused a significant decrease in total solids, total protein, ash, lactose and acetaldehyde contents of yoghurt as well as scores of organoleptic properties. While it did not affect the fat content of yoghurt. Addition of gelatin to yoghurt milk improved the organoleptic properties and increased acetaldehyde content and titratable acidity, while decreased pH values. Adding of gelatin did not affect significantly total solids, total protein, fat, ash and lactose contents of yoghurt for each type of milk. Total solids, ash and titratable acidity increased as storage period progressed, while total protein, lactose and pH values decreased. Scores of organoleptic properties did not change as storage period advanced up to the sixth day, thereafter, decreased up to the end of storage period. Syneresis of whey decreased up to the sixth day of storage, then it increased as storage period progressed. Total bacterial, streptococci and lactobacilli counts increased during the first three days of storage period, then decreased up to the end of storage period.

Keywords: Yoghurt, non-fat dry milk, gelatin, sensory evaluation.

INTRODUCTION

Yoghurt is the most popular fermented milk produced in Egypt and worldwide. Its consumption in Egypt has been increased tremendously. The value of yoghurt in human nutrition is based, non only on the nutritive value of the milk from which it is made and increased digestibility, but also on the beneficial effect of intestinal microflora, prophylactic and healing effects (Rasic & Kurman, 1978 and Buttriss, 1997).

Many health benefits have been attributed to yoghurt such as improved lactose tolerance, protection against gastrointestinal infections, effective treatment for specific types of diarrhea, relief of constipation, improved immunity, cholesterol reduction and protection against cancer (Rasic & Kurman, 1978; Agerbaek *et al.*, 1995; Schaafsma, 1996; Tvede, 1996 and Buttriss, 1997).

It is well known that making yoghurt from cow's milk has a weak body and texture. Therefore it has been recommended to fortify cow's milk with non-fat dry milk, stabilizers, whey products and ropy culture to

El-Salam *et al.*, 1996; Hassan *et al.*, 1999; Harby & El-Sabie, 2001 and Zedan *et al.*, 2001). Because of the higher price of non-fat dry milk and its availability at any time, other ingredients have been tried.

The objectives of this study were to evaluate the possibility of making a good quality yoghurt from cow's milk fortified with gelatin and to investigate the effects of the rate of replacing non-fat dry milk with gelatin on the physical, rheological, chemical and sensory evaluation of yoghurt.

MATERIALS AND METHODS

Starter culture:

Streptococcus salivarius subsp. *thermophilus* EMCC 1043 and *Lactobacillus delbrueckii* subsp. *bulgaricus* EMCC 1102 were obtained from Cairo Mircen (Ain Shams University, Egypt). These strains were activated by three successive transfers in sterile 10% reconstituted non-fat dry milk.

Manufacture of yoghurt:

Preliminary study was carried out to choose the best amount of non-fat dry milk that should be added to cow's milk to get the best yoghurt. The preliminary obtained results revealed that yoghurt made from cow's milk fortified with 3% non-fat dry milk was the most acceptable yoghurt.

Fresh cow's milk was obtained from the herd of Tokh Tanbisha Farm belonging to Fac. of Agric., Minufiya Univ. Milk was standardized to 3% fat and divided into eleven batches. One portion was used as a control and the milk was fortified with 3% non-fat dry milk (4).

Non-fat dry milk was added to five portions at the rate of 1.5%, one of them was used as control (C₂), while gelatin was added to the other four portions at each of them at the rate of 0.30, 0.35, 0.40 and 0.45% (T₁, T₂, T₃ and T₄), respectively. The other five portions were not fortified with non-fat dry milk, one of them was used as control (C₃), while the other four ones were fortified with gelatin at the rate of 0.45, 0.50, 0.55 and 0.60% (T₅, T₆, T₇ and T₈), respectively.

Non-fat dry milk was added to warm milk and all batches were heated to 85°C for 20 min. gelatin which was dissolved in some milk was added to heated milks. Heated milks were cooled to 40 ± 2°C, then inoculated with 1% of *S. thermophilus* subsp. and 1% of *Lactobacillus delbrueckii* subsp. *bulgaricus*. The inoculates batches were packed in plastic cups and incubated at 40 ± 2°C for 3.0-3.5 hrs. until the desirable acidity was evaluated. All batches were stored at 6 ± 1°C for 12 days and sampled for analysis were picked at days 1, 3, 6, 9 and 12 days. These experiments were done in triplicated.

Bacteriological analysis:

Total viable bacterial counts were enumerated on standard plate count agar (Messer *et al.*, 1985). MRS medium was used to enumerate *Lactobacillus delbrueckii* subsp. *bulgaricus* (de Man *et al.*, 1960). *S. salivarius* subsp. *thermophilus* was enumerated on yeast lactose agar (Skinner and Quesnel, 1978).

Rheological analysis:

Syneresis was determined according to the method of Dannenberg and Kessler (1988) with slight modification. Hundred grams yoghurt in plastic cup was cut into four sections and transferred into a funnel fitted with 120 mesh metal screen. The whey was drained into a graduated cylinder. The amount of whey drained off was measured after 15, 30, 45, 60, 90 and 120 min. at room temperature ($20 \pm 1^\circ\text{C}$) on yoghurt stored for one day and after 2 h on yoghurt stored for 3, 6, 9 and 12 days.

Chemical analysis:

Yoghurt samples were analyzed at the first and twelfth days for total solids, total nitrogen, fat, ash contents, while analyzed during storage for titratable acidity and pH and according to Ling (1963). Lactose was determined according to the method described by Nickerson *et al.* (1976), while acetaldehyde was determined according to Lees and Jago (1969).

Sensory evaluation:

Yoghurt samples were evaluated for flavour, appearance, acidity and body and texture by 15 panelists of the staff members of Dairy Sci. and Tech. Dep., Fac. of Agric. Minufiya Univ. Results were recorded in a score sheet described by Kebary and Hussein (1999). Samples were presented to judges in plastic cups in random order. Judges were provided with room temperature rinse water, plastic spoons and score sheets.

Statistical analysis:

2×3 factorial design were used to analyze the data. Duncan's test was used to make the multiple comparisons (Steel and Torrie, 1980). Significant differences were determined at $p \leq 0.05$.

RESULTS AND DISCUSSION

Syneresis of all yoghurt batches was determined at different times on the first day of storage. Whey separation from all yoghurt treatments increased by increasing the time up to 2 hours (Fig. 1). Similar results were reported by Harwalkar & Kalab (1983); Kebary & Hussein (1999) and Harby & El-Sabie (2001). Therefore, it has been decided to measure whey syneresis during storage of all yoghurt treatments for 2 hours. Decreasing the rate of fortifying cow's milk with non-fat dry milk caused a significant increase of whey syneresis (Fig. 2 and Table 5) (C_1 , C_2 and C_3). Fortification of cow's milk with gelatin even a accompanied by decreasing non-fat dry milk caused a significant ($p \leq 0.05$) reduction of whey syneresis (Fig. 2 and Table 5). This reduction of whey separation was proportional to the amount added of gelatin to each type of milk (T_1 , T_2 , T_3 and T_4) and (T_5 , T_6 , T_7 and T_8). Although, T_2 , T_3 and T_4 fortified with only 1.5% non-fat dry milk, they exhibited lower whey separation than C_1 that fortified with 3.0% non-fat dry milk. Similar results were observed for treatments T_7 and T_8 which were not fortified with non-fat dry milk, but fortified with 0.55 and 0.60% gelatin,

Table (1): Gross composition of yoghurt fortified with gelatin.

Yoghurt treatments*	Total solids (%)		Total protein (%)		Fat (%)		Lactose (%)		Ash (%)	
	1 day	12 day	1 day	12 day	1 day	12 day	1 day	12 day	1 day	12 day
C ₁	14.68	15.06	4.52	4.48	3.30	3.20	4.72	4.09	0.99	1.07
C ₂	12.31	12.46	4.04	3.99	3.20	3.10	3.96	3.18	0.92	0.96
T ₁	12.58	12.71	4.29	4.24	3.20	3.10	3.94	3.21	0.95	0.96
T ₂	12.61	12.75	4.33	4.30	3.20	3.10	3.95	3.20	0.95	0.97
T ₃	12.70	12.78	4.38	4.32	3.20	3.10	3.93	3.22	0.96	0.97
T ₄	12.74	12.84	4.43	4.37	3.20	3.10	3.94	3.22	0.96	0.98
C ₃	11.72	12.18	3.72	3.68	3.10	3.00	3.11	2.51	0.83	0.86
T ₅	11.99	12.48	3.99	3.94	3.10	3.00	3.13	2.53	0.84	0.86
T ₆	12.04	12.56	4.04	4.01	3.10	3.00	3.11	2.53	0.84	0.88
T ₇	12.44	12.63	4.08	4.05	3.10	3.00	3.14	2.52	0.85	0.89
T ₈	12.49	12.70	4.13	4.07	3.10	3.00	3.12	2.53	0.85	0.90

* C₁ = yoghurt made from cow's milk fortified with 3.0% non-fat dry milk.

C₂, T₁, T₂, T₃ and T₄ = yoghurt treatments made from cow's milk fortified with 1.5% non-fat dry milk and 0.0, 0.30, 0.35, 0.40 and 0.45% gelatin respectively.

C₃, T₅, T₆, T₇ and T₈ = yoghurt treatments made from cow's milk without fortification with non fat dry milk, but fortified with 0.45, 0.50, 0.55 and 0.60% gelatin respectively.

Table (2): Changes in titratable acidity and pH values during storage of yoghurt fortified with gelatin.

Yoghurt treatments*	Titratable acidity (%)					pH value				
	Storage period (days)					Storage period (days)				
	1	3	6	9	12	1	3	6	9	12
C ₁	0.96	1.08	1.13	1.22	1.26	4.63	4.33	4.18	4.04	3.98
C ₂	0.88	0.96	1.04	1.16	1.18	4.68	4.36	4.20	4.12	4.10
T ₁	0.90	0.97	1.05	1.18	1.20	4.61	4.33	4.17	4.12	4.10
T ₂	0.93	1.01	1.08	1.20	1.22	4.56	4.32	4.16	4.08	4.00
T ₃	0.97	1.03	1.10	1.22	1.24	4.49	4.30	4.16	4.05	3.98
T ₄	1.00	1.06	1.12	1.22	1.24	4.37	4.26	4.12	4.02	3.96
C ₃	0.83	0.91	0.98	1.05	1.13	4.70	4.43	4.28	4.19	4.13
T ₅	0.85	0.94	1.02	1.09	1.17	4.68	4.38	4.25	4.17	4.11
T ₆	0.88	0.96	1.06	1.11	1.20	4.66	4.35	4.21	4.13	4.10
T ₇	0.92	1.00	1.09	1.13	1.24	4.62	4.30	4.19	4.09	4.06
T ₈	0.94	1.01	1.12	1.15	1.26	4.60	4.28	4.18	4.04	4.00

* See Table (1).

respectively (Fig. 2 and Table 5). These results are in agreement with those reported by Abd El-Salam *et al.* (1996); Hassan *et al.* (1999); Harby & El-Sabie (2001). They found that addition of stabilizers to yoghurt milk decreased the whey separation from yoghurt. This reduction of whey syneresis might be due to increasing the water binding capacity of yoghurt curd. Whey separation of all yoghurt treatments decreased as storage period progressed up to the sixth day of storage period then increased. This increase might be due to the increasing of acidity during storage. Similar results were reported by Abd El-Salam *et al.* (1996); Hassan *et al.* (1999) and Kebary & Hussein (1999).

Gross composition of yoghurt is presented in Table (1). Total solids of yoghurt were significantly decreased ($p \leq 0.05$) by decreasing the rate of fortification of cow's milk with non-fat dry milk (Tables 1, 5). Control yoghurt that made with adding 3% non-fat dry milk had the highest total solids content followed with C₂ and C₃ (Tables 1, 5). On the other hand, fortification of cow's milk with gelatin did not affect significantly ($p > 0.05$) the total solids content of yoghurt either made with adding 1.5% non-fat dry milk (T₁, T₂, T₃ and T₄) or treatments which made without adding non-fat dry milk (T₅, T₆, T₇ and T₈) (Tables 1, 5). Total solids content of all yoghurt treatments increased significantly ($p \leq 0.05$) during storage period, which might be due to the evaporation of moisture. These results are in agreement with those reported by El-Nagar and Brennan (2001).

Table (3): Acetaldehyde content of yoghurt fortified with gelatin.

Yoghurt treatments*	Storage period (days)				
	1	3	6	9	12
C ₁	33.1	35.2	34.3	27.1	24.1
C ₂	30.0	32.1	33.1	26.8	22.6
T ₁	31.3	32.9	29.2	23.9	20.4
T ₂	32.9	35.0	32.4	27.4	23.2
T ₃	33.8	35.8	31.6	27.6	23.9
T ₄	34.2	36.2	33.8	28.2	24.1
C ₃	28.6	30.1	33.1	25.9	23.2
T ₅	29.3	31.8	28.5	22.6	19.9
T ₆	30.4	32.6	29.3	23.4	21.3
T ₇	31.0	33.9	30.1	26.6	23.6
T ₈	31.9	35.0	32.2	28.2	24.2

* See Table (1).

Total protein content of yoghurt decreased significantly ($p \leq 0.05$) by decreasing the rate of fortification of cow's milk with non-fat dry milk (Tables 1, 5). Control yoghurt made with adding 3.0% non-fat dry milk contained the highest total solids followed by treatments C₂, T₁, T₂, T₃ and T₄ those made with adding 1.5% non-fat dry milk, then treatments C₃, T₅, T₆, T₇ and T₈ those made without fortification with non-fat dry milk (Tables 1, 5). Treatments C₂, T₁, T₂, T₃ and T₄ were not significantly ($p < 0.05$) different from each other. Also, treatments C₃, T₅, T₆, T₇ and T₈ were not significantly different from each other, which means that adding gelatin did not affect the protein content of yoghurt (Tables 1, 5) similar results were obtained by Mehanna and Mehanna (1989).




Fig. (1): Effect of replacing non-fat dry milk with gelatin on syneresis of yoghurt.

* See Table 1.




Fig. (2): Effect of replacing non-fat dry milk with gelatin and storage on syneresis of whey (%) of yoghurt (test was carried out for 2h).

* See Table 1.




Fig. (3): Effect of replacing non-fat dry milk with gelatin and storage on total bacterial counts of yoghurt.

* See Table 1.




Fig. (4): Effect of replacing non-fat dry milk with gelatin and storage on streptococci counts of yoghurt.

* See Table 1.




Fig. (5): Effect of replacing non-fat dry milk with gelatin and storage on lactobacilli counts of yoghurt.

* See Table 1.

Table (4): Organoleptic scores of yoghurt fortified with gelatin.

Yoghurt treatments*	Flavour (out of 45)					Body and texture (30)					Appearance (15)					Acidity (10)					Total score (out of 100)				
	Storage period (days)					Storage period (days)					Storage period (days)					Storage period (days)					Storage period (days)				
	1	3	6	9	12	1	3	6	9	12	1	3	6	9	12	1	3	6	9	12	1	3	6	9	12
C ₁	43	42	40	40	39	27	27	26	25	24	13	12	12	11	10	9	9	8	8	7	92	90	86	84	80
C ₂	40	40	39	39	37	25	25	24	23	22	11	11	10	10	9	8	8	7	7	7	84	84	80	79	75
T ₁	40	40	40	39	38	25	25	24	24	23	12	12	11	10	9	8	8	7	7	7	85	85	82	80	77
T ₂	41	41	40	40	39	26	25	25	24	24	13	12	11	11	10	9	9	8	8	7	89	87	84	83	80
T ₃	42	42	41	40	39	27	27	26	25	24	13	13	12	12	11	9	9	8	8	7	91	91	87	85	81
T ₄	41	41	41	39	38	26	26	25	24	23	12	12	11	11	10	8	9	8	8	7	87	88	85	82	78
C ₃	39	38	38	37	35	23	23	22	21	20	10	10	9	8	8	7	7	6	6	6	79	78	75	72	69
T ₅	40	40	39	39	36	24	24	23	22	20	11	10	9	8	8	8	7	7	6	6	83	81	78	75	70
T ₆	42	42	41	40	38	25	25	24	23	22	11	11	10	10	10	8	8	8	7	6	86	86	83	80	76
T ₇	41	41	40	40	37	26	26	25	24	24	12	12	11	10	10	9	9	8	7	7	88	88	84	81	78
T ₈	41	41	40	38	36	25	25	25	24	22	11	11	10	10	9	8	8	8	7	7	85	85	83	79	74

* See Table (1).

2684

Table (5): Statistical analysis of yoghurt properties.

Properties of yoghurt	Mean squares	Effect of treatments						Mean square						Effect of storage (days)					
		Multiple comparisons*												Multiple comparisons*					
		C ₁	C ₂	T ₁	T ₂	T ₃	T ₄	C ₃	T ₅	T ₆	T ₇	T ₈				1	3	6	9
T.S. (%)	16.425*	A	B	B	B	B	B	C	C	C	C	C	12.131*	B					A
Protein (%)	3.125*	A	B	B	B	B	B	C	C	C	C	C	3.424*	A					B
Fat (%)	6.431	A	A	A	A	A	A	A	A	A	A	A	1.021*	A					A
Ash (%)	3.025*	A	B	B	B	B	B	C	C	C	C	C	5.467*	B					A
Lactose (%)	0.942*	A	B	B	B	B	B	C	C	C	C	C	9.321*	A					B
Acetaldehyde (ppm)	59.312*	A	C	AB	AB	A	A	D	C	C	AB	AB	113.231*	C	A	B	D	E	
pH value	35.118	C	AB	B	CD	CD	D	A	AB	AB	C	C	0.932*	A	B	C	D	E	
Titratabel acidity (%)	36.121*	AB	BC	BC	AB	AB	A	CD	CD	BC	AB	AB	1.239*	E	D	C	B	A	
Syneresis after 2h (%)	53.463*	D	B	C	CD	CD	D	A	A	AB	B	C	76.432*	A	C	D	C	B	
Organoleptic properties:																			
Flavour	69.12*	A	B	B	AB	A	AB	C	AB	A	AB	AB	93.126*	A	A	AB	B	C	
Body and texture	23.809*	A	C	C	AB	A	B	E	D	C	AB	C	36.241*	A	A	AB	B	C	
Appearance	9.427*	A	C	B	A	A	B	D	C	C	AB	C	58.432	A	A	AB	B	C	
Acidity	7.362*	A	B	B	A	A	B	C	B	B	B	B	26.782	A	A	AB	B	C	
Total score	296.321	A	D	D	AB	A	C	E	D	C	AB	D	934.635	A	A	AB	B	C	

2685

* Each different letters (in the same row) means that the multiple comparisons are different from each other, letter A is the highest mean followed by B, C, ... etc.
 * Significant at 0.05 level.
 † See legend Table (1).

Protein content of all yoghurt treatments decreased slightly ($p \leq 0.05$) during storage period. Similar results were reported by El-Nagar and Brennan (2001).

Concerning fat content, all yoghurt treatments were not significantly ($p > 0.05$) different from each other, which means neither the rate of fortification with non-fat dry milk nor the rate of fortification of cow's milk with gelatin affected significantly ($p > 0.05$) the fat content of the yoghurt treatments (Tables 1, 5) (Mehanna and Mehanna, 1989). The fat content of yoghurt treatments did not change during storage period (Mehanna & Mehanna, 1989 and Kebary & Hussein, 1999).

Lactose content of yoghurt treatments decreased significantly ($p \leq 0.05$) by decreasing the rate of fortification of cow's milk with non-fat dry milk. Treatment C_1 contained the highest lactose content followed by treatments C_2 , T_1 , T_2 , T_3 and T_4 then treatments C_3 , T_5 , T_6 , T_7 and T_8 , which might be due to the lactose content of added non-fat dry milk. Either treatments C_2 , T_1 , T_2 , T_3 and T_4 or treatments C_1 , T_5 , T_6 , T_7 and T_8 were not significantly ($p > 0.05$) different from each other in lactose content, which means that the rate of adding gelatin did not affect significantly ($p > 0.05$) the lactose content of yoghurt treatments (Tables 1, 5). Lactose content of yoghurt treatments decreased ($p \leq 0.05$) throughout storage period, which might be due to the consumption of lactose by lactic acid bacteria (El-Nagar and Brennan, 2001).

Ash content of yoghurt decreased ($p \leq 0.05$) by decreasing the amount added of non-fat dry milk, which might be due to the ash content of non-fat dry milk (Tables 1, 5). Supplementation of cow's milk with gelatin did not affect significantly ($p > 0.05$) the ash content of yoghurt. Ash content of all yoghurt treatments increased slightly during storage (El-Nagar and Brennan, 2001).

Slight differences were noticed among yoghurt treatments in titratable acidity of yoghurt (Tables 2, 5) (Mehanna and Mehanna, 1989). On the other hand, titratable acidity of all yoghurt treatments increased significantly ($p \leq 0.05$) as storage period progressed (Tables 2, 5). These results are in agreement with those reported by Mehanna & Mehanna (1989); Abd El-Salam *et al.* (1996); Kebary & Hussein (1999); Harby & El-Sabie (2001) and Zedan *et al.* (2001).

pH values as affected by the rate of fortification with non-fat dry milk or gelatin and storage period followed opposite trends of those of titratable acidity (Tables 2, 5). Similar results are reported by Badawi & El-Sonbaty (1997); Hassan *et al.* (1999); Harby & El-Sabie (2001) and Zedan *et al.* (2001).

Acetaldehyde content of yoghurt decreased ($p \leq 0.05$) by decreasing the rate of fortification with non-fat dry milk (Tables 3, 5), while increased slightly by increasing the rate of adding gelatin (Tables 3, 5). Acetaldehyde content of all yoghurt treatments increased up to the 3rd day of storage, then decreased up to the end of storage period. Similar results were reported by Salama (2001).

The changes of total, lactobacilli and streptococci bacterial counts are illustrated in Figs. (3, 4, and 5). These bacteria followed almost similar trends. The count of total, lactobacilli and streptococci bacteria decreased by decreasing the rate of adding non-fat dry milk, while they increased slightly by adding gelatin. Total lactobacilli and streptococci bacterial counts increased as storage period progressed up to the third day, then decreased up to the end of storage period. These results are in agreement with those reported by Kebary *et al.* (1996); Hussein & Kebary (1999) and Harby & El-Sabie (2001).

Scores of sensory evaluation of yoghurt are presented in Table (4). Scores of organoleptic properties (flavour, body and texture, appearance, acidity and total scores) followed similar trends. Scores of sensory evaluation decreased by decreasing the rate of fortification of cow's milk with non-fat dry milk. Scores of organoleptic properties increased by adding gelatin and this increase was proportional to the rate of adding gelatin (Tables 4, 5). Yoghurt treatment that made with adding 1.5 and 0.4% of non-fat dry milk and gelatin, respectively was not significantly ($p > 0.05$) different from control yoghurt that made with adding 3.0% non-fat dry milk and T₇ which made from unfortified milk and adding 0.55% gelatin (Tables 4, 5). Scores of organoleptic properties of all yoghurt treatments did not change during storage period up to the sixth day, then decreased as storage period proceeded up to the end of storage period (Tables 4, 5).

It could be concluded that it is possible to make a good quality yoghurt from cow's milk by using non-fat dry milk and gelatin and saving about 50% of the non-fat dry milk used in making the control yoghurt.

REFERENCES

- Abd El-Salam, M. H.; H. M. El-Etriby and N. M. Shahein (1996). Influence of some stabilizers on some chemical and physical properties of yoghurt. *Egyptian J. Dairy Sci.* 24: 25.
- Abou-Dawood, A.E.; E.L. Ghita; I.A. Abd El-Gawad; H.A. El-Gazzar and S.A. Ibrahim (1984). Utilization of unsalted whey protein in some dairy products. III. In yoghurt manufacture. *Annals of Agriculture Science, Moshtohor*, 21: 719.
- Agerbaek, M.; L.V. Gerdes and B. Richelsen (1995). Hypocholesterolaemic effect of new fermented milk products in healthy middle-aged men. *Europ. J. Clinical Nutr.* 49: 346.
- Badawi, R.M. and A.H. El-Sonbaty (1997). Viability of *Staphylococcus aureus* and *Escherichia coli* in Zabady made with bifidobacteria. *Egyptian J. Dairy Sci.* 25: 217.
- Buttriss, J. (1997). Nutritional properties of fermented milk products. *Int. J. Dairy Technol.* 50: 21.
- Dannenberg, F. and H. G. Kessler (1988). Effect of denaturation of β -lactoglobulin on texture properties of set-style non-fat yoghurt. I. *Syneresis. Milchwissenschaft* 43: 632.
- De Man J. C.; M. Rogosa and M. E. Sharp (1960). A medium for the cultivation of lactobacilli. *J. Appl. Bacteriol.* 23: 130.

- El-Nagar, G. F. and C. S. Brennan (2001). The influence of fiber addition on the texture and quality of stirred yoghurt. Proc. 8th Egyptian conf. Dairy Sci. and Techn. pp. 505 – 523.
- Harby, S. and W. El-Sabie (2001). Studies on skim milk yoghurt using some stabilizers. Proc. 8th Egyptian Conf. Dairy Sci. and Techn. pp. 537.
- Harwalkar, V. R. and M. Kalab (1983). Susceptibility of yoghurt to syneresis. Comparison of centrifugation and drainage methods. *Milchwissenschaft* 38: 517.
- Hassan, F. A. M.; W. A. Helmy and A. E. Enab (1999). Utilization of some local polysaccharide in manufacture of yoghurt. *Egyptian J. Dairy Sci.* 27: 281.
- Hussein, S. A. and K. M. K. Kebary (1999). Improving viability of bifidobacteria by microentrapment and their effect on some pathogenic bacteria in stirred yoghurt, *Acta Alimentaria* 28: 113 – 131.
- Kebary, K. M. K. and S. Hussein (1999). Manufacture of low fat Zabady using different fat substitutes. *Acta Alimentaria* 28: 1-14.
- Kebary, K. M. K.; A. M. Moussa; A. I. Hamed and E. T. Yousef (1996). Quality of zabady made with *Bifidobacterium bifidum* D1. The proceeding of 36th Week of Science, Alepo Univ. Syria, Book 3, Part 1: 79 – 97.
- Lees, G. J. and G. R. Jago (1969). Methods for the estimation of acetaldehyde in cultural dairy product. *Aust. J. Dairy Techn.*, 24: 181.
- Ling, E. R. (1963). *A Text Book of Dairy Chemistry*. Vol. 2, Practical 3rd Ed. Chapman and Hall, Ltd., London.
- Mehanna, N. M. and A. S. Mehanna (1989). On the use of stabilizer for improving some properties of cow's milk yoghurt. *Egypt J. Dairy Sci.*, 17: 289.
- Messer, J. W.; J. Behney and L. O. Leudecke (1985). *Standard Methods for the Examination of Dairy Products*. 14th ed., Am. Publ. Health Assoc., Washington.
- Nickerson, T. A.; I. F. Vujcic and A. Y. Lin (1976). Calorimetric estimation of lactose and its hydrolytic products. *J. Dairy Sci.*, 59 (3): 386 – 390.
- Rasic, J. L. and J. A. Kurman (1978). *Yoghurt, Scientific Grounds, Technology, Manufacture and Preparations*. Technical Dairy Publishing House, Copenhagen, Denmark, 445 pp.
- Salama, F. M. M. (2001). Preparation of high protein therapeutic acidophilus yoghurt. *Egyptian J. Dairy Sci.*, 29: 277 – 284.
- Schaafsma, G. (1996). State of the art concerning probiotic strains in milk products. *Newsletter Int. Dairy Federation* 145: 23-24.
- Skinner, F. A. and L. B. Quesnel (1978). *Streptococci*. Academic Press, New York, p. 390.
- Steel, R. G. D. and J. H. Torrie (1980). *Principles and Procedures of Statistics. Biometrical approach*. 2nd Ed. McGraw-Hill Book Co., New York.
- Tvede, M. (1996). Potential of probiotic strains in stabilizing intestinal microflora to prevent gastrointestinal infection. *Newsletter Int. Dairy Federation* 145: 30.

Zedan, M. A.; A. N. Zedan; K. M. K. Kebary and S. F. Mahmoud (2001).
Effects of fortification of cows milk with acetylated whey protein
concentrates on the quality of set yoghurt. Egyptian J. Dairy Sci. 29:
285.

تأثير استبدال اللبن الفرز المجفف بواسطة الجيلاتين على صفات اليوجورت المصنوع من اللبن البقرى إبراهيم إبراهيم بدران قسم علوم وتكنولوجيا الألبان - كلية الزراعة - جامعة المنوفية - مصر

في هذا البحث تم تصنيع عشرة معاملات من اليوجورت من اللبن البقرى المدعم أو الغير مدعم بلبن فرز مجفف مع إضافة نسب مختلفة من الجيلاتين بالإضافة إلى العينة القياسية التي صنعت من اللبن البقرى المدعم بإضافة ٣% من اللبن الفرز المجفف (تصبح إجمالي المعاملات إحدى عشر معاملة) ، أما الخمس معاملات الأولى فقد تم تصنيعها من لبن بقرى مدعم بـ ١.٥% لبن فرز مجفف وأضيف إلى كل منها الجيلاتين بنسب صفو ، ٠.٣٠ ، ٠.٣٥ ، ٠.٤٠ ، ٠.٤٥ ، ٠.٥٠ ، ٠.٥٥ ، ٠.٦٠ . وقد تم أخذ عينات من كل المعاملات في الأيام ١ ، ٢ ، ٣ ، ٦ ، ٩ ، ١٢ من التخزين بالثلاجة وذلك لإجراء التحليلات الربولوجية والكيموية والحسية .

ولقد أوضحت النتائج المتحصل عليها بعد تحليلها إحصائياً ما يلي :

- أدى خفض تدعيم اللبن البقرى باللبن الفرز المجفف إلى خفض نسب الجوامد الكلية، البروتين الكلى ، الرماد ، اللاكتوز والأسيتالدهيد كما أدى إلى خفض درجات التقييم الحسى لليوجورت بينما لم تؤثر على نسبة الدهن فى اليوجورت .
- أدى إضافة الجيلاتين إلى اللبن المصنوع منه لليوجورت إلى تحسين الخواص الحسية وكانت أحسن المعاملات هى تلك المعاملة من لبن مدعم بـ ١.٥% لبن فرز مجفف وإضافة ٠.٤% جيلاتين وكانت غير مختلفة معنوياً عن المعاملة المصنعة من لبن بقرى فقط والمضاف له ٠.٥٥% جيلاتين وكذلك المعاملة القياسية المصنعة بإضافة ٣% من اللبن الفرز المجفف وكذلك أدى إضافة الجيلاتين إلى زيادة نسبة الأسيتالدهيد والحموضة بينما أدى إلى إنخفاض قيم الـ pH لليوجورت فى حين لم يؤثر إضافة الجيلاتين معنوياً على نسب الجوامد الكلية والبروتين الكلى والدهن والرماد واللاكتوز .
- ازدادت نسب الجوامد الكلية والرماد والحموضة بتقدم فترة التخزين بينما انخفضت نسب البروتين الكلى واللاكتوز وقيم الـ pH .
- لم تتغير درجات التقييم الحسى أثناء الستة أيام الأولى من التخزين بينما انخفضت بتقدم التخزين بعد ذلك
- انخفض انفصال الشرش بتقدم فترة التخزين حتى اليوم السادس ثم ازداد بزيادة فترة التخزين
- زابت أعداد البكتريا الكلية والـ Streptococci والـ Lactobacilli حتى اليوم الثالث من التخزين ثم انخفضت أعدادها بتقدم فترة التخزين .