

Determination of formalin in animal and poultry inactivated vaccines using different methods

N. A. Y. Sherif

Central Laboratory for Evaluation of Veterinary Biologics, Abbasia, Cairo, Egypt

In this study, two methods for determination of formalin amount were applied on samples of inactivated vaccines representing local or foreign companies. The first method; matching method was conducted by using phenyl hydrazine while in the other method is; spectrophotometry phloroglucinol was used. Spectrophotometrical method was found more sensitive and more accurate than the matching one. At the same time, the spectrophotometry method could be used for determination of formalin in all inactivated vaccines either bacterial or viral vaccines.

During vaccine production, several control tests are performed to ensure that vaccines have been made under optimal circumstances; the raw and intermediate products follow the required standard. At the end of the production process, many tests must be done on the bulk product and the final vaccine. Such tests include safety, potency, sterility, identity as well as residual chemical constituents particularly formalin which is used as inactivating agent during manufacture. Formalin is considered as toxicogenic or carcinogenic material for both animals and human beings. Tests have been formulated for each vaccine according to (British Pharmacopoeia, 1973; Quality Control of Vaccine, 1983 and European Pharmacopoeia, 1997) which are generally accepted all over the world.

Brandly *et al.* (1946) showed that 0.025% diluted formalin could destroy the infectivity of Newcastle disease virus (NDV).

King (1991) evaluated different methods for inactivation of NDV and avian influenza viruses (AIV). He found that NDV or AIV were inactivated by binary ethyleneimine (BEI) (0.01M) with no adverse effect on haemagglutinating or hemolytic activities. The effect of formalin (0.1%) was variable and depressed HI titres of antisera.

Soliman *et al.* (1996) found that binary ethyleneimine, used as inactivating agent for NDV, had no effect on the antigenicity of the virus when used in a concentration of 0.01 or 0.03 M. These results compared favourably with those obtained when using formaldehyde solution as an inactivating agent at concentration of 0.1 or 0.2%, which significantly reduced antigenicity of the haemagglutination titer of 1024 to that of 32.

The widespread use of formaldehyde and the reports of adverse effects have created the need for specific, sensitive and simple method for its determination.

The present work aimed to compare between two currently used methods for determination of formalin: the first one is by using phenyl hydrazine and the other is by using phloroglucinol (spectrophotometry).

Material and Methods

Tested batches. Total number of (136) batches of inactivated vaccines were used in this study. These batches represent local and imported vaccines: (88) viral poultry inactivated vaccines, (29) bacterial poultry inactivated vaccines, (10) viral large animal inactivated viral vaccines and (9) inactivated bacterial vaccines of large animal.

Phenyl hydrazine method. It was carried out according to Quality Control of Vaccines (1983). Standard formaldehyde solution was prepared by measuring out 1 ml of aqueous formaldehyde solution (40%) into a 100 ml standard flask and diluting to the mark with distilled water. The concentration of the solution was found to be 3800 ppm. One ppm of formaldehyde solution was prepared freshly prior to use. In five test tubes represented five concentrations (1.0, 0.50, 0.1, 0.05 % and 0.01 %) of the formaldehyde (El-Gomhoria Company for Chemicals, Drugs and Medical Supplies, Egypt) standards. 1ml in each tube and another test tube represented the sample (1ml of the tested inactivated vaccine plus 100 ml distilled water). The following reagents were put in order: 0.1 ml of phenyl hydrazine 1% (Riedel-de Haen Allied Signal, Germany), 0.1 ml of potassium ferricyanide 5% (BDH Chemicals Ltd., Poole, England) and Three drops of concentrated hydrochloric

Table (1): Calibration graph was drawn using the concentration (ppm) against absorbance (optical density OD).

Concentration (ppm)	Absorbance (Optical Density)
0.05	0.1031
0.10	0.2059
0.15	0.3178
0.20	0.4413
0.25	0.6147




Fig. (1): Calibration graph was drawn using the concentration (ppm) against absorbance (optical density OD).

acid (HCl) (The Egyptian Company for Chemicals and Drugs (ADWIA), Egypt). After 5 min. the colour of both tubes was matched.

2. Phloroglucinol method. It was performed according to the method described by (Gayathri and Balasubramanian, 2000):

Preparation of Calibration Graph. In five test tubes containing 1 ml of 1% phloroglucinol (BDH Chemicals Ltd., Poole, England), 0.5, 1.0, 1.5, 2.0 and 2.5 ml of 1 ppm standard formaldehyde solution was added separately. 4 ml concentrated sulphuric acid (ADWIA Company, Egypt.) was added carefully to each tube using a long stem funnel. The solution was allowed to stand for 20 min. to attain room temperature. The solution was then transferred into 10 ml standard flask, washed with 1ml of 9 M sulphuric acid and diluted to the mark with the same acid. Absorbance was measured at 435 nm against reagent blank prepared according to the same procedure (Table 1, Fig. 1).

Determination of formaldehyde in inactivated vaccines. One ml of the inactivated vaccine was diluted to 100 ml with distilled water. The diluted solution was filtered and used for analysis. One ml of the sample solution was taken and analyzed for formaldehyde content following the procedure described under calibration graph.

Calculation. Calculation of formaldehyde concentration (ppm) was obtained from the

calibration graph based on the following equation:

$$\frac{\Delta \text{ OD (Sample)}}{\Delta \text{ OD (Standard)}} \times \text{Concentration of standard}$$

Results and Discussion

Several methods for determination of formaldehyde have been developed, e.g. high performance liquid chromatography (HPLC) method (Zegota, 1999; Sandner *et al.*, 2001 and Possanzini and Di Pola, 2003), gas chromatography (GC) method (Ren and Guo, 1997; Suliman and Soma, 2000 and Shiraishi *et al.*, 2001), laser spectrometer (Rehle *et al.*, 2001 and Richter *et al.*, 2002) and by spectrophotometry (Lodge, 1989; Lancaster *et al.*, 2000; Ross *et al.*, 2002 and Mason *et al.*, 2004).

In this study, two methods were used; the principle of the first method is base on the reaction of formaldehyde with phenyl hydrazine solution (1%), potassium ferricyanide (5%) in acid solution (HCl) forming a red to faint pink coloured compound, the intensity of which can be matched visually with 1, 0.50, 0.1, 0.05 and 0.01% of formaldehyde standard solutions (Quality Control of Vaccines, 1983). The second method is based on the reaction of formaldehyde with phloroglucinol in acidic solution (Gayathri and Balasubramanian, 2000).

Table (2): Determination of formaldehyde concentration (%) in random batches of inactivated vaccines using visual method (Matching).

Poultry inactivated vaccines						Large animal inactivated vaccines					
Viral Vaccines			Bacterial vaccines			Viral Vaccines			Bacterial vaccines		
Type	No. of batches	Mean+SE	Type	No. of batches	Mean+SE	Type	No. of batches	Mean+SE	Type	No. of batches	Mean+SE
ND	44	0.044+0.002	Cholera	9	0.049+0.007	RVF	3	0.05+0.0	Ovine & caprine HS	2	0.05+0.0
IBD	8	0.05 + 0.0	Coryza	10	0.03+0.006	AHS	2	0.05+0.0	Clostridia	3	0.03+0.02
EDS	5	0.038+0.007	FRHS	6	1.0+0.0	Entero-3	2	0.01+0.0	Black leg	2	0.05+0.0
RVHS	4	0.05+0.0	MG	2	0.01+0.0	FMD	3	0.01+0.0	Cattle HS	2	0.1+0.0
ND+TRT	2	0.53+0.48	E. coli	2	0.01+0.0						
ND+IB	2	0.05+0.0									
ND+IBD	5	0.034+0.009									
ND+EDS	2	0.1+0.0									
ND+IB+EDS	4	0.05+0.0									
ND+IB+SHS	2	0.02+0.0									
ND+IB+IBD	3	0.05+0.0									
ND+IB+IBD+Reo	3	0.04+0.01									
PPMV	2	0.01,0.04									
AE	2	0.025+0.015									
Total	88	0.12+0.03		29	0.24+0.07		10	0.039+0.007		9	0.057+0.009

Mean + Standard Error (SE)

Table (3): Determination of formaldehyde concentration (%) in random batches of inactivated vaccines using spectrophotometry method.

Poultry inactivated vaccines						Large animal inactivated vaccines					
Viral Vaccines			Bacterial vaccines			Viral Vaccines			Bacterial vaccines		
Type	No. of batches	Mean±SE	Type	No. of batches	Mean±SE	Type	No. of batches	Mean±SE	Type	No. of batches	Mean±SE
ND	10	0.0465±0.001	Cholera	5	0.0522±0.001	RVF	3	0.0518±0.003	Ovine & caprine HS	2	0.0535±0.001
IBD	3	0.0521±0.002	Coryza	5	0.0352±0.001	AHS	2	0.0507±0.0003	Clostridia	3	0.0311±0.003
EDS	3	0.0395±0.001	FRHS	3	1.0209±0.02	Enter-3	2	0.0171±0.002	Black leg	2	0.0514±0.001
RVHS	3	0.0532±0.001	MG	2	0.0159±0.0002	FMD	3	0.0016±0.0003	Cattle HS	2	0.1088±0.002
ND+TRT	2	0.0534±0.003	E. coli	2	0.0151±0.0003						
ND+IB	2	0.0563±0.002									
ND+IBD	5	0.0323±0.001									
ND+EDS	2	0.1058±0.001									
ND+IB+EDS	3	0.0507±0.001									
ND+IB+SHS	2	0.0269±0.002									
ND+IB+IBD	3	0.0516±0.001									
ND+IB+IBD+Reo	3	0.0446±0.001									
PPMV	2	0.0155±0.001									
AE	2	0.0301±0.0002									
Total	45	0.0462±0.002		17	0.2095±0.09		10	0.0296±0.007		9	0.0579±0.01

Mean ± Standard Error (SE)

Table (4): Comparison between determination of formaldehyde concentration (%) in random batches of inactivated vaccines using visual method (Matching) and spectrophotometry method.

Type of vaccine	Visual method (matching)		Spectrophotometry method	
	No. of batches	Mean ± SE	No. of batches	Mean ± SE
Viral poultry inactivated vaccines	88	0.12 ± 0.03	45	0.0462 ± 0.002
Bacterial poultry inactivated vaccines	29	0.24 ± 0.07	17	0.2095 ± 0.09
Viral large animal inactivated vaccines	10	0.039 ± 0.007	10	0.0296 ± 0.007
Bacterial large animal inactivated vaccines	9	0.057 ± 0.009	9	0.0579 ± 0.01

Mean ± Standard Error (SE)

The obtained results were illustrated in (Tables 2-4) which showed determination of formaldehyde percent in 136 random batches of inactivated vaccines used in this study, which were collected during the period of (2004-2005). These batches represent local and imported vaccines including 88 viral poultry inactivated vaccines, 29 bacterial poultry inactivated vaccines, 10 viral large animal inactivated vaccines and 9 bacterial large animal inactivated vaccines.

From the obtained results, it was noted that the mean percentage of formaldehyde of the viral poultry inactivated vaccines was (0.0462 %) by spectrophotometrical method and (0.12 %) by the visual method. On the other hand, formaldehyde mean percentages in bacterial poultry inactivated vaccines were 0.2095 and 0.24% in the spectrophotometrical and in visual methods respectively. The viral large animal inactivated vaccines show mean percentage of 0.0296 % and 0.039 % by spectrophotometrical and visual methods respectively. Bacterial large animal inactivated vaccines showed formaldehyde mean percentage of 0.0579 and 0.057 % by spectrophotometrical and visual methods respectively.

Although the two methods gave nearly the same values, it is very clear that the spectrophotometry method is more sensitive than the visual one, where the former gave very accurate percentage of formaldehyde while the later gave approximate values. The obtained results are in agreement with those obtained by (Gayathri and Balasubramanian, 2000) who said that the determination of formaldehyde spectrophotometrically using phloroglucinal is a simple, accurate and very sensitive method, also with Ross *et al.* (2002) who conducted an international collaborative study of quantitative colorimetric method for determination of formaldehyde in veterinary vaccines products by 15 laboratories in North America, Europe and Japan. Moreover, Amer (2004) revealed that the developed spectrophotometry method using Rosaniline was conveniently applied to the determination of traces of formaldehyde in veterinary biological products.

In conclusion, the determination of formaldehyde spectrophotometrically using phloroglucinol method is more simple, accurate and sensitive than the visual (matching) method. The spectrophotometry method could be used for determination of formaldehyde not only in the poultry viral inactivated vaccines but also in

all veterinary inactivated vaccines either bacterial or viral.

Acknowledgement

The author is deeply grateful to Prof. Dr. Mohamed M. Taha, Director of Central Laboratory for Evaluation of Veterinary Biologics, Abbasia, Cairo, Egypt for his kind and great help.

References

- Brandly, C. A.; Moses, M. E.; Jones, E. E. and Jungherr, E. L. (1946): Immunization of chickens against ND. *Amer. J. Vet. Res.*, 7: 307-332.
- British Pharmacopoeia (1973)**: London Her Majesty's Stationery Office.
- European Pharmacopoeia (1997)**: 3rd ed. Published in accordance with the convention on the Elaboration of a European Pharmacopoeia. Council of Europe, Strasbourg.
- Gayathri, N. and Balasubramanian, N. (2000): Spectrophotometric determination of formaldehyde. *Analyt. Lett.*, 33 (14): 3037-3050.
- King, D. J. (1991): Evaluation of different methods of inactivation of NDV in egg fluid and serum. *Avian Dis.*, 35 (35): 505.
- Lancaster, D. G.; Fried, A.; Wert, B.; Henry, B. and Tittel, F. K. (2000): Difference-frequency-based tunable absorption spectrometer for detection of atmospheric formaldehyde. *Appl. Opt.*, 39 (24): 4436-4443.
- Lodge, J. P. (1989): *Methods of air sampling and analysis*. 3rd ed. Lewis Publishers, Michigan, pp. 274-278.
- Mason, D. J.; Sykes, M. D.; Panton, S. W. and Rippon, E. H. (2004): Determination of naturally occurring formaldehyde in raw and cooked mushrooms by spectrophotometry. *Food Addit. Contam.*, 21 (11): 1071-1082.
- Amer, N. A. A. (2004): Determination of quantity of some additives in veterinary biologics. *Fac. Sci., Ain Shams Univ., Egypt*.
- Possanizi, M. and Di Pola, V. (2003): Simultaneous determination of formaldehyde in ambient air by hydrazine reagent and HPLC. *Ann. Chim.*, 93 (1-2): 149-156.
- Quality Control of Vaccines (1983)**: Rijksinstituut Voor Volksgezondheid En Milieuhygiene Bilthoven. The Netherlands. Egypt. Org. Biol. Prod. Vaccines Library, Agouza, Giza, Egypt.
- Rehle, D.; Leleux, D.; Erdelyi, M.; Tittel, F.; Fraser, M. and Friedfeld, S. (2001): Ambient formaldehyde detection with a laser spectrometer based on difference frequency generation in PPLN. *Appl. Phys. B*, 72 (8): 947.
- Ren, Q. and Guo, Y.J. (1997): Rapid determination of low molecular weight aldehydes in air by gas chromatography. *Se Pu.*, 15 (4): 356-357.
- Richter, D.; Fried, A.; Wert, B. P.; Walega, J. G. and Tittel, F. K. (2002): Development of a tunable mid-IR difference frequency laser source for highly sensitive airborne trace gas detection. *Appl. Phys. B*, 75 (2-3): 281-288.
- Ross, P. F.; Draayer, H. and Itoh, O. (2002): In international collaborative study on a method for determination of formaldehyde in veterinary vaccines. *Biologicals*, 30 (1): 37-41.
- Sandner, F.; Dott, W. and Hollender, J. (2001): Sensitive indoor air monitoring of formaldehyde and other carbonyl compounds using the 2,4-dinitrophenyl hydrazine method. *Int. J. Hyg. Environ. Health*, 203 (3): 275-279.
- Shiraishi, T.; Soma, Y.; Ishitani, O. and Sakamoto, K. (2001): Application of an integrated prepsation-GC-NPD

system to automated continuous measurement of formaldehyde and acetaldehyde in the atmosphere. *J. Environ. Monit.* 3 (6): 654-660.

Soliman, S. M.; Hamdy, A.; Zaghloul, W. A. and El-Bordiny, F. (1996): *Assiut Vet. Med. J.*, 35 (69): 157.

Suliman, F. E. and Soma, Y. (2000): The determination of carbonyl compounds in air using a robotic sampling

preparation system integrated to a gas chromatograph with a nitrogen-phosphorus detector. *J. Env. Monit.* 2 (5): 470-475.

Zegota, H. (1999): High-performance liquid chromatography of methanol released from pectins after its oxidation to formaldehyde and condensation with 2,4-dinitrophenyl hydrazine. *J. Chromat. A.*, 863 (2): 227-233.