

EFFECT OF LOW PROTEIN DIETS SUPPLEMENTED WITH LYSINE AND METHIONINE ON THE PERFORMANCE OF MUSCOVY DUCKLINGS

Sonbol, S.M.¹, G. A. Abd El-Rahman¹, R.E. Khidr² and Mona, M. Hassan²

¹Faculty of Agriculture, Zagazig University, Zagazig, Egypt.

²Desert Research Center, Mataria, Cairo, Egypt.

ABSTRACT

This experiment was performed to assess response of muscovy ducklings (0-13 weeks of age) to low protein diets supplemented with lysine and methionine. The corn – soybean meal based experimental diets contained two levels of crude protein were supplemented with lysine and methionine. The influence of supplemental amino acids was studied within each protein level.

The results showed that Muscovy ducklings received low protein(20%) diet with high lysine (1.20%) and 0.83% methionine+ cystine, had significantly ($P < 0.05$) the highest value of average live body weight (876.60 g) during the starting period (0-4 w.k). However, during the finishing and the total period ducklings fed on diet contained high protein diet with low lysine and low methionine+cystine, had a highly significant ($P < 0.01$) increase in body weight, weight gain and improved the feed conversion values. Higher percentage of feather, giblets were recorded in group fed on a diet contained low protein with low lysine and high methionine+cystine%. However, higher percentage of carcass (72.41) was recorded for group fed high protein with low lysine and high methionine+cystine%. The lowest feed cost/kg. gain was (62.50%) recorded by group fed high protein with low lysine and low methionine+cystine levels. It could be recommended in the diet 22% protein, 1.10% lysine and 0.83% methionine+cystine during the starting period (0-5 wk.) followed by diet contained 18% protein, 0.90% lysine and 0.70% methionine+cystine, during the finishing period (5-13 wk.) of age.

Keywords: Ducklings, protein, lysine and methionine.

INTRDUCTION

The protein cost is considered mainly the most expensive item in ducks production. Animal protein sources expensive as compared to plant protein sources, but; plant protein are nutritionally imbalanced, therefore some of essential amino acids should be supplemented. Methionine and lysine are generally the first and second limiting amino acids in corn-soybean diets (D'Mello, 1994) and protein requirement may be reduced in corn-soybean meal-type diets by supplementation with methionine and lysine (Waibel *et al.*, 1995). So, using the optimum level of crude protein and supplementing corn soybean diets with the essential amino acids would cause an increase in the net profit.

The present work aimed to study the response of muscovy ducklings to low protein diets supplemented with lysine and methionine.

MATERIALS AND METHODS

The experimental work of the present study was carried out at Ras Suder Research Station (Desert Research Center) in southern Sinai Governorate. The field work started on 3rd September 2003 and terminated 13 weeks later.

Three hundred and twenty Muscovy ducks of genotype ST₁₂ were randomly divided into eight experimental groups, equal in number (40 each). Each group was allocated to replicates (20 each), in a factorial design 2 x 2 x 2, which contained two dietary protein levels; P1 (22 and 18%), P2 (20 and 16%) during the starting and finishing period, respectively. Within each protein level; ducklings were divided into two sub-groups that fed on diets contained two lysine levels, Lys1 (1.20 and 1.05%) or Lys2 (1.10 and 0.90%) during the starting and finishing period, respectively. Within each lysine groups; ducklings were divided into two sub-group which contained two levels of total sulfur amino acids, TSAA.1 (0.93 and 0.80%), or TSAA.2 (0.83 and 0.70%) during the starting and finishing period, respectively. The xperimental design of the experiment are presented in table 1, while the composition of the experimental diets during the starting and finishing periods are presented in table 2.

Table (1): Experimental design

Treatment Group	E			Finisher		
	Protein %	Lysine %	Methionine+cystine %	Protein %	Lysine %	Methionine +cystine%
1	22	1.20	0.93	18	1.05	0.80
2			0.83			0.70
3		1.10	0.93		0.90	0.80
4			0.83			0.70
5	20	1.20	0.93	16	1.05	0.80
6			0.83			0.70
7		1.10	0.93		0.90	0.80
8			0.83			0.70

Feed and water were offered *ad libitum* during the whole experimental period. Individual bird weights and feed consumption were recorded at different intervals. Feed conversion was calculated as g feed/g gain. At the end of the experimental period, 8 ducks (4 males and 4 females) were randomly from each group, fasted for 12 hours, then slaughtered to estimate the dressing weight percentage as well as the other carcass traits relatively to the live body weight.

Statistical analysis was carried out according to Snedecor and Cochoran (1982). The statistical model which used in the analysis was:

$$Y_{ijk1} = U + P_i + L_j + M_k + PL_{ij} + PM_{ik} + LM_{jk} + PLM_{ijk} + e_{ijkl}$$

Where:

Y_{ijk1} = An observation,

U = the overall mean,

P_i = the protein level in the diets ($i = 1$ and 2),

Table (2): Composition and chemical analysis of the experimental diets (0- 13 weeks old)

Ingredients %	Starter (0- 5 weeks)								Finisher (5- 13 weeks)							
	1	2	3	4	5	6	7	8	1	2	3	4	5	6	7	8
Yellow corn	58.71	58.81	57.50	57.50	63.00	63.00	63.00	63.00	69.14	69.24	69.20	69.30	72.00	72.10	72.10	72.20
Soy bean meal (44%)	35.50	35.50	30.34	30.44	31.60	31.80	31.70	31.80	26.80	26.80	26.80	26.80	20.72	20.72	20.72	20.72
Corn gluten (60%)	2.00	2.00	5.00	5.00	-	-	-	-	-	-	-	-	-	-	-	-
Wheat bran	-	-	3.40	3.40	1.40	1.30	1.40	1.40	-	-	-	-	3.00	3.00	3.00	3.00
Di-calcium phosphate	1.90	1.90	1.80	1.80	1.90	1.90	1.90	1.90	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00
Limestone	1.10	1.10	1.20	1.20	1.10	1.10	1.10	1.10	1.20	1.20	1.20	1.20	1.20	1.20	1.20	1.20
Sodium chlorid	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30
Vit. & Min premix*	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30
DI- Methionine lysine	0.19	0.09	0.16	0.06	0.28	0.18	0.28	0.18	0.20	0.10	0.20	0.10	0.26	0.16	0.26	0.16
	-	-	-	-	0.12	0.12	0.02	0.02	0.06	0.06	-	-	0.22	0.22	0.12	0.12
Total	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100
Chemical omposition**																
Crude protein (%)	22	22	22	22	20	20	20	20	18	18	18	18	16	16	16	16
Kcal (ME)/ kg feed	2840	2840	2839	2836	2838	2847	2835	2838	2920	2920	2920	2920	2920	2920	2920	2920
Calorie/protein (ratio)	129	129	129	129	142	142	142	142	162	162	162	162	182	182	182	182
Lysine	1.20	1.20	1.10	1.10	1.20	1.20	1.10	1.10	1.00	1.00	0.94	0.94	1.01	1.01	0.90	0.90
Methionine+cystin %	0.93	0.83	0.93	0.83	0.93	0.83	0.93	0.83	0.80	0.70	0.80	0.70	0.80	0.70	0.80	0.70
Calcium %	1	1	1	1	0.99	0.94	0.94	0.94	0.98	0.98	0.98	0.98	0.97	0.97	0.97	0.97
Total phosphorus	0.74	0.74	0.74	0.74	0.74	0.76	0.76	0.76	0.74	0.74	0.74	0.74	0.75	0.75	0.75	0.75

* Vitamin and Minerals premix 0.30 of the diet supplies the following per kg of the diets: vit.A 1200 IU, vit. D. 2500 IU, vit. E. 10 mg, vit K. 3mg, vit B₁. 1mg, vit B₂. 4mg pantothenic acid 10mg, folic acid 1mg , niacine 40mg, vit, B₆ 3mg, vit B₁₂ 20mcg, Mn. 62 mg, fe 44mg, Zn. 56mg, cu 5mg and se 100mg.

** Calculated according to NRC (1994).

- L_j = the lysine level in the diets ($j = 1$ and 2),
 M_k = the methionine level in the diets ($K = 1$ and 2),
 PL_{ij} = the interaction between protein level and lysine level ($ij = 1, 2, \dots, 4$),
 PM_{ik} = the interaction between protein level and methionine level ($ik=1, 2, \dots, 4$), LM_{jk} = the interaction between lysine level and methionine level.
 PLM_{ijk} = the interaction between protein, lysine level and methionine level ($ijk = 1, 2, \dots, 4$),
 e_{ijk1} = Random error.

Duncan's new multiple range test (Duncan, 1955) was used to detected the significance between means.

RESULTS AND DISCUSSION

Live Body Weight (BW)

The effects of dietary treatments on body weight are illustrated in Table 3. At 3 weeks of age, Ducklings fed diets contained P2 (20%) with Lys₂ (1.10%) and TSAA₁ (0.93%) had significantly ($P < 0.05$) the heaviest body weight (522.10 g). At 4 weeks of age, ducklings fed diet P2 (20%), Lys₁ (1.20%) and TSAA₂ (0.83%) had significantly ($P < 0.05$) the heaviest body weight (876.60g); however, at 9 and 13 weeks of age; ducklings fed on diet contained P1 (18%), L2 (0.90%) and TSAA₂ (0.70%) had significantly ($P < 0.01$) the heaviest body weight meanwhile; the lowest body weight was recorded by group fed on diet obtained P1 (18%) , lys₁ (1.05%) and TSAA₁(0.80%). This may due to that free amino acid supplements are absorbed more rapidly than protein-bound amino acids resulting in an imbalanced supply of amino acids, imbalanced diets depressed feed intake followed by a retardation of growth. Similar results were obtained by Wang *et al.* (2004) who found that methionine supplementation of diets meat ducks. significantly improved body weight at 21 days of age. Also, Leclercq and De. Carville (1989) observed that live body weight of male muscovy ducklings from 8 to 12 weeks of age improved as the protein level increased from 15 to 17%.

Body Weight Gain

Ducklings fed on diet contained P1(20%) with Lys₂ (1.10%) and TSAA₁ (0.93%) had significantly ($P < 0.01$) the highest weight gain (473.42 g) during the period 0-3 weeks of age. However, there were no significant differences between treatments during the periods 0-4 and 0-5 weeks of age. During the periods; 5-9 and 9-13 weeks of age; ducklings received P1 (18%) with Lys₂ (0.90%) and TSAA₂ (0.70%) showed a significant ($P < 0.01$) increase in weight gain, being (839.1g). Ducklings fed on P1 with Lys₂ and TSAA₂ during the whole experiment had significantly ($P < 0.01$) the highest value of weight gain, followed by group fed p₂, Lys₁ and TsAA₂ while the lowest value of weight gain was recorded by group fed diet contained p₁, Lys₁ and TSAA₁.(Table 4).

Table (3): Effects of protein, lysine and TSAA. levels on average body weight ($\bar{X} \pm SE$) of ducklings

Age (weeks)	P1				P2			
	Lys1		Lys2		Lys1		Lys2	
	TSAA1	TSAA2	TSAA1	TSAA2	TSAA1	TSAA2	TSAA1	TSAA2
0	49.1±0.7	49.3±0.7	48.7±0.8	48.0±0.6	49.9±0.9	48.9±0.7	48.7±0.5	47.9±0.5
3	433.2±18.7 ^d	470.6±16.9 ^c	427.9±19.1 ^d	495.5±14.1 ^b	479.9±16.3 ^{bc}	500.4±16.4 ^b	522.1±13.9 ^a	496.6±12.8 ^b
4	788.0±29.8 ^{de}	811.5±31.6 ^{cd}	763.4±36.2 ^{ef}	836.7±21.2 ^{b^c}	781.4±23.2 ^e	876.6±25.1 ^a	862.6±22.5 ^{ab}	799.2±23.7 ^{de}
5	1142.5±33.9	1120.9±39.6	1152.6±53.0	1273.7±32.0	1185.3±30.7	1214.6±30.8	1147.6±31.4	1174.0±34.8
9	2826.5±76.2 ^e	2861.9±93.0 ^{de}	2960.4±100.8 ^{cd}	3371.0±98.8 ^a	3053.1±97.1 ^{bc}	3106.2±96.6 ^b	2849.8±95.1 ^{de}	3091.0±86.9 ^b
13	3501.2±77.7 ^d	3559.0±94.0 ^d	3683.9±100.1 ^c	4185.1±91.0 ^a	3749.8±91.5 ^{bc}	3862.6±98.1 ^b	3530.4±96.9 ^d	3857.5±87.3 ^b

a, b means in the same row with different superscripts differ significantly at ($p < 0.05$)

Table (4): Effect of protein, lysine and TSAA levels on weight gain ($\bar{X} \pm SE$) of ducklings

Age (weeks)	P1				P2			
	Lys1		Lys2		Lys1		Lys2	
	TSAA1	TSAA2	TSAA1	TSAA2	TSAA1	TSAA2	TSAA1	TSAA2
0-3	384.0±18.1 ^d	421.3±16.2 ^c	379.1±18.3 ^d	447.5±13.5 ^b	430.0±15.5 ^{bc}	451.4±15.6 ^b	473.2±13.4 ^a	448.7±12.3 ^b
0-4	738.9±29.2	762.2±31.0	715.0±36.2	464.8±37.1	731.5±22.6	827.7±24.4	813.9±22.0	799.2±23.7
0-5	1093.4±33.4	1072.4±39.3	1103.4±52.2	1227.8±31.8	1135.4±30.9	1165.7±30.8	10.98.9±31.4	1162.1±34.8
5-9	1684.0±60.8 ^d	1728.1±68.8 ^{cd}	1807.9±71.6 ^{bc}	2096.6±90.7 ^a	1867.8±83.9 ^b	1891.9±81.4 ^b	1702.1±97.3 ^d	1917.1±81.5 ^b
9-13	674.7±10.0 ^e	702.2±11.9 ^{cd}	723.5±14.7 ^c	839.1±20.8 ^a	746.6±23.3 ^b	755.9±18.7 ^b	680.6±17.6 ^{de}	766.5±19.7 ^b
0-13	3455.6±78.2 ^d	3509.6±93.7	3635.0±99.5 ^c	4612.1±91.9 ^a	3749.9±91.2 ^{bc}	3813.7±98.1 ^b	3481.7±97.0 ^d	3809.6±87.2 ^b

a, b means in the same row with different superscripts differ significantly ($P < 0.01$).

Similar results were obtained by Lai-Mingkueipolasek *et al.* (2003) who found that the highest weight gain of Peking ducks was obtained when they fed on low protein diets supplemented with lysine and methionine during the period from 0 to 3 weeks of age. Also, Leeson *et al.* (1982) reported that 18% crude protein during the finishing period was adequate for maximum weight gain of ducks.

Feed Conversion ratio (g feed/g gain)

Statistical analysis showed that feed conversion of ducklings was significantly ($P < 0.01$) better (1.34) when the diet contained P2 (20%) with Lys2 (1.10) and TSAA1 (0.93%) during the period 0-3 weeks of age. However, feed conversion of ducklings fed on diets contained P1 with Lys2 and TSAA2 was significantly ($P < 0.01$) the best (3.13) as compared to the other treatments during the whole experimental period (0-13 wk.) as shown in Table 5. Similar results were obtained by leclercq and De-carville (1989) who found an improvement in feed conversion of male Muscovy ducklings fed on a diet contained 17% as compared to those fed on 13% crude protein during the finishing period. Ibrahim (2004) showed that feed conversion of Muscovy ducklings was improved when diet contained low methionine (0.60%) level as compared with high methionine level (0.70%) during the finishing period.

Protein efficiency ratio (PER)

Data reflecting the average PER values are shown in Table (6). The PER values of ducklings was increased (2.13) significantly ($P < 0.05$) when ducklings fed on a diet contained P2 with Lys2 and TSAA2 during the period from 0 to 4 weeks of age. However, TSAA1 (0.93%) had improved significantly ($p < 0.01$) the PER of ducklings during the period 0 to 5 weeks of age. During the periods 5-9, 9-13 and 0-13 weeks of age, ducklings fed on diets contained P2 (16%) with Lys2 and TSAA2 had significantly ($P < 0.01$) the highest PER as compared to the other treatments group, it may reflect the effects of amino acids imbalances which reduced the efficiency of protein utilization, (D'Mello, 1994). Similar results were obtained by Sonbol, et al., (2001). Who reported that ducklings fed lower protein diets during the starting and finishing periods had significantly higher PER.

Sonbol and Habeeb (1991) showed that PER was improved when broiler diet supplemented with 0.10% methionine.

Carcass traits

Results in Table 7 showed that higher significant ($P < 0.05$) blood percentage was recorded for ducklings that fed on a diet contained P2 with Lys1 and TSAA2. The highest significant ($p < 0.01$) feather and giblets percentage was recorded for ducklings fed on a diet contained P2 with Lys2 and TSAA1. However, ducklings fed on diets contained P1 with Lys2 and TSAA1 had significantly ($p < 0.05$) the highest carcass percentage being 72.41. Similar results were obtained by Dean (1977) who reported that adding methionine to a corn-soya based diet, improved feathering of pekin ducks during the finishing period. Mazanowski *et al.* (1991). showed that increasing the protein level from 13 to 16% improved the carcass quality of ducks.

Table (5): Effects of protein, lysine and TSAA levels on feed conversion ($\bar{X} \pm SE$) of ducklings

Age (weeks)	P1				P2			
	Lys1		Lys2		Lys1		Lys2	
	TSAA1	TSAA2	TSAA1	TSAA2	TSAA1	TSAA2	TSAA1	TSAA2
0-3	1.83 ± 0.04 ^a	1.84 ± 0.04 ^a	1.82 ± 0.06 ^a	1.75 ± 0.02 ^b	1.79 ± 0.01 ^{ab}	1.58 ± 0.12 ^a	1.34 ± 0.01 ^d	1.51 ± 0.04 ^c
0-4	1.28 ± 0.01 ^d	1.42 ± 0.01 ^c	1.36 ± 0.04 ^c	1.25 ± 0.01 ^d	1.56 ± 0.02 ^d	1.49 ± 0.08 ^b	1.26 ± 0.02 ^d	1.18 ± 0.04 ^e
0-5	2.24 ± 0.02 ^d	2.76 ± 0.03 ^a	2.62 ± 0.12 ^b	2.12 ± 0.03 ^e	2.48 ± 0.02 ^c	2.49 ± 0.01 ^c	2.22 ± 0.01 ^d	2.28 ± 0.03 ^d
5-9	4.82 ± 0.15 ^a	4.10 ± 0.01 ^b	3.85 ± 0.04 ^c	3.23 ± 0.01 ^f	3.47 ± 0.11 ^{de}	3.63 ± 0.01 ^d	4.25 ± 2.23 ^b	3.40 ± 0.14 ^{ef}
9-13	6.30 ± 0.06 ^a	5.79 ± 0.02 ^c	5.33 ± 0.01 ^d	4.40 ± 0.03 ^g	4.69 ± 0.02 ^f	5.06 ± 0.10 ^e	5.98 ± 0.16 ^b	4.72 ± 0.05 ^f
0-13	4.29 ± 0.07 ^a	4.07 ± 0.05 ^b	3.77 ± 0.01 ^c	3.13 ± 0.01 ^f	3.42 ± 0.05 ^e	3.56 ± 0.01 ^d	3.95 ± 0.15 ^b	3.34 ± 0.07 ^e

a, b means in the same row with different superscripts differ significantly ($P < 0.01$).

Table (6): Effects of protein, lysine and TSAA levels on protein efficiency ratio ($\bar{X} \pm SE$) of ducklings

Age (weeks)	P1				P2			
	Lys1		Lys2		Lys1		Lys2	
	TSAA1	TSAA2	TSAA1	TSAA2	TSAA1	TSAA2	TSAA1	TSAA2
0-3	1.25 ± 0.05	1.25 ± 0.05	1.25 ± 0.05	1.30 ± 0.01	1.40 ± 0.01	1.57 ± 0.09	1.47 ± 0.38	1.66 ± 0.03
0-4	1.78 ± 0.01 ^c	1.61 ± 0.01 ^d	1.68 ± 0.06 ^d	1.83 ± 0.01 ^c	1.61 ± 0.02 ^d	1.68 ± 0.09 ^d	1.99 ± 0.03 ^b	2.13 ± 0.06 ^a
0-5	2.03 ± 0.01 ^d	0.65 ± 0.02 ^f	1.74 ± 0.08 ^e	2.15 ± 0.03 ^c	2.02 ± 0.01 ^a	2.01 ± 0.01 ^d	2.25 ± 0.01 ^a	2.19 ± 0.03 ^b
5-9	1.16 ± 0.04 ^e	1.36 ± 0.01 ^d	1.45 ± 0.02 ^c	1.73 ± 0.01 ^b	1.80 ± 0.06 ^{ab}	1.73 ± 0.01 ^b	1.48 ± 0.08 ^c	1.85 ± 0.08 ^a
9-13	0.88 ± 0.01 ^e	0.97 ± 0.01 ^d	1.05 ± 0.01 ^c	1.27 ± 0.01 ^b	1.34 ± 0.01 ^a	1.24 ± 0.03 ^b	1.05 ± 0.03 ^c	1.33 ± 0.02 ^a
0-13	1.25 ± 0.02 ^g	1.31 ± 0.02 ^f	1.41 ± 0.01 ^e	1.70 ± 0.01 ^{bc}	1.73 ± 0.03 ^b	1.67 ± 0.01 ^c	1.52 ± 0.05 ^d	1.79 ± 0.03 ^a

a, b means in the same row with different superscripts differ significantly ($p < 0.01$)

Table (7): Effect of protein, Lysine and TSAA levels on some carcass traits ($\bar{X} \pm SE$) of ducklings at 13 weeks of age

Carcass traits	P1				P2			
	Lys1		Lys2		Lys1		Lys2	
	TSAA1	TSAA2	TSAA1	TSAA2	TSAA1	TSAA2	TSAA1	TSAA2
Preslaughter weight (g)	3500.0 ± 318.9	3478.8 ± 355.9	3620.0 ± 362.0	3292.5 ± 198.9	3452.5 ± 268.6	3747.5 ± 390.8	3416.3 ± 319.9	3648.8 ± 490.6
Blood %	5.24 ± 1.10 ^b	4.83 ± 0.89 ^b	2.81 ± 0.02 ^c	5.24 ± 0.20 ^b	6.65 ± 0.26 ^a	6.98 ± 0.70 ^a	6.59 ± 0.95 ^a	6.38 ± 0.14 ^a
Feather %	5.80 ± 0.78 ^b	4.88 ± 0.47 ^c	5.14 ± 0.01 ^{bc}	3.88 ± 0.05 ^d	5.68 ± 0.64 ^b	3.57 ± 0.16 ^{de}	6.91 ± 0.72 ^a	3.07 ± 0.05 ^e
Giblets %	2.74 ± 0.10 ^e	3.39 ± 0.007 ^c	2.75 ± 0.02 ^e	3.93 ± 0.14 ^a	3.66 ± 0.04 ^b	3.51 ± 0.14 ^{bc}	3.97 ± 0.31 ^a	3.12 ± 0.21 ^d
Liver %	1.90 ± 0.09	2.01 ± 0.11	2.41 ± 0.01	2.40 ± 0.20	2.55 ± 0.13	2.07 ± 0.32	2.19 ± 0.31	2.05 ± 0.05
Carcass %	69.72 ± 1.08 ^c	70.23 ± 1.62 ^{bccc}	72.41 ± 0.14 ^a	65.00 ± 1.20 ^f	68.53 ± 0.91 ^d	66.89 ± 1.85 ^e	71.64 ± 1.68 ^{ab}	69.18 ± 2.08 ^{cd}

a, b means in the same row with different superscripts differ significantly (P < 0.01).

Feed cost

It could be concluded from Table (8) that highest feed cost "LE"/ kg gain during the experimental period was recorded by group fed P1 with Lys1 and TSAA1, while, the lowest feed cost/kg gain was recorded by group fed P1 with Lys2 and TSAA2. Similar results were obtained by Torkia (2002) who reported that feed cost significantly increased in hens fed a diet contained high protein level with higher lysine and methioine supplementation as compared to hens feed a diet contained high protein level with the lowest level of methioine and lysine. Rukmangadhan, *et al.* (1990) found that for optimal costs of Khalki Campbell ducks; diets must containe 22 and 18% crude protein in the starting and finishing period, respectively.

Table(8):Effects of protein lysine and TSAA. levels on feed cost/ kg gain

Treatments group	Feed cost /kg. gain	%
1	1.37	100
2	1.22	88.89
3	1.06	77.89
4	0.85	62.50
5	0.95	69.86
6	1.03	75.52
7	1.15	84.31
8	0.88	64.67

Conclusion

It could be concluded that; feeding Muscouy ducklings on diet contained low protein level (20%) supplemented with 0.10 % of methionine was sufficient to obtaine maximume growth during the period 0-4 weeks of age .At the5th week of age feeding Muscouy ducklings on diet contained 22% protein, 1.10% lysine and 0.83% TSAA. followed by 18% protein, 0.90% lysine and 0.70 TSAA. during the finishing period (5-13 wk.), improved body weight, weight gain and feed efficiency.

REFERENCES

Dean, W.F. (1977). Effect of feeding low protein grower- finisher rations, with and without supplemental lysine and methionine or weight gain, feed conversion, carcass composition and Feathering in Ducks. Duck Res. Progr. Rpt., Cornell Univ. Duck. Res. Lab. P. 20.

D'Mello. (1994). Amino acids imbalances, antagonisms, and toxicities. Page 63- 97 in: Amino acids in farm animal Nutrition. J.P.F. D'Mello, ed. CAB international, walling ford, oxon, ox10. 8 DE, UK.

Duncan, D.B. (1955). Multiple range and multiple F test Biometric 11: 1- 24.

Ibrahim,W.G. (2004). Studies on the nutritional requirements of Ducks. ph. D, Faculty of Agriculture, Zagazig University.

Lai-Mingkueipolasek, L.; J. Huang and K. Lin (2003). Effects of low deitary crude protein supple-mented with methionine and lysine on the growth performace and carcass traits of peking ducks. Journal of Taiwan Livestock research, 36: 39, 273 – 282, 19 ref.

- Leclercq, B. and H. De Carville (1989). Effects of protein concentration of finisher on growth and carcass quality of Muscovy ducklings. Water Fowl Production Proceeding the satellite conference for the XVIII world's poultry congress September 11 – 18 China 1989, 181 – 184.
- Leeson, S.; J.D. Summers and J. Proulx (1982). Production and carcass characteristics of the Duck. *Poultry Science*, 61 : 2456 – 2464.
- Mazanowski, A.; Z. Bernacki and D. Kokoszynski. (1991). Effect of feed mixtures containing different crude protein and metabolizable energy levels on meat traits of Ducks. *Zeszyty Naukowe, Zootechnix* 22 (177) 61 – 72. [Nutr. Abs. & Rev. 63 (4) 1983 (267)].
- National Research Council (1994). Nutrition requirement of poultry .9th, National Academy press. Washington DC., USA.
- Rukmangadhan, S.; R. Rajini and N. Ramamurthy (1990). Crude protein requirement of female khaki campbel ducklings *Indian Journal of Poultry Science* 25 (4) 301 – 303 [Nutr. Abs. & Rev. 62 (4) 1992 (258)].
- Snedecor, G. and W.G. Cochoran (1982). *Statistical methods* 6th Ed. Iowa state college press Ames, Iowa, U.S.A.
- Sonbol, S.M. and A. Habeeb (1991). Effect of dietary energy, methionine, choline and sulfate levels on productive performance and some blood constituents of broiler chicks during the growing periods. *Egypt. J. Appl. Science*. 6 (3): 310 – 328.
- Sonbol, S.M.; R.E. Khidr and Mona M. Hassan. (2001). Effect of feed mixtures containing different protein and Energy levels on Growth and Carcass characteristics of Ducks *Desert Inst. Bull.*, Egypt, 51, No. 2, 361 – 376.
- Torkia, A.A. (2002). Effect of dietary protein, lysine and methionine levels on the performance of laying hen ph. D., Faculty of Agriculture, Zagazig Univerity.
- Waibel, P.E.; G.W. Carlson and S.L. Noll (1995). Replacing protein in corn-soybean turkey diets with methionine and lysine. *Poultry sci.* 74: 1143 – 1158.
- Wang, Y.Z.; Z.R. Xus and J. Feng (2004). The effect of betaine and DL-methionine on growth performance and carcass characteristics in meat ducks. institute of feed science, Zhejiang University, No. 164. Qiatao North Road, Hangzho 4310029, China.

تأثير إضافة الليسين والميثيونين للعلائق منخفضة البروتين على معدل أداء البيط المسكوفي

شريف محمد سنبل^١ - جمال الدين عبد الرحمن^١ - رأفت السيد خضرا^١ و
منى محمد على حسن^٢

١- كلية الزراعة - جامعة الزقازيق.

٢- قسم تغذية الحيوان والدواجن - مركز بحوث الصحراء - المطرية - القاهرة.

أجريت هذه الدراسة على ٣٢٠ كتكوت بط مسكوفي عمر يوم ولمدة ١٣ أسبوع حيث قسمت الطيور عشوائيا إلى ٨ مجاميع تجريبية في تصميم احصائي ٢×٢×٢ (بكل مجموعة ٤٠ طائر قسمت إلى ٢ مكررة بكل مكررة ٢٠ طائر).

اشتملت المجاميع على أحد مستويين رئيسيين من البروتين مستوى مرتفع (٢٢ ، ١٨%) أو مستوى منخفض (٢٠ ، ١٦%) خلال مرحلتى (البادئ والناهي) على التوالي. كل مستوى من البروتين تم تقسيمه إلى مجموعتين فرعيتين تحوى إما مستوى ليسين مرتفع (١,٢٠ و ١,٠٥%) أو ليسين منخفض (١,١ و ٠,٩٠%) خلال مرحلتى (البادئ والناهي) على التوالي، ثم قسمت مجاميع الليسين التجريبية إلى تحت مجاميع اشتملت على إما مستوى مرتفع من الميثيونين+السستين (٠,٩٣ و ٠,٨٣%) أو مستوى ميثيونين+ سستين منخفض (٠,٨٠ و ٠,٧٠%) خلال مرحلتى (البادئ والناهي) على التوالي.

وتشير النتائج إلى أن تغذية البيط المسكوفي خلال مرحلة البادئ على عليقة منخفضة البروتين مع انخفاض فى محتواها من الليسين وارتفاع محتواها من الاحماض الامينية الكبريتية أدى إلى حدوث زيادة معنوية ($P < 0.01$) فى الوزن الحى، كما أنه عند التغذية على عليقة مرتفعة فى محتواها من البروتين ، منخفض فى الليسين ومنخفض فى محتواها من الاحماض الامينية الكبريتية أنت للحصول على أفضل وزن حى، مع أعلى زيادة فى الوزن، أعلى كفاءة من تحويل الغذاء خلال مرحلة الناهي ٥-٩ ، ٩-١٣ و ٠-١٣ أسبوع من العمر.

أوضحت المعاملات زيادة معنوية فى وزن الدم، الريش والأحشاء المأكولة ($p < ٠,٠١$) للمجموعة المغذاة على عليقة احتوت على ٢٠% بروتين، ١,١% ليسين و ٠,٩٣ ميثيونين+ سستين فى البادئ ثم ١٦% بروتين، ٠,٩٠ ليسين و ٠,٨٠ ميثيونين فى الناهي.

نستخلص من النتائج المتحصل عليها من هذه التجربة أن أفضل تركيبة لعلائق البيط المسكوفي والتي تحقق أعلى معدل نمو مع أقل تكلفة هي:

عليقة مرحلة البادئ (٠ - ٥ أسابيع):

وتحتوى على ٢٢% بروتين، ١,١% ليسين و ٠,٨٣ ميثيونين + سستين

عليقة مرحلة الناهي (٥- ١٣ أسبوع) :

وتحتوى على ١٨% بروتين، ٠,٩٠ ليسين ، ٠,٧٠ ميثيونين + سستين