

EFFECT OF SUBSTITUTING CONCENTRATE FEED MIXTURE BY CORN GRAINS FED WITH AMMONIATED RICE STRAW AND BENTONITE SUPPLEMENT ON:

1- FEED INTAKE, DIGESTION COEFFICIENTS AND SOME RUMEN PARAMETERS OF LACTATING FRIESIAN COWS.

Maklad Eman, H.M¹; H.M. Ead²; M.M. El-Shinnawy¹; A.M. Abdel-Khabir² and M.E. Sayed-Ahmed²

- 1. Animal Production Department, Faculty of Agriculture, Mansoura University, Egypt.**
- 2. Animal Production Research Institute, Agriculture Research Center, Dokki, Giza, Egypt.**

ABSTRACT

Twenty lactating Friesian cows with average live body weight ranging from 450 to 520 kg were used. All animals were in the 2nd to 4th lactation season. Animals were randomly distributed into five similar groups (G) (four animals each), to be fed formulated experimental rations which were as follows: G 1 : Control : 70 % concentrate feed mixture (CFM) + 30 % rice straw (RS), G 2 : ration 1 : 70 % CFM + 30 % ammoniated rice straw (ARS) + bentonite (B), G 3 : ration 2: 65 % CFM +5% ground corn grain (GCG) + 30 % ARS +B, G 4 : ration 3 : 60 % CFM +10% GCG + 30 % ARS +B and G 5: ration 4 : 55 % CFM +15% GCG + 30 % ARS + B. These proportions were chosen to achieve approximately isonitrogenous rations containing about 12-13% CP. The results obtained showed that digestibility of DM was significantly ($P<0.05$) higher with control and ration 4 than the other ones, while OM digestibility was significantly ($p<0.05$) higher with ration 4 than the other rations. The digestibility of CP, NFE, NDF, hemi. and cell. were significantly ($p<0.05$) higher when animals were fed the control and ration 4 than the other rations, while the CF was significantly ($p<0.05$) improved with increasing the replacement of corn grains to the highest level with ration 4. Nutritive values as total digestible nutrients (TDN), metabolizable energy (ME), net energy (NE) and relative feeding value RFV were significantly ($p<0.05$) higher with ration 4 than the other rations. Based on the mean value of ruminal pH, buffering capacity (BC), $\text{NH}_3\text{-N}$ concentration, the results showed that ruminal pH was decreased ($p<0.05$) when feeding ration 4 compared with ration 1 (7.09 and 7.38, respectively). The BC was higher ($p<0.05$) when feeding ration 2, 3 and 4 than the control or ration 1. The $\text{NH}_3\text{-N}$ concentration was significantly ($p<0.05$) increased by the replacement with corn grains. The VFA concentration was significantly ($p<0.05$) lower by increasing the replacement with corn grain in ration 3 and ration 4 than the other rations. The calculated effective neutral detergent fiber (eNDF) values ranged from 39.41 to 46.15 with different rations. The lowest eNDF value was recorded ($p<0.05$) for ration 4 compared with the other ones.

Keywords: Cows, ammoniated rice straw, effective NDF, relative feeding values, rumen fermentation.

INTRODUCTION

Fiber-rich, low protein forages and crop residues are the most abundant and appropriate feeds for ruminants in the tropics. Strategies to improve the utilization of these feeds should aim to provide supplements to

rumen microbes by selective feeding or chemical treatment usually with urea (Manda *et al.* 1999).

In ruminant nutrition it has long been recognized that the structural carbohydrates are essential to proper functioning of that the rumen and its microbes fermentation. Furthermore developments have been characterized as neutral detergent fiber (NDF), acid detergent fiber (ADF) and acid detergent lignin (ADL). Nowadays non-fibrous carbohydrates (NFC) receive attention and are divided and characterized in terms of soluble sugars and starches of different origin that differing in rumen degradation rates and hence, also in rumen resistance. It was also realized that the microbial population in the foregut may have requirements for nutrient and energy different from that of host animal. Until now, our knowledge of how we can stimulate a beneficial microflora in the gastro intestinal tract is very limited. It can be expected that some carbohydrates are the primary sources of proiotics as alternatives to antibiotics (Williams *et al.* 2001).

Nutritionists often go beyond and the measures of carbohydrates nutrition to include nutrients such as non-fiber carbohydrates (NFC), starch, effective NDF (eNDF), (Mertens,1997). Each of these nutrients look at a slightly aspect of carbohydrates nutrition. The total intake of rapidly fermentable carbohydrates is probably more important than the percentage of the carbohydrates (Oetzel and Nordlund, 1998). Ruminal adaptation to diets high in fermentable carbohydrates apparently has two key aspects-microbial adaptation (particularly the lactate-producing bacteria) and ruminal papillae length (longer papillae probate greater VFA absorption and thus lower ruminal pH) (Dirksen *et al.*, 1985).

Although it is clear that increased starch digestion in the total tract improves performance (Nocek and Tamminga, 1991), the optimal site of starch digestion is still unclear. Much of the work on alteration of the site of starch digestion has been done to compare different grain sources. Although this information is useful and practical, more specific questions about starch digestion in those studies are difficult to address because of confounding factors, such as protein and fiber concentration and their levels and source.

The main objective of this study was to evaluate, the effect of partially substituting of concentrate feed mixture (CFM) by yellow corn grains fed together with ammoniated rice straw supplemented with bentonite on nutrient or grains digestibility coefficients, nutritive values and rumen liquor parameters in lactating Friesian cows.

MATERIALS AND METHODS

The experimental work of this study was conducted at El-Karada Animal Production Research Station, Animal Production Research Institute, Agricultural Research Center, Ministry of Agriculture and Department of Animal Production, Fac. of Agric., Mansoura University during the years 2003-2005.

Twenty lactating Friesian cows from the herd station with body weight ranging from 450 to 620 kg were used. All animals were in the 2nd to 4th lactation season. Cows were randomly distributed into five similar groups (four each) . All groups were individually fed according to NRC (1984)

recommendations, based on their live body weight and milk yield (requirements for maintenance were 1% of LBW concentrate +1% of LBW roughage and requirements for lactation were ½ Kg concentrate per 1Kg milk yield). The experimental period lasted for 140 days (20 weeks).

The five experimental rations were formulated as follows:

- G 1 : Control : 70 % concentrate feed mixture CFM + 30 % rice straw RS.
- G 2 : ration 1 : 70 % CFM + 30 % ammoniated rice straw ARS +3%bentonite B.
- G 3 : ration 2 : 65 % CFM +5% ground corn grain GCG + 30 % ARS +3%B.
- G 4 : ration 3 : 60 % CFM +10% GCG + 30 % ARS +3%B.
- G 5 : ration 4 : 55 % CFM +15% GCG + 30 % ARS +3% B.

The experimental rations were formulated to be almost iso-nitrogenous and contained about 12.5-13.5% crude protein as recommended by Ørskov *et al.* (1972) to ensure maximal rate of fermentation in the rumen.

Bales of unchopped rice straw (30 tons) were arranged in three stacks, each consisting of three layers in cement pit. The stacks were covered with plastic sheet, leaving a free margin of fifty cm. plastic on each side to be covered with the soil. The stacks were injected by ammonia at the rate of 3% NH₃ weight basis through a hold metal pipe, then good covered each side of the plastic by the soil and left for three weeks to accomplish the reaction with ammonia. Then the stacks were opened and aerated before starting the feeding trial.

The CFM used contained wheat bran, undecorticated cotton seed meal, yellow corn, molasses, rice bran, limestone, soybean meal and salt. The yellow corn grains were spread to dry in the sun, it was coarsely ground using a local chopper usually available in the villages for grinding wheat grain.

Bentonite was obtained from El-Fayoum governorate, Egypt. It was added at the rate of 30 g/kg DMI according to Abd El-Baki *et al.* (1988) recommendation.

Management of feeding

The CFM was fed with or without ground corn grain (GCG) was offered to animals firstly at morning, while untreated or treated rice straw was given after consumption of the CFM + GCG mixture. Drinking fresh and clean water was available at all times.

Experimental animals and rations:

Five digestibility trials were conducted using three cows chosen randomly from each group to determine nutrients digestibility coefficients and nutritive values of the experimental rations. Each digestibility trial consisted of 15 days preliminary period followed by 7 days collection period. During the digestion trials, cows were fed their allowances according to the experimental assignment of each group. Acid insoluble ash (AIA) was used as a natural marker (Van Keulen and Young, 1977). Nutrients digestibility was calculated from the equations stated by Schneider and Flatt (1975) according to the following formula :

$$\text{Nutrient digestibility \%} = 100 - \frac{\text{AIA \% in feed}}{\text{AIA \% in feces}} \times \frac{\text{nutrient \% in feces}}{\text{nutrient \% in feed}} \times 100$$

Where: AIA is acid insoluble ash.

Feces samples were taken from the rectum of each cow twice daily with 12 hours interval at during the collection period of each trial and dried in a forced air oven at 65°C for 48 hours. Dried samples were composted for each cow and representative samples were taken, ground and kept for chemical analysis.

Samples of CFM, GCG, RS, ARS were taken at the beginning, middle and the end of each trial. At the end of the collection period composite samples were dried in a forced air oven at 65°C for 48 hours, then ground and kept for chemical analysis.

Chemical analysis and rumen parameters :

Proximate chemical analysis of CFM, GCG, RS, and ARS and feces were carried out as well as fiber fractions (NDF,ADF ADL, Hemic. and Cell.) according to the methods of AOAC (1990). Acid insoluble ash was determined according to method of VanKeulen and Young (1977).

Ruminal fluid samples were taken using rubber stomach tube before morning feeding and at 2, 4 and 8 hrs post- feeding from three animals of each treatment. The collected rumen fluid samples were filtered through three layers of gauze without squeezing for the determination of pH, buffering capacity, ammonia-N and total volatile fatty acids (TVFAs) concentration. Ruminal pH was estimated by pH meter (Orion Research, model 201 digital pH meter). Rumen liquor buffering capacity was determined according to Nicholson *et al.* (1963). Ruminal ammonia-N was determined according to Conway (1957). The TVFAs were determined by the steam distillation method as described by Warner (1964).

Statistical analysis:

The statistical analysis was performed using the least squares method described by Likelihood programme of SAS (1994). The obtained data for nutrient digestibility, nutritive value, effective NDF (eNDF), were subjected to one way analysis of variance according to the following model:

$$Y_{ij} = \mu + T_i + e_{ij}$$

Where: Y = Observation of the tested factor

μ = Overall mean

T_i = Treatment effect

e_{ij} = Error

The data of rumen liquor parameters were subjected to tow way analysis of variance according to the following model:

$$Y_{ijk} = \mu + T_i + P_j + TP_{ij} + e_{ijk}$$

Where: Y = Observation of the tested factor

μ = Overall mean

T_i = Treatment effect

P_j = Time effect

TP_{ij} = Interaction effect of the treatment x time

e_{ijk} = Error

The differences among means were carried out according to Duncan's New Multiple Range Test (Duncan, 1955).

RESULTS AND DISCUSSION

The CP content of ARS was practically doubled (8.19 vs 4.08 % respectively) compared with the untreated RS (Table 1). Generally, the summative proximate chemical analysis of the CFM, RS, ARS and GCG (Table 1) used to formulate the experimental rations were within the normal published ranges (Toykawa *et al.*, 1982; Yoon *et al.*, 1983; Cheva-Isarakul *et al.*, 1984, Ibrahim, 1987, Maklad, 1996 and Sayed-Ahmed, 2001). The CP of rice straw was marginal to deficient and not adequate for ruminal microbial breakdown of ingested forage. Ammonia treatment caused a reduction in neutral detergent fiber (NDF) content (65.5 Vs 73.3 %) and increased in neutral detergent soluble (NDS) (34.52 and 26.7%), compared with the untreated material. The proportion of fiber to cell soluble is a major determinant of energy availability in forages (Buxton and Redfearn, 1997). Also the energy contribution from structural and nonstructural carbohydrate must be considered (Nocek and Russell, 1988).

Corn grain was higher in neutral detergent soluble (NDS) % and rumen available carbohydrate (RAC) % than CFM (82.63 vs 59.66% and 89.38 vs 81.88% respectively).

The average daily dry matter intake by cows (Table 2) of each roughage and concentrate were in accordance with those of Hagemeister *et al.*, (1981) who reported that the greatest utilization of energy for microbial protein synthesis has been suggested when diets contained approximately 30 % roughage and 70 % concentrate. The average daily intake of total concentrate (concentrate feed mixture +corn grains) as the % of body weight (BW) was ranged from 2.6 to 3.29 of BW, while the roughage ranged was from 1.16 to 1.41% of BW. The total DM intake ranged from 3.82 to 4.70 % of BW. Such differences among groups may be due to the differences in animals average body weight and average daily milk yield. When cow eats more corn grains, ammoniated rice straw consumption was tended to be usually under 2% of BW (Wheeler , 2003) .

Concerning nutrient digestibility coefficients, the digestibility of DM was significantly ($P<0.05$) the highest with control and ration 4, but the differences among control, R1 and R2 were not significant while OM digestibility was significantly ($p<0.05$) the highest ration 4 composed with the other rations (Table 3) without significant difference among control , R1, R2 and R3 rations. These results are in agreement with those recorded by Fouad (2002) when feeding ammoniated rice straw with corn grain .

The highest CP digestibility was noticed for control ration and it did not significantly differ when compared with ration 1, 2 and 4, but the lowest value ($p<0.05$) was recorded with ration 3. The highest NFE digestibility ($p<0.05$) was with R4, compared with R1, 2 and 3 , but the differences were not significant among control , R1, R2 and R3 in this respect . Ration 4 showed the highest ($p<0.05$) CF digestibility compared with control, R1 and R3, but the differences among latter rations were not significant. The NDF digestibility was significantly ($p<0.05$) highest in R4 ration, compared with R1, R2 and R3, but the differences among latter rations were not significant. Likewise,

Table (1): The chemical composition of the ingredients and experimental rations.

Item	DM	Chemical composition (% as DM)															
		OM	CP	EE	CF	NFE	Ash	NDF	ADF	Hemi.	Cellu.	ADL	NFC*	UNDF ¹	ANDF ²	NDS ³	RAC ⁴
Ingredients																	
Concentrate feed mixture (CFM)	90.86	92.72	16.48	2.14	15.17	58.93	7.28	40.34	20.65	19.69	13.07	7.58	33.76	7.33	33.01	59.66	81.88
Corn grain (CG)	90.43	98.30	11.53	3.77	2.50	80.50	1.70	17.37	3.55	13.82	2.14	1.41	65.63	0.58	16.79	82.63	89.38
Rice straw (RS)	91.56	82.38	4.08	1.82	28.48	48.00	17.62	73.32	56.06	17.26	39.45	16.61	3.14	29.22	44.1	26.68	62.05
Ammoniated rice straw (3% ARS)	91.89	80.65	8.19	1.65	27.70	43.11	19.35	65.48	57.19	8.29	40.27	16.92	5.33	26.59	38.89	34.52	63.45
Experimental rations																	
30% RS+70%CFM	89.04	89.61	12.74	2.04	19.19	55.64	10.39	50.30	31.34	18.96	21.03	10.31	24.53	12.44	37.86	49.70	76.86
30%ARS+B+70%CFM	89.45	89.07	13.97	1.99	18.96	54.15	10.93	47.93	31.68	16.25	21.28	10.40	25.18	11.96	35.97	52.07	77.19
30%ARS+B+65%CFM+5%CG	89.46	89.35	13.73	2.08	18.40	55.14	10.65	46.79	30.84	15.95	20.74	10.10	26.75	11.34	35.45	53.21	77.87
30%ARS+B+60%CFM+ 10%CG	89.48	89.62	13.58	2.20	17.85	55.99	10.38	45.66	30.00	5.66	20.21	9.79	28.23	10.72	34.94	54.34	78.54
30%ARS+B+55%CFM+ 15%CG	89.49	89.96	13.27	2.24	17.50	56.95	10.04	44.42	29.02	15.40	19.57	9.45	30.03	10.07	34.35	55.58	79.27

* Non fiberous carbohydrates%= OM% - (CP%+NDF%+EE%), (Calsamiglia et al., 1995).

(1) UNDF : Unavailable NDF = NDF x 0.01 x ADL x 2.4 (Fox et al., 2000) .

(2) ANDF : Available NDF = NDF – UNDF

(3) NDS : Neutral detergent solubles = 100 – NDF

(4) RAC: Rumen available carbohydrate =
$$\frac{[0.9 (NDS - (\text{Protein} + \text{Lipid})) + (NDF \times \text{NDF availability})]}{[(NDS - (\text{Protein} + \text{Lipid})) + NDF]}$$
 (Nocek and Russell, 1988)

ration 4 showed the highest hemicellulose and cellulose digestibility coefficients, compared with R1, R2 and R3 rations. Similar result was found by Hill and West (1991).

In this respect, Mehrez (1995) reported that the addition of readily available substrates (in the form of cereals) to poor quality roughage, increased its rate of fermentation in the rumen through more colonization of rumen bacteria.

The nutritive values expressed as TDN, ME, NE and RFV were significantly ($p < 0.05$) the highest with ration 4, compared with the other rations. Higher starch intakes may promote rapid rates of passage from the rumen, causing a "washout" of grain from the rumen to the lower tract. The highest DCP values ($p < 0.05$) were noticed with ration 1, 2 and 4, without significant difference among them, compared with the values of control and R3 rations

Table (2): Average daily dry matter intake of concentrate, rice straw, ammoniated rice straw with or without bentonite supplementation by dairy cows

Items	Control	Ration 1	Ration 2	Ration	Ration
Average body weight	476.7	530.0	606.7	540.0	513.3
Roughage : concentrate	30 : 70	30 : 70	30 : 70	30 : 70	30 : 70
Bentonite g/h/d	0.00	630	690	700	740
Intake of DM from :					
Concentrate feed mixture (CFM) :					
Kg/h/d	13.30	14.56	15.03	13.77	13.24
As % BW	2.79	2.75	2.48	2.55	2.58
Corn grains (CG) :					
Kg/h/d	0.0	0.0	1.15	2.29	3.60
As % BW	0.0	0.0	0.19	0.43	0.71
Total concentrate :					
Kg/h/d	13.30	14.56	16.18	16.05	16.84
As %BW	2.79	2.76	2.67	2.99	3.29
Untreated rice straw (RS) :					
Kg/h/d	5.75	0.0	0.0	0.0	0.0
As % BW	1.21	0.0	0.0	0.0	0.0
3%Ammoniated rice straw (ARS) :					
Kg/h/d	0.0	6.30	7.01	6.95	7.20
As % BW	0.0	1.19	1.16	1.29	1.41
Total dry matter intake:					
Kg/h/d	19.05	20.87	23.18	23.01	24.04
As % BW	4.00	3.95	3.82	4.28	4.70

Table(3): Effect of feeding the experimental rations on the digestion coefficients and feeding values by cows.

Items	Control	Ration 1	Ration 2	Ration 3	Ration 4
Nutrient digestibility (%):					
DM	76.59 ^{ab} ± 0.93	72.24 ^{bc} ± 0.68	76.07 ^{bc} ± 2.69	71.93 ^c ± 1.31	80.86 ^a ± 0.76
OM	79.91 ^b ± 1.25	75.88 ^b ± 0.72	79.32 ^b ± 2.77	75.93 ^b ± 1.20	85.06 ^a ± 0.57
CP	85.56 ^a ± 0.65	83.49 ^a ± 0.68	83.47 ^a ± 2.23	77.05 ^b ± 1.75	85.05 ^a ± 0.66
EE	86.71 ± 0.28	83.46 ± 3.07	80.04 ± 2.04	80.90 ± 2.11	84.65 ± 1.13
CF	70.67 ^b ± 0.98	71.71 ^b ± 1.25	74.88 ^{ab} ± 1.52	71.31 ^b ± 2.03	77.78 ^a ± 1.08
NFE	81.56 ^{ab} ± 1.85	75.10 ^b ± 0.87	79.67 ^b ± 3.49	76.78 ^b ± 1.92	87.12 ^a ± 1.07
NDF	69.50 ^{ab} ± 2.09	61.28 ^c ± 1.20	67.08 ^{bc} ± 4.61	61.11 ^c ± 1.86	75.57 ^a ± 1.08
ADF	63.92 ± 1.38	59.70 ± 0.88	62.00 ± 4.41	60.46 ± 1.69	68.83 ± 1.36
Hemi.	78.71 ^{ab} ± 3.28	64.36 ^c ± 3.14	76.89 ^b ± 5.05	62.34 ^c ± 3.35	88.27 ^a ± 1.97
Cell.	85.44 ^{ab} ± 0.46	75.50 ^c ± 1.43	78.40 ^{bc} ± 4.82	76.67 ^c ± 1.37	86.19 ^a ± 0.91
ADL	20.01 ± 3.24	27.37 ± 1.81	28.32 ± 3.92	27.00 ± 3.13	32.88 ± 2.27
NFC	97.79 ± 0.69	98.87 ± 0.56	98.55 ± 0.23	98.92 ± 0.10	99.14 ± 0.13
Feeding value (%)					
TDN	73.81 ^b ± 1.13	69.67 ^b ± 0.65	72.78 ^b ± 2.49	69.89 ^b ± 1.10	78.36 ^a ± 0.47
DCP	10.90 ^{bc} ± 0.08	11.67 ^a ± 0.10	11.46 ^{ab} ± 0.30	10.39 ^c ± 0.24	11.27 ^{ab} ± 0.08
NFCI/DCPI	2.66 ^b ± 0.02	2.46 ^c ± 0.02	2.69 ^b ± 0.09	3.07 ^a ± 0.07	2.97 ^a ± 0.02
ME(Mcal/k	2.63 ^b ± 0.04	2.48 ^b ± 0.02	2.59 ^b ± 0.09	2.49 ^b ± 0.04	2.79 ^a ± 0.01
ME(Mj/Kg)	10.99 ^b ± 0.16	10.37 ^b ± 0.10	10.83 ^b ± 0.36	10.41 ^b ± 0.15	11.67 ^a ± 0.08
gDCP/ME	41.53 ^c ± 0.79	47.08 ^a ± 0.77	43.84 ^b ± 1.14	41.74 ^{bc} ± 0.36	40.40 ^c ± 0.23
NE(Mcal/K	1.69 ^b ± 0.03	1.59 ^b ± 0.02	1.66 ^b ± 0.06	1.59 ^b ± 0.03	1.80 ^a ± 0.02
DDM% ^{***}	68.20 ^{ab} ± 0.83	64.62 ^b ± 0.61	68.06 ^b ± 2.41	64.35 ^b ± 1.17	72.35 ^a ± 0.69
RFV ^{***}	211.58 ^b ± 6.43	197.96 ^b ± 5.91	201.63 ^b ± 7.46	213.45 ^b ± 6.47	264.02 ^a ± 12.4

a, b and c : Means within the same raw with different superscripts are significantly different (P<0.05).

* NE (Mcal / kg) = (TDN% x 0.0245) - 0.12 (NRC, 2001)

** DDM% of DM = 88.9 - 0.779 x (ADF% of DM) (Schroeder , 1996)

*** RFV = DMI x DDM / 1.29 (Schroeder , 1996)

The NFCI/DCPI ratio was significantly (p<0.05) higher with ration 3 and 4 than the other rations, but the difference was not significant between the former rations in this respect. The DDM% was higher (p<0.05) with ration 4 than R1, R2 and R3 rations. In this field, Knowlton *et al* (1998) reported that nitrogen digestibility in total tract was lower for diets containing dry corn than for diets containing higher moisture corn. Increasing in microbial N flow in feces and in microbial N produced in the large intestine for dry ground corn diets support the observation that starch digestion in the large intestine increases for dry ground corn diets. The increase in N lost as starch fermentation increased in the large intestine. Also, N from urea recycled from the blood to the large intestine may be incorporated into microbial N and

excreted (Ørskov, 1982). Recent studies with lactating dairy cows indicate that from 1 kg to nearly 5 kg of starch may disappear post ruminally in cows on high starch diets (Knowlton *et al* , 1998).

In general, increased flow of starch to the small intestine increased the quantity of starch digested. The grinding of corn may be decreased fecal starch flow and increased starch digestion in the total tract because of increased starch disappearance from the large intestine (Oliveira *et al*, 1995).

McAllister *et al* (1992) suggested that increasing the supply of starch post-ruminally to the small intestine may improve the feed efficiency by reducing the loss of energy through methane or heat . This would support the suggest of an advantage to increasing the availability of starch present in the small intestine . However , this may be offset by decreases in efficiency of starch digestion and absorption of glucose in the small intestine with increasing post ruminal starch passage (Owens *et al* 1986) .

Regarded rumen parameters, based on the mean value of ruminal pH (Table 4) the obtained results showed that it was decreased ($p < 0.05$) when feeding ration 4 compared with ration 1 (7.09 and 7.38 , respectively). In this field, Øroskov (1987) reported that the supplements fed with either untreated or treated straw diets are very important, since rumen bacteria which ferment or digest cellululosic feed are very sensitive to low rumen pH caused by supplementation . The pH values obtained in present study were within a normal range of 6-7 . Such range is suitable for the growth and activity of cellulytic bacteria (Prasad *et al* 1972) .

The mean value of BC was higher ($p < 0.05$) when feeding ration 2 , 3 and 4 than the control or ration 1. Maekawa *et al* (2002) showed that when feeding corn diets the total amount of fermentated acids produced in rumen is lower and more importantly, the rate of fermentation of acids produced could be considerably slower .

The mean values of the $\text{NH}_3\text{-N}$ concentrations was significantly ($p < 0.05$) increased by the replacement corn grains. In this respect, Bunting *et al* (1989) showed the amount ranging from non to over 80% of NH_3 from urea degradation is incorporated into bacteria N and the availability of energy is the major determinant of that percentage.

The mean values of the VFA concentrations, showed that the concentration was significantly ($p < 0.05$) decreased by increasing the replacement of corn grain in ration 3 and ration 4 than the other ones. Krause *et al* (2002) showed that the quantity of organic matter fermented in the rumen drives VFA production.

Nocek and Tamminga (1991) showed that the starch degradability in the rumen, rate and extent of VFA production, post-ruminal starch flow as digestion of starch can vary in response of the physiological status of the cow, grains type, growing conditions and both physical and chemical processing methods . The starch in similarly processed wheat, oats and barley is generally more degradable than the starch in corn. Most published reports on the site of starch digestion in lactating cows have involved duodenally but not ilealy cannulated cows. Some of the very high post-ruminal starch disappearances were observed when dry grains were fed (Oliveira *et al* 1995) might have been more associated with fermentations of

starch in the large intestine than digestion in the small intestine. Fermentation of starch in the large intestine result in loss of microbial N in feces, although the VFA produced are available for absorption and use by the cow.

Table (4) : Effect of feeding experimental rations on some rumen liquor parameters at different times of sampling.

Items		Control	Ration 1	Ration 2	Ration 3	Ration 4
Parameters	Hours					
PH-Values	0	7.39 ± 0.04	7.78 ± 0.13	7.63 ± 0.14	7.38 ± 0.06	7.49 ± 0.13
	2	7.22 ± 0.11	7.47 ± 0.18	7.40 ± 0.26	6.97 ± 0.19	6.81 ± 0.13
	4	7.20 ± 0.13	7.20 ± 0.10	7.35 ± 0.30	7.23 ± 0.09	7.31 ± 0.38
	8	7.49 ± 0.19	7.06 ± 0.15	6.88 ± 0.18	7.67 ± 0.30	6.76 ± 0.19
	Means	7.32 ^{ab} ± 0.07	7.38 ^a ± 0.10	7.32 ^{ab} ± 0.13	7.32 ^{ab} ± 0.11	7.09 ^b ± 0.14
Buffering capacity BC (ml eq/100ml)	0	9.23 ± 0.29	10.70 ± 0.06	14.87 ± 0.69	14.07 ± 0.90	12.50 ± 1.00
	2	11.43 ± 0.35	11.43 ± 0.30	14.03 ± 0.32	13.23 ± 0.97	12.47 ± 1.98
	4	9.23 ± 0.07	8.23 ± 0.38	12.47 ± 1.03	10.97 ± 0.89	11.47 ± 0.44
	8	7.80 ± 0.26	8.37 ± 0.44	10.83 ± 0.55	11.37 ± 1.49	10.03 ± 0.61
	Means	9.43 ^c ± 0.41	9.68 ^c ± 0.45	13.05 ^a ± 0.55	12.41 ^{ab} ± 0.61	11.62 ^b ± 0.58
NH ₃ -N (mg/100ml)	0	14.00 ± 1.01	12.04 ± 1.64	15.12 ± 2.80	17.27 ± 2.05	11.39 ± 3.09
	2	14.93 ± 0.81	15.68 ± 2.39	17.73 ± 1.31	17.17 ± 1.73	15.96 ± 0.97
	4	19.00 ± 0.73	18.67 ± 1.08	21.28 ± 1.32	23.80 ± 1.70	24.27 ± 1.82
	8	20.65 ± 1.46	18.67 ± 0.67	21.28 ± 2.26	21.75 ± 3.67	24.92 ± 1.54
	Means	17.14 ^{bc} ± 0.94	16.26 ^c ± 1.06	18.85 ^{abc} ± 1.17	20.00 ^a ± 1.35	19.13 ^{ab} ± 1.9
Total VFA's (ml eq/100ml)	0	9.68 ± 0.04	7.03 ± 0.43	8.52 ± 0.20	7.05 ± 1.08	5.35 ± 1.24
	2	8.95 ± 0.10	7.62 ± 0.72	8.22 ± 1.61	7.37 ± 0.88	7.75 ± 1.08
	4	7.88 ± 1.01	8.18 ± 0.89	7.95 ± 1.20	7.72 ± 0.86	6.07 ± 0.16
	8	8.67 ± 0.84	6.50 ± 0.35	7.67 ± 0.98	5.52 ± 1.04	7.42 ± 0.10
	Means	8.80 ^a ± 0.34	7.33 ^{bc} ± 0.33	8.09 ^{ab} ± 0.49	6.91 ^{bc} ± 0.48	6.65 ^c ± 0.57
%eNDF*		44.91 ± 0.57	46.15 ± 0.72	44.71 ± 1.66	44.69 ± 3.46	39.41 ± 1.81

a, b and c : Means within the same raw with different superscripts are significantly different (P<0.05).

* % eNDF = (pH - 5.425) / 0.04229 (Fox et al., 2000)

The calculated effective neutral detergent fiber eNDF values ranged from 39.41 to 46.15 with different rations. The lowest eNDF value was recorded (P<0.05) in ration 4 compared with the other rations. Effective NDF (eNDF) is the percentage of the NDF effective in stimulating chewing and salivation, rumination as rumen motility (Russell et al, 1992).

From the foregoing results, it could be concluded that corn grains can be replaced at the rate of 0.7% of body weight would supply adequate amounts of both digested dry matter (DDM) and CP. Therefore, corn would be the preferred supplement if the low quality forage supply was limited in diets of lactating cows. This replacement showed appropriate similar results in terms of nutrient digestibility coefficients, feeding values and rumen parameters as that obtained with the control ration

REFERENCES

- Abd El-Baki, S.M.S.; E.A. Hassona; M.S. Nowar and A.M. Aiad (1988). Improvement of urea utilization by the addition of bentonite, Kaolin and tafla clays in ruminants rations. *Zagazig Univ. J. Agric. Res.*, 15: 656.
- AOAC (1990). Association of Official Analytical Chemists. Official Methods of Analysis. 15th Ed., Washington DC .
- Bunting, L.D. ; J. A. Boling and C.T. Mackown (1989). Effect of dietary protein level on nitrogen metabolism in the growing bovine : 1. Nitrogen recycling and intestine protein supply in Calves . *J. Anim. Sci.*, 67 :810.
- Buxton, D.R. and D.D. Redfearn (1997). Plant limitation to fiber digestion and utilization . *J. Nutr.* 127 (5 supp. 1) :8145.
- Calsamiglia, S.; M.D. Stern and J.L. Firkins (1995). Effects of protein source on nitrogen metabolism in continuous culture and intestinal digestion *in vitro*. *J. Anim. Sci.*, 73: 1819.
- Cheva-Isarakul, Boonlom and Cheva- Isarakul , Boonserm (1984). Comparison of the intake and digestibility of different crop residues by sheep, cattle and buffaloes . In " The Utiliaztion of Fibrous Agicultural Resides as Animal Feeds " , pp.88-97 , editor P.T. Doyle . (School of Agriculture and Foresty , University of Melbourne , Parkville, Victoria) .
- Clark, J.H.; T.H. Klusmeyer and M.R. Cameron (1992). Symposium nitrogen metabolism and amino acid nutrition in dairy cattle, microbial protein synthesis and flows of nitrogen fractions to the duodenum of dairy cows . *J. Dairy Sci.*, 75 : 2304 .
- Conway, E.F. (1957). Microdiffusion Analysis and Volumetric Error. Rev. Ed. Lock wood, London.
- Dirksen, G.U.; H.G. Liebich and E. Mayer (1985). Adaptive changes of the ruminal mucosa and their functional and clinical significance. *Bovine Practitioner* 20:116.
- Duncan, D.B. (1955). Multiple Range and Multiple F Test. *Biometrics*, 11:10.
- Fouad, R.T. (2002). Effect of replacing corn by barley grains in concentrate feed mixture on the performance of sheep . *J. Agric. Sci. Mansoura Univ.*, 27(3): 1371-1380.
- Fox, D.G.; T.P. Tylutki; M.E. Van Amburgh; L.E. Chase; A.N. Pell; T.R. Overton; L.O. Tedeschi; C.N. Rasmussen and V.M. Durbal (2000). The net carbohydrate and protein system for evaluating herd nutrition and nutrient excretion. *Animal Science Mimeo* 213, Department of animal science, Cornell university, 130 Morrison Hall, Ithaca, New York 14853.
- Hagermeister, H.; W. Luppig. and W. Kaufmann. (1981). Microbial protein synthesis and digestion of the high yielding cows. (C.F. Clark *et al.*, 1992; *J. Dairy Sci.*, 75: 2304).
- Hill, G.M. and J.W. West (1991). Rumen protected fat in kline barley and corn diets for beef cattle : Digestibility, physiological and feed lot responses. *J. Anim. Sci.*, 69:3376.
- Ibrahim, B.K.M. (1987). Improving the feeding value of roughages .Ph. D. Thesis, Fac. of Agric. Mansoura univ.
- Knowlton , K.F. ; B.P. Glenn and R.A. ERDman (1998). Performance , ruminal fermentation and site of starch digestion in early lactation cows

- fed corn grain harvested and processed differently . J . Dairy Sci ., 81 : 1972.
- Krause, K.M.; D.K. combs and K.A. Beauchemin (2002). Effect of Forage particle size and grain fermentability in midlactation cows . 11. Ruminant PH and Chewing activity . J. Dairy Sci . 85 : 1947 .
- Maekawa, M.; K.A. Beauchemin and D.A. Christensen (2002) . Effect of concentrate level and feeding management on chewing activities , saliva secretion , and ruminal pH of lactating dairy cows . J . Dairy Sci. 85 : 1165.
- Maklad, E.H.M. (1996). Use of untraditional feed ingredients in feeding animals. Ph.D. Thesis, Fac. of Agric., Mansoura University.
- Manda, T.; M. Murai and A. Yamazaki (1999). Safe ammonia treatment system of cereal straws for ruminant feeding .Department of research planning and coordination, Hokkaido National Agricultural Experiment Station, Japanese.
- McAllister, T.A. ; L.M. Rode ; K. J. Cheng and J.G. Buchanan-Smith (1992). Effect of Formaldehyde treated barley or escape protein on ruminal environment and digestion in steers . Can . J. Anim . Sci . 72 : 317 .
- Mehrez, A.Z. (1995) Fermentation of feeds in the rumen. Proc. 5th Conf. Animal Nutrition, 69–68, Ismailia , Egypt .
- Mertens, D.R.(1997). Creating a system for meeting the fiber requirements of dairy cows. J. Dairy Sci.80:1463.
- Nicholson, J.W.G.; M.M. Cunningham and D.W. Friend (1963). The addition of buffers to ruminant rations. 4. The effect of additions of sodium bicarbonate, sodium propionate, limestone and cod liver oil on interrumen environment. Canadian J. Anim. Sci., 42: 309.
- Nocek, J.E. and J.B. Russell (1988). Protein and energy as an integrated system, relationship if ruminal protein and carbohydrate availability to microbial synthesis and milk production . J. Dairy Sci. 71 : 2070 .
- Nocek, J.E. and S. Tamminga (1991). Site of digestion of starch in the gastro intestinal tract of dairy cows and its effect on milk yield and composition. J.Dairy Sci. 74: 3598.
- NRC (National Research Council), (2001). Nutrient Requirements of Dairy Cattle, 7th rev. ed. National Academy Sci., Washington, DC.
- Oetzel, G.R. and K.V. Nordlund (1998). Effect of dry matter intake and feeding frequency on ruminal pH in lactating dairy cows. J.Dairy Sci.81(suppl.1) : 297. (Abstr.).
- Oliveira , J.S. ; J.T. Huber; J. M. Simas; C.B. Theurer and R.S. Swingle (1995) . Effect of sorghum grain processing on site and extent of digestion of starch in lactating dairy cows . J . Dairy Sci . 78 : 1318 .
- Ørskov, E.R.; C. Fraser and I. McDonald (1972). Digestion of concentrates in sheep. 4. The effects of urea on digestion, nitrogen retention and growth in young lambs. Br. J. Nutr., 27:491.
- Ørskov, E.R. (1982). Protein nutrition in ruminants. Acad. Press , New York,NY.
- Ørskov, E.R. (1987). The feeding of ruminants, principles and practice. First Published in U.K. by Chalcombe Publication.

- Owens, F.N. , R.A. Zinn and T.K. Kim (1986). Limits to starch digestion in the ruminant small intestine. *J. Anim. Sci.*, 63 : 1634 .
- Prasad, J.; S.S. Ahluwalia and B.P. Joshi (1972). Clinico-Biochemical studies in digestion in cattle and buffaloes. *Indian J. of Animal Sci.*, 42: 911.
- Russell, J.B. ; J.D. O'Connor ; D.G. Fot ; P.J. Van Soest and C.J. Sniffen (1992). A net carbohydrate and protein system for evaluating cattle diets .I. Ruminant fermentation . *J. Anim. Sci.* 70 : 3551 .
- SAS Institue (1994). *SAS / STAT R User's Guide : Statistics. Ver. 6.04, Fourth Edition SAS Institute Inc, Cary, NC.*
- Sayed-Ahmed. M. E. (2001). Effect of addition bentonite to ruminant feeds on utilization NPN. M. Sc. Thesis Fac. of Agric. Mansoura Univ.
- Schneider, B.H. and W.P.Flatt (1975). The evaluation of feeds through digestibility experiments. The university of Georgia Press Athens 30602.
- Schroeder. J.W. (1996). Quality forage for maximum production and return . NDSU Extension service , North Dakota State University .
- Toyokawa, Kojo. Takayasu. Ichior and Tsubomatsu, Kaizo (1982). Studies on the utilization of rice straw. Estimation of nutritive value by the method of artificial rumen . *Journal of Japanese Grassland Science* 28, 225–233 .
- Van Keulen. J. and B.A.Young (1977). Evaluation of acid insoluble ash as a natural marker in ruminant digestibility studies. *J. Anim. Sci.*, 44: 282.
- Warner, A.C.I. (1964). Production of volatile fatty acids in the rumen, methods of measurements. *Nutr. Abst. & Rev.*, 34. 339.
- Wheeler, B. (2003). Guidelines for feeding dairy cows. Ministry of Agriculture and Food , Government of Ontario , Canada .
- Williams, B.A.; M.W.A. Verstegen and S. Tamminga (2001). Fermentation in the monogastric large intestine its relation to animal health .*Nutrition Research Reviews* (in press).
- Yoon, C.S.; E.S. Choi; T.K. Oh; N.H. Lee; C.W. Kim and C.S. Kin (1983). Effect of aqueous ammonia treated rice straw on feed intake, nutritive value and rumen characteristics. *Korean J. Anim. Sci.*, 25: 613.

تأثير إحلال الأذرة جزئيا محل مخلوط العلف المصنع مع التغذية على قش الأرز المعامل بالأمونيا وإضافة البنتونيت على ١- الهضم وبعض قياسات الكرش في إبقار الفريزيان الحلابة

إيمان حنفي محمود مقلد* ، حسين محمد الشافعي عيد** ، محمد محمد الشناوي* ،
عبد الخبير محمد عبد الخبير** ومحمد السيد سيد احمد**
* قسم إنتاج الحيوان ،كلية الزراعة ، جامعة المنصورة.
** معهد بحوث الانتاج الحيواني ، مركز البحوث الزراعية

اجرى هذا البحث بهدف دراسة تأثير زيادة الطاقة المتاحةفى كرش الإبقار الحلابة بتغذيتها على
حبوب الأذرة المجروشة بإحلالها بنسب مختلفة (١٠ ، ١٥ ، ٢٠ %) محل مخلوط العلف المصنع على
كل من معاملات الهضم والقيمة الغذائية وبعض مقاييس الكرش حيث استخدم عشرون بقرة فريزيان تراوح
موسم حليبها بين الثاني والرابع وزعت فى خمس مجموعات لتغذي على احدى العلائق الآتية :

(مجموعة المقارنة) ٧٠ % علف مصنع + ٣٠% قش أرز غير معامل.
(عليقة أولى) ٧٠ % علف مصنع + ٣٠% قش أرز معامل بـ ٢% أمونيا+٣% بنتونايت.
(عليقة ثانية) ٦٥ % علف مصنع + ٥% حبوب أذرة مجروشة + ٣٠% قش أرز معامل بـ ٢% أمونيا +
٣% بنتونايت.
(عليقة ثالثة) ٦٠ % علف مصنع + ١٠% حبوب أذرة مجروشة + ٣٠% قش أرز معامل بـ ٢% أمونيا +
٣% بنتونايت.
(عليقة رابعة) ٥٥ % علف مصنع + ١٥% حبوب أذرة مجروشة + ٣٠% قش أرز معامل بـ ٢% أمونيا
+ ٣% بنتونايت.

وفى تم تكوين الخططات حتى تكون متماثلة تقريبا فى محتواها من البروتين الخام (١٢,٥-١٣,٥%) وهى
النسبة التى تلبي احتياجات البكتريا وتوفير الظروف المناسبة لتخمير المواد الخشنة بالكرش
وكانت أهم النتائج المتحصل عليها كما يلى:

١. أشارة نتائج التحليل الكيماوى لحبوب الأذرة مقارنة بمخلوط العلف المصنع إلى ارتفاع محتواه من
المركبات الكربوهيدراتية غير الليفيه (NFC) الكربوهيدرات المتاحة بالكرش (RAC) ، المركبات
النذائية (NDS) (٦٥,٢٣ ، ٨٩,٢٨ ، ٨٢,٢٣%) ، (٢٣,٧٦ ، ٨١,٨٨ ، ٥٩,٦٦) لكل من الأذرة
ومخلوط العلف المصنع على التوالى ، و كان محتوى العلف المصنع من كلا من البروتين الخام (CP
) ، مستخلص الألياف المتعادل المستفاد منه (ANDF) مرتفع مقارنة بحبوب الأذرة حيث كان فى
العلف المصنع ١٦,٤٨ ، ٢٣,٠١% بينما كان فى حبوب الأذرة ١١,٥٣ ، ١٦,٧٩% على التوالى، و
أيضا تشير نتائج التحليل الكيماوى إلى زيادة نسبة كلا من RAC ، NDS ، CP و انخفاض نسبة
الألياف الخام (CF) فى قش الأرز بت معاملته بالأمونيا ٣% .

٢. كان متوسط المأكول اليومي من المادة الجافة كنسبة مئوية من وزن الجسم الحى يختلف بين
المجموعات المختلفة حيث كان ٤ ، ٣,٩٥ ، ٣,٨٢ ، ٤,٢٨ ، ٤,٧٠% لكل من عليقة المقارنة وعليقة
١ و ٢ و ٣ على الترتيب .

٣. تحسنت معنويا معاملات هضم المادة العضوية CF ، NFC فى العلائق التى تحتوى على نسب
مختلفة من حبوب الأذرة وخاصة مع العليقة الأخيره (١٥%حبوب أذرة) وبالنسبة لمعاملات هضم
البروتين الخام فلم يكن هناك فروق معنويه باستثناء المجموعة الرابعة (١٠% حبوب أذرة) فلقت
إنخفاض بها بالمقارنة بالمجموعات الأخرى ، وبالنسبة لمعاملات هضم الهيميسيليلوز والدهن الخام
والنجنين فلم يظهر معها تحسن معنوى مقارنة بالمجموعات الأخرى.

٤. تحسنت معنويا قيمة مجموع المركبات الغذائية المهضومة (TDN) مع العليقة الأخيره (١٥%حبوب
أذرة) بالمقارنة بباقى المجموعات الأخرى.

٥. تحسنت قيم البروتين الخام المهضوم DCP (معنويا على مستوى ٥%) مع المجموعة الأولى بالمقارنة
بمجموعة المقارنة بينما إنخفضت معنويا مع المجموعة الثالثة بالمقارنة بباقى المجموعات الأخرى ،

- وكذلك فإن الطاقة الممتلئة والطاقة الصافية و الطاقة المتاحة (RFV) تزداد معنوياً مع العليقة الأخيرة (١٥%حبوب أذره) بالمقارنة بباقي المجموعات الأخرى.
٦. ازدادت قيمة الموضوم من المادة الجافة (DDM) مع العليقة الأخيرة (١٥%حبوب أذره) أو مجموعة المقارنه بالمقارنة بباقي المجموعات الأخرى .
 ٧. كانت متوسطات قيم تركيز أبون الأيدروجين (pH) فى سائل الكرش فى الحدود المناسبة لنشاط البكتريا وتأثرت معنوياً عند الأوقات المختلفة (قبل التغذية مباشرة وبعدها بـ ٢، ٤، و٨ ساعات) ولقد نقصت قيمة (pH) بالتقدم فى الوقت حتى وصلت لأقل قيمة عند الساعة الرابعة ثم ارتفعت مرة أخرى عند الساعة الثامنة مع كل المجموعات.
 ٨. وجد أن تركيز الأمونيا ($\text{NH}_3\text{-N}$) كان متشابها بين كل المجموعات وذلك قبل الأكل مباشرة ثم بعد ذلك وبمرور الوقت زاد التركيز حتى وصل إلى ذروته فى الساعة الثامنة مع كل المجموعات ، وكان المتوسط العام لتركيز الأمونيا ($\text{NH}_3\text{-N}$) يزداد مع العلائق التى تحتوى على نسب مختلفة من حبوب الأذره بالمقارنة بباقي المجموعات الأخرى.
 ٩. إنخفض تركيز الأحماض الدهنية الطيارة (TVFA's) معنوياً مع مجموعات إحلال الأذره وعلى وجه الخصوص مع العليقة الأخيرة (١٥%حبوب أذره) بالمقارنة بباقي المجموعات الأخرى.
 ١٠. كانت متوسطات قيم السعة التنظيمية للكرش (BC) منخفضه معنوياً مع مجموعة المقارنه و السجموعه الأولى مقارنة بالمجموعات الأخرى .
 ١١. إزدادت قيمة الألياف القابلة للتخمر (%eNDF) معنوياً عند التغذية على عنيقة المجموعة الأولى بالمقارنة بالمجموعة الرابعة وكانت القيم ٤٦,١٥ ، ٣٩,٤١ % على الترتيب .
- يستخلص من هذه الدراسة إنه يمكن فى الظروف المماثلة لهذه التجربة تغذية أبقار الفريزين الحلابة على علائق تحتوى على حبوب الأذره المجروشة بنسبة ١٥% من المادة الجافة الكلية الساكنة بجانب العلف المصنع وذلك عند التغذية على القش المعامل بالأمونيا مع إضافة البنتونايت أثناء التغذية حيث أن هذه الخلطات كانت الأفضل غذائياً فى صورة تحسن ملحوظ فى معاملات الأيضم والقيمة الغذائية وقياسات الكرش.