

EFFECT OF WATER REGIME AND POTASSIUM FERTILIZER LEVELS ON YIELD OF SOYBEAN AND WATER USE EFFICIENCY AT NORTH DELTA

Saied, M.M.; H.A. Shams El-Din; E.H. Omar and A.A.S. Gendy
Soils, Water and Environment Res. Inst., Agric. Res. Center, Egypt

ABSTRACT

Two field experiments were carried out at Sakha Agric. Res. Station in 2004 and 2005 seasons to study the effect of different soil moisture depletion, 35, 50 and 65% and potassium fertilizer levels; untreated 24 and 48 kg K₂O/fed. on soybean yield, some water relations and N, P, K concentrations of seeds.

The obtained results revealed that the irrigation at 50% soil moisture depletion surpassed the other two depletions (35 and 65%) in increasing the seed yield of soybean by 6.37 and 14.29% in the 1st season and 9.09 and 16.59 in the 2nd season. Application of 48, kg K₂O/fed. increased seed yield by 21.53 and 13.23% than application of 0.0 and 24 kg K₂O/fed. in the 1st season. While, in the 2nd season the seed yield increased by 21.43 and 10.36%.

The greatest amounts of water consumptive use were found to be 2007.43 and 1981.83 m³/fed. in the 1st and 2nd seasons respectively under 35% depletion.

The most suitable water applied throughout the two seasons are 2646.4 and 2490.7 m³/fed. in the 1st and 2nd seasons under irrigation 50% depletion of available soil moisture which produced the greatest seed yield.

The highest values of crop water use and field water use efficiencies were achieved from interaction between irrigation at 50% depletion of available soil moisture and application of 48 kg K₂O/fed. in both seasons.

INTRODUCTION

Soybean is the world's cultured plant as edible crop which is grown for its numerous utilizations and considered as important source of oil and protein. In Egypt increasing soybean productivity is urgently needed to meet the demands of both human and livestock. Proper irrigation and potassium fertilization are often critical factors in the success of the crop. As our share from Nile water is limited and the necessity of expanding cultivation in the newly reclaimed soil, there is a great need to make better use of our limited water resources.

The main problem have been observed during the summer season is the severe competition between the main crops growing in such period regarding irrigation water duty. It is of considerable importance to study the effect of rationalize the amount of irrigation water added to the unit cultivated area of land for significant without causing great reduction in yield. The effect of applied irrigation water on yield of soybean was studied by other investigators. Among them Sourour *et al.* (1988), Dornbos *et al.* (1989), El-Borai *et al.* (1993) and Shams El-Din *et al.* (1995), they reported that water stressed soybean plants produced fewer pods and lower weight of both 100 seed weight and seed yield.

Moreover El-Borai *et al.* (1993) measured 53 to 57% yield reductions in soybean resulting from water stress.

El-Awag *et al.* (1993), El-Refaie and El-Kabbany (1995) and El-Naggar *et al.* (1995) stated that the greatest yield obtained the most suitable water applied (increasing soil moisture tension) decreased consumptive use and water use efficiency which increased by decreasing the depletion of available soil moisture.

Eisa and Hanna (1997), concluded that the recommended treatment (seven irrigations) recorded both the highest amount of irrigation water applied and water consumptive use, while the lowest values were obtained from application of three irrigations in the two seasons. The best water use efficiency for seed production was obtained from irrigating soybean five times at 15, 30, 60, 75 and 105 days from sowing.

Also, substantial increases in soybean growth, yield quantity and seed quality were achieved by adding potassium fertilizer. In order to face the K-needs of grain legumes, soil and fertilizers might be able to supply K at a sufficient rate through all growth stages (Faucannier, 1980 and Abd El-Hady *et al.*, 1987).

Application of potassium fertilizer up to 36 kg K₂O/fed. had significant effect on the number of pods/plant, plant height and seed and straw yield of soybean (Derar and Saleep, 1997).

The objective of the present study was to investigate the effect of irrigation at different soil moisture levels and potassium fertilizer levels on soybean yields and its components, mineral composition in seeds and some water relations.

MATERIALS AND METHODS

Two field experiments were conducted at Sakha Agricultural Research Station in North Nile Delta region, during summer season of 2004 and 2005 respectively to study the effect of water regime and level of potassium fertilizer applied on seed yield of soybean, mineral composition of seeds and some water relations.

The soil of the experimental field was clayey in texture, non saline and non alkaline as shown in Table (1a). Soil chemical analysis in soil paste extract was determined according to Page (1982) and physical properties of soil were determined according to Klute (1986). While Table (1b) illustrates some soil-water features.

Table (1a): Some physical and chemical properties of the experimental field.

| Depth, cm | Particle size distribution | | | Texture class | EC dS/m | SAR | pH | Cations meq/L | | | | Anions meq/L | | | |
|-----------|----------------------------|--------|--------|---------------|---------|------|------|-----------------|----------------|------------------|------------------|------------------------------|-------------------------------|-----------------|------------------------------|
| | Sand % | Silt % | Clay % | | | | | Na ⁺ | K ⁺ | Ca ⁺⁺ | Mg ⁺⁺ | CO ₃ ⁻ | HCO ₃ ⁻ | Cl ⁻ | SO ₄ ⁻ |
| 0-30 | 28.0 | 29.1 | 42.9 | Clayey | 3.38 | 4.28 | 7.81 | 13.4 | 0.8 | 12.5 | 7.1 | 0.0 | 3.3 | 11.9 | 21.0 |
| 30-60 | 27.1 | 29.6 | 43.1 | Clayey | 3.6 | 4.91 | 7.93 | 16.1 | 0.7 | 13.0 | 8.5 | 0.0 | 3.3 | 20.4 | 14.6 |
| 60-90 | 27.9 | 28.7 | 43.4 | Clayey | 3.79 | 4.31 | 7.96 | 14.6 | 0.7 | 15.8 | 6.8 | 0.0 | 3.0 | 14.5 | 20.4 |

Table (1b): Soil moisture characteristics of the experimental field.

| Depth, cm | Field capacity % | Wilting point % | Available water % | Bulk density, g/cm ³ |
|-----------|------------------|-----------------|-------------------|---------------------------------|
| 0-15 | 43.9 | 23.96 | 20.04 | 1.24 |
| 15-30 | 39.0 | 21.2 | 17.8 | 1.36 |
| 30-45 | 37.0 | 20.11 | 16.89 | 1.39 |
| 45-60 | 36.2 | 19.67 | 16.53 | 1.47 |
| Average | 39.03 | 21.21 | 17.82 | 1.37 |

A split plot design with four replicates was used and the experimental treatments were as follows:

The main plots: (Water depletion)

D₃₅ : Irrigation at 35% depletion of soil available water.

D₅₀ : Irrigation at 50% depletion of soil available water.

D₆₅ : Irrigation at 65% depletion of soil available water.

Sub plots-(potassium fertilizer levels).

K₀ Untreated

K₁ Application of 24 kg K₂O/fed.

K₂ Application of 48 kg K₂O/fed.

Soybean (*Glycine max* L.) variety Giza 21 was sowed in the first week of June and harvested on the 20 and 25th Sep. 2004 and 2005 for the two seasons, respectively.

Fertilizers were applied to all plots as recommended practices at the rate of 20 kg N/fed. as urea (46% N) and 15 kg P₂O₅/fed. as calcium superphosphate (15.5% P₂O₅). Potassium fertilizer was added in the form of potassium sulphate (48% K₂O). Irrigation scheduling was based on soil moisture content which was gravimetrically determined in soil samples taken from successive layers before and after each irrigation of 15 cm depth down to 60 cm for scheduling irrigation.

Studied characters:

1. Yield and its components: The total plot area (10.5 m²) was harvested, 100 seeds weight and seed yield (ton/fed.) was determined . All the data were statistically analyzed according to Snedecor and Cochran (1967).
2. Plant height, number of tillers and pods per plant.
3. Mineral composition of seeds: Nitrogen, phosphorus and potassium were determined according to methods introduced by Jackson (1967).
4. Amount of irrigation water applied (W_a) to each treatments was measured using cutthroat flume 10 x 90 cm (Early, 1975).
5. Water consumptive use by growing crop was calculated in m³/fed. from sowing until harvesting according to the following equation:

$$Cu = \sum_{i=1}^{i=n} \frac{\theta_2 - \theta_1}{100} \times DB \times \frac{60}{100} \times 4200 \text{ (Israelsen and Hansen, 1962)}$$

Where:

Cu : Water consumptive use m³/fed.

n : Number of irrigations

- θ_2 : Soil moisture content (%) after irrigation
 θ_1 : Soil moisture content (%) before the next irrigation
DB : Bulk density (g/cm^3).
4200 : Area of feddan 0.42 ha.
6. Field water use efficiency (kg/m^3) = yield ($\text{kg}/\text{fed.}$)/amount of water applied ($\text{m}^3/\text{fed.}$), (Jensen, 1983).
7. Crop water use efficiency (kg/m^3) = yield ($\text{kg}/\text{fed.}$)/actual water consumption use ($\text{m}^3/\text{fed.}$), Jensen 1983.

RESULTS AND DISCUSSIONS

Seed yield of soybean:

A. Effect of soil moisture depletion levels:

Data in Table (2) show the effect of irrigation treatments on seed yield of soybean and yield components during the two successive seasons. The obtained results revealed that all estimated characters were significantly affected by the tested depletion treatments except for plant height which was higher under 35% depletion. Irrigation at 50% soil moisture depletion surpassed the other two depletion treatments (35 and 65%) in increasing the seeds of soybean by 6.37 and 14.29% in the 1st season and 9.09 and 16.59% in the 2nd season. The highest average values (1062.94 and 1096.23 $\text{kg}/\text{fed.}$) for seed yield, (4.42 and 5.25) for number of tillers, (88.75 and 77.75) for number of pods/plant and (18.39 and 17.33 gm) for weight of 100 seeds in the 1st and 2nd seasons, respectively were obtained under irrigation at 50% depletion of available soil moisture. The more droughted soybean plants which irrigated at 65% depletion of available water produced the lowest average seed yield and yield parameters in the two growing seasons. These results are in agreement with those of El-Borai *et al.* (1993) and Shams El-Din *et al.* (1995), they reported that water stressed soybean plants produced lower pods and lower weight of both 100 seed and seed yield.

The increase in seed yield under irrigation at 50% depletion of available soil water is mainly attributed to the suitable soil moisture in the root zone and to the presence of water supply continuously to plants in the different growth stages.

B. Effect of potassium levels:

Results in Table (2) show significant differences among levels of potassium fertilization for seed yield, plant height, number of tillers, number of pods per plant and weight of 100 seeds in both growing seasons. Increasing the K rate from 24 to 48 $\text{kg K}_2\text{O}/\text{fed.}$ increased seed yield and yield components in the two studied seasons. The high level of potassium fertilizer had the highest average values of these traits followed by the recommended level of 24 $\text{kg K}_2\text{O}/\text{fed.}$ These results show that 48 $\text{kg K}_2\text{O}/\text{fed.}$ may be gained the balance nutrition in the plants and reflected on the yield and its components. In this respect, Derar and Saleep (1997) stated that application of potassium fertilizer up to 36 $\text{kg K}_2\text{O}/\text{fed.}$ had significant effect on the number of pods/plant, plant height and seed and straw yields of soybean.

Table (2): Soybean yield and yield components as influenced by moisture depletion and potassium fertilization.

| Treatments | Seed yield, kg/fed. | Plant height, cm | Number of tillers | Number of pods/plant | Weight of 100 seeds, gm | Seed yield, kg/fed. | Plant height, cm | Number of tillers | Number of pods/plant | Weight of 100 seeds, gm |
|------------------------|---------------------|------------------|-------------------|----------------------|-------------------------|----------------------|------------------|-------------------|----------------------|-------------------------|
| | First season (2004) | | | | | Second season (2005) | | | | |
| Depletion level | | | | | | | | | | |
| D ₃₅ | 995.24 | 97.75 | 3.75 | 75.58 | 17.26 | 996.62 | 100.01 | 3.83 | 57.58 | 16.32 |
| D ₅₀ | 1062.94 | 92.01 | 4.42 | 88.75 | 18.39 | 1096.23 | 95.98 | 5.25 | 77.75 | 17.33 |
| D ₆₅ | 911.05 | 91.83 | 3.00 | 47.08 | 16.45 | 914.32 | 91.08 | 3.33 | 51.33 | 15.12 |
| F-test | ** | * | ** | ** | ** | ** | * | ** | ** | ** |
| L.S.D. 0.05 | 9.52 | 5.14 | 0.32 | 8.96 | 0.79 | 8.75 | 8.39 | 0.75 | 8.09 | 0.73 |
| 0.01 | 13.18 | | 0.48 | 13.09 | 1.13 | 12.18 | | 1.14 | 11.30 | 1.05 |
| Potassium fert. | | | | | | | | | | |
| K ₀ | 878.47 | 83.83 | 3.17 | 47.33 | 16.92 | 880.62 | 83.08 | 3.25 | 58.02 | 16.11 |
| K ₂₄ | 971.32 | 89.08 | 3.67 | 78.47 | 17.29 | 1005.19 | 87.83 | 4.25 | 60.58 | 16.22 |
| K ₄₈ | 119.44 | 108.67 | 4.33 | 85.17 | 17.88 | 1121.36 | 115.75 | 4.92 | 68.08 | 16.45 |
| F-test | ** | ** | ** | ** | ** | ** | ** | ** | * | * |
| L.S.D. 0.05 | 10.96 | 4.22 | 0.37 | 6.43 | 0.72 | 9.77 | 8.79 | 0.31 | 8.83 | 0.33 |
| 0.01 | 15.02 | 5.78 | 0.51 | 8.81 | 0.95 | 13.39 | 12.05 | 0.42 | | |
| Interaction | | | | | | | | | | |
| D x K | ** | ** | ns | ** | ** | ** | ** | ns | ** | ** |

The interaction between irrigation and potassium fertilizer levels gave a high significant effects on seed yield and yield components except number of tillers in the two studied seasons.

These results are in coincide with those of Faucannier (1980), Abd El-Hady *et al.* (1987) and Derar and Saleep (1997).

Some water relations:

1. Water consumptive use:

The water consumptive use values for different treatments of soil moisture depletion and potassium fertilizer levels are presented in Table (3). The greatest amount of water consumptive use were found to be 2007.43 and 1981.83 m³/fed. in the first and the second season, respectively under 35% depletion of available soil moisture. It was followed by 50% depletion, while the lowest values (1785.8 and 1780.1 m³/fed.) for both seasons were obtained under 65% depletion available soil moisture. As for the effect of potassium fertilizer levels, data in Table (3) show that as potassium fertilizer increased from 24 to 48 kg K₂O, the water consumptive use was increased under different soil moisture depletion. Increasing water consumptive use as application of potassium fertilization increased is mainly due to its effect on increasing transpiration.

Obtained results agreed with those of El-Naggar *et al.* (1996) who reported that increasing K₂O from 0 to 72 kg/fed. increased water consumptive use in maize and soybean.

2. Amount of water applied:

The amount of irrigation water delivered to each treatment was measured and recorded in Table (3). The data indicated that the irrigation at 35% depletion of available soil moisture under application of 48 kg K₂O/fed. received the highest amount of irrigation water followed by interaction between irrigation at 50% depletion and application 48 kg K₂O/fed. for both seasons. On the other hand, the irrigation at 65% depletion of available soil moisture under untreated treatments (K₀). received the lowest amount of irrigation water. It can be seen that from Table (3) the most suitable water applied throughout the two seasons are 2646.6 and 2490.7 m³/fed. in the first and the second seasons which produced the greatest seed yield. The listed values are obtained as irrigation events was based on 50% depletion from available water and 48 kg K₂O/fed. In addition, these values of irrigation water consists of watering excluding the sowing one. Similar results were obtained by El-Awag *et al.* (1993) and El-Naggar *et al.* (1996).

3. Field and crop water use efficiencies:

Data in Table (3) indicated that the values of field water use efficiency were lower than that of crop water use efficiency. Both efficiencies are dealing with the capability of each unit of both water consumed and water applied in producing the crop yield.

It was noticed that the highest values of crop water use and field water use efficiencies were achieved from combination between irrigation at 50% depletion of available soil moisture and application 48 kg K₂O/fed. in both seasons.

Table (3): Some water relations for soybean as affected by soil moisture depletion levels and potassium fertilization.

| Treatments | First season, 2004 | | | | | | Second season, 2005 | | | | | |
|-----------------|-------------------------|------------------------|---------------------|---|-------------------------------------|-------------------------|-------------------------|---------------------|---------------------------|--------------------------|-------------------------|-------------------------|
| | Soil moisture depletion | Potassium fert. levels | Seed yield, kg/fed. | Water consumptive use, m ³ /fed. | Water applied, m ³ /fed. | CWUE, kg/m ³ | FWUE, kg/m ³ | Seed yield, kg/fed. | WCU, m ³ /fed. | Wa, m ³ /fed. | CWUE, kg/m ³ | FWUE, kg/m ³ |
| D ₃₅ | K ₀ | 875.43 | 1948.4 | 2824.6 | 0.45 | 0.31 | 878.73 | 1939.6 | 2492.4 | 0.45 | 0.35 | |
| | K ₂₄ | 964.3 | 1990.3 | 2905.4 | 0.48 | 0.33 | 965.45 | 1981.5 | 2546.23 | 0.49 | 0.38 | |
| | K ₄₈ | 1146.0 | 2083.6 | 3041.2 | 0.55 | 0.38 | 1145.68 | 2024.4 | 2601.4 | 0.57 | 0.44 | |
| | Mean | 995.24 | 2007.43 | 2923.73 | 0.5 | 0.34 | 996.62 | 1981.83 | 2546.6 | 0.5 | 0.39 | |
| D ₅₀ | K ₀ | 975.57 | 1871.5 | 2507.1 | 0.52 | 0.39 | 976.45 | 1846.6 | 2372.9 | 0.53 | 0.41 | |
| | K ₂₄ | 1031.25 | 1910.2 | 2578.5 | 0.54 | 0.4 | 1128.28 | 1902.5 | 2444.7 | 0.59 | 0.46 | |
| | K ₄₈ | 1182.0 | 1959.7 | 2646.6 | 0.6 | 0.45 | 1183.95 | 1938.3 | 2490.7 | 0.61 | 0.48 | |
| | Mean | 1062.94 | 1913.8 | 2577.4 | 0.55 | 0.41 | 1096.23 | 1895.8 | 2436.1 | 0.58 | 0.45 | |
| D ₆₅ | K ₀ | 784.4 | 1748.4 | 2272.0 | 0.45 | 0.35 | 786.68 | 1731.2 | 2224.6 | 0.45 | 0.35 | |
| | K ₂₄ | 918.4 | 1785.8 | 2356.2 | 0.51 | 0.39 | 921.85 | 1788.4 | 2298.1 | 0.51 | 0.40 | |
| | K ₄₈ | 1030 | 1823.2 | 2424.6 | 0.56 | 0.42 | 1034.45 | 1820.7 | 2339.6 | 0.57 | 0.44 | |
| | Mean | 910.93 | 1785.8 | 2350.93 | 0.51 | 0.3 | 914.33 | 1780.1 | 2287.4 | 0.51 | 0.4 | |

While the lowest values were obtained from interaction between irrigation at 35% depletion of available soil-moisture without K application

The clear increase in crop and field water use efficiencies values could be attributed to the higher yield of soybean.

The present results are in general agreement with those obtained by El-Refaie and El-Kabbany (1995) and El-Naggar *et al.* (1995).

Mineral composition of seeds:

Data in Table (4) show the effect of available soil moisture depletion levels and potassium fertilizer levels on mineral composition of seeds.

Table (4): Contents of NPK (%) in seeds of soybean as affected by soil moisture depletion levels and potassium fertilizer rates.

| Irrigation level | Potassium fert. | First season | | | Second season | | |
|------------------|-----------------|--------------|------|------|---------------|------|------|
| | | N | P | K | N | P | K |
| D ₃₅ | K ₀ | 3.00 | 0.63 | 0.78 | 3.14 | 0.61 | 0.76 |
| | K ₂₄ | 3.10 | 0.66 | 0.90 | 3.18 | 0.67 | 0.93 |
| | K ₄₈ | 3.20 | 0.66 | 0.93 | 3.23 | 0.69 | 0.96 |
| | Average | 3.10 | 0.65 | 0.87 | 3.18 | 0.66 | 0.88 |
| D ₅₀ | K ₀ | 3.08 | 0.58 | 0.86 | 3.11 | 0.53 | 0.84 |
| | K ₂₄ | 3.16 | 0.60 | 0.98 | 3.14 | 0.62 | 1.02 |
| | K ₄₈ | 3.22 | 0.64 | 1.01 | 3.26 | 0.66 | 1.13 |
| | Average | 3.15 | 0.61 | 0.95 | 3.17 | 0.60 | 1.00 |
| D ₆₅ | K ₀ | 3.22 | 0.54 | 0.90 | 3.25 | 0.58 | 0.88 |
| | K ₂₄ | 3.30 | 0.58 | 1.06 | 3.35 | 0.63 | 1.11 |
| | K ₄₈ | 3.34 | 0.62 | 1.12 | 3.38 | 0.65 | 1.16 |
| | Average | 3.29 | 0.58 | 1.03 | 3.33 | 0.62 | 1.05 |
| K average | K ₀ | 3.10 | 0.58 | 0.85 | 3.17 | 0.57 | 0.83 |
| | K ₂₄ | 3.19 | 0.61 | 0.98 | 3.22 | 0.64 | 1.02 |
| | K ₄₈ | 3.25 | 0.64 | 1.02 | 3.29 | 0.67 | 1.08 |

Data indicated that nitrogen and potassium percentage in seeds of soybean increased with increasing the water stress for both seasons. On the other hand, phosphorus percentage decreased with decreasing water stress in both seasons.

The highest mean value of nitrogen (3.29 and 3.33%) and (1.03 and 1.05%) for potassium was obtained at irrigation 65% depletion of available soil moisture in the 1st and 2nd seasons respectively. On the contrary to this, the highest mean values of phosphorus percentage 0.65 and 0.66% were obtained with irrigation at 35% depletion of available soil moisture in both seasons.

The increase in phosphorus percentage in seeds may be due to the effect of high soil moisture content on the solubility of phosphorus compounds in soil. Concerning the effect of potassium fertilizer levels on N, P and K, data revealed that application of K up to 48 kg K₂O/fed. increased the N, P and K in seeds of soybean.

Barbariuck (1985) pointed out that the increase of N uptake as a result of K application was not nutritional response to K but resulted from the suppression of N uptake.

Similar results were obtained by Abd El-Mawly (1987) who reported that K addition increased NPK in soybean plant.

Finally, to achieve the highest seed yield of soybean and its water use efficiency, it could be recommended by irrigation at 50% depletion of soil available water and application of 48 kg K₂O/fed.

REFERENCES

- Abd El-Mawly, S.E. (1987). Effect of potassium and phosphate application and water regime on yield and chemical composition of soybean. M.Sc. Thesis, Fac. Agric. Minia Univ.
- Abdel-Hady, A.H.; K.G. Asy and Y.H. Mohamed (1987). Effect of K fertilizer on the production of *Trifolium alexandrium* in Egyptian soils. Mediterranean Potash News No. 3.
- Barbarick, K.A. (1985). Potassium fertilization of alfalfa grown on a soil high in potassium. Agron. J. 77. 442.
- Derar, R.A. and S.R. Saleep (1997). Response of soybean plants to potassium and iron application. Menofiya J. Agric. Res. Vol. 22 (1997), No. 4: 1187-1196.
- Dornbos, D.L.; Jr., R.E. Mullen and R.M. Shibles (1989). Drought stress effects during seed fill period on soybean seed germination and vigor. Crop Sci., 29. 476-480.
- Early, A.C. (1975). Irrigation scheduling for wheat in the Punjab CENTO Scientific programme on the optimum use of water in agriculture, report No. 17, Lyallpur, Pakistan 3-5 March pp. 115-127.
- Eisa, M.S. and A.M. Hanna (1997). Frequency and timing of irrigation effect on yield and its components of two soybean cultivars. J. Agric. Sci. Mansoura Univ., 22(12): 4105-4113.
- El-Awag, T.I.; A.M. Hanna and I.M. El-Naggar (1993). Influence of bio- and mineral phosphate fertilization on soybean production and some water characters under different levels soil moisture. Egypt. J. Appl. Sci., 8(11): 575-594.
- El-Borai, M.A.; M.M. Radi; M.B. Habeeb; F.H. Shalaby and Nadia, A. El-Aidy (1993). Effect of some water stress regimes and plant population on yield and seed quality of soybean. J. Agric. Sci., Mansoura Univ., 18(8): 2198-2205.
- El-Naggar, I.M.; A.M. Hanna; E.A.Y. El-Kabbany and A.. El-Fawail (1995). Effect of soil moisture levels and different sources of nitrogen on physical and chemical properties of soil and maize production. M. J. Agric. Res., 20(3): 1237-1263.
- El-Naggar, I.M.; T.I. El-Awag and A.M. Hanna (1996). Production of some field crops as influenced by water regime and potassium application. Egypt. J. Appl. Sci., 11(9): 211-225.
- El-Refaie, M.M.A. and E.A.Y. El-Kabbany (1995). Effect of water deficient and nitrogen fertilization rate on some water relations and wheat yield. Menofiya. Agric., Res., 20(6): 2437-2454.
- Faucannier, D. (1980). Potassium requirements of grain legumes. Proc. 11th Conger. Inter. Potash Inst. Res. Topics No. 7.
- Israelsen, O.W. and V.E. Hansen (1962). Irrigation principles and practices. 3rd Ed. John. Wiley Sons, New York, p. 240-272.

- Jackson, M.L. (1967). Soil chemical analysis. Constable and Co. Ltd., London.
- Jensen, M.C. and J.E.E. Middleton (1965). Scheduling irrigation from pan evaporation. Circular 386 Washington Agricultural Experiment Station.
- Jensen, M.E. (1983). Design and operation of farm irrigation system. Am. Soc. Agric. Eng. Michigan USA.
- Klute, A. (1986). Methods of soil analysis. Part 1: Physical and mineralogical methods (2nd ed). Am. Soc. Agron. Inc. in Soil Sci. of Agron. Inc. Madison, Wisconsin, USA.
- Page, A.L. (1982). Methods of soil analysis. Part 2: Chemical and microbial properties (2nd ed.). Am. Soc. Agron. in Soil Sci. Soc. Am. Inc. Madison, Wisconsin, USA.
- Shams El-Din, H.A.; M.M. Saied; R.Y. Rizk and S.A. Abd El-Wahab (1995). Effect of soil moisture, salinity and inoculation on water use efficiency, nodulation, yield and seed mineral composition of soybean. J. Agric. Sci. Mansoura Univ., 20(9): 4177-4186.
- Snedecor, G.W. and W.G. Cochran (1967). Statistical methods, 6th ed. Iowa Univ. Press, U.S.A.
- Sorour, S.Gh.R.; M.H. Ibrahim and M.A. El-Borai (1988). Effect of irrigation regimes during different growth stages of soybean on yield, seed composition and water relations. Proc. 3rd Egyptian Conf., Agron. Kafr El-Sheikh, 5-7 Sept. Vol. 11: 203-213.

تأثير مستويات استنفاد الرطوبة الأرضية ومعدلات التسميد البوتاسي على محصول فول الصويا وكفاءة استخدام المياه بشمال الدلتا محمود محمد سعيد ، حسن على شمس الدين ، السعيد حماد عمر ، الجندي عبدالرازق سليمان جندي

معهد بحوث الأراضي والمياه والبيئة - مركز البحوث الزراعية - الجيزة - مصر

أقيمت تجربتان حقليتان بمحطة البحوث الزراعية بسخا في الموسمين الزراعيين ٢٠٠٤ ، ٢٠٠٥م لدراسة تأثير ثلاثة مستويات من استنفاد الرطوبة الأرضية الميسرة وهي ٣٥ ، ٥٠ ، ٦٥% وثلاثة معدلات من التسميد البوتاسي صفر ، ٢٤ ، ٤٨ كجم بوب/أفدان على محصول فول الصويا ومكوناته وبعض العلاقات المائية وعلى تركيز النيتروجين ، الفوسفور ، والبوتاسيوم في بذور فول الصويا. وقد أظهرت النتائج المتحصل عليها أن معاملة الري عند ٥٠% استنفاد من الرطوبة الأرضية الميسرة تفوقت على معاملة الري ٣٥% ، ٦٥% في زيادة محصول بذور فول الصويا بنسبة ٦,٣٧% ، ١٤,٢٩% في الموسم الزراعي الأول ، زيادة بنسبة ٩,٠٩% ، ١٦,٥٩% في الموسم الزراعي الثاني. أيضا إضافة البوتاسيوم بمعدل ٤٨ كجم بوب/أفدان أدى إلى زيادة محصول البذور بنسبة ٢١,٥٣% ، ١٣,٢٣% مقارنة بالكنترول بدون إضافة ومعدل الإضافة ٢٤ كجم بوب/أفدان في الموسم الزراعي الأول بينما كانت الزيادة في الموسم الزراعي الثاني هي ٢١,٤٧% ، ١٠,٣٦% مقارنة بنفس المعاملات المذكورة.

اتضح من النتائج أيضا أن أعلى القيم للاستهلاك المائي كانت ٢٠٠٧,٤٣ ، ١٩٨١,٨٣م/أفدان في الموسمين الأول والثاني مع المعاملة التي تروى عند فقد ٣٥% من الرطوبة الميسرة ، وأن أنسب كمية نضياء الري النضاف في الموسمين كانت ٢٦٤٦,٤ ، ٢٤٩٠,٧م/أفدان مع المعاملة التي تروى عند استنفاد ٥٠% من الرطوبة الأرضية الميسرة وهي التي أعطت أعلى محصول من بذور فول الصويا وأن أعلى القيم لكفاءات استخدام المياه تحققت من التفاعل بين الري عند ٥٠% استنفاد من الرطوبة الأرضية الميسرة وإضافة ٤٨ كجم بوب/أفدان في كلا الموسمين.