

## THE COMPOST QUALITY AND ITS USE AS A GROWING MEDIA

Mahmoud, E. K.\* and H. A. Salem\*\*

\* Soil and Water Sciences Dept., Faculty of Agriculture, Tanta University, Egypt.

\*\* Floriculture Dept., Faculty of Agriculture, Alex. University, Egypt.

### ABSTRACT

Compost quality is very important when composted material is used as a growing media. In this study, six different Compost types were collected from Delta Bio-Tec Company, Nile Company and Mahmoud (2003) to determine the physical and chemical characteristics, stability/maturity; and investigate the suitability of the Delta Bio-Tec composts as a growing media for vinca growth. The physical properties of the different Compost types except compost biosolid Compost (BC) were not close to those of an "ideal substrate" for plant growth. The compost (BC) was the highest in N and K contents, and the lowest in P content. The Delta Bio-Tec composts and Nile compost were not stabilized and still phytotoxic to cress seeds. Plant growth, indicated that vinca yields were greater for peat than for composts alone or composts mixed with peat. Plants were died in the compost chicken manure at the ratio of 1:1, 3:1 and 4:0 (compost : peat, v/v). The high salt concentration and low stability/maturity of the all compost types except compost (BC) were a major limitations for compost use in growing media. Application strategies for these composts must be designed to minimize salt and phytotoxic damage to the plants in growing media .

### INTRODUCTION

Composting is one method to increase the recycling of organic by-products generated in cities and the farm. Composting transforms organic by-products with undesirable properties into a final product, compost, that is a valuable commodity for soil improvement. High-quality composts are suitable for potting mixes, home gardens and amendments for food chain crop growth. High-quality composts are highly mature and stable and have consistent chemical and physical characteristics (Sikora and Sullivan, 2000). Compost stability and maturity are important factors affecting the successful application of compost for agricultural purposes (Inbar *et al.*, 1990 and Mathur *et al.*, 1993). Compost stability and maturity are comprehensive properties indicating the degree of organic matter decomposition and potential of phytotoxicity caused by insufficient composting (Wu and Ma, 2001). Application of unstable or immature compost may cause slow plant growth and damage crops by competing for oxygen or causing phytotoxicity to plants due to insufficient biodegradation of organic matter (Keeling *et al.*, 1994 and Hue *et al.*, 1995).

Peat is commonly used as a growing medium in horticulture. Increasing demand and rising costs for peat as potting medium in horticulture have led to search for high-quality, low cost substrates, i.e., organic waste materials such as bark, spent mushroom compost, leaf mould, ... etc. These organic materials have been introduced as an alternative to peat substrate in potting media after proper composting (Cull, 1981; Bik, 1983; Lohr *et al.*,

1984; Chen *et al.*, 1988 and Mahmoud, 2003). The objective of this study is to determine physical and chemical characteristics, stability, maturity of selected different types form compost; and assess the suitability of Delta Bio-Tec composts as a growing media for vinca growth.

## MATERIALS AND METHODS

Compost samples were collected from the Delta Bio-Tec Company, Nile Company; and Mahmoud (2003). The composition of feed-stock of different types form compost used in this study are shown in Table (1). Three to five samples were collected from each type of compost, placed in polyethylene bags to Lab. A subsample was air-dried and ground to pass through a 2-mm screen for chemical analyses.

**Table 1. Production and feed stocks of different types form compost used in this study.**

Compost	Production	Feed stock
VW	Delta Bio-Tec Company	Vegetable wastes
CW	Delta Bio-Tec Company	Cow wastes
CHW	Delta Bio-Tec Company	Chicken wastes
OSW	Delta Bio-Tec Company	Ostrich wastes
BC	Mahmoud	Biosolids + crop residue
CR	Nile Company	Crop residues

### Chemical and Physical Analyses

Ash content of air-dried samples was determined by heating (at 480°C for 8 h) in a muffle oven (NEYM-525). Organic matter was estimated by loss upon ignition at 480°C for 8 h (50% from O.M was considered organic C) (AFNOR, 1991). Dried and ground samples were wet digested with concentrated HNO<sub>3</sub>+HClO<sub>4</sub> (Jones and Case, 1990). Total heavy metals, Ca<sup>++</sup> and Mg<sup>++</sup>, were determined using atomic absorption (Perkin Elmer AA model 2380) and total phosphorus was measured colorimetrically using spectrophotometer. Total nitrogen was determined by using Kjeldahl method (Stevenson, 1982). Electrical conductivity and pH values were determined in a compost-water extract solution (1:10, v/v) using CMD 830 conductivity meter and pH Controller model 5997, respectively. Bulk density, particle density, total porosity, container capacity, and air space were determined for selected composts by methods of Inbar *et al.*(1986).

### Carbon Dioxide Evolution

Microbial respiration of compost sample, based on CO<sub>2</sub> evolution, was measured using a modified procedure of Iannotti *et al.*(1994). Approximately 5 g of screened sample with moisture content adjusted to 60% was sealed in a 0.5 L vessel along with a beaker containing a known volume of 1M NaOH solution. The samples were incubated at room temperature (22±3°C). Excess NaOH was titrated with 0.5M HCl through precipitation with barium chloride as barium carbonate.

### **Seed Germination Test:**

Germination test on compost samples was measured in triplicate using cress seeds (*Lepidium sativum*). Compost layers of 5 mm thickness from each type were laid in Petri dishes, covered with filter paper, and then soaked to water saturation for the percent germination test as described by Mathur *et al.* (1986).

### **Compost as growing Media**

A greenhouse trial was conducted to evaluate Delta Bio-Tec composts as growing media in Green Oasis nursery: 2.5 km after Abeer gate - Khorsheed - Agriculture Road, Alexandria. Vinca (*Catharanthus roseus*) growing in trays (cell size 5x5x3.2 cm deep) for peat substrate and transplanted into (13.5 cm x 12 cm deep) pots using chicken, cow, and vegetable composts with peat at a volume ratio as potting substrate: (compost : peat) of 0:1 (0% compost), 1:3 (25% compost), 1:1 (50% compost), 3:1 (75% compost), 1:0 (100% compost) and arranged in a randomized complete block with 9 replications. All treatments were fertilized and watered equally by PGMix (14+16+18). After 40 days, plants were removed from media and washed, dried at 70°C, and weighted.

## **RESULTS AND DISCUSSION**

### **Chemical Properties of Compost**

The average values of composition and micro-nutrient elements (Fe, Cu, Zn and Mn) are shown in Table (2) for the different types of compost. The composts pH values ranged from 7.3 to 8.35, which are considered suitable for supporting growth of many plant species (Dick and McCoy, 1993). Electrical conductivity (EC) values in water extracts from composts were variable, and are higher than 3 dS/m in the chicken and crop residues composts. Such values could pose problems for seed germination, particularly if used in potting media (He *et al.*, 1995). The C/N ratio was above 15 : 1 of all different types the composts, except the ostrich and chicken composts, that is higher than C/N ratio of 15 which temporary net N immobilization of nitrogen may be induced after soil application (Gilmour, 1998). The compost (BC) was the highest in N and K contents, and the lowest in P content.

Chicken compost contained much higher concentrations of all elements except Fe, 60 mg/kg (Cu), 397 mg/kg (Mn) and 187.5 mg/kg (Zn). This may be due to using these elements in feed materials for the chickens. On the other hand, the crop residues compost had the lowest contents of these elements. The average values of micro-elements of selected composts was lower than the pollutant concentrations (PC) regulated by USEPA (1993) for sewage sludge. For example, Zn in the present study ranged between (32.5-187.5 mg/kg) was lower than the 2800 mg/kg, and Cu (14-60 mg/kg) was below the 1500 mg/kg.

**Table 2: The average values of chemical composition and micro-nutrients for the selected composts.**

Parameter	OSW	VW	CW	CHW	CR	BC
pH (1:10)	8.35	7.30	7.80	8.00	7.50	7.40
EC (1:10)	1.91	0.21	2.43	6.00	4.11	0.60
dS/m	1.31	1.01	1.23	1.60	1.44	1.85
N %	0.51	0.49	0.56	1.83	0.63	0.48
P %	0.35	0.15	0.40	1.55	1.50	2.60
K %	8.10	46.30	29.2	13.47	17.30	16.40
C/N	21.26	93.47	71.94	43.11	49.90	61.00
O.M %	78.74	6.53	28.06	56.89	50.10	39.00
Ash %						
<b>Micro-nutrient elements (mg/kg)</b>						
Fe	2837.5	1155.0	2972.5	3115.0	3057.5	3415.0
Mn	145.0	12.5	62.5	397.5	277.5	141.0
Zn	52.5	32.5	35.0	187.5	65.0	107.5
Cu	42.5	50.0	22.5	60.0	40.0	14.0

**Physical Properties**

The bulk density and particle density of the compost (BC) were much lower than other types of compost (Table 3). The particle density of the ostrich and chicken composts had greater than other composts due to the presence of sand which was used as a bedding material. Total porosity decreased and as a result the bulk density/particle density increased in all compost types. The physical properties of all compost, types except compost (BC), were not close to those of an "ideal substrate" for plant growth and peat as suggested by de Boodt and Verdonck (1972) (Table 3).

**Table 3. Physical properties of peat and different types of compost**

Composts	Bulk density	Particle density	Total porosity	Container capacity	Air space
	g/cm <sup>3</sup>		%		
OSW	0.726	2.305	68.47	33.30	35.20
VW	0.473	1.593	70.30	36.80	33.40
CW	0.647	1.754	63.11	38.20	24.90
CHW	0.646	2.029	68.16	34.50	33.70
BC	0.192	1.841	89.50	74.30	15.20
Peat	0.177	1.563	88.67	73.10	15.60
Ideal substrate*	-	-	85.00	55.75	20.30

\* From de Boodt and Verdonck (1972).

**Compost Stability and Maturity**

Compost stability refers to the degree to which composts have been decomposed to more stable materials. Stability was tested by CO<sub>2</sub> evolution. Compost maturity refers to the phytotoxicity associated with compost. Compost phytotoxicity test was based on the seed germination.

### Carbon Dioxide Evolution

The average values of accumulated CO<sub>2</sub> evolution after 6 days of incubation for selected composts are given in Table (4). The accumulated CO<sub>2</sub> evolution values of Delta Bio-Tec composts were higher than Nile and Mahmoud composts. The compost chicken manure was higher than other composts in the accumulated CO<sub>2</sub> evolution. Wu *et al.* (2000) found that the CO<sub>2</sub> evolution was markedly reduced as compost stabilized. The results showed that Delta Bio-Tec composts and Nile compost were higher in microbial activity than Mahmoud compost. Based on CO<sub>2</sub> evolution, all the Delta and Nile composts were not on obvious stabilization process. The much greater differences in values imply that source material and composting time have a great impact on the compost stabilization process.

**Table 4. The average accumulated CO<sub>2</sub> evolution after 6 days of incubation and germination rate after 3 days.**

Compost	CO <sub>2</sub> evolution* g/kg	Germination rate* %
OSW	38.6 ± 1.6	10 ± 8
VW	35.8 ± 0.8	30 ± 6
CW	78.2 ± 4.2	20 ± 3
CHW	137.0 ± 1.0	0 ± 0
CR	20.6 ± 0.2	40 ± 5
BC	2.5 ± 0.2	100 ± 3

\* Mean of three replicates ±S.D.

### Seed Germination

The average germination rate for selected composts is presented in Table (4). Mahmoud compost had high germination rate (100%) which corresponded well to their overall low CO<sub>2</sub> evolution and EC (Table 2,4). In contrast, the germination rates for the Delta Bio-Tec and Nile composts were low (10-40%), indicating that the Delta Bio-Tec composts and Nile compost were still phytotoxic to cress seeds despite their high CO<sub>2</sub> evolution and EC values.


Many of the substances found in immature compost can result in the reduction in the seed germination rate with its magnitude depending on the source of waste material and composting process. For example, a variety of organic compounds, including short (Shiralipour *et al.*, 1997) and long-chain fatty acids (Sesay *et al.*, 1997), phenolic acids, salts, ammonia, and heavy metals (BBC Labs, 1999).

### Vinca Growth

Generally, vinca yields were greater for peat than for composts alone or composts mixed with peat (Fig. 1). Plant fresh and dry weights decreased with increasing compost decomposition rate. Vinca fresh weight and dry weight were greatest when composts the were mixed with peat at the ratio of 1:3, 1:1, 3:1 and lowest at 4:0 the ratio of (compost : peat, v/v). Plants were died in the compost of chicken manure at ratio 4:0, 3:1, 2:1 (compost : peat).

It possibly due to the effect of increased EC in the chicken manure. EC values above 3.5 dS/m are, generally, considered to be high to support plant growth in containers (Eames, 1977 and Lemaire *et al.*, 1985). Results obtained with vinca indicate that the Delta Bio-Tec composts were not superior to peat as a growing media. Flynn *et al.* (1995) and Mahmoud (2003) showed that the compost was superior to peat as a substrate, but the compost + peat mixture gave better results; especially at the ratio of 2:1 (v/v).

Fig. (1): Fresh weight and dry weight of the vinca (*Catharanthus roseus*) plants grown in peat, compost and compost + peat.



## Summary

In Egypt, the companies have begin in production of large quantities of composts, which are marketed as a growing media . The high salt concentration and low stability/maturity of composts were a major limitation for compost use in growing media. Application strategies for these composts must be designed to minimize salt hazard and phytotoxic damage to plants in potting media.

## REFERENCES

- AFNOR (1991). Matieres fertilisants et supports de culture, Recueil des norms Francaises, AFNOR, Paris, p. 713.
- BBC Laboratories, Inc. (1999). <http://www.bbc.labs.com>.
- Bik, A.R. (1983). Substrates in floriculture. Proc. 21<sup>st</sup> Int. Hortic. Congr., Hamburg, 11: 811-822.
- Chen, Y.; Y. Inbar and Y. Hadar (1988). Composted agricultural wastes as potting media for ornamental plants. Soil Sci., 145: 298-303.
- Cull, D.C. (1981). Alternatives to peat as container media. Organic resources in the U.K. Acta Hort., 126: 69-81.
- de Boodt, M. and O. Verdonck (1972). The physical properties of the substrates used in horticulture. Acta Hortic., 26: 37-44.
- Dick, W.A. and E.L. McCoy (1993). Enhancing soil fertility by addition of compost. p. 622-644. In: H.A.J. Hoitink and H.M. Keener (eds.). Science and engineering of composting: Design, environmental, microbiological and utilization aspects. Renaissance Publications, Worthington, Ohio.
- Eames, A.G. (1977). Could spent mushroom compost be used for container shrubs? Mushroom J., 52: 114. (Abstr.)
- Flynn, R.P.; C.W. Wood and E.A. Guertal (1995). Lettuce response to composted droiler litter as a potting substrate component. J. Amer. Soc. Hort. Sci., 120 (6): 964-970.
- Gilmour, J.T. (1998). Carbon and nitrogen mineralization during co-utilization of biosolids and composts. p. 89-112. In S.L. Broen *et al.*(ed.) Beneficial co-utilization of agricultural, municipal and industrial by-products. Kluwer Academic Publishers, Dordrecht, The Netherlands.
- He, X.T.; T.J. Logan and S.J. Traina (1995). Physical and chemical characteristics of selected U.S. municipal solid waste composts. J. Environ. Qual., 24: 543-552.
- Hue, N.V. and J. Liu (1995). Predicting compost stability. Compost Sci. Util., 3: 8-15.
- Iannotti, D.A.; M.E. Grebus; B.L. Toth; L.V. Madden and H.A.J. Hoitink (1994). Oxygen respirometry to assess stability and maturity of composted municipal solid waste. J. Environ. Qual., 23: 1177-1183.
- Inbar, Y.; Y. Chen and Y. Hadar (1986). The use of composted separated cattle manure and grape more as peat substitute in horticulture. Acta Hortic., 178: 147-154.

- Inbar, Y.; Y. Chen and Y. Hadar (1990). Humic substances formed during the composting of organic matter. *Soil Sci. Soc. Am. J.*, 54: 1316-1323.
- Jones, J.B. and V.W. Case (1990). Sampling handling and analyzing plant tissue samples. p. 389-427. In R.L. Wasterman (ed.) *Soil testing and plant analysis*. 3<sup>rd</sup> ed. SSSA Book Series 3. SSSA, Madison, WI.
- Keeling, A.A.; I.K. Paton and J.A.J. Mullett (1994). Germination and growth of plants in media containing unstable refuse-derived compost. *Soil Biol. Biochem.*, 26: 767-772.
- Lemaire, F.; A. Dartigues and L.M. Riviere (1985). Properties of substrate made with spent mushroom compost. *Acta Hortic.*, 172: 13-29.
- Lohr, V.I.; R.G. O'Brien and D.L. Coffey (1984). Spent mushroom compost in soilless media and its effects on the yield and quality of transplants. *J. Am. Soc. Hortic. Sci.*, 109: 693-697.
- Mahmoud, E.K. (2003). Low technology co-composting of municipal sludge and crop residues as soil improver and peat substitute. Ph.D. Thesis, Alexandria Univ., Egypt.
- Mathur, S.P.; J.Y. Daigle; M. Levesque and H. Diné (1986). The feasibility of preparing high quality composts from fish scarp and peat with seaweeds or crop scrap. *Biological Agriculture & Horticulture*, 4: 27-38.
- Mathur, S.P.; G. Owen; H. Diné and M. Schnitzer (1993). Determination of compost biomaturity. 1. Literature review. *Biological Agriculture & Horticulture*, 10: 65-85.
- Sesay, A.A.; K. Lasaridi; E. Stentiford and T. Budd (1997). Controlled composting of paper pulp sludge using the aerated static pile method. *Compost Sci.*, 5: 82-96.
- Shiralipour, A.; D.B. McConnell and W.H. Smith (1997). Phytotoxic effects of a short-chain fatty acid on seed germination and root length of *Cucumis sativus* cv. "Poinset". *Compost Sci. Util.*, 5: 47-52.
- Sikora, L.J. and D.M. Sullivan (2000). Case studies of municipal and on-farm composting in the United States of America. In USA Land application of agricultural, industrial and municipal by-products. SSS A Book & Series no. 6, p. 605-625.
- Stevenson, F.J. (1982). *Humus chemistry: genesis, composition, reactions*. John Wiley & Sons, New York, NY.
- U.S. Environmental Protection Agency (U.S. EPA) (1993). Standards for use or disposal of sewage sludge. *Fed Regist.*, 58 (32): 9248-9415.
- Wu, L. and L.Q. Ma (2001). Effect of sample storage on biosolids compost stability and maturity evaluation. *J. Environ. Qual.*, 30: 222-228.
- Wu, L.; L.Q. Ma and G.A. Martiney (2000). Comparison of methods for evaluating stability and maturity of biosolids compost. *J. Environ. Qual.*, 29: 424-429.



## جودة الكمبوست وإستخدامه كبيئة نمو للنباتات

عيسوى قاسم محمود\* ، حسنى عبد الله سالم\*\*

\* قسم علوم الأراضى والمياه، كلية الزراعة، جامعة طنطا، مصر.

\*\* قسم الزهور ونباتات الزينة، كلية الزراعة، جامعة الإسكندرية، مصر.

جودة الكمبوست مهمة عند إستخدامه كبيئة نمو. فى هذه الدراسة أختير ستة أنواع مختلفة من الكمبوست من إنتاج شركة دلتا Bio-Tec وشركة النيل، الكمبوست المنتج بواسطة محمود (٢٠٠٣) لتقدير الخصائص الفيزيائية والكيميائية ودرجة ثبات ونضج الكمبوست ودراسة مدى امكانية استخدام الكمبوست المنتج بواسطة شركة دلتا Bio-Tec كبديل للبيت بزراعة نباتات الونكا *Vinca sp.*

وأوضحت النتائج مايلي:-

أن الخصائص الفيزيائية لاتكون مثالية لنمو النباتات ماعدا الكمبوست (BC).

التركيز العالى من الأملاح فى كمبوست الداغنى ومخلفات المحاصيل المحدد الرئيسى عند إستخدامه كبيئة نمو. الكمبوست (BC) محتواها عالى من البوتاسيوم والنتروجين ومنخفض فى الفوسفور. كمبوست شركة دلتا Bio-Tec وشركة النيل لم يصل إلى درجة الثبات والنضج التام مما أدى ذلك إلى سمية نبات *Cress*. محصول الونكا كان أكبر فى بيئة البيت عن الكمبوست أو مخلوط الكمبوست مع البيت. موت النباتات النامية فى الكمبوست الداغنى عند نسبة ٤ : صفر ، ٣ : ١ ، ١ : ١ (الكمبوست : البيت (حجما)). التركيز العالى من الأملاح وقلة درجة الثبات والنضج للكمبوست المختبر ماعدا كمبوست (BC) هما المحددان لمستخدم الكمبوست كبيئة نمو بدل البيت.

والإستراتيجية لإستخدام هذا الكمبوست يجب أن تقنن لتقليل أثر الأملاح والسمية للنباتات النامية فى بيئة النمو.