

THE EFFECT OF USING DATE PALM WASTE (MID RIB AND PINNE OF DATE PALM) AS INGREDIENTS IN BROILER DIETS

EL-Faham, A. I.

Poultry Prod .Dept. Faculty of Agric., Ain Shams Univ., Cairo – Egypt.

ABSTRACT

An experiment was conducted to determine the effect of addition date palm waste (mid rib of date palm, MDP and pinnae of date palm, PDP) to broiler diets (starter/grower/ finisher or grower/ finisher or Finisher) on the productive performance, carcass characteristics and economical evaluation of broiler chicks. A total number of 480 one- week old male Hubbard broiler chicks were distributed equitably into 8 dietary treatments in 3 replicates of 20 birds each .Four experimental diets in each period (starting, growing and finishing) were formulated in which (control diet) was 0.0% date palm waste, in the other diets wheat bran WB , MDP and PDP were incorporated at level of 5% to obtain basal diet 1, basal diet 2 and basal diet 3, respectively. Control diet and basal diet 1were used in the whole experimental period (control and T1), while basal diet2 and basal diet 3 were used in whole experimental period (T2 and T5) or in growing/ finishing period (T3 and T6) or in finishing period (T4 and T7) respectively.

The results indicated that:

- 1- There were no significant differences in body weight, feed intake , feed conversion and mortality rate between chicks fed control diets, wheat bran diets (Basal 1), MDP diets (Basal 2) and PDP diets (Basal 3) in different growth periods.
- 2- Carcass characteristics indicated that added MDP in broiler chicks significantly decreased dressing % (T3) and breast % (T4), but drumstick% significantly increased (T2).
- 3- The results of chemical composition of white breast meat showed that, broiler chicks fed MDP (T3) or PDP (T7) gave significantly lower protein % compared with those fed control diets.
- 4- The results of economical evaluation showed that mid rib and pinnae of date palm could be included at 5 % in broiler diets and the best feeding cost per kg gain was recorded for broiler chicks fed pinnae of date palm (T5).

Keywords: Date palm, pinnae of date palm, broiler diet.

INTRODUCTION

In recent years, considerable attention has been given to search for more available and cheaper feed ingredients particularly agricultural by-products for animal feeding. Furthermore, 21 million tons of agricultural crop residues are produced in Egypt annually (Deraz, 1996). However, only 4.0 to 4.3 million ton of the crop residues are used for feeding animals (El-Shinnawy, 1990; Hathout and El-Nouby, 1990 and Khorsned, 2000).

Therefore, conclusive research has been conducted to evaluate the effect of using some agricultural by products as untraditional feedstuffs on the performance of animals and poultry.

The main shortcoming of agriculture by- products as untraditional feedstuffs in poultry diets lies in high fiber content and their poor palatability. Moreover, fiber sources may contain phytic acid, which has been shown to reduce calcium, phosphorus and zinc availability to chicks (Lease and Williams, 1967 and lease, 1968). The fiber sources themselves may reduce

trace mineral availability by binding the mineral to fiber matrix (Ismail- Beigi *et al.* 1977)

On the other hand Miller *et al.* (1979) found that various fractions isolated from wood hemicellulose extract, a chemically complex material high in phenolics, resulted in improvements in performance of chicks when incorporated into practical corn soybean meal diet. Moreover, Ricke *et al.* (1982) concluded that the improvements due to addition of cell wall carbohydrates is caused by change in a microbial

metabolism in the digestive tract rather than a change in post absorptive (tissue) metabolism. Also, the high fiber diets increased the digestive passage rate and increased bacterial synthesis of amino acids in the large intestine and caeca. (Parsons *et al.* 1983). Moreover, many successful trails by using some components of rice milling , wheat bran ,corn bran, soy bran, alfalfa meal, sawdust, prickley pears, oat hulls and date stone meal in broiler , layer, quail and ducklings diets (Sutton *et al.*, 1981; Thompson and Weber, 1981; Ricke *et al.*, 1982; Ghazalah., 1990; Attia, *et al.* , 1995; Jorgensen *et al.*, 1996; Ensaf El-full *et al.* , 2000; El-Nagmy *et al.*, 2001; and Abaza *et al.*, 2004).

Wheat bran, is usually not included in appreciable quantities in the poultry diets. However, Lee *et al.*(1985) supplemented isonitrogenous diet with 5or 10% wheat bran to fed chickens from 3 to 28 day of age . they found that increasing amounts of wheat bran in the diet decreased feed intake and there was a significant interaction of bran with dietary salt or calcium. wheat bran contains an active phytase in germinated grain, the enzyme may improve the utilization of phytase phosphorus in diets when fed to animals and chicks (Nelson, 1980; McGillivray, 1978.

It was generally known that date have a fundamental economic importance in many desert areas especially in sinai and El-Wadi El-Gedid governorate in Egypt, where inedible dates, date pits, pinnae of date palm and mid rib of date palm are not utilized for human consumption. However, several experiments by many investigators revealed that date stone meal can be included in broiler, layer, quail and duckling rations. (Zumbado *et al.*, 1987; Panigrahia and Powell., 1991; El-Bogdady *et al.*, 1995; Osman *et al.* 1995; Soliman., 1996 and El-Nagmy *et al.*,2000).

No trail has been reported to use pinnae of date palm or mid rib of date palm as untraditional feed ingredients in poultry diets. So, the present investigation was undertaken to assess the effects of use pinnae of date palm and mid rib of date as untraditional feed ingredients in broiler diets.

MATERIALS AND METHODS

The present study was carried out at the Poultry Farm of the Poultry Production Department Faculty of Agriculture, Ain Shams University, Shoubra El-Kheima. The present study was designed to study the possibility for using pinnae of date palm (PDP) and mid rib of date palm (MDP) as untraditional feedstuffs in broiler diets. Such materials are produced in abundant quantities in El-Wadi El-Gedid Governorate.

Fifty Kilograms. from each crop-residue were sun dried until complete drying and fine ground to pass through a 1.0-mm screen.

Chemical analysis of PDP and MDP were carried out according to A.O.A.C (1980) and results are presented in (Table1).

Four experimental diets were formulated in which (control diet) was 0.0 % date palm waste, in the other diets wheat bran (WB), (MDP) and (PDP) were incorporated at level of 5% to obtain basal diet 1, basal diet 2 and basal diet 3 respectively. Starter, grower and finisher experimental diets (table2) were formulated to meet the nutrient requirements of the broiler chicks according to Egyptian Ministry of Agriculture decree (1996).

Table (1): Chemical composition percent of wheat bran, mid rib and pinnae of date palm on dry matter basis.

Items %	Wheat bran	Mid rib of date palm	Pinnae of date palm
Dry matter	88.17	88.10	93.70
Crude protein	15.21	2.69	5.85
Crude fiber	9.95	34.02	27.64
Ether extract	4.44	2.98	0.78
Nitrogen free extract	52.94	41.70	49.86
Ash	5.63	6.71	9.57
Organic matter	94.37	93.29	90.43
ME (Kcal/kg) *	1300	695	627

Each value represents pooled results of 3 samples per each tested ingredient.

* The metabolizable energy (ME)for wheat bran was calculated according to feed composition tables for animal and poultry feedstuffs used in Egypt (2001)., and for MDP and PDP were calculated on the basis of its chemical composition according to Carpenter and Clegg (1956) .

A total number of 480 day-old male commercial broiler chicks (Hubbard) were fed starter diet (22.61% CP and 2996 Kcal, ME/Kg) for the first week of age. AT the 7th day of age, chicks were wing-banded, weighed and randomly assigned to 8 dietary treatments of 60 chicks each, in three replicates (20 chicks per replicate). Chicks were randomly allocated into battery brooders with raised wire floors equipped with electric, thermostatically controlled heaters. Chicks were given starter diets

(1-2weeks), grower diets (3-4weeks) and finisher diets until the end of experimental period (7 weeks). The eight experimental dietary treatments were as follows:

treatments	Starter diet	Grower diet	Finisher diet
Control	Control diet	Control diet	Control diet
T1	WB - diet	WB - diet	WB - diet
T2	MDP - diet	MDP - diet	MDP - diet
T3	Control diet	MDP - diet	MDP - diet
T4	Control diet	Control diet	MDP - diet
T5	PDP - diet	PDP - diet	PDP - diet
T6	Control diet	PDP - diet	PDP - diet
T7	Control diet	Control-diet	PDP - diet

Feed and water were available (ad. Libitum) all time during experimental period. Body weights and feed consumption were weekly recorded and body weight gain and feed conversion were calculated.

Table (2). Composition and calculated analysis of the control and basal diets.

Ingredients	Starter diets		Grower diets		Finisher diets	
	Control	Basal 1*	Control	Basal 1*	Control	Basal 1*
Yellow corn	61	54	63	56	66	59
Soybean meal (44%)	27	27	27	27	27	27
Meat and bone meal (60%)	10	10	6	6	2	2
Wheat bran	-	5	-	5	-	5
Vegetable oil	0.93	2.93	2.04	4.04	2.42	4.42
Bone meal	0.45	0.45	1.30	1.30	1.9	1.9
Salt	0.30	0.30	0.30	0.30	0.30	0.30
Premix **	0.30	0.30	0.30	0.30	0.30	0.30
DL Methionine	0.02	0.02	0.06	0.06	0.08	0.08
Total	100	100	100	100	100	100
Calculated analysis						
Crude protein %	22.61	22.82	20.42	20.63	18.30	18.51
ME Kcal/ kg	2996	2932	3053	3054	3080	3081
Lysine%	1.29	1.31	1.16	1.17	1.03	1.03
Methionine + cystine %	0.72	0.72	0.71	0.72	0.70	0.70
Calcium %	1.01	1.01	0.96	0.95	0.85	0.85
Available phosphors %	0.50	0.51	0.47	0.48	0.43	0.44
Price of ton feed L.E ***	1171	1191	1172	1192	1154	1174

* Starter, grower and finisher basal diets 2 & 3 are the same as (basal diet 1) but wheat bran substitution by mid rib of date palm (basal 2) or by pinnae of date palm (basal 3).
 ** Premix, vitamin and mineral mixture supplied each kg diet: Vit A 12000 IU, Vit D3 2500 IU, Vit E 12mg, Vit k3 3mg, Vit B1 1mg, Vit B2 6mg, Vit B6 3mg, Vit B12 13mg , Niacin 30mg, Pantothenic acid 12 mg , Folic acid 1mg, Biotin 75 mg, Choline chloride 600mg, Copper 5mg, Manganese 70mg, Zinc 50mg, iron 60 mg, Selenium 0.1mg and cobalt 0.1mg .
 *** The price of ton feed for basal diets which content mid rib or pinnae of date palm were 1176, 1177 and 1159 L.E for starter, grower and finisher diets respectively.

At the end of experiment (7 week of age) six chicks of each dietary treatment were randomly taken and slaughtered to complete bleeding, followed by plucking the feathers. After the removal of head, viscera, shanks, spleen, gizzard, liver, heart and abdominal fat, the rest of the body was weighed to determine the dressed weight. Weights of different parts of carcass (breast, thigh, drumstick and back) were recorded to the nearest gm. Representative samples of white breast meat were analyzed for moisture, ether extract , crude protein and ash content using standard methods of A.O.A.C. (1990). Mortality rate was daily recorded and feed cost /Kg gain (L.E) was calculated.

The data was examined statistically using the computerized analysis of variance and Duncan's multiple range test procedures within the statistical analysis system, SAS (1996).

RESULTS AND DISCUSSION

Results of proximate analysis (on dry weight basis) of MDP and PDP used in this research in comparison with WB was summarized in table (1). The experimental data showed that WB was the highest in crude protein

(15.21%), while MDP was the lowest (2.69%) and PDP was in the middle (5.85%). The crude protein content of PDP is comparable to that found in some grains commonly used in poultry diets (Salem and Hegazi 1971). Ether extract was relatively higher in WB (4.44%) and MDP (2.98%) than those found in PDP (0.78%). While, nitrogen free extract (NFE) was higher in WB (52.94%) than those found in PDP (49.86%) and MDP (41.70%). On the other hand, MDP contained the highest value in crude fiber (34.02%) followed by PDP (27.64%) and WB had the lowest (9.95%).

This high fiber percentage may limit the diet inclusion level of date palm wastes, especially MDP in poultry diets (El Boushy and Van Der Poel 1994).

The metabolizable energy (ME) was calculated to be 695 and 627kcal/kg in MDP and PDP on the basis of its chemical composition according to Carpenter and Clegg (1956), respectively. These figures for ME were much lower than WB (1300 kcal/kg) according to Egyptian Ministry of Agriculture feed composition Tables (2001).

However, there are numerous inconsistent and conflicting findings concerning the chemical composition and metabolizable energy for date waste.

The metabolizable energy was calculated by Hermes and Al-Homidan (2004) to be 2512 and 1126 kcal/kg in Sukarry date waste and date pits meal. While Al-Yousef (1985) reported that ME of Khudri² date was estimated to be 2409 kcal/kg. On the other hand, Radwan *et al.*, (1997) and El-Nagmy *et al.* (2001) calculated ME of date stone meal on the basis of its chemical composition and found it to be 1746 and 2757 kcal ME/kg, respectively.

Productive Performance.

Growth performance of broiler chicks fed experimental diets (starter, grower and finisher) of control; WB, MDP and PDP in different periods are shown in Table 3. It is clear that initial live body weight at 7 days of age did not differ among the experimental groups and the overall mean being 103.39. Chicks fed experimental diets during hall experimental period (1-7 weeks of age) reflect that the lowest value in body weight were for chicks fed diets containing MDP (T2) followed by that fed control diet while chicks fed diets containing PDP (T5) were the highest; differences among treatments were not significant. It was obvious from Table 3 that no significant differences in body weight were detected among the control diet, WB, MDP, PDP in different treatments.

The results of feed intake showed that birds fed the experimental diets containing PDP (T5, T6 and T7) had the lowest feed intake being 4.5%, 6.0% and 5.7% lower than that of birds on control diet, respectively during the period from 1 to 7 weeks age; differences were statistically not significant. During the different experimental periods (1-2, 1-4 and 1-7 weeks of age) the birds fed different experimental diets went down and the differences in feed intake failed to be not significant.

However, birds fed control diets or diets containing MDP (T3) consumed more feed than the other dietary treatments.

Table (3): Effects of feeding different experimental diets on the growth performance of broiler chicks at different age intervals.

Treatment age wks	control	T1	T2	T3	T4	T5	T6	T7	SE	Significant Of differences
Body weight (gm)										
1	108.7	106.0	104.3	100.0	101.3	102.7	102.3	101.0	1.99	NS
2	258.3	248.0	249.0	235.3	240.3	246.7	239.7	234.7	5.68	NS
4	886.7	879.3	900.3	897.0	885.7	889.7	834.0	840.0	21.72	NS
7	1887.3	1917.3	1873.7	1927.0	1921.7	1956.0	1917.3	1903.7	29.05	NS
Feed intake (gm)										
1-2	249.0	233.7	224.3	209.3	210.3	238.7	229.0	224.7	11.19	NS
1-4	1379.0	1368.7	1409.3	1425.0	1383.6	1382.7	1322.7	1333.0	46.54	NS
1-7	4064.3	3961.3	3909.3	4225.7	3967.3	3882.7	3822.7	3833.0	91.75	NS
Feed conversion										
1-2	1.66	1.65	1.55	1.55	1.51	1.66	1.67	1.68	0.058	NS
1-4	1.77	1.77	1.77	1.79	1.76	1.77	1.81	1.80	0.058	NS
1-7	2.29	2.19	2.21	2.32	2.18	2.10	2.11	2.13	0.059	NS

NS = not significant

The results of feed conversion (g feed/ g gain) showed the same trend since the differences were not significant during starting, starting/ growing, as well as during the whole experimental period and the birds fed PDP in experimental diets (T5, T6 and T7) were the best followed by those fed MDP (T4), while those fed control diets or PDP (T3) were the worst. These results are in agreement with those reported by many investigators (Yeong *et al.*, 1981; Sawaya *et al.*, 1984; Onwudikee, 1986 and El-Boushy and Van Der Poel 1994).

They concluded that high crude fiber percentage would limit the diet inclusion level of date waste, especially date pits to 10-15% in poultry diets. Moreover, the date wastes are poorly digested by poultry and the lower levels of critical essential amino acids in date waste are not enough to meeting the requirements for birds.

Similar observation were reported by other investigators, Osei and Amo(1987) found that no significant differences in final body weight of broilers by using 5.10 and 15% date meal in their diets. El-Nagmy (2001) reported that partial substitution of yellow corn either by date stone meal or prickley pear peels as sources of energy in quail diets had beneficial effect on the quails performance. Also, Holder (1980) and Lee *et al.*(1985) found that no significant difference in body weight gain and feed efficiency for chicks fed diets containing up to 7.5% clover or wheat bran up to 10% compared to the control.

On the other hand, these findings are in contrast with the results obtained by Radwan *et al.*, (1997), who reported that raising level of date stone meal in quail, diets up to 24% tended to improve body weight gain of grower quail.

Also Abaza, *et al.* (2004) reported that wheat bran can be used up to 35% in Matrouh laying hen diets with no detrimental effect could be found on feed conversion or egg quality.

Carcass characteristics.

Data for carcass characteristics and carcass parts of slaughtered broilers from different treatments are presented in Table 4. It is worth to note that broiler chicks fed MDP (T3) and WB (T1) reflected the lowest dressing percent compared with those fed control diets and the corresponding values were 57.65%, 58.84% and 62.04% respectively and in most cases differences were significant.

In addition, broiler chicks fed MDP (T4) gave the lowest breast% compared to those fed control diets being 28.26 % VS 31.18% respectively.

In contrast the drumstick% of broiler chicks fed MDP (T2) gave the highest percent, differences failed to be significant. These findings are in contrast with the results obtained by Ramdan *et al.* (1997) in grower chicks and layer, Hermes and Al-Homidan (2004) in layer, and Ramdan *et al.* (1997) in growing quail, they reported that date waste (whole dates, date pits and date stone meal) could be replaced yellow corn without any adverse effect on productive performance, quality characteristics and carcass characteristics. In general, no significant differences were obtained among dietary treatments with regard to thigh %, wing % abdominal fat % and giblets % (Table 4).

Chemical composition of breast meat.

Data for chemical composition of white breast meat are summarized in Table 5. A significant ($P < 0.05$) decrease in fat content of white breast meat for broiler chicks fed WB (T1) compared to those fed control diets being 0.45% and 0.75% respectively. Also, broiler chicks fed MDP (T3) or fed PDP (T7) gave significantly lowest protein% compared with those fed control diets and the corresponding values were 21.48%, 21.95% and 23.86% respectively. These findings are in agreement with those of Patel *et al.* (1981) and McDougall *et al.* (1996).

They found that feeding wheat bran, cellulose or alfalfa meal resulted in significant reductions in hepatic lipid deposition, liver fat and serum cholesterol this may explain and support our findings concerning the reduction of breast fat content.

Economical evaluation.

Data for economical evaluation are summarized in (Table 6) .These results are based on recent prices of local market for feed ingredients. Because the calculated energy (ME- Kcal/ kg) value of wheat bran was much lower than yellow corn, it was necessary to increase the level of fat by about 2% in (basal 1) diets, to keep all diets iso- caloric (Table 2). The average cost/ton of final experimental diets (starter, grower and finisher) were : 1171,1172 and 1154 L.E for control diets, and 1191,1192 and 1174 L.E for basal 1 diets and 1176 , 1177 , and 1159 L.E for basal 2 or basal 3 diets respectively . Fat supplementation (20kg vegetable oil/ ton final diet) to(basal 1) diets raised the cost/ ton of final feed by about 20 L.E as compared to control diets.

Table (4). Effect of feeding different experimental diets on carcass traits (%) of broiler chicks

Treatment Criteria	control	T1	T2	T3	T4	T5	T6	T7	SE	Significant Of differences
Dressing %	62.04 BA	58.84 BC	60.39 BAC	57.65 C	62.63 A	61.68 BA	62.50 A	63.79 A	1.12	**
Breast %	31.18 CB	32.14 AB	30.21 BCD	30.16 BCD	28.26 D	33.34 A	29.01 CD	29.26 BCD	0.67	**
Thigh %	20.39	19.85	20.67	19.44	20.13	19.87	22.44	20.23	1.03	NS
Drumstick %	18.54 CB	18.95 CB	20.86 A	16.25 D	16.24 D	18.32 B	16.13 D	17.03 D	0.67	**
Back %	16.97 B	16.73 B	15.92 B	22.26 A	23.10 A	15.83 B	20.36 A	21.31 A	1.07	**
Wings %	12.43	12.52	12.54	11.87	12.25	12.91	12.11	12.15	0.453	NS
Abdominal fat %	1.20	1.03	1.70	1.36	1.24	1.07	1.41	1.47	0.176	NS
Giblets %	4.74	4.23	4.74	4.70	4.37	4.93	4.87	4.63	0.196	NS

Means in each row having different letter (s) are significantly different ($P < 0.05$); NS= not significant

* $P < 0.05$; ** $P < 0.01$.

Table (5) . Effect of feeding different experimental diets on chemical composition (%) of broiler chicks breast meat.

Treatment Criteria	control	T1	T2	T3	T4	T5	T6	T7	SE	Significant Of differences
Moisture	73.90	74.33	71.97	73.40	73.93	74.33	73.87	74.30	0.94	NS
Ether extract	0.75 A	0.45 B	0.63 BA	0.60 BA	0.64 BA	0.58 BA	0.57 BA	0.56 BA	0.07	*
Protein	23.86 BA	22.45 BDC	21.48 D	24.61 A	23.19 BDAC	23.35 BAC	23.14BDAC	21.95 DC	0.54	*
Ash	1.49	1.77	1.92	1.40	1.59	1.39	1.45	1.18	0.06	NS

Means in each row having different letter (s) are significantly different ($P < 0.05$); NS= not significant * $P < 0.05$.

It was clear that using WB, MDP except (T3) and PDP reduced relatively the feeding cost of broiler chickens compared with those fed control diets. The best feeding cost per kg gain was calculated for chicks fed PDP (92.1%) compared with those fed control diets (T5). Such improvement was due to the superior body weight and feed conversion for broiler chicks fed PDP in whole experimental periods.

Table (6). Effect of feeding different experimental diets on feed cost/ kg gain of broilers at marketing age.

Items	Control	T1	T2	T3	T4	T5	T6	T7
Feed cost/bird,(L.E*)								
2- week	0.292	0.278	0.264	0.245	0.246	0.281	0.268	0.263
4- week	1.616	1.630	1.659	1.676	1.621	1.627	1.555	1.562
7- week	4.715	4.675	4.557	4.922	4.616	4.525	4.453	4.460
Feed cost/kg gain L.E)								
2- week	1.95	1.96	1.83	1.81	1.77	1.95	1.95	1.97
4- week	2.08	2.11	2.08	2.10	2.07	2.07	2.13	2.11
7- week	2.65	2.58	2.58	2.69	2.54	2.44	2.45	2.74
Relative FC,(%)**	100	97.4	97.4	101.5	95.9	92.1	92.5	103.4

* Based upon the price of feed.

** Relative to the FC of the control at 7- week.

Health condition and mortality rate.

Under the condition of the present study all chicks appeared healthy and the total mortality rate was 2.5 % during the whole experimental period without any clear differences among treatments .

CONCLUSION

From the present results, it could be concluded that it is possible from the economic point of view to inclusion MDP or PDP by 5% level in broiler diets (starter, grower and finisher) .

Further studies must be carried on the use of such waste especially with PDP in order to discover more information about these material in feeding livestock's.

REFERENCES

- Abaza, I.M; Ali, M.N. and Hassan, M.S. (2004) Nutritional and physiological studies on improving the utilization of wheat bran in laying hen diets. Egypt, Poul. Sci. 24:1015-1031.
- Al-Yousef, Y.M.(1985) , Dates as feedstuff for poultry and the effect of Alkali on date Pit fiber. Ph. D. Thesis presented to Faculty of graduated school of the University of Missouri Columbia, USA.
- A.O.A.C (1990) Association of Official Analytical Chemists. Methods of Analysis, 15th edition .Arlington, USA.

- Attia, Y.A., Mona Osman and Merfat Salah El-Din (1995). Effect of enzyme mixture Treatment on performance and nutrient digestibility of Alexandria layers fed diets containing different levels of whole rice milling by product. First Egyptian Hungarian Poul. Conf., 17-19 Sept. 1995, Alex, Egypt.
- Carpenter, K.J and Clegg, K.M.(1956). The metabolizable energy of poultry feeding Stuffs in relation to their chemical composition. J.Ci. food Agric., 7:45.
- Deraz, T.A (1996). The production of microbial protein from some agricultural wastes and its utilization in ruminant. Ph.D. Thesis Faculty of Agric. Ain Shams University.
- El-Bogdady, A.H., Osman, A.M.A and Toson, M.A. (1995). Performance of growing Japanese quail fed different levels of date stone meal. J. Agric., Sci. Mansoura Univ., 20:2191-2149.
- El-Boushy, A.R.Y. and Van Der Poel A.F.B.(1994): Poultry feed from waste processing and use. Department of Animal Nutrition, Wageningen Agricultural University, the Netherlands. Published by Chapman Hall, 2-6 Boundary row, London SE 18 HN, UK. PP: 276-286.
- El-Nagmy, K.Y; Ali, A.M.; and Abdel-Malak, M.S.(2001). The effect of using some untraditional feedstuffs on the performance of Japanese quails in north Sinai. Egypt. Poul. Sci. 21:701-717.
- El-Shinnawy M.M.(1990). The role of fibrous residues in residues in feeding ruminants. 3rd Intern. Symp. On feed Manufac. and quality control., PP.321-326.
- Ensaf A.El-full; Asker, N.E.A; Ali, M.M.M; Hala M.Abdel Wahab and Omar, E.M.(2000). The use of rice bran in broiler and layer diets with reference to enzyme supplementation. Egypt. Poul. Sci. 20:517-543.
- Feed composition Tables for Animal and poultry feedstuffs used in Egypt.(2001). Technical Bulletin Nr. I, Central Lab for feed and food, Ministry of Agriculture Egypt.
- Ghazalah, A.A., Arafa, S.A. and Abd- El Ghany, A.L.(1990). Nutritional value of rice bran in laying hen diets. Egypt. Poul. Sci. 10: 365-383.
- Hathout, M.K. and H.M. El-Nouby (1990). Practical application of crop residues treatment in Egypt. 3rd Intern. Symp. On feed. Manufac. and quality control. Pp. 337-347.
- Hermes, I.H. and Al-Homidan, A.H.(2004). Effects of using date waste (whole dates and date pits) on performance, egg components and quality 14 characteristics of Baladi Saudi and Leghorn laying hens. Egyptian J. Nutrition and Feeds 7:223-241.
- Holder, D.P.(1980). Use of clover in broiler starter rations. Poul. Sci, 59:2355-2357.
- Ismail-Beigi, F. Faradji, B. and Reinhold, J.G. (1977). Binding of Zinc and iron to wheat breed, wheat bran and their components. Amer. J. Clin. Nutr. 30:1721-1725.
- Jorgensen, H., X. Zhao, K.E. Bachknudsn, and B.O. Egyum (1996). The influence of dietary source and level on development of the gastrointestinal tract, digestibility and energy metabolism in broiler chickens. Br. J. of Nutr. 75:379-395.

- Khorshed, M.M.(2000) Different treatments for improving quality of some crop residues used in ruminant nutrition. Ph.D. Thesis.Fac. of Agricultural Ain Shams University.
- Lease,J.G.(1968). Effect of a soluble fraction of oil seed meals on Zn65 from Ca. Mg Zn65 phytate complexes by the chick. *J.Nutr.* 96:126-138.
- Lease,J.G. and Williams,W.P.(1967). Availability of Zinc and comparison of in Vitro and Vivo Zinc uptake of certain oil seed meals. *Poultry Sci.*, 46:233-241.
- Lee, J.H., Han, I.K. and Paik, I.K (1985). The effect of interaction of wheat bran, calcium and salt on the performance of broiler chicks. *Korean Journal of Animal science* 27(10), 655-662.
- McDougall, G.J. Morrison,I.M. Stewart, D. and Hillman, J.R.(1996). Plant cell walls as dietary fiber range structure, processing and function. *J.Sci. food Agric* 70:133-150.
- McGillivray, J.J.(1978). Biological availability of phosphorus sources P.73-85 in first Annu. Int. Miner. Conf.
- Miller, L., Faheamed, GC. Schussler, S.L (1979). Isolation of biologically active fractions from wood hemi cellulose extracts. *J.Anim. Sci.* 48:1129-1134.
- Nelson, T.S.(1980).Phosphorus availability in plant origin feed stuffs for poultry and Swine. pages 58-84 in 3rd Annu. Int. Miner conf.
- Onwudikee, O.C.(1986a): Palm kernel meal as feed for poultry 1.Composition of palm kernel and availability of its amino acids to chicks. *Anim.Feed Sci. and Technology*, 16:179-186.
- Osai, S.A. and Amo, J.(1987). Palm kernel cake as broiler feed ingredient. *Poultry Sci.*, 66: 1870- 1873.
- Osman,A.M; Toson, M.A. and El- Bogdady, A.H.(1995). Effect of replacing part of starter and grower diets by date stone meal on performance of Peking Duckling. *J. Agric. Sci. Mansoura Univ.* 20(5): 2151-2159.
- Panigraha, S. and powell,O. (1991). Effect of high rates of inclusion of Palm kernel meal in broiler chick. *Anim. Feed Sci. and Techno.*, 34:37-47.
- Parsons, C.M., Potter, L.M. and Brown, JR (1983). Effects of dietary carbohydrate and of intestinal microflora on excretion of endogenous amino acids by poultry. *Poult. Sci.* , 62:483-489.
- Patel, M.B., McGinnis, J. and pulbols, M.H (1981). Effect of dietary cereal grain, citrus pectin, and guar gum on liver fat in laying hens and young chicks. *Poult. Sci.*, 60: 631-636.
- Radwan, M.A.H., Ghazalah, A.A., Ghoneim, S.I, Shetawi, M.M, and Ali, A.M.(1997). Studies on growing quail nutrition in North Sinai using untraditional rations. Second Hungarian Egyptian Poultry Conference, 16- 19 september 1997 Godollo. Hungary 143-158.
- Radwan, M.S.M.; Hataba, N.A.; El Faham A.I. and Ibrahim S.A.(1997). Using date stone meal in grower chicks and layer diets. *Egyptian J.of Nutrition and feeds (special Issue)* : 321-333.
- Ricke, S.C., Vanderaar, P.J. Fahey, J.R. and Berger, L.L. (1982). Influence of dietary fibers on performance and fermentation characteristics of gut. contents from growing chicks. *Poult. Sci.*, 61:1335-1343.

- S.A.S(1996) SAS procedure Guide. Version 6.12 Edition. SAS Institute INC., Cary, NC,USA.
- Sawaya. W.N, Khalil, J.K. and Safi, W.J. (1984): chemical composition and nutritional quality of date seeds. J. food Sci 49:617-619.
- Salem, S.A., and Hegazi, S.M.(1971). Chemical composition of the Egyptian dry dates. J.Sci. food Agric.22:632.
- Soliman (1996) Use of date stone meal in broiler chicken diets. Egypt. Poultr.Sci. 16(1): 187-198.
- Sutton, C.D., Muir ,W.M. and Begin, J.J. (1981). Effect of fiber on cholesterol metabolism in the quail.Poultr. Sci., 60: 812-817.
- Thompson, S.A.and.Weber,C.W. (1981). Effect of dietary fiber sources on tissue mineral levels in chicks. Poultr. Sci., 60:840-845.
- Yeong, S.W., Mukherlee ,T.K. and Hutagalung, R.I (1981): The nutritive value of palm kernel cake as a feedstuff for poultry. Proceedings of a Nutritional workshop on oil palm By-products utilization, Kualalumpur, Malaysia. P.P: 100-107.
- Zumbado,M.E.,L.R Zumbado, and Muriilb, M.J. (1987). Composition and feeding value of whole palm kernel meal for laying hen. Poultry Sci., 66:196.

تأثير استخدام مخلفات النخيل (جريد وزعف النخيل) كخامات اعلاف فى علائق

بىدارى التسمين

أحمد إبراهيم سليمان الفحام

قسم إنتاج الدواجن - كلية الزراعة - جامعة عين شمس

أجريت هذه التجربة للتعرف على تأثير استخدام مخلفات النخيل (جريد للنخيل وزعف النخيل) فى علائق بدارى التسمين (بى/ نامى/ نامى او نامى/ نامى او نامى) على الاداء الانتاجى وصفات الذبيحة والتقييم الاقتصادى لبدارى التسمين. استخدم ٤٨٠ كتكوت هيرد عمر اسبوع فى التجربة وزعت الكتاكيت على ٨ معاملات غذائية بثلاثة تكرارات وكل مكرر ٢٠ كتكوت. كونت أربعة علائق لكل فترة (بى/ نامى ونامى ونامى) بحيث عليقة الكنترول احتوت على صفر% مخلفات نخيل بينما العلائق الأخرى ادخلت فى مكونات عليقة الردة وجريد النخيل وزعف النخيل بمعدل ٥% للحصول على عليقة قاعدية ١ وعليقة قاعدية ٢ وعليقة قاعدية ٣ على التوالى .

استخدمت عليقة الكنترول وعليقة القاعدية طوال الفترة التجريبية (كنترول ومعاملة رقم ١) بينما عليقة قاعدية ٢ وعليقة قاعدية ٣ استخدموا طوال الفترة التجريبية(معاملة رقم ٢ و ٥) أو فترة نامى ونامى (معاملة رقم ٣ و ٦) أو فترة الناهى(معاملة رقم ٤ و ٧) على التوالى.

أوضحت النتائج ما يلى :

- ١- لا يوجد فروق معنوية فى وزن الجسم واستهلاك العلف والاستفادة من الغذاء ومعدلات النفوق بين كتاكيت التسمين المغذاه على علائق الكنترول وعلائق الردة (عليقة قاعدية ١) وعلائق جريد النخيل (عليقة قاعدية ٢) وعلائق زعف النخيل (عليقة قاعدية ٣) فى مختلف فترات النمو.
- ٢- صفات الذبيحة أوضحت النتائج أن استخدام جريد النخيل فى تغذية بدارى التسمين يخفض معنويا نسبة التصافى (معاملة رقم ٣) ونسبة الصدر (معاملة رقم ٤) ويزيد معنويا % للساك (معاملة رقم ٢).
- ٣- التركيب الكيمايى للحم الصدر أوضحت النتائج ان استخدام جريد النخيل (معاملة رقم ٣) او زعف النخيل (معاملة رقم ٧) فى تغذية بدارى التسمين يخفض معنويا % للبروتين فى لحم الصدر مقارنة بالكنترول .
- ٤- نتائج التقييم الاقتصادى أوضحت انه يمكن استخدام جريد النخيل وزعف النخيل بمعدل ٥% فى علائق بدارى التسمين وافضل تكلفة غذائية لكل كيلو جرام نمو حسب لبدارى التسمين المغذاه على زعف النخيل (معاملة رقم ٥).