

Aerobic Sporeforming Bacteria in Some Foods of Animal Origin

Ahmed; A. M. Saad, A.H. and Takwa H. Ismail*

Dept. of Food Hygiene and Control, Faculty of Vet. Med., Suez Canal University

** Dept. of Food Hygiene & Control, Animal Health Research Institute, Israeli

ABSTRACT

The incidence of aerobic sporeforming bacteria in some foods of animal origin obtained from Ismailia city markets were investigated. A total of 150 food samples, 25 each of frozen beef, frozen minced beef, frozen sausage, cow milk, soft Dommietta cheese and full cream milk powder were subjected to microbiological analyzed. Aerobic sporeforming organisms were detected with 100% in all examined frozen meat samples, while 23 (92%), 14 (56%) and 10 (40%) of cow milk, soft dommietta cheese and milk powder samples were positive for the incidence of aerobic sporeformer bacteria. The mean values of total aerobic sporeforming counts for frozen meat, minced beef and frozen sausage were 6×10^3 , 2.2×10^6 and 4×10^5 10^2 CFU/g respectively, while for cow milk, soft Dommietta cheese and milk powder samples were 3.5×10^4 , 5.1×10^3 and 2×10^2 cfu/ml or g respectively. Nine Bacillus strains represented by 203 isolates were identified from the food samples. Genus of *B. cereus*, *B. flexus* and *B. pumilus* had both proteolytic and lipolytic activities.

INTRODUCTION

Foods of animal origin, specially meat and milk, represent a part of the daily diet of many individuals and considered as an important economic component of the industrial activities. Meat, milk and their products are an excellent source of protein, good sources of vitamin B-complex and minerals especially calcium, iron and phosphorus (1, 2). Food contaminations by aerobic sporeformer during various stages of production and processing are inevitable. Air, water, soil, intestine, lymph nodes, processing equipment, utensils and human being, in addition to further processing and handling techniques share the responsibility of microbial contamination of the meat during slaughtering processes (3). On the other hand, milk may be contaminated via fecal contamination of the udder. Sporeformer bacteria present in milk not only survive in their end products but may also activate causing products spoilage and/or constitute pathogens as *Bacillus cereus*, which has raised considerable health concern (4).

Aerobic sporeforming bacteria, genus Bacillus, are ubiquitous and widely distributed in nature. Genus Bacillus is described as gram positive, aerobic, mostly catalase positive rods forming endospores very resistant to many adverse conditions (5). The Bacillus spores are resistant to heat, desiccation, disinfectants, ionizing radiation and ultraviolet light (6). *Bacillus spp.* is among the main spoilage organisms in food due to their versatile

metabolism and heat-resistant spores. Except for *B. cereus*, no specific identification method is available, although information about *B. spp.* would be useful in monitoring good manufacturing practice (7). Thus, the objective of this work was to evaluate the incidence and enzymatic activity of aerobic sporeforming bacteria in the meat and milk samples randomly collected from the retail markets at the Ismailia city, Egypt.

MATERIALS AND METHODS

Samples Collection

A total of 150 samples of meat, milk and their products were microbiological analyzed in this study allover of 8 months. Twenty five samples (200 ± 15 g) each of frozen beef, frozen minced beef, frozen sausage, raw cow milk, soft dommietta cheese and full cream milk powder were randomly purchased from more than fifteen-retail foods markets located in various regions of Ismailia city, Egypt. Sterile techniques were used during samples collection, packaging and microbiological analysis. Samples were transferred in ice box to the laboratory where they were subjected to microbiological analysis for detection of the total sporeforming bacterial counts, and identification of the most isolates and their proteolytic and lipolytic activities.

Samples Preparation

The technique recommended by APHA (8) was used to prepare the food samples. Frozen beef and their products were kept over

night in refrigerator at 8-12°C in its original case for thawing. Beef, minced beef, sausage and soft Dommietta cheese samples were minced in a separate sterile food processor (*Ts/12/Omas/UK*). 25g or ml of each sample was added to 225 ml of 0.1% sterile buffer peptone water (*Oxoid*) in sterile stomacher bags of approximately 500-mL capacity. Samples were blended in a Seward stomacher (*400R/UK*) for 2 min to prepare a 1:10 dilution. One ml of the original dilution was transferred serially into sterile test tubes containing 9 ml of 0.1% sterile peptone water to obtain a final dilution of 10^7 .

Determination of Sporeforming Counts

The technique recommended by Austin (9) was used for enumeration of aerobic sporeforming counts. A loopful from each dilution was streaked duplicate onto trypticase soy agar plates (*Oxoid*), and were incubated at $35 \pm 0.5^\circ\text{C}$ for 48 ± 2 hr. The average number of colonies per countable plates was determined and the total number of colonies per gram was then expressed as cfu/g.

Identification of Isolates

Five colonies were randomly picked from each plate. Each colony was isolated for further purification, identification and determinations the proteolytic and Lipolytic activities according to MacFadyean (10).

RESULTS AND DISCUSSION

Total Aerobic Sporeforming Counts

The obtained results in the present study revealed that all frozen meat samples from the Ismailia markets (100%) were positive for aerobic sporeforming organisms, while 23 (92%), 14 (56%) and 10 (40%) of cow milk, soft dommietta cheese and full cream milk powder respectively were positive for the presence of aerobic sporeformer bacteria Table, 1. The mean values of total aerobic sporeformer counts for meat, minced meat and sausage were 6×10^3 , 2.2×10^6 and 4×10^5 CFU/g respectively. For raw milk, soft domiati cheese and full cream milk powder samples were 3.5×10^4 , 5.1×10^3 and 2×10^2 CFU/ml or g respectively Table 1,2. In 1999, Nassar (11) showed that the counts of the *B. spp.* in the locally manufactured meat products ranged between 104 to 105 CFU/g. Other investigators recorded variable data

concerning the incidence of aerobic sporeformer bacteria in locally manufactured meat, milk and their products from Egypt (12, 13). The variable values of total aerobic sporeformer counts in this study reflected the level of hygienic and sanitary measures applied during the production and processing of meat and milk in Egyptian markets. Minced meat usually made from the less desirable cuts that may contain higher surface bacterial counts and consequently distributed inside the meat during mincing. Khalafalla et al. (14) confirmed that minced meat manufactured from the frozen meat had a higher microbial load when compared with that produced from fresh one; they confirmed that the additional contaminations were related to technique and operation used for mincing of meat. Aerobic sporeforming bacteria form spores during unfavorable growth conditions. These spores are heat-resistant and can survive processing, pasteurization and cooking (15). Generally, if foods are cooled slowly or kept warm before serving for suitable time and under room temperature, sporeformer may germinate and multiply rapidly at such temperatures and produce their exotoxin which is heat stable and may cause a severe public health hazard (16).

The Egyptian Organization for Standardization and Quality Control (17) does not establish a microbial standard limit for aerobic sporeformer bacteria in frozen meat, minced meat and sausage, meanwhile Food Safety Authority of Ireland (18) established that the acceptable limit should not be more than 105 CFU/g for *B. spp.* in meat and meat products. Accordingly the presence of *B. spp.* counts of 2(8%), 7(28%) and 5(20%) meat, minced meat and sausage are considered unfit for human consumption and constitute a public health hazard. Minced meat seem to be the most hazard meat products on consumer's health in the retail markets, which may be due to their locally manufacture under inadequate hygienic measures from low quality meat. On the other hand it is worthy to mention that, there is no clear microbial standard limit for the presence of the *B. spp.* in raw milk and their products. Whatever, the poor microbiological quality of milk and their products may associate with the preparation on the premises, premises type and little or no food hygiene control.

The frequency distributions of aerobic sporeforming strains isolated from food samples were shown in Fig. 1 and Table 3. Nine *Bacillus* strains out of 203 isolates were identified from meat and milk samples. Sporeformer bacteria commonly isolated from frozen minced meat (46 strains) followed by frozen sausage (42), cow milk (35) and soft dommietta cheese (28) then frozen meat and full cream milk powder (26 strains). Sporeformers in frozen meat and minced meat samples mainly consisted of *B. subtilis* (30.8%, 23.9%) and *B. brevis* (19.2%, 10.9%), followed by *B. flexus* (15.4%, 13%) respectively, while in frozen sausage samples mainly consisted of *B. brevis* (23.8%) followed by *B. cereus* and *B. subtilis* (21.4%). On the other side, sporeformers in cow milk, soft dommietta cheese full cream milk powder samples consisted mainly of *B. cereus* (28.5%, 39.3% and 42.4%) followed by *B. subtilis* (14.3% and 34.6%) respectively. The identification with classical methods based on morphological and biochemical criteria showed *B. cereus* to be the most frequently occurring *Bacillus* spp. which isolated from examined food samples Figure, 2.

It is assumed that raw meat or milk directly obtained from healthy animals is free of microorganisms. Microbial contamination of foods during processing is undesirable but unavoidable. The initial microflora on the surfaces of the raw foods of animal origins mainly constitutes of aerobic sporeforming organisms (19). The non hygienic methods of milk production and high ambient temperatures with the lack of prompt cooling after milking are the main reason affecting microbial contamination of raw milk (20). *B. spp.* is the most predominated foods contaminant which capable of existing in two forms: active vegetative cells and dormant spores. These two forms often differ in their resistance properties to heat, chemicals, irradiation and other processing stresses. Similarly, spores are typically more resistant than vegetative cells to the alternative food processing technologies which interpret the presence of these organisms in the meat and milk products.

Aerobic sporeformer are considered either considered food spoilage microorganisms as genera of *B. flexus* and *B.*

pumilus or food-borne pathogens as *B. cereus* and *B. subtilis* (21). Of those most frequently isolated from the food of animal origin, *B. brevis*, *B. cereus* and *B. subtilis* are important for food hygiene because of their hydrolytic activities on food components and the ability of some strains to produce toxins or to grow at refrigerated temperatures. The high frequency of *B. cereus* (62.5%) was recorded in the meat products (22). It is clear from the obtained results that a high incidence of *B. cereus* was observed in the frozen sausage samples. *Asplund et al.* (23) showed that the growth of *B. cereus* is a problem in the sausage especially when are stored at high temperatures. A dried milk product is known to be frequently contaminated with *B. cereus* (24). *B. cereus* is known as producer of emetic and enterotoxin, it is described as causative agent in food-borne illness, most of *B. spp.* produces emetic toxins and cytotoxins, but none of those cytotoxic strains produced enterotoxin like *B. cereus* (25).

The proteolytic and lipolytic activities of *B. spp.* isolated were shown in Table 4. Genus of *B. cereus*, *B. flexus* and *B. pumilus* had both proteolytic and lipolytic activities. Proteolytic bacillus strains had variable degree of proteolysis on the examined plates. On the other hand, genus of *B. cereus*, *B. flexus*, *B. coagulans*, *B. pumilus* and *B. subtilis* had lipolytic activity, meanwhile genus of *B. megaterium*, *B. polymyxa* and *B. mycoides* not shown lipolytic activity in this study. *B. cereus* and *B. subtilis* had strong lipolytic activities by 31.2% and 28.6% respectively.

Food spoilage required contamination of foods by proteolytic and lipolytic bacterial strains have the ability to produce extracellular protease and lipases enzymes that can decomposed the protein and fat to low molecular weight substances. The most *Bacillus* strains which had proteolytic activity were *B. subtilis* (23.9%) followed by *B. cereus* (17.9%) and *B. brevis* (14.9%), while *B. coagulans* had the lowest proteolytic activity by 1.5%. Aerobic sporeformer organisms have the ability to produce thermostable protease, especially from *B. cereus* and *B. polymyxa* that can survive ultra-heat treatment of foods (26). Protease enzymes can attack the nitrogen molecules naturally occur in meat causing, severe deteriorative changes in the colour and

odour of foods even under refrigerated or frozen storage (27). Proteolysis is the most important process recovered during cheese storage. It contributes cheese off-flavor, off odor and abnormal texture through the breakdown of the released proteolytic products of amino acids and peptides into amines and acids (13).

Meat contains significant amount of fat that is susceptible to hydrolytic and oxidative

rancidity. The growths of lipolytic bacteria in meat and their products play a role in quality losses and render the foods unwholesomeness (28). Growth of aerobic sporeformer bacteria in raw milk produces extracellular lipase enzyme which adsorbed on milk fat globules and concentrated in the manufactured cheese. During storage, the enzyme causes bitter flavor by hydrolysis of fats into fatty acids and glycerides (29).

Table 1. The Incidence No. (%) of Aerobic Sporeforming Bacteria in Food Samples.

	Frozen Beef and Beef Products			Milk and Milk Products		
	Raw	Minced	Sausage	Raw	*S.D. Cheese	Powder
Positive No. (%)	25 (100)	25 (100)	25 (100)	23 (92)	14 (56)	10 (40)
Negative No. (%)	0 (0)	0 (0)	0 (0)	2 (8)	11 (44)	15 (60)
Total	25 (100)	25 (100)	25 (100)	25 (100)	25 (100)	25 (100)

*S.D. Soft dommietta Cheese

Table 2. Statistical Analysis of Total Aerobic Sporeformer Counts in Frozen Beef, Cow Milk and some of their Products.

		Aerobic Sporeforming Count (CFU/g)			
		Minimum	Maximum	Mean	* ± S.E.
Frozen Beef	Raw	< 10 ²	1.4 x 10 ⁶	6 x 10 ⁵	2 x 10 ³
	Minced	2 x 10 ²	6.7 x 10 ⁸	2.2 x 10 ⁶	2.1 x 10 ⁴
	Sausage	< 10 ²	2 x 10 ⁶	4 x 10 ⁵	5 x 10 ⁴
Cow Milk	Raw	2 x 10 ²	4.2 x 10 ⁶	3.5 x 10 ⁴	3 x 10 ³
	S.D. Cheese	< 10 ²	1.6 x 10 ⁵	5.1 x 10 ³	2 x 10 ²
	Powder	< 10 ²	3.8 x 10 ⁴	2 x 10 ²	3 x 10 ²

*Sample No. = 25

* ± S.E. means ± Standard Error

Figure 1. Frequency Distribution of B. spp. among some of Foods of Animal Origin




Table 3. Frequency Distribution of Aerobic Sporeforming Species Isolated from some Foods of Animal Origin

Sporeformer Strains	Meat			Cow Milk			Isolates No.
	Raw	Minced	Sausage	Raw	D. Cheese	Powder	
	*F(%)	F(%)	F(%)	F(%)	F(%)	F(%)	
<i>B. brevis</i>	4(19.2)	5(10.9)	10(23.8)	3(8.6)	2(7.1)	0(0)	24
<i>B. cereus</i>	2(3.8)	7(15.2)	9(21.4)	10(28.5)	11(39.3)	11(42.4)	50
<i>B. coagulans</i>	0(0)	2(4.4)	0(0)	5(14.3)	1(3.6)	1(3.9)	9
<i>B. flexus</i>	4(15.4)	6(13)	2(4.8)	1(2.8)	3(10.7)	0(0)	16
<i>B. megaterium</i>	3(11.6)	4(8.7)	2(4.8)	3(8.6)	1(3.6)	2(7.6)	15
<i>B. polymyxa</i>	4(15.4)	3(6.5)	4(9.5)	2(5.7)	1(3.6)	0(0)	14
<i>B. mycoides</i>	0(0)	8(17.4)	6(14.3)	3(8.6)	2(7.1)	1(3.9)	20
<i>B. pumilus</i>	1(3.8)	0(0)	0(0)	3(8.6)	3(10.7)	2(7.6)	9
<i>B. subtilis</i>	8(30.8)	11(23.9)	9(21.4)	5(14.3)	4(14.3)	9(34.6)	46
Total	**26(100)	46(100)	42(100)	35(100)	28(100)	26(100)	203

Figure 2. Incidence of *B. cereus* in Compared with Other *B. spp.* Isolated from some Foods of Animal Origin




Table 4. Proteolytic and Lipolytic Activities for *B. spp.* isolated from some Foods of Animal Origin

Sporeformer Strains	Isolates (No.)	Proteolytic Activity		Lipolytic Activity	
		+ve	%	+ve	%
<i>B. brevis</i>	24	20	14.9	7	9.1
<i>B. cereus</i>	50	24	17.9	24	31.2
<i>B. coagulans</i>	9	2	1.5	1	1.3
<i>B. flexus</i>	16	11	8.2	11	14.3
<i>B. megaterium</i>	15	9	6.7	0	0
<i>B. polymyxa</i>	14	7	5.3	0	0
<i>B. mycoides</i>	20	17	12.7	0	0
<i>B. pumilus</i>	9	12	8.9	12	15.5
<i>B. subtilis</i>	46	32	23.9	22	28.6
Total	203	134	100	77	100

CONCLUSION

It could be concluded that the presence of aerobic sporeforming bacteria in the meat, milk and their products at the retail level in Ismailia city confirmed the original microbial contamination of raw meat and milk as well as the additional contamination occurred during foods processing. Application of Hazard Analysis and Critical Control Point (HACCP) systems in foods production establishments' aids in identifying and evaluating the food safety hazards. HACCP provides a quality control tools designed to replace or supplement traditional food inspection.

REFERENCES

1. Price, J.F., Schweigert, B.S. (1987): The science of meat and meat products. 3rd ed., Food and Nutrition Press, Westport.
2. Varnam, A.H., Sutherland, J.P. (1994): Milk and milk products: Technology, Chemistry and Microbiology, 3rd Ed., Chapman and Hall, London, UK.
3. Bell, R.G. (1997): Distribution and sources of microbial contamination on beef carcasses. J. Appl. Microbiol., 82: 292-300.
4. Te Giffel, M.C., Wagendrop, A., Herrewegh, A., Driehurs, F. (2002): Bacterial spores in silage and milk raw. Antonie Van Leeuwenhoek, 81: 625-630.
5. Claus, D., Berkeley, R. (1986): Genus *Bacillus*. Cohn 1872, 174^{al}. In: Sneath, P.H.A., Mair, N.S., Sharpe, M.E., Holt, J.G. (eds.) Bergey's Manual of Systematic Bacteriology, vol. 2, Baltimore, Williams and Wilkins, pp. 1105-1139.
6. Pirttijarvi, T.S., Andersson, M.A., Salkinoja-Salonen, M.S. (2000): Properties of *Bacillus cereus* and other bacilli contaminating biomaterial-based industrial processes. Int. J. Food Microbiol., 60(2-3): 231-239.
7. Deak, T., Timar, E. (1988): Simplified identification of aerobic sporeformers in the investigation of foods. Int. J. Food Microbiol., 2: 115-125.
8. APHA (1992): American Public Health Association: Compendium Methods for the Microbiological Examination of Foods. 2nd Ed., Washington D.C, USA.
9. Austin, J.W. (1998): Determination of aerobic and anaerobic sporeforming. Polyscience publications, P.O. Box 1606, Station St-Martin, Laval, Quebec, Canada H7V 3P9.
10. MacFadyean, J.F. (1988): Biochemical tests for identification of medical bacteria. 2nd Ed., Willams and Wilkins Baltimore, London.
11. Nassar, A.M. (1999): Bacteriological quality of locally manufactured and imported beef luncheon. Assiut Vet. Med. J., 42, 83: 191-199.
12. Ahmed, A.M. (1997): Quantitative and qualitative studies on proteolytic and

- Master Dissertation, Faculty of Vet. Med., Suez Canal University, Ismailia, Egypt.
13. **Salwa, A. Aly, Gala, E.A. (2002):** Effect of milk pretreatment on the keeping quality of domiati cheese. *Pakistan J. of Nutrition*, 1, 3: 132-136.
 14. **Khalafulla F, Gergis AF, El-Sherif A. (1993):** Effect of freezing and mincing technique on microbial load of minced meat. *Nahrung*, 37, 5: 422-427.
 15. **Novak, J.S., Call, J., Tomasula, P., Luchansky, J.B. (2005):** An assessment of pasteurization treatment of water, media, and milk with respect to *Bacillus* spores. *J. Food Prot.*, 68, 4: 751-757.
 16. **S.C.D.C. (2003):** *Bacillus* food poisoning. S.C.D.C., South Cambridgeshire District Council, Environmental Health, 9-11 Hills Road, Cambridge, CB2 1PB.
 17. **EOS (2005):** Egyptian Standards: Frozen meat, mined meat and sausage. Egyptian organization for standardization and quality control, E.S No. : 1522, 2097 and 1972.
 18. **FSAI (2001):** Guidelines for the interpretation of results of microbiological analysis of some ready-to-eat foods sampled at point of sale. Food Safety Authority of Ireland, Abbey Court, Lower Abbey Street, Dublin 1, pp. 8.
 19. **Doyle, M. (2002):** Survival and growth of bacterial pathogens on raw meat during chilling. Review Article, Food Research Institute, University of Wisconsin, Madison, US.
 20. **Sharaf, O.M., Hewedy, M.M., Tawfek, N.F. (1989):** Studies of the bacterial flora of raw milk in Fayoum markets, Egypt. *J. Food Sci.*, 17, 1-2: 153.
 21. **Turnbull, B., Kramer, M. (1995):** *Bacillus*. In *Manual of clinical microbiology*. Murray, P.R.; Baron, E.J.; Pfaller, M.A.; Tenover, F.C.; Tenover, R.H. (eds), ASM Press. Washington D.C., pp. 349-356.
 22. **Fang, T.J., Wei, Q.K., Liao, C.W., Hung, M.J., Wang, T.H. (2003):** Microbiological quality of 18 degrees C ready-to-eat food products sold in Taiwan. *Int. J. Food Microbiol.*, 80, 3: 241-250.
 23. **Asplund, K., Nurmi, E., Hill, P., Hirn, J. (1988):** The inhibition of the growth of *Bacillus cereus* in liver sausage. *Int. J. Food Microbiol.*, 7, 4: 349-352.
 24. **Becker, H., Schaller, G., von Wiese, W., Terplan, G. (1994):** *Bacillus cereus* in infant foods and dried milk products. *Int. J. Food Microbiol.*, 1: 1-15.
 25. **From, C., Pukall, R., Schumann, P., Hormazabal, V., Granum, P.E. (2005):** Toxin-producing ability among *Bacillus* spp. outside the *Bacillus cereus* group. *Appl. Environ. Microbiol.*, 71, 3: 1178-1183.
 26. **Matta, H., Punj, V. (1998):** Isolation and partial characterization of a thermostable extracellular protease of *Bacillus polymyxa* B-17. *Int. J. Food Microbiol.*, 42 (3): 139-145.
 27. **Abd El-Rahman, H.A., Marriott, N.G., Yassein, M.A., Ahmed, A.M. (1998):** Quantitative studies of proteolytic and lipolytic bacteria in frozen minced beef and hamburger. *J. of Muscle Foods*, 9, 2: 144-148.
 28. **Ahmed, A.M. (1997):** Control of proteolytic and lipolytic bacteria in comminuted meat. Ph.D. Dissertation, Faculty of Vet. Med., Suez Canal University, Ismailia, Egypt.
 29. **Beresford, T., Fitzsimons, N., Cogan, T., Condon, S. (1998):** Non starter lactic acid bacteria: Growth in cheese, contribution to flavor development, In: *Proceedings of the Internat. Dairy Symposium*. 19 October Rhodia Stockport, UK, pp: 98-134.

الملخص العربي

الميكروبات الهوائية المتجترثة في بعض الأغذية ذات الأصل الحيواني
 علي معوض احمد - أحمد حسن علي - تقوي حسين اسماعيل*
 الرقابة الصحية علي الأغذية - كلية الطب البيطري - جامعة قناة السويس
 * قسم الرقابة الصحية علي الأغذية - مركز بحوث صحة الحيوان - الاسماعيلية

تم إجراء الفحص الميكروبيولوجي علي عدد ١٥٠ عينة من الأغذية بواقع عدد ٢٥ عينة من مجمدات اللحم البقري واللحم المفري والسجق وكذلك من اللبن البقري الخام والجبن الأبيض الدمياطي واللبن البودرة المعبأ كامل الدسم وذلك لأستبيان احتمالية تواجد الميكروبات الهوائية المتجترثة وأيضا مقدرة تلك الجراثيم علي تحليل البروتينات والدهون في تلك الأغذية وتبين من الفحص الميكروبيولوجي أن الجراثيم الهوائية المتجترثة متواجدة بنسبه ١٠٠% في كل عينات اللحوم بينما متواجدة في عينات اللبن البقري والجبن الدمياطي واللبن البودرة بنسبة ٢٣(٩٢%)، ١٤(٥٦%)، ١٠(٤٠%) علي التوالي. وقد كانت قيمة متوسط العدد الكلي للجراثيم الهوائية المتجترثة في عينات اللحم البقري واللحم المفري والسجق هي $٦,٢ \times 10^2$ و $٢,٢ \times 10^2$ و ٤×10^1 خلية لكل جرام علي التوالي وكانت في عينات اللبن البقري والجبن الدمياطي واللبن البودرة $٣,٥ \times 10^1$ و $٢,٢ \times 10^1$ و ٤×10^1 خلية لكل جرام علي التوالي. هذا وقد تمكن من عزل عدد ٢٠٣ عزله جرثومية صنفت إلى ٩ عترات من جرثومة الباسيلاي وكانت أعلي عدد من العترات معزولة من عينات اللحم المفري المجدد (٤٦) يليها عينات السجق لمجدد (٤٢) ثم اللبن البقري (٣٥) والجبن الدمياطي (٢٨) واخيرا اللحم البقري المجدد واللبن البودرة (٢٦). وأثبت معمليا قدرة ميكروبات الباسيلاي سريس والباسيلاي فليكسس والباسيلاي بيوميليس علي أفرز إنزيمات هاضمه لها القدرة علي تحليل البروتينات والدهون هذا وقد أوصت الدراسة علي ضرورة تطبيق نظام تحديد النقاط الحرجه وتحليل المخاطر في جميع أماكن إنتاج وتجهيز وتخزين الأغذية من أصل حيواني وذلك للتأكد من سلامتها للاستهلاك الأدمي.