

## EFFECT OF TILLAGE TREATMENTS AND INTERCROPPING PATTERNS ON WATER USE EFFICIENCY AND YIELD COMPONENTS OF SOYBEAN AND MAIZE

[11]

Sherif, Sahar<sup>1</sup>. A.; Wafae Kh. Mohamed<sup>1</sup>; Sahar T. Ibrahim<sup>1</sup>;  
H.E. Osman<sup>2</sup> and S.I. El-Khatib<sup>2</sup>

### ABSTRACT

Two field experiments were performed at Sids Agriculture Farm Research Station, Bani Swif Governorate during 2003 and 2004 seasons, to study the suitability of different tillage treatments, i.e. chisel plough two and three passes at 10, 15 and 20cm depths and intercropping patterns of maize and soybean i.e. two ridges of maize : two ridges of soybean (2:2) and two ridges of maize : four ridges of soybean (2:4) on water consumption, growth characters, yield and quality of maize (cv. T.W.C. 310) intercropped with soybean (cv. Clark). The results indicated that using chisel plough 3 passes decreased the value of mean weight diameter (M.W.D) by 33.00%, 27.92% and 31.87% as compared when using chisel plough 2 passes for 10, 15 and 20cm depths respectively. On other hand, yield and quality of maize, as well as, yield and quality of soybean were significantly increased by using tillage with chisel plough 3 passes. Both yield of maize and soybean per feddan. in pure stand were always higher than those within any intercrop combination, these results were true in both seasons. The data also revealed that maize yield in (2:2) pattern and using chisel plough 3 passes at 15cm depth gave the highest yield whereas, the highest yield of soybean per feddan was obtained when soybean plants grown in (2:4) pattern with 3 passes of chisel plough at 15cm depth. On other hand the highest values of water use efficiency (W.U.E) in the two seasons were recorded when (2:4) pattern was applied and using chisel plough 3 passes at 15cm depth. The highest values of land equivalent ratio (LER) in the two seasons were 1.54 and 1.53 respectively when (2:4) pattern was applied.

**Keywords:** Intercropping patterns, Tillage systems, Maize, Soybean, Water applied

---

1- Field Crops Res. Inst. A.R.C., Giza, Egypt.

2- Agric. Eng. Res. Inst., A.R.C., Dokki, Giza, Egypt

(Received July 9, 2005)

(Accepted November 14, 2005)

## INTRODUCTION

Before sowing seeds it is necessary to prepare a suitable seedbed for seed germination. Tillage will ensure the adequate moisture and air quantity needed for plant. In addition the seedbed should be as free as possible from weeds and applied fertilizer be incorporated eventually with the soil. **El-Nakib and Fouad, (1990)** showed that the mean weight diameter (MWD) increased by increasing of working depth and forward speed because less breakdown would be produced at higher speed and depth. **Abo-Habaga, (1992)** concluded that decreasing the percentages of less than ( $\phi > 50\text{mm}$ ) in the seedbed increased the main distance between adjacent seeds in row for a given number of seeds per unit area and then the crop yield was increased, on the other hand plant deviation percentage increased with large aggregates ( $\phi 750\text{mm}$ ). Also, he found that the best aggregate size diameters for drill machine was ( $\phi 20\text{-}50\text{ mm}$ ) for good germination and distribution. **El-Hanafy et al (1995)** reported that using the chisel plough followed by rotary plough for barley preparation land is considered the best combination to get the highest barley yield. **Sherif et al (1995)** indicated that tillage treatments significantly affected all characters studied i.e plant height, ear length, stem diameter, ear weight, weight of grains per ear, grain yield per plant, 100 grains weight and grain yield per fad except percentage of barren plants, number of ears per plant and number of rows per ear. He further added that no tillage treatment reduced grain yield as compared with conventional tillage. **El-Douby et al (1996)** concluded that the highest maize grain yield was obtained when

(4:2) intercropping pattern applied and maize was spaced at 20 cm apart in both seasons. Whereas, the highest soybean yield was produced with (2:4) intercropping pattern which included 33% maize + 133% soybean in both seasons. They also added that intercropping increased "LER" values by 19% and 23% when (2:4) and (4:2) patterns when maize was sown at 20 cm apart in the first and second seasons, respectively. **Shalaby, (1988)** indicated that the effects of using chisel plough two passes followed by scraper, followed by wooden level 20 cm and chisel plough two passes followed by rotary plough once 20 cm gave high soil surface roughness of 35.30 and 32.60 % and clod mean weight diameter of 40.12 and 36.48 mm respectively. While using chisel plough two passes followed by rotary plough 15 cm depth gave roughness of 25.20% and M.W.D. of 22.00 mm. He also concluded that number of plants per  $\text{m}^2$  was affected greatly by soil profile and roughness. **El-Khatib, (2000)** summarized that the mean weight diameter (M.W.D.) after tillage with chisel plough one pass, chisel plough two passes and chisel plough two passes + disk harrow were 76.21, 60.23 and 43.35 mm respectively.

## MATERIAL AND METHODS

Two field experiments were carried out at Sids Agricultural Farm Research Station Bani-Swif Governorate during 2003 and 2004 growing seasons, to study the effect of tillage systems and intercropping patterns on the amount of applied water, water use efficiency and yield of maize and soybean. A split plot design with three replicates was used, the main plots were allocated for tillage treatments whereas the sub plots were

devoted for intercropping patterns. The sub-plot area was 42 m<sup>2</sup> (7m x 6m) i.e 1per100 feddan. The treatments were as follows:

#### A. Tillage treatments

- 1- Chisel plough 2 passes (10, 15 and 20cm depths).
- 2- Chisel plough 3 passes (10, 15 and 20cm depths).

#### B. The intercropping treatments

The intercropping patterns were

- 1- (2: 2) maize was grown on two ridges alternated with two ridges of soybean
- 2- (2: 4) maize was grown on two ridges alternated with four ridges of soybean
- 3- solid planting for both crops.

Soybean (cv. Clark) was seeded immediately after inoculation with Rhizobium bacteria to stimulate nodulation and irrigated at once. Seedling was carried out on 10<sup>th</sup> May, 2003 and 15<sup>th</sup> May, 2004 seasons. It was planted at 10 cm on both sides of all ridges and thinned to two plants per hole in all treatments. Maize (cv., Three Way Cross 310) was seeded at the first irrigation of soybean. It was seeded on 1<sup>st</sup> and 10<sup>th</sup> June, 2003 and 2004 seasons, at 30cm apart and thinned to two plants / hill, whereas plants grown in pure stand were spaced at the same distances, but thinned to one plant / hill.

#### C- The irrigation treatments

The developed surface irrigation used in this investigation is a new technique to transmit irrigation water, by means of a pump, from the main source (open canal)

to soil surface at the upper part of the field carrying water to be applied in the furrows through perforated 4-inch Aluminum line. Water meters of 0.1 cubic meter accuracy (to measure the amount of water applied) were attached to the network. The capacity of the used pump was up to 120 m<sup>3</sup>/feddandan head. The pump was connected to the main line by flexible quick hitch hose. Irrigation was applied based on 40% of soil moisture content and evapotranspiration "ET<sub>p</sub>". Frequency of irrigation was estimated described by Cuenca, (1989) as follows:

- 1- Total available moisture (TAW)

$$TAW = FC - CEW \quad (1)$$

where

TAW is the total available moisture (mm per m).

FC is field capacity

CEW is crop extractable water

- 2- Available moisture (AM) at 40% depletion

$$(AM_{40}) = \% \text{ depletion} \times TAW \times \text{root zone depth} \quad (2)$$

- 3- Frequency of irrigation (I<sub>fr</sub>)

$$I_{fr} = \frac{Am_{40}}{Et_{mgs}} \quad (3)$$

where: Et<sub>mgs</sub> is evapotranspiration at the midpoint of the growing season.

The quantity of water applied was estimated using the class A pan evaporation equation:

$$ET_p = K_p E_{pan} \quad (4)$$

Where

ET<sub>p</sub> = Evapo-transpiration of grass reference crop, mm per d

K<sub>p</sub> = pan coefficient

E<sub>pan</sub> = pan evaporation, mm per d.

The irrigation water was calculated on 100%  $ET_p$  basis and 100% water application efficiency, due to the even distribution of water within the strips and non-water losses, as a result to precision land leveling by laser technology on the following basis:

- 1- The measured evaporation from the A pan between irrigation rounds.
- 2- A pan coefficient = 0.8 for dry regions.
- 3- Average crop coefficient =1 for all stages of growth.
- 4- Evapo-transpiration Potential ( $ET_p$ ) =100%

Before starting the experimental, soil analysis was done. Table (1) shows the results of the mechanical analysis and the bulk density of the soil. Field capacity was 39.6 % by weight and the wilting point was 18 % by weight.

Each plot was fertilized with calcium superphosphate (15.5%  $P_2O_5$ ) at a rate of 150kg per feddan. it was applied during land preparation and before planting. Potassic fertilizer was added in the form of potassium sulphate (48%  $K_2O$ ) at the second irrigation at a rate of 50kg  $K_2O$  per feddan. N fertilizer as ammonium sulphate (20.5% N) was applied at a rate of 100 kg N per feddan. in 2 equal doses at the first and second irrigations. All the experimental treatments received the same agricultural practices as recommended.

### Measurement

At harvest time ten plants were taken at random from the middle rows of each treatment to estimate growth characters,

and quality of maize: plant height cm, height of first ear cm, ear length cm, ear diameter cm, average number of rows per ear, average number of kernels per row, weight of 100- grains.

Soybean: plant height cm, average number of fruiting branches per plant, average number of pods per plant, weight of pods per plant, weight of 100-seed.

Maize grain yield in kg per feddan as well as, yield of soybean in ton per feddan were calculated on hole plot basis.

### Methods of calculations

#### 1-Water use efficiency "WUE" (kg per $m^3$ )

WUE = yield (kg/feddan) per total applied water ( $m^3$ /feddan)

#### 2-Clod size distribution

The mean weight diameter "MWD" was determined according Ashery, (1985) as follow

$$M.W.D = \frac{W_1 S_1 + \dots + W_n S_n}{W}$$

$$M.W.D. = \frac{\Delta W_1 \Delta I}{W}$$

Where: M.W.D. is mean weight diameter (mm),  $W_i$  is soil weight on  $i$ th sieve,  $S_i$  is sieve number (at first),  $W_n$  is soil weight on  $S_n$ ,  $S_n$  is sieve number (at last),  $W$  is total weight of soil sample,  $I$  is sieve number and  $\Delta I$  is  $I^{th}$  sieve mesh (mm).

Table 1. Some physical properties of the experimental spoil

Depth cm	Coarse sand %	Fine sand %	Silt %	Clay %	Texture class	Organic matter %	CaCO <sub>3</sub> %	Bulk density cm <sup>3</sup>
(0-15)	4.67	15.96	18.50	60.48	Clay	5.50	3.50	1.10
(15-30)	4.50	13.50	19.00	63.00	Clay	5.00	4.0	1.09
(30-45)	4.90	14.00	18.60	62.50	Clay	2.00	3.90	1.15
(45-60)	3.50	15.50	16.00	65.00	Clay	2.00	3.50	1.15

### 3- Competitive relationships

#### 1- Land equivalent ratio (LER)

LER is determined as the sum of the fractions of the yield of the intercrops relative to their sole crop yields (Willey and Osiru, 1972). Land equivalent ratio LER was determined according to the following formula:

$$LER = \frac{Y_{ab}}{Y_a} + \frac{Y_{ba}}{Y_b}$$

Where:  $Y_{aa}$  is pure stand yield of species a,  $Y_{bb}$  is pure stand yield of species b,  $Y_{ab}$  is mixture yield of a (when combined with b) and  $Y_{ba}$  is mixture yield of b (when combined with a).

#### 2- Relative crowding coefficient (RCC)

This was proposed according to Hall (1974). It assumes that mixture treatment forms a replacement series. Each series has its own coefficient (K) which gives a measure to indicate that series has pro-

duced more, less or equal yield to that expected. Relative crowding coefficient (RCC) was determined according to the following formula: for species (a) in a mixture with species (b).

$$K_{ab} = \frac{Y_{ab} \times Z_{ba}}{(Y_{aa} - Y_{ab}) \times Z_{ab}}$$

Where:  $Z_{ab}$  is sown proportion of species a (in a mixture with b) and  $Z_{ba}$  is sown proportion of species b (in a mixture with a).

$$K_{ba} = \frac{Y_{ba} \times Z_{ab}}{(Y_{bb} - Y_{ba}) \times Z_{ba}}$$

If a species has a coefficient less than, equal to, or greater than 1, it means it has produced less yield, the same yield, or more yield than the "expected", respectively. The component crop with the higher coefficient is the dominant one. To determine if there is a yield advantage of mixing, the product of the coefficient of mixing, the product of the coefficient is formed by multiplying  $K_{ab} \times K_{ba}$ . If  $K > 1$ , there is a yield advantage, if  $K = 1$  there is no difference and if  $K < 1$  there is a yield disadvantage.

### 3- Aggressively (A)

This parameter was proposed by Mc Gilchrist, (1960). It gives a simple measure of how much the relative yield increase in species (a) is greater than that of species (b). Aggressivity "A" is determined according to the following formula:

$$A_{ab} = \frac{\text{Mixture yield of a} - \text{Expected yield of a}}{\text{Expected yield of a}} - \frac{\text{Mixture yield of b} - \text{Expected yield of b}}{\text{Expected yield of b}}$$

$$A_{ab} = \frac{Y_{ab}}{Y_{aa} \times Z_{ab}} - \frac{Y_{ba}}{Y_{bb} \times Z_{ba}}$$

An aggressively value of zero indicates that the component species are equally competitive. For any other situation, both species will have the same numerical value but the sign of the dominant species will be positive and the dominated negative. The greater the numerical value the bigger the difference in competitive abilities and the bigger the difference between actual and "expected" yield.

## RESULTS AND DISCUSSION

### 1- Effect of tillage passes and ploughing depth on soil clods size distribution

Data in Table (2) showed that the highest percentage (39%) of the largest soil clods size (more than 50 mm) was obtained by tillage 2 passes with chisel plough at 20 cm depth. While the lowest percentage (18 %) was obtained by tillage

3 passes at 10 cm depth this may be due to increase the depth gave big aggregates and 2 passes gave less breakdown. The data revealed that using chisel plough 3 passes increased the percentage of small soil clods size (less than 50 mm). Also, the data indicated that the differences in M.W.D. were affected by tillage depth and tillage system. Using chisel plough 3 passes decreased the value of M.W.D. compared with using chisel plough 2 passes by 33.00 %, 27.92% and 31.87% for 10, 15 and 20cm depths, respectively.

### 2- Effect of tillage passes and ploughing depth on water applied, water use efficiency, yield, and quality of maize intercropped with soybean

Data on yield and quality as well as growth traits of maize are presented in Table (3). The data indicated significant differences, except in case of plant height (cm), height of first ear (cm), ear length (cm), number of rows per ear in both seasons.

Tillage 2 passes with chisel plough at 15cm depth gave lower quality as well as lower grain yield compared with tillage 3 passes at 15cm depth in both seasons

On other hand, yield of maize increased by 10.50, 12.28 and 13.82 % when chisel plough was done for 3 passes compared with 2 passes at 10, 15 and 20cm depths respectively, in 2003 season and 6.46, 12.07 and 9.49 % in 2004 season respectively. The higher yield was obtained by tillage 3 passes at 15cm depth. These results were supported by those obtained by El-Sayed, (1983) who revealed that tillage using one pass of chisel plough was not suitable for mechanized planting where the surface of soil

Table 2. Effect of tillage treatments on soil clods size distribution

Tillage treatments	Ploughing Depth, cm	Soil clods size distribution, %				M.W.D. mm
		5-10 mm	10-20 mm	20-50 mm	> 50 mm	
Chisel	10	17	21	37	25	30.12
plough	15	15	19	34	32	36.35
2 passes	20	10	16	35	39	41.80
Chisel	10	29	23	30	18	20.18
plough	15	17	28	33	22	26.20
3 passes	20	12	24	35	29	28.48

Table 3. Effect of tillage passes and depth of ploughing on water applied, yield and quality, and water use efficiency of maize intercropped with soybean in 2003 and 2004 seasons

Treatments <sup>a</sup> characters	2003 season						LSD at 0.05
	Chisel plough 2 passes			Chisel plough 3passes			
Ploughing depth (cm)	10	15	20	10	15	20	
Plant height (cm)	293.56	296.68	301.33	299.45	303.64	309.74	N. S
Height of 1 <sup>st</sup> ear (cm)	80.37	86.07	88.83	91.93	93.43	97.33	N. S
Ear length (cm)	18.49	20.78	16.72	20.40	22.78	16.70	N. S
Ear diameter (cm)	4.15	4.05	4.34	4.59	5.30	4.07	0.45
Av. no. of rows per ear	12.03	12.57	11.50	13.00	13.47	12.50	N. S
Av. no. of kernels per row	40.96	45.98	37.94	44.99	48.91	41.40	1.61
Wt. of 100-grain (gm)	30.27	32.70	29.73	35.40	38.93	31.53	2.28
Yield (kg per fed.)	2443.33	2525.00	2306.67	2730.0	2878.33	2676.67	128.800
Water applied (m <sup>3</sup> per fed)	2332.78	2366.11	2374.44	2377.78	2399.44	2418.89	11.30
W.U.E (kg per m <sup>3</sup> )	1.05	1.07	0.97	1.15	1.20	1.11	0.035

Table 3. Cont.

Treatments characters	2004 season						LSD at 0.05
	Chisel plough 2 passes			Chisel plough 3passes			
Ploughing depth (cm)	10	15	20	10	15	20	
Plant height (cm)	285.22	289.77	293.96	290.54	298.11	303.50	N.S
Height of 1 <sup>st</sup> ear (cm)	77.80	83.07	85.87	87.93	90.70	93.30	N.S
Ear length (cm)	18.91	20.60	15.62	19.67	21.17	17.53	N.S
Ear diameter (cm)	4.53	4.80	3.85	4.55	5.27	4.13	0.79
Av. no. of row per ear	11.30	11.90	10.67	11.83	12.17	11.23	N.S
Av. no. of kernelsper row	39.43	43.08	34.40	43.03	47.60	38.99	0.363
Wt. of 100- grain (gm)	27.43	30.03	25.90	31.57	35.50	27.43	1.15
Yield (kgperfad)	2365.00	2448.33	2226.67	2528.33	2784.33	2460.00	95.28
Water applied (m <sup>3</sup> perfad)	2338.33	2350.11	2371.50	2380.33	2443.44	2480.89	17.59
W.U.E (kg perm <sup>3</sup> )	1.01	1.04	0.94	1.06	1.14	0.99	0.031

being big clods and seed were not surrounded by uncompacted soil and lease bulk density.

Data in the same table, indicated that water applied (m<sup>3</sup> per fad) of maize were reduced by (1.93, 1.41 and 1.87%) when chisel plough was done for 2 passes compared with 3 passes at depth at 10, 15 and 20cm respectively, in 2003 whereas, in 2004 season it was (1.79, 3.97 and 4.61%) for the same respective treatments. These results are in agreement with those obtained by El-Sayed, (1983) who found that the plant root system was small to catch up with water percolation through the soil. This lead to increase of

water consumption in the deeply ploughed system. Also the W.U.E select the same trend and reduced by 9.52, 12.15 and 14.43% in 2003 season whereas, in 2004 season it was 4.95, 9.62 and 5.34 %.

### 3- Effect of intercropping patterns on water use efficiency, yield and its components of maize intercropped with soybean

Data in Table (4) showed that growth of maize in monoculture was significantly higher than in other intercropping combinations. These results were supported by



Table 4. Effect of intercropping patterns on WUE, yield, and its components of maize intercropped with soybean in 2003 and 2004 seasons

characters	Intercropping patterns			LSD at 0.05
	2003 season			
	2:2	2:4	solid	
Plant height (cm)	295.20	301.23	305.81	N.S
Height of 1 <sup>st</sup> ear (cm)	81.80	90.20	97.18	3.1
Ear length (cm)	16.84	19.13	21.96	0.632
Ear diameter (cm)	4.28	4.50	4.74	0.708
Av. no. of rowsper ear	12.42	12.53	13.03	N.S
Av. no. of kernelsper row	39.56	46.83	48.50	1.97
Wt. of 100-grain ( gm)	29.25	32.53	36.90	3.30
Yield (kgperfad)	2577.50	2260.83	2941.67	502.0
Water applied (m <sup>3</sup> perfad)	2449.44	2397.11	2338.67	15.75
W.U.E (kg per m <sup>3</sup> )	1.05	0.94	1.26	0.033

Table 4. Cont.

characters	Intercropping patterns			LSD at 0.05
	2004 season			
	2:2	2:4	solid	
Plant height (cm)	288.52	294.34	297.69	N.S
Height of 1 <sup>st</sup> ear (cm)	78.58	86.72	93.43	2.75
Ear length (cm)	17.69	18.42	20.73	0.607
Ear diameter (cm)	4.25	4.44	4.88	0.519
Av. no. of rowsper ear	10.97	11.58	12.15	N.S
Av. no. of kernelsper row	37.99	40.69	44.58	0.942
Wt. of 100-grain (gm)	26.37	29.30	33.27	4.61
Yield (kgperfad)	2500.50	2122.50	2791.67	421.40
Water applied (m <sup>3</sup> perfad)	2456.67	2372.00	2348.57	20.33
W.U.E (kg per m <sup>3</sup> )	1.02	0.89	1.19	0.022

those obtained by Kamel *et al* (1990) and El-Douby, (1992). The detrimental effect of intercropping on growth characters of maize plants may be due to the increase in plant density per unit area of both components. Maize density was estimated to 67% of maize Population in solid planting when maize was intercropped with soybean in (2:4) pattern. The adverse effects appeared more conspicuous when maize grown in (2:2) pattern. This is attributed more to inter and intra competition between plants as a result of the heavy density of plants per unit area.

Data on maize quality indicted clearly that ear length, number of rows per ear, number of kernels per row and weight of 100 grains of solid planting were superior to those in all intercrop associations, except, in case of average number of kernels per row in (2:4) pattern in 2003 season. However, estimated values for all traits of maize plants grown in (2:4) pattern were higher than those plants grown in (2:2) pattern. Kamel *et al* (1990) found that yield of maize grown in (2:2) pattern was higher than those grown in (2:4) pattern. It seemed that maize yield in the intercrop combination was closely parallel to maize density, interpreting superiority of maize yield in (2:2) pattern over that in (2:4) pattern.

Data on yield of grains per Fadden showed that none of the intercropping patterns was able to give yield equal to or more than solid maize treatment. The estimated excesses in yield of solid maize treatment over (2:2) and (2:4) patterns were 12.38 and 23.15 % in 2003 season and 10.43 and 23.97 % in 2004 season respectively. Several investigators verified these results Kamel *et al* (1990) and El-Douby (1992).

On other hand, the data revealed that the water applied increased by 4.52 and 2.44% over those in pure stand when intercropping patterns (2:2 and 2:4) were applied in 2003 season respectively whereas, in 2004 season it was 4.40 and 1.00 % for the same respective patterns. When the highest water use efficiency was obtained when maize grown in pure stand than those grown under different intercropping combinations it was reduced by (20.00 and 34.04%) when (2:2 and 2:4) patterns were applied in 2003 season. In 2004 season the reduction was (16.67 and 33.71%) for the same selective patterns. These results were supported by El-Khatib and Sherif, (2003).

#### 4- Interaction effect of tillage treatments, depth of ploughing and intercropping patterns on water applied, yield and quality of maize intercropped with soybean

The interaction effect of tillage treatments and intercropping patterns on growth, yield and quality and water applied of maize is presented in Table (5). Data indicated that differences between treatments were not great enough to reach the 5% level of significance, except in case of ear diameter and water applied in 2003 and 2004 seasons and ear length in 2004 season. Moreover, maximum plant height and height of first ear were obtained when maize plants were grown in pure stand and ploughed 3 passes with the chisel plough at 20cm depth, whereas minimum values were obtained in (2: 2) pattern and chisel plough was done 2 passes at 10cm depth..

The interaction effect on maize yield per feddan was relatively influenced by maize population. However, none of the

Table 5. Interaction effect of tillage treatments and intercropping patterns on water applied, yield and quality of maize intercropped with soybean in 2003 and 2004 seasons

Characters		2003 season										
		Plant height (cm)	H. of first ear (cm)	Ear Length (cm)	Ear diameter (cm)	No of rows per ear	No of kernels per row	Wt. of 100 grain (gm)	Yield (Kg per fad)	Water applied (m <sup>3</sup> per fad)	WUE (kg per m <sup>3</sup> )	
Treatments												
Chisel plough 2 passes	2:2	10	287.67	75.50	14.93	4.30	11.50	37.27	27.10	2350.00	2401.60	0.98
		15	290.87	78.30	16.83	4.50	12.00	42.53	29.50	2420.00	2430.00	0.99
		20	296.23	80.10	13.37	4.00	11.00	35.43	25.00	2150.00	2453.33	0.88
	2:4	10	294.33	80.50	19.13	4.50	12.10	41.10	30.10	2150.00	2350.67	0.92
		15	298.67	86.60	22.03	4.90	12.60	47.10	32.50	2275.00	2386.33	0.96
		20	300.43	90.30	16.30	4.10	11.40	38.30	28.10	1975.00	2410.00	0.82
	Solid maize	10	298.67	85.10	21.40	3.65	12.50	44.50	33.60	2830.00	2270.00	1.25
		15	300.50	93.30	23.47	4.35	13.10	48.30	36.10	2880.00	2290.00	1.26
		20	307.33	96.10	20.50	4.91	12.10	40.10	32.50	2795.00	2330.10	1.20
Chisel plough 3 passes	2:2	10	291.70	85.10	19.00	4.15	12.50	39.13	30.30	2810.00	2453.33	1.15
		15	296.50	88.50	21.93	5.10	12.80	45.47	35.50	2975.00	2471.67	1.20
		20	308.20	83.30	15.00	3.61	12.00	37.50	28.10	2796.00	2486.67	1.12
	2:4	10	300.33	9.60	20.10	4.52	13.00	45.37	35.60	2380.00	2400.67	0.99
		15	305.53	94.70	22.10	5.29	13.60	48.63	38.2	2510.00	2410.00	1.04
		20	308.10	98.50	15.10	3.70	12.50	41.70	30.70	2275.00	2425.00	0.94
	Solid maize	10	306.33	98.30	22.10	5.10	13.50	50.47	40.30	3000.00	2295.30	1.25
		15	309.10	100.1	24.30	5.50	14.00	52.63	43.10	3150.00	2323.30	1.30
		20	312.93	110.20	20.00	4.90	13.00	45.00	35.80	2995.00	2623.33	1.14
1.SD5%		N. S	N. S	N. S	0.775	N. S	N. S	N. S	N. S	7.516	N. S	

Table 5. Cont.

Treatments		2004 season										
		Plant height (cm)	H. of first ear (cm)	Ear Length (cm)	Ear diameter (cm)	No of rows per ear	No of kernels per row	Wt. of 100 grain (gm)	Yield (Kg per fad)	Water applied (m <sup>3</sup> per fad)	WUE (kg per m <sup>3</sup> )	
Chisel plough 2 passes	2:2	10	280.50	73.10	18.43	4.23	10.70	34.53	24.50	2310.00	2366.67	0.98
		15	287.67	75.00	19.03	4.44	11.30	40.83	26.10	2385.00	2411.67	0.99
		20	290.50	77.90	14.20	3.61	10.10	30.50	22.80	2125.00	2433.33	0.87
	2:4	10	287.20	77.30	18.00	4.36	11.30	40.07	27.10	2035.00	2325.0	0.88
		15	290.13	8per4.10	20.17	4.83	11.80	43.30	30.50	2185.00	2363.33	0.92
		20	292.27	87.20	15.17	3.78	10.60	35.10	25.10	1955.00	2380.00	0.82
	Solid	10	287.95	83.00	20.30	4.99	11.90	43.70	30.70	2750.00	2293.33	1.20
	maize	15	291.50	90.10	22.60	5.13	12.60	45.10	33.50	2775.00	2275.00	1.22
		20	299.10	92.50	17.50	4.16	11.30	37.60	29.80	2600.00	2301.10	1.13
Chisel plough 3 passes	2:2	10	285.23	80.10	18.90	4.30	11.30	40.10	28.50	2650.00	2483.33	1.07
		15	290.60	85.50	20.00	4.94	11.80	47.00	31.30	2953.00	2518.33	1.17
		20	296.60	79.90	15.00	3.96	10.60	35.00	25.00	2580.00	2526.67	1.02
	2:4	10	290.63	86.50	19.10	4.56	11.90	42.00	30.70	2135.00	2356.67	0.98
		15	300.60	90.10	21.00	5.13	12.60	45.50	35.10	2350.00	2397.00	0.98
		20	305.20	95.10	17.10	4.00	11.30	38.17	27.30	2025.00	2410.00	0.84
	Solid	10	295.75	93.60	21.00	4.78	12.30	47.00	35.50	2800.00	2301.00	1.22
	maize	15	303.13	96.50	22.50	5.75	13.00	50.30	40.10	3050.00	2415.00	1.26
		20	308.70	104.90	20.50	4.44	11.80	43.80	30.00	2775.00	2506.00	1.10
LSD 0.05			N. S	N. S	0.624	0.083	N. S	N. S	N. S	N. S	12.92	N. S

intercropping patterns exceeded the pure stand. The excess in yield of maize grown in pure stand and ploughed 3 passes at 15cm depth over those grown in (2:2) pattern was 5.56 and 3.28 % in 2003 and 2004 seasons respectively, whereas, the excess over those grown in (2:4) pattern was 15.40 and 22.95 % in 2003 and 2004 seasons, respectively.

In addition, maximum value of WUE was obtained when maize plants were grown in pure stand and ploughed 3 passes at a depth of 15cm whereas, the minimum values were obtained when plants were grown in (2:4) pattern and ploughed 2 passes at a depth of 30cm in the both seasons .

#### **5- Effect of tillage systems treatments on water applied, water use efficiency, yield and quality of soybean intercropped with maize**

Data in Table (6) showed that soybean growth characters, yield and quality were affected by increasing tillage passes and ploughing depth. These results were true in cases of plant height, number of branches per plant, water applied in both seasons. Ploughing 3 passes at 15cm depth gave higher quality as well as higher grain yield compared with other treatments in both seasons, except in case of number of pods per plant in both seasons.

On other hand, data revealed that the yield of soybean increased by 15.18, 12.48 and 13.32 % when ploughing 3 passes with chisel ploughs compared with 2 passes at 10, 15 and 20cm depths in 2003 season, respectively and 18.64, 15.30 and 15.72 % in 2004 season respectively. It is clear that higher yield of soybean was obtained when soybean plants

were grown under 3 passes with chisel plough at 15cm depth in both seasons.

Data in the same table indicated that the water use efficiency (kg per m<sup>3</sup>) of soybean has the higher value when ploughing 3 passes at 15cm depth in both seasons. These results are in agree with those obtained by El-Sayed, (1983).

#### **6- Effect of intercropping patterns on yield and quality of soybean intercropped with maize**

Data in Table (7) showed significant effects on plant height, average number of pods per plant in both seasons. However data analysis showed that soybean growth in pure stand was significantly higher than on other any intercrop combinations. In addition, values of growth characters of soybean grown in (2:4) pattern were mostly higher than those obtained in (2:2) pattern. Growing two rows of maize alternated with four rows of soybean had the highest values, whereas in case of two rows of maize alternated with two rows of soybean possessed the least values. These results are in agreement with those obtained by Kamel *et al* (1990). They revealed a general tendency towards more growth vigor and weight when plants grow in row strip alternated with two rows of maize. However, the general increase in growth characters of soybean plants grown in (2:4) pattern may be due to more light intercepted by foliage as well as the low below and above ground competition between both components in the mixture. On other hand, the minimum growth values were associated with (2:2) pattern may be due to low light intensity owing to the shade of maize plants.

Table 6. Effect of tillage treatments on water applied, WUE, yield and quality of soybean intercropped with maize in 2003 and 2004 seasons

characters	Treatments						L.S.D
	Chisel plough 2 passes			Chisel plough 3 passes			
	10	15	20	10	15	20	
Ploughing depth (cm)							0.05
Plant height (cm)	63.63	66.41	70.07	65.28	69.47	72.89	N. S
No. of branches per plant	2.26	2.53	3.14	2.52	2.65	3.28	N. S
No. of pods per plant	44.81	47.40	52.04	48.52	51.61	52.18	0.323
Weight of pods per plant (gm)	89.30	93.07	89.60	96.10	102.80	94.43	N.S
Weight of 100 seeds (gm)	16.96	17.31	17.13	18.28	18.61	18.03	N. S
Yield (kg per feddan)	885.44	970.26	809.29	1043.87	1108.61	933.67	22.45
Water applied (m <sup>3</sup> per feddan)	2269.45	2312.78	2368.89	2286.67	2325.00	2366.67	9.142
WUE (kg per m <sup>3</sup> )	0.390	0.420	0.340	0.461	0.480	0.400	0.006
	2004 season						
Plant height (cm)	58.53	64.24	68.12	62.27	65.08	69.66	N. S
No. of branches per plant	2.21	2.43	2.74	2.31	2.54	2.95	N. S
No. of pods per plant	42.93	45.22	48.42	45.98	48.70	52.15	N. S
Weight of pods per plant (gm)	85.97	91.30	84.70	97.13	103.37	89.97	N.S
Weight of 100 seeds (gm)	16.51	16.98	16.43	17.96	18.30	17.71	0.108
Yield (kg per fed.)	824.21	915.45	773.62	1013.03	1070.43	917.97	90.36
Water applied (m <sup>3</sup> per fed.)	2237.22	2285.00	2325.00	2293.56	2328.00	2394.45	14.14
W.U.E (kg per m <sup>3</sup> )	0.368	0.401	0.333	0.442	0.460	0.383	0.002

Table 7. Effect of intercropping patterns on yield and quality of soybean intercropped with maize in 2003 and 2004 seasons

Intercropping patterns Characters	2003 season				2004 season			
	2:2	2:4	solid	L.S.D.	2:2	2:4	solid	L.S.D.
Plant height (cm)	63.74	67.66	72.47	2.13	61.78	64.57	67.59	0.78
No. of branches per plant	2.53	2.71	2.96	N.S	2.37	2.44	2.79	N.S
No. of pods per plant	46.67	48.53	54.53	0.852	45.46	46.79	49.45	0.603
Weight of seeds per plant (gm)	82.32	95.70	104.63	11.0	78.50	92.45	105.32	9.03
Weight of 100 seeds (gm)	17.23	17.69	18.25	0.262	16.86	17.38	17.70	0.283
Yield (kg per feddan)	677.66	953.73	1244.18	250.49	626.93	924.67	1205.76	200.03
Water applied (m <sup>3</sup> per feddan)	2449.44	2385.28	2130.00	11.60	2456.67	2388.61	2086.34	17.32
W.U.E (kg per m <sup>3</sup> )	0.277	0.399	0.584	0.004	0.255	0.387	0.578	0.002

Data revealed that yield of soybean in pure stand in the two seasons were significantly higher than that grown in (2:2) pattern as well as those grown in (2:4) pattern.

In this respect, Kamel *et al* (1990) reported that increases in yield of soybean were closely parallel with the increases of soybean ratio in the intercrop pattern. Increases in soybean yield associated with (2:4) may be related to the increase in soybean population as compared with (2:2) pattern.

The data also indicated that the highest WUE were obtained when soybean was grown in pure stand than in (2:2 and 2:4) patterns in both seasons. Whereas the highest value of water applied was obtained when soybean was grown in (2:2) pattern. These results are in agreement with those obtained by El-Khatib and Sahar (2003).

#### 7- Interaction effect of intercropping patterns and tillage treatments on water applied, WUE, yield and quality of soybean

The interaction effect of tillage treatments and intercropping patterns on WUE, water applied, yield and quality of soybean were not significant except in cases of plant height and yield per feddan in the second season, as well as, water applied and WUE in both seasons presented in Table (8). Values of yield and quality of soybean plants in (2:4) pattern were relatively superior to that in (2:2) pattern. None of the intercrop combinations exceeded the solid planting of soybean under any treatment of tillage system. It is also clear that the highest yield

of soybean in the intercrop combinations was obtained with (2:4) pattern and ploughing with chisel plough 3 passes at 15cm depth, whereas the least yield was associated with (2:2) pattern and ploughing with chisel plough 2 passes at 20cm depth.

Moreover, it is evident that maximum value of WUE was obtained when soybean plants were grown in pure stand and ploughing with chisel plough 3 passes whereas the minimum values were obtained when plants were grown in (2:2) pattern and ploughing with chisel plough 2 passes.

#### 8- Competitive relationships

Intercropping patterns exhibited effects on the relative yield of maize, as well as, the RY of soybean (Table 9). Highest RY value for maize was obtained in (2:2) pattern. These results seemed coincided with maize densities in the mixture. Kamel *et al* (1990) came to a similar conclusion. Highest RY value of soybean was obtained with (2:4) pattern. This increase in RY was associated with the increase of soybean population in the mixture. Results indicated that the highest LER value was obtained when both crops were oriented in (2:4) pattern. The reduction in LER associated with (2:2) pattern were 7.14 and 7.12 below LER values for (2:4) in both seasons respectively.

Relative crowding coefficient (RCC) followed a similar trend as in LER (Table 9). K and the total RCC values were superior when both components were oriented in (2:4) pattern. Results hold true in both seasons. Several investigators supported these results such as Kamel *et al* (1990) and Sherif (1993).



Table 8. Interaction effect of intercropping patterns and tillage treatments on water applied, WUE yield and quality, of soybean intercropped with maize in 2003 and 2004 seasons

Treatments		Characters		2003 season						
		Plant height (cm)	No. of branches perplant	No. of pods perplant	Wt. of pods perplant (gm)	Wt. of 100-seeds (gm)	Yield (Kgperfeddan)	Water applied (m <sup>3</sup> perfeddan)	W.U.E (kg perm <sup>3</sup> )	
Chisel plough 2 passes	2:2	10	59.23	2.17	42.80	81.10	16.13	603.70	2401.67	0.25
		15	62.37	2.43	44.73	87.70	16.67	750.27	2430.00	0.31
		20	63.90	2.63	48.77	76.10	17.13	550.47	2453.33	0.22
	2:4	10	63.63	2.10	43.97	92.10	17.07	825.50	2326.67	0.36
		15	66.47	2.27	46.13	94.0	17.30	885.50	2378.33	0.37
		20	70.03	3.40	49.53	88.70	16.90	750.27	2400.00	0.31
	Solid soybean	10	68.03	2.50	47.67	94.70	17.67	1227.13	2080.00	0.59
		15	70.40	2.90	51.33	97.50	17.97	1275.0	2130.00	0.60
	20	76.27	3.40	57.83	104.00	17.37	1127.13	2253.33	0.50	
Chisel plough 3 passes	2:2	10	61.87	2.53	45.37	82.00	17.77	730.50	2453.33	0.30
		15	65.97	2.63	47.17	90.00	18.07	830.67	2471.67	0.34
		20	69.10	2.77	51.17	77.00	17.63	600.33	2486.67	0.24
	2:4	10	64.70	2.47	48.50	99.50	18.27	1100.7	2356.67	0.47
		15	69.33	2.60	51.03	103.90	18.60	1185.17	2403.33	0.49
		20	71.80	3.43	52.03	96.00	17.97	975.25	2446.67	0.40
	Solid soybean	10	69.27	2.57	51.70	106.80	18.80	1300.40	2050.00	0.63
		15	73.10	2.73	56.33	114.50	19.17	1310.00	2100.00	0.62
	20	77.77	3.63	62.33	110.30	18.50	1225.43	2166.67	0.57	
LSD5%		N.S	N.S	N. S	N.S	N.S	N.S	75.16	0.002	

Table 8. Cont.

Treatments	Characters		2004 season							
	Plant height (cm)	No. of branches perplant	No. of pods perplant	Wt. of pods perplant (gm)	Wt. of 100-seeds (gm)	Yield (Kgperfeddan)	Water applied (m <sup>3</sup> perfeddan)	W.U.E (kg perm <sup>-3</sup> )		
Chisel plough 2 passes	2:2	10	55.47	2.03	42.27	77.30	16.10	550.25	2366.67	0.23
		15	60.83	2.20	43.43	81.50	16.53	635.70	2411.67	0.26
		20	63.20	2.53	46.70	70.30	16.00	500.10	2433.33	0.21
	2:4	10	58.50	2.07	42.83	90.10	16.43	785.50	2325.00	0.34
		15	63.70	2.37	44.93	92.60	16.90	875.50	2363.33	0.37
		20	68.53	2.67	47.57	85.50	16.50	700.50	2380.00	0.29
	Solid soybean	10	61.63	2.53	43.70	90.50	17.00	1136.87	2020.00	0.56
		15	68.20	2.73	47.30	99.80	17.50	1235.14	2080.00	0.59
		20	72.63	3.03	51.00	98.30	16.80	1120.26	2161.67	0.52
Chisel plough 3 passes	2:2	10	61.20	2.27	45.03	80.30	17.50	700.00	2483.33	0.28
		15	63.50	2.43	46.07	85.70	17.77	775.50	2518.33	0.31
		20	66.50	2.73	49.27	75.60	17.27	600.00	2526.67	0.24
	2:4	10	61.97	2.23	46.83	95.30	18.13	1085.50	2386.67	0.54
		15	65.23	2.47	48.70	98.70	18.47	1150.70	2415.00	0.48
		20	69.50	2.83	49.87	92.50	17.87	950.30	2461.67	0.39
	Solid soybean	10	63.63	2.43	46.08	115.80	18.25	1253.60	2010.67	0.62
		15	66.50	2.73	51.33	125.70	18.67	1285.10	2050.67	0.63
		20	72.97	3.30	57.30	101.80	18.00	1203.60	2195.00	0.55
LSD5%		0.758	N.S	N.S	N.S	N.S	15.60	129.20	0.005	

Table 9. Effect of intercropping patterns on competitive relationships in 2003 and 2004 seasons

Treatments	2003 season							
	RY		LER	K		RCC	Agg	
	Maize	soybean		maize	soybean		maize	soybean
2:2	0.88	0.55	1.43	7.08	1.20	8.50	+0.66	-0.66
2:4	0.77	0.77	1.54	6.74	1.62	10.92	+1.19	-1.19
2004 season								
2:2	0.90	0.52	1.42	8.59	1.08	9.28	+0.75	-0.75
2:4	0.76	0.77	1.53	6.44	1.62	10.43	+1.16	-1.16

Data on aggressively (Agg). revealed that the least value was associated with (2:2) pattern. Moreover, maize was always the dominant component while soybean was the dominated in all intercrop combinations in both seasons. These results were concordant with those obtained by Attia and El-Bially, (1990)

#### REFERENCES

- Abo-Habaga, M.M. (1992). Influence of Soil Clods Size Distribution in Seedbed on the Applicability of a Grain Drill Machine for Wheat. *Misr J. Agric. Eng. 9(1): 58-66.*
- Ashery, E.S.R (1985). *Development of the Locally Made Chisel Plough* pp. 73-74. M.Sc. Thesis., Fac. Agric., Saba Basha, Alex. Univ., Egypt.
- Attia, A.M. and M.E. El-Bially (1990). Interspecific competition and yield advantage of some summer crops as affected by some intercropping patterns and nitrogen rates. *Proc. 4<sup>th</sup> Conf. Agron. Cairo, Vol. II: 613-625.*
- Cuenca, R.H. (1989). *Irrigation System Design: An Engineering Approach*. pp. 120-125. Prentice-Hall, Englewood Cliffs, NJ.
- El-Douby, A.K. (1992). *Responses of Maize Grain Yield to Different Intercropping Patterns* pp. 192-197. Ph. D. Thesis, Fac. Agric., Moshtohor, Zagazig Univ., Egypt.
- El-Douby, A.K.; K.E. El-Habbak; H.E. Khalil and Zahira M. Attia (1996). Effect of some intercropping patterns on growth and yield of maize and soybean. *Annals of Agric. Sci., Moshtohor, 34(3): 919-933.*
- El-Hanafy, E.H.; H. Abd-Alla and S.M. Radwan (1995). Effect of Some Seedbed Preparation on Seed Drill Performance, Soil Physical Properties and Barely Yield. *Misr J. Agric. Eng., 12(3): 591-600.*
- El-Khatib, S.I. (2000). Effect of Tillage System on Laser Land Leveling Efficiency. *Arab Univ. J. Agric Sci., Ain Shams Univ., Cairo, 8(1): 31-40.*
- El-Khatib, S.I. and Sahar A.M. Sherif (2003). Effect of Soybean – Maize Inter-

- cropping and Laser Land Leveling on Water Use Efficiency, Plant Growth, Yield and Total Net Return, *Egypt, J. Agric. Res.*, 81(2): 699-717.
- El-Nakib, A.A. and H.A. Fouad (1990). Effect of Minimum Tillage With Conditioner Implement On Soil Physical Properties. *Miser J. Agric. Eng.* 7(2):121-131.
- El-Sayed, G.H. (1983). *Mechanized Techniques for Suitable Seedbed Preparation and Planting of Maize*, pp. 60-73 M.Sc. Faculty of Agric. Agric Eng. Department, Zagazig Univeristy, Egypt
- Hall, R.L. (1974). Analysis of the nature of niterfrenc between plants of different species. *Aust. J. Agric. Res.*, 25:749-756.
- Kamel, A.S.; M.N. Sherif; M.A. El-Masry; S.K. Badr and M.A. Abd-El-Aziz (1990). Intercropping maize with soybean as influenced by different patterns. *Agric. Rev.* 28(1): 1-19.
- Mc Gilchrist, C.A. (1960). Analysis and competition experiments *Biometrics*, 21: 975-985.
- Shalaby, S.A. (1988). *Effect of Seed bed Preparation Systems on Wheat Transplanting*, pp. 105-109. M.Sc. Thesis, Faculty of Agric., Mansoura Univ., Egypt.
- Sahar, A.M. Sherif (1993). *Studies on Fertilizer Requirements under Different Patterns of Soybean – Maize Intercropping under Some African Environmental Conditions* pp. 88-96. M.Sc. Institute of African research and studies. Natural Resources Department Cairo University, Cairo.
- Sherif, M.N.; S.K. Badr and A.M. Aly (1995). Effect of Some Preceding Winter Crops And Tillage on Growth, Yield And Quality of Maize. *Egypt. J. Applied Sci.*; 10 (12): 127 – 138.
- Willey, R.W. and S.O. Osiru (1972). Studies on mixture of maize and beans (*phasolus vulgaris*) with particular reference to plant populations. *J. Agric. Sci. Cambridge*, 79: 519-529.

مجلة نشاء الجامعات العربية للدراسات والبحوث الزراعية ، جامعة عين شمس ، القاهرة ، ١٤ (١) ، ١٧٣-١٩٣ ، ٢٠٠٦

## دراسة تأثير نظم الحرت ونظم تحميل فول الصويا مع الذرة الشامية على كفاءة استخدام مياه الري و مكونات المحصول

111

سحر على محمود شريف<sup>١</sup> - وفاء خميس محمد<sup>١</sup> - سحر طلعت ابراهيم<sup>١</sup> -  
حسن البنا عثمان<sup>٢</sup> - صلاح الدين اسماعيل الخطيب<sup>٢</sup>

١- قسم انتكثيف المحصولي - معهد بحوث المحاصيل الحقلية - مركز البحوث الزراعية - الجيزة- مصر  
٢- معهد بحوث الهندسة الزراعية - مركز البحوث الزراعية - النقى - الجيزة- مصر

كما أشارت البيانات الى أن محصول فول الصويا ومكوناته سلك نفس الاتجاه بالنسبة لمعاملات الخدمة بينما كان أفضل محصول ناتج تحت ظل نظامي التحميل كان عند زراعة فول الصويا في نظام تبادلي (٤:٢) مقارنة بنظام (٢:٢).

أوضحت البيانات أن معدل كفاءة استخدام المياه قد اختلف باختلاف المعاملات فكان أعلا قيمة سجلت عند تحميل فول الصويا مع الذرة الشامية تحت نظام (٤:٢) والحرت بثلاثة أوجه مع عمق ١٥ سم

كما أوضحت البيانات أن معاملات التجربة قد أدت الى زيادات في قيم معدل كفاءة استخدام الأرض وكذا معامل الحشد النسبي. ولم تزدى هذه المعاملات إلى خلق ضغوط تنافسية محسوسة وذلك عند قياس قيم العدوانية. من ناحية أخرى فإن زراعة الذرة محملا مع فول الصويا في نظام تبادلي (٤:٢) قد أدى الى أعلا قيمة في معدل كفاءة استغلال الأرض.

أجريت تجربتان حقليتان بمحطة بحوث سدس بمحافظة بني سويف خلال عامي ٢٠٠٣ ، ٢٠٠٤ وذلك لدراسة تأثير عدد أوجه الحرت وأعماق مختلفة (١٠ ، ١٥ ، ٢٠ سم) في ظل نظم تحميل مختلفة للذرة الشامية وفول الصويا ٢ خط ذرة شامية : ٢ خط فول صويا (٢:٢) ، ٢ خط ذرة شامية: ٤ خط فول صويا (٤:٢) على المحصول ومكوناته . كذلك دراسة تأثير المعاملات السابقة على معدل كفاءة استغلال الأرض وكفاءة استخدام مياه الري.

أشارت النتائج المتحصل عليها أن مكونات محصول الذرة والمحصول قد تأثرت بعدد أوجه الحرت وعمقه حيث نجد أن أفضل محصول تم الحصول عليه عندما تم الحرت ثلاثة أوجه على عمق ١٥ سم. كان لنظام الزراعة (٢:٢) تأثير واضح على محصول الذرة/فدان ومكوناته حيث كانت الزيادة معنوية في المحصول ومكوناته مقارنة بنظام (٤:٢) في حين لم ينجح أي من النظامين في التفوق على محصول الزراعة النقية .

تحكيم: أ.د عبد الغنى الجندي / أ.د محمد عباس المصرى