

**EFFECT OF NITROGENOUS AND PHOSPHATIC
FERTILIZERS ON PERFORMANCE OF RAINFED PEARL
MILLET (*PENNISETUM GLAUCUM* L.) GROWN ON
CLAY LOAM SOIL**

[12]

Eltilib¹, A.M.A.; T.E.M. Hago¹; A.H. Awad Elkarim^{1,a}
and S.A.M. Ali²

ABSTRACT

An experiment was conducted for two summer seasons (2000 and 2001) to study the effect of nitrogenous and phosphatic fertilizers on performance of rainfed pearl millet (*Pennisetum glaucum* L.) grown on clay loam soil at University of Zalingei site, Western Darfour states, Sudan. The experimental design used was split-plot design with five replications. Four nitrogen treatments (0, 30, 60 and 80 kg N/ha) and four phosphorus treatments (0, 15, 30 and 60 kg P/ha) were used. Nitrogen treatments were assigned to the main plots and phosphorus levels were allotted to the sub-plots. Urea (46% N) and triple superphosphate (48% P₂O₅) were used as sources of nitrogen and phosphorus, respectively. Phosphorus treatments were applied at sowing, whereas nitrogen fertilizer was split into two equal doses, one was added four days after emergence and the second was three weeks after sowing. Results of the experiments showed that application of nitrogen increased the plant height over the control. The increments were in range of 5.7% - 16.1%. Leaf area index (LAI), grain and stover yields were significantly affected by nitrogen application. The increments over control were in the range of 39% - 92.3%, 10.1% - 112.7% and 34.1% - 126.9% for LAI, grain yield and stover yield, respectively. Regarding phosphorus, the results revealed that phosphorus addition had a significant effect on plant height, LAI, grain and stover yields. The increments over control ranged from 2.4% - 11.3%, 6.3% - 17.9%, 6.7% - 24.6% and 6.8% - 35.4% for plant height, LAI, grain and stover yields, respectively. The interaction between nitrogen and phosphorus had a significant effect only on LAI and plant height at the sixth week measurement in the first and second seasons, respectively. The tallest plants were obtained from 3N × 3P treatment whereas the greatest LAI was obtained by 3N × 2P treatment.

Keywords: Phosphatic fertilizers, Pearl millet, Plant height, Leaf area index, Grain yield

-
- 1- Department of Soils & Environment Sciences, Faculty of Agriculture, University of Khartoum, Shambat 13314, Sudan
2- Faculty of Agriculture, University of Zalingie, Western Darfour State, Sudan
a- E-mail: akarim61@yahoo.com

(Received September 23, 2005)

(Accepted December 12, 2005)

INTRODUCTION

Soils, particularly in the arid and semiarid regions, rarely contain adequate amounts of essential plant nutrients to sustain high yields of most crops. This is particularly true for nitrogen and mostly for phosphorus. Use of chemical fertilizers has been practiced for a long time in modern agriculture to compensate these deficiencies in nitrogen and phosphorus.

In Sudan, pearl millet is an important cereal crop, next to sorghum. It constitutes the staple food of the majority of Western Sudan (Darfur and Kordofan) where it occupies an area of 1.2 – 2.9 million hectares. Most of the area is cultivated and harvested under traditional rainfed agriculture (Abuelgasim and Jain, 1987; FAO, 1999). It is also cultivated on alluvial soil along wadis. Recently, some farmers in the mechanized clay plain of central Sudan started to grow pearl millet instead of sorghum which is susceptible to *Striga hermonthica* (Elmahi, et al 1995). Very little research work was conducted in this sector compared with irrigated agriculture. In particular, pearl millet did not receive the appropriate attention. Therefore, the present study was conducted to investigate the effect of nitrogen and phosphorus fertilization on performance and yield of pearl millet grown on clay loam soil under Western Sudan conditions.

MATERIAL AND METHODS

An experiment was carried out for two summer seasons (2000 and 2001) to investigate the effect of nitrogen and phosphorus fertilizers on growth and yield of rainfed pearl millet (*Pennisetum glaucum*). The study site was at the Uni-

versity of Zalingei, Western Darfur state (latitude 12° 45' N and longitude 23° 29' E with an attitude of 900m above mean sea level).

The soil of the site is clay loam (41% clay), pH range from 6.4 to 7.0 with low available soil phosphorus (0.83 – 0.9 ppm). According to Black *et al* (1969) standards, the soil organic carbon is low (0.25 – 0.32%) and, hence, low soil nitrogen content (0.029– 0.035%). The experimental design used was spilt-plot design replicated five times with nitrogen in the main plots and phosphorus in the subplots. The area of main plots and subplots was 108m² and 27m², respectively. A local pearl millet cultivar (Darmasa) was sown on the 12th and 15th of July in the first and second seasons, respectively. Traditional hand sowing was used by using local implement. Nitrogen treatments consisted of four N levels: 0, 30, 60 and 80 kg N/ha designated as 0N, 1N, 2N and 3N, respectively, phosphorus treatments consisted of four P levels: 0, 15, 30 and 60 kg P/ha designed as 0P, 1P, 2P and 3P, respectively. Urea (46% N) and triplesuperphosphate (48% P₂O₅) were used as sources of nitrogen and phosphorus, respectively. Triplesuperphosphate was applied at sowing in bands of 5cm depth and at a distance of 5cm from the seeds. Urea was split into equal doses, one dose added four days after germination and the other was three weeks after sowing by hand placement method. Thinning was done when the plant were four weeks old to end up with three plants per hole. Growth parameters, plant height and leaf area index (LAI) were measured at an interval of two weeks starting a month after sowing until harvest. The crop was harvested at 120 and 112 days from sowing in the first and

second seasons, respectively. Grain and stover yields were determined in an area of 6.25m² in the centre of the middle three rows in each plot.

RESULTS AND DISCUSSION

Nitrogen application had a significant effect ($P \leq 0.05$) on plant height (Table, 1). The plant height was significantly increased over the control by nitrogen treatment. The increases in plant height produced by 1N, 2N and 3N treatments over the control at 12th week after sowing were 7.1%, 10.5%, 9.3% and 5.7%, 11.0% and 16.1% for the first and second seasons, respectively. These findings were in conformity with results obtained by Ibrahim, (1996).

Nitrogen application had a significant effect on LAI in both seasons (Table, 2) where LAI index increased with increasing nitrogen rate up to the 8th week of the sowing but this disappeared as the season proceeded. The increments in LAI over the control due to nitrogen application at the rate of 1N, 2N and 3N at the 8th week measurement were 39.0%, 52.4% and 60.0%, respectively, in the first season and 39.6%, 78.6% and 92.3%, respectively, in the second season. Mourad *et al* (1986) found that LAI values were significantly increased with increasing nitrogen rate up to 90 kg N/ha, which was almost equivalent to the rate used in this study.

Phosphorus application had a significant effect on plant height (Table, 3). The increments in plant height produced by P treatments ranged, on average, between 2.4% - 10.0% over the control. Similar findings were reported by Okalebo (1984). The added phosphorus increased

LAI significantly over the control (Table, 4) for the first three measurements but later this disappeared. These results were in line with those obtained by Elmedani, (1997) who quoted similar results.

Grain and stover yields were significantly affected by nitrogen and phosphorus application (Table, 5). The increments due to nitrogen fertilization were in the range of 10.1% - 112.7% and 34.1% - 126.9% of grain and stover yields, respectively, whereas, the respective increments in grains and stover yields due to phosphorus application ranged between 6.7% - 24.6% and 6.8% - 35.8%.

These results were in line with those obtained by many research workers (Mason, 1989 and Singh & Sharma, 1995).

The yield of the first season was higher than that of the second season. This was mainly due to the fluctuating rainfall which is a critical factor that determines yield in rainfed agriculture. The rainfall in the first season was favorable for crop growth in terms of total amount (777.8 mm) and distribution throughout the growth period whereas in the second season the rainfall was low (421 mm), less than the twenty years mean (488.4 mm). These results were similar to those reported by Bationo *et al* (1989) who reported that low rainfall can adversely affect crop productivity.

However, the interaction of N × P had exerted a non significant effect on most measured parameters. Exception of this, LAI at the 6th week after sowing in the first season was significantly affected by treatment interaction (Table, 6). Similarly, plant height recorded at the 6th week measurement in the second season was significantly affected by N × P interaction (Table, 7).

Table 1. Effect of nitrogen fertilizer on plant height (cm)

Nitrogen level	Time (weeks after sowing)				
	4	6	8	10	12
First season					
0 N	43.56 b	96.97 c	159.48 c	198.15 b	199.98 b
1 N	48.45 ab	123.70 bc	189.72 b	212.78 a	214.22 a
2 N	53.38 a	141.24 a	203.67 a	220.03 a	220.89 a
3 N	51.61 a	133.09 b	200.70 ab	218.33 a	218.51 a
Second season					
0 N	40.80 c	73.87 d	110.28 c	159.89 c	161.11 c
1 N	45.91 b	97.06 c	133.58 b	169.84 bc	170.23 bc
2 N	50.72 a	105.71 b	144.58 b	178.57 ab	178.80 ab
3 N	52.39 a	116.30 a	156.23 a	187.73 a	187.73 a

Means within the same column followed by the same letter are not significantly different at $P \leq 0.05$ according to Duncan's Multiple Range Test (DMRT).

0N = 0 kg N/ha 1N = 30 kg N/ha 2N = 60 kg N/ha 3N = 80 kg N/ha

Table 2. Effect of nitrogen fertilizer on leaf area index (LAI)

Nitrogen level	Time (weeks after sowing)			
	4	6	8	10
First season				
0 N	0.76 b	2.49 c	3.13 c	2.55 a
1 N	1.19 a	3.41 b	4.35 b	2.65 a
2 N	1.35 a	3.93 a	4.77 ab	2.72 a
3 N	1.26 a	3.81 ab	5.01 a	2.83 a
Second season				
0 N	0.65 c	1.09 c	1.54 c	1.10 a
1 N	0.91 b	1.67 b	2.15 b	1.12 a
2 N	1.17 a	2.14 a	2.75 a	1.43 a
3 N	1.27 a	2.29 a	2.97 a	1.50 a

Means within the same column followed by the same letter are not significantly different at $P \leq 0.05$ according to Duncan's Multiple Range Test (DMRT).

Symbols as shown in Table 1

Table 3. Effect of phosphorus fertilizer on plant height (cm)

Phosphorus level	Time (weeks after sowing)				
	4	6	8	10	12
First season					
0 P	44.68 b	111.04 c	125.90 c	205.79 c	207.31 c
1 P	50.06 a	124.24 b	188.00 bc	211.34 b	212.35 bc
2 P	51.34 a	125.59 b	197.44 a	218.70 ab	219.03 a
3 P	50.91 a	134.13 a	192.25 ab	213.46 a	214.93 ab
Second season					
0 P	43.01 c	83.35 c	120.28 c	162.58 c	163.00 c
1 P	47.40 b	99.77 b	136.67 b	173.97 b	174.08 b
2 P	49.34 a	103.63 ab	142.29 a	178.55 ab	179.36 ab
3 P	50.07 a	106.19 a	145.43 a	180.94 a	181.43 a

Means within the same column followed by the same letter are not significantly different at $P \leq 0.05$ according to Duncan's Multiple Range Test (DMRT).

0P = 0 kg P/ha 1P = 15 kg P/ha 2P = 30 kg P/ha 3P = 60 kg P/ha

Table 4. Effect of phosphorus fertilizer on leaf area index (LAI)

Phosphorus level	Time (weeks after sowing)			
	4	6	8	10
First season				
0 P	0.84 b	2.99 b	4.11 a	2.61 a
1 P	1.25 a	3.53 a	4.37 a	2.67 a
2 P	1.23 a	3.58 a	4.37 a	2.76 a
3 P	1.23 a	3.53 a	4.41 a	2.71 a
Second season				
0 P	0.74 b	1.54 b	2.13 b	1.28 a
1 P	1.03 a	1.82 a	2.35 a	1.28 a
2 P	1.11 a	1.91 a	2.47 a	1.31 a
3 P	1.12 a	1.93 a	2.45 a	1.28 a

Means within the same column followed by the same letter are not significantly different at $P \leq 0.05$ according to Duncan's Multiple Range Test (DMRT).

Table 5. Effect of nitrogen and phosphorus fertilizers on grain and stover yields

Treatments	Yield (ton/ha)	
	Grain	Stover
First season		
0 N	1.652 b	5.16 c
1 N	1.819 b	6.92 b
2 N	2.152 a	7.97 a
3 N	2.184 a	8.07 a
Second season		
0 N	0.512 c	1.6 c
1 N	0.653 c	2.49 b
2 N	0.909 b	3.08 ab
3 N	1.089 a	3.63 a
First season		
0 P	1.781 c	6.46 b
1 P	1.900 bc	6.90 ab
2 P	2.018 ab	7.39 a
3 P	2.109 a	7.38 a
Second season		
0 P	0.668 b	2.29 c
1 P	0.799 a	2.59 bc
2 P	0.819 a	2.82 ab
3 P	0.857 a	3.10 a

Means within the same column followed by the same letter are not significantly different at $P \leq 0.05$ according to Duncan's Multiple Range Test (DMRT).

Table 6. Effect of treatment interaction on LAI at the six week after sowing (1st season)

Phosphorus level	Nitrogen level			
	0N	1N	2N	3N
0 P	1.81 h	3.08efg	3.68abcde	3.38de
1 P	3.11 ef	3.63abcde	3.83abcd	3.55bcde
2 P	2.57fg	3.40de	4.13ab	4.22a
3 P	2.48g	3.51cde	4.07abc	4.07abc

Mean within same column followed by the same letters are not significantly different at $P < 0.05$ according to Duncan's Multiple Range Test (DMRT)

Table 7. Effect of treatment interaction on plant height (cm) at the six week after sowing (2nd season)

Phosphorus level	Nitrogen level			
	0N	1N	2N	3N
0 P	66.07h	79.70g	88.00f	99.63e
1 P	69.53h	98.70e	110.63cd	120.20ab
2 P	79.20g	105.67cd	108.23cd	121.43ab
3 P	80.67fg	104.17cd	115.97bc	123.93a

Mean within same column followed by the same letters are not significantly different at $P < 0.05$ according to Duncan's Multiple Range Test (DMRT)

The highest value of LAI was obtained from 3N × 3P treatment which the tallest plants were recorded from 3N × 3P treatment.

REFERENCES

- Abuelgasim, H.E. and R.P. Jain (1987). Breeding pearl millet for drought in the Sudan. Proceedings of the International Pearl Millet Workshop (7-11 April, 1986). *ICRISAT Center India* : 307 – 308.
- Batino, A.; C.B. Christianson and A.U. Mokwunye (1989). Soil fertility management of pear millet – producing sandy soil of Sahelian West Africa: The niger experience, soil, crop and water management systems for rainfed agriculture in the Sudano Sahelian zone: Proceedings of an International Workshop (11 – 16 Jan. 1987). *ICRISAT Sahelian Center Niamey, Niger*: 159 – 167.
- Batino, A.; C.B. Christianson and W.E. Baethgen (1990). Plant density and nitrogen fertilizer effects on pearl millet production in Niger. *Agron. J.* 82: 290 - 294.
- Black, C.A.; D.D. Evans; J.E. Ensminger; J.L. White and F.E. Clark (eds.) (1969). *Methods of Soil Analysis. Part 2. Chemical and Microbiological Properties*. Madison, Wisconsin, USA.
- Elmahi, H.A.; M.E.H. Omer and M.O. Idris, (1995). Dawney mildew of pearl millet in Sudan. *International Sorghum and Millet Newsletter (ISMN) No. 36*: 50-55.
- Elmedani, E.H.Y. (1997). *The Effect of Phosphate Rocks, Triple Superphosphate and Their Combinations on the Performance of Sorghum bicolor var. dura L. and Sorghum sudanese (piper) (stapf)*. pp. 27-35. M.Sc. Thesis Faculty of Agriculture, University of Khartoum.
- FAO, (1999). *Food and Agricultural Organization of the United Nations, Year Book*, No. 53 Rome.
- Gomez, K.A. and A.A. Gomez (1984). *Statistical Procedures for Agricultural*

Research. 2nd Edition. pp. 75-165. John Wiley and Sons Inc. New York.

Ibrahim, A.E.S. (1996). Growth and forage yield of pearl millet (*Pennisetum americanum* L.) as affected by irrigation frequency, seed rate and nitrogen in the United Arab Emirates. *U. of K. J. of Agric. Sci.* 4(1): 45-60.

Mason, S.C. (1989). Agronomy and cropping systems project (Nu-113) nitrogen contribution of intercropped cowpea. *INTOSORMIL Annual Report*: 31-36.

Mourad, S.B.; M.A. El-lakany; A.I. El-Agamy and F.H. Soliman (1986). Response of some varieties of hybrids of

maize to plant densities and nitrogen fertilization. I. Growth and flowering traits. *Al-Azhar J. Agric. Res.* 4: 401-411.

Okalebo, J.R. (1984). Bulursh millet (*Pennisetum typhoides*) responses to nitrogen and phosphorus fertilizers at Ithookwe, Kitui, Kenya. *East Afr. Agric. For. J.* 49(3): 62-77.

Singh, K.N. and D.P. Sharma (1995). Effect of subsurface drain spacing and nitrogen level on yield and growth of upland cotton (*Gossypium hirsutum*) and pearl millet (*Pennisetum glaucum*) in saline soil. *Indian, J. of Agric. Sci.* 65 (1):27-33.

مجلة اتحاد الجامعات العربية للدراسات والبحوث الزراعية، جامعة عين شمس، القاهرة، ١٤٤ (١)، ١٩٥-٢٠٣، ٢٠٠٦

أثر إضافة أسمدة النتروجين والفسفور علي نمو وإنتاجية الدخن المطري في التربة الطينية الطميية

[١٢]

عبد المنعم محمد أحمد التلب^١ - تاج الدين الشيخ موسى هجو^١ -

عبد الكريم حسن عوض الكريم^١ - صديق عبد العزيز محمد علي^٢

١- قسم علوم التربة و البيئة - كلية الزراعة - جامعة الخرطوم - شمبات ١٣٣١٤ - السودان

٢- كلية الزراعة - جامعة زاننجي - ولاية غرب دارفور - السودان

محلي من الدخن (درمسا)، وأستخدم أربعة معاملات من النتروجين هي: صفر و ٣٠ و ٦٠ و ٨٠ كجم نتروجين للهكتار، وأربعة معاملات من الفسفور هي: صفر و ١٥ و ٣٠ و ٦٠ كجم فسفور للهكتار. استعمل اليوريا كمصدر للنتروجين (٤٦%)

أجريت تجربة حقلية لموسمين متتاليين (٢٠٠٠ و ٢٠٠١) بموقع جامعة زاننجي بولاية غرب دارفور بالسودان، لدراسة تأثير إضافة سمادي النتروجين والفسفور علي نمو وإنتاجية الدخن المزروع في التربة الطينية الطميية. تم زراعة صنف

بالنسبة للحبوب ، بينما تراوحت نسبة الزيادة بين ٣٤,١% - ١٢٦,٩% بالنسبة للقصب.

أما الفسفور فقد كان له تأثير معنوي علي طول النبات ودليل مساحة سطح الورقة وإنتاجية الحبوب والقصب ، وقد تراوحت نسبة الزيادة مقارنة بالشاهد بين ٢,٤% - ١١,٣% و ٦,٣% - ١٧,٩% لمساحة سطح الورقة و ٦,٧% - ٢٤,٦% للحبوب و ٦,٨% - ٣٥,٤% للقصب.

وكانت العلاقة بين الفسفور والنتروجين ايجابية علي طول النبات فقط في الاسبوع السادس في الموسم الأول وعلي مساحة سطح الورقة في الاسبوع السادس من الموسم الثاني ، وقد تحصل علي أكثر النباتات طولاً من المعاملة $P_3 \times N_3$ وسجلت أكبر مساحة لسطح الورقة من المعاملة $P_2 \times N_3$.

(N والسيوبر فوسفات الثلاثي كمصدر للفسفور (٤٨% P_2O_5). وأتبع في تنفيذ التجربة تصميم القطع المنشقة في خمس مكررات وكانت معاملات النتروجين في القطع الرئيسية ومعاملات الفسفور في القطع الفرعية. السماد الفسفوري أضيف مع الزراعة اما النتروجين فقد أضيف علي جرعتين الأولى بعد أربعة أيام من الإنبسات والثانية بعد ثلاثة أسابيع من الزراعة.

أوضحت النتائج أن إضافة النتروجين أدت إلي زيادة معنوية في طول النبات ، حيث تراوحت الزيادة بين ٥,٧% - ١٦,١% مقارنة بالشاهد . كما أدت إضافة النتروجين إلي تأثير معنوي علي دليل مساحة الورقة، وكل من إنتاجية الحبوب والقصب حيث تراوحت الزيادة بين ٣٩% - ٩٢,٣% لدليل مساحة سطح الورقة و ١٠,١% - ١١٢,٧%

تحكيم: أ.د عبد المنعم محمد الجلا