

Dept. of Poultry Diseases,  
Fac. Vet. Med., Assiut Univ.

## **SOME STUDIES ON THE INCIDENCE OF ORNITHOBACTERIUM RHINOTRACHEALE INFECTION IN CHICKEN EMBRYOS AND LAYERS**

(With One Table and 8 Figures)

By

**M.A. SHAHATA; T.Y. ABD EL- MOTELIB  
and HEBAT-ALLAH A. MOHAMED\***

\* Animal Health Research Institute

(Received at 18/6/2006)

**بعض الدراسات عن نسبة الإصابة ببكتريا الأورنيثو  
فى أجنة البيض والدجاج البياض**

**مصطفى عبد المطلب شحاته ، طلبه يونس عبد المطلب ،  
هبة الله عبد الحليم محمد**

تم جمع ٨٠ عينة من كتاكيت فاقسة وناقفة حديثاً و ١٤٠ عينة من دجاج بياض وناقف حديثاً عمر ٨ - ١٢ شهراً (٣٥ عينة من المفصل ، ٣٥ عينة من قناة البيض و ٣٥ عينة من المخ، ٣٥ عينة من الرئة والقصبه الهوائية والأكياس الهوائية) وذلك من مزارع مختلفة بمحافظة أسيوط. وقد أمكن عزل ميكروب الأورنيثو بنسبة ٤٠% من الكتاكيت الفاقسة حديثاً وبنسبة ١٧% ، ٢٢,٨% ، ٨,٥% ، ٣٤% من كل من المفصل وقناة البيض والمخ والرئة والقصبه الهوائية والأكياس الهوائية على التوالي. وقد تم إجراء عدوى صناعية بالميكروب المعزول على كتاكيت عمر ٣٠ يوم عن طريق الحقن فى الوريد والحقن فى الأكياس الهوائية والذى أدى إلى إنكماش الطيور ونقص فى الوزن وتورم العين وصعوبة التنفس وإفرازات بالأنف والبعض أظهر بعض الحركات العصبية قبل النفوق الذى وصلت نسبته إلى ٣٠% وذلك فى اليوم الثالث والرابع بعد إجراء العدوى الصناعية أما فى الإِسبوع الثانى من الحقن ظهرت بعض الحالات بها عرج وعدم القدرة على الوقوف أما الصفة التشريحية أظهرت التهابات بالأكياس الهوائية وغشاء فيبريني على الكبد والقلب بالإضافة إلى التهابات بالرئة وبعض الطيور كانت بها مواد صفراء متجينة بالأكياس الهوائية فى منطقة البطن والبعض الآخر أظهر احتقانات بالمخ. وقد أظهرت الكتاكيت التى حقنت عن طريق الأكياس الهوائية نفس الأعراض والصفة التشريحية السابقة ولكن بعد فترة أطول. تم إعادة عزل الميكروب من الحالات المعدة صناعياً. بإجراء اختبار الحساسية لميكروب الأورنيثو المعزول وجد أن الأموكسيسيلين والنتراسيكلين والأنروفلوكساسين هم الأدوية الأكثر تأثيراً.

## SUMMARY

Eighty samples were collected from freshly dead hatched chicks, also one hundred and forty freshly dead layers 8 –12 month old age were collected (35 samples from joint, 35 from oviduct, 35 from brain and 35 from lung, trachea and air sac) from different farms in Assiut Governorate. *Ornithobacterium rhinotracheale* organism was isolated at a rate of 40% from newly hatched chicks and at a rate of 17%, 22.8%, 8.5% and 34% from joint, oviduct, brain and lung, trachea and airsac respectively. Experimental infection of 30 days old chicks by isolated organism through intravenous and intraairsac inoculation was done. Intravenous inoculation leads to depression, ruffling feather of the infected birds and decreased body weight, eye affection, difficult in breathing, nasal secretion and some birds showed nervous signs before death which reached to 30% in the 3<sup>rd</sup> and 4<sup>th</sup> day after infection but in the 2<sup>nd</sup> week postinfection some birds revealed arthritis and could not stand. Postmortem examination revealed airsacculitis, perihepatitis, pericarditis, pneumonia and oedema of the lung. Some birds showed caseous, yellow material on the abdominal airsacs, and others revealed congestion in the brain. Birds which inoculated intra airsac showed also the same previous signs and lesions but it appeared at longer time. Reisolation of the organism from experimentally infected birds was succeeded. Sensitivity test revealed that amoxycillin, tetracycline and enrofloxacin were the most effective drugs.

**The aim of this work is designed to cover the following points:**

- Isolation and identification of the *Ornithobacterium rhinotracheale* organism and its incidence in the newly hatched chicks and layers.
- Experimental infection of the isolated organism to 30 days-old chicks.
- In vitro sensitivity test of the isolated organism to some antimicrobial drugs.

*Key words: Ornithobacterium rhinotracheale, hatched chicks and layers*

## INTRODUCTION

*Ornithobacterium rhinotracheale* (ORT) has been repeatedly found to be involved in cases of respiratory disease in broiler chickens (Charlton *et al*; 1993 – Hafez *et al*; 1993 and Leorat *et al*; 1994).

ORT is a gram – negative, non motile, pleomorphic rod, non sporulating bacterium of the rRNA super family V, in the taxonomic

neighbourhood of the genera *Cytophaga*, *Riemerella*, *Flavobacterium*, *Sporocytophaga* and *Capnocytophaga*.

Osteitis, meningitis and joint infections, can be induced by intravenous application with ORT. The infection can be transmitted horizontally by aerosol, as well as vertically through eggs, which probably accounts for its rapid and worldwide spread. Twelve serotypes can be distinguished within the species ORT, of which serotype A is the most prevalent, therapeutic treatment of the disease can be difficult because acquired resistance against the regular antibiotics is very common within the genus (Vandamme *et al.*, 1994).

Travers *et al.* (1996) were able to induce the disease by inoculation ORT isolates via the coudal abdominal airsac into 28-days old broiler. They noted that the ORT isolates were capable of causing primary disease, with statistically significant resultant mass loss, respiratory and arthritic symptoms were reproduced but no sinusitis was observed. Also they observed a highly significant reisolation of ORT from the brains of broilers challenged with the isolates.

Van Empel *et al.* (1997) could isolated the ORT from the respiratory tract, liver, joints and brains of diseased birds as well as from egg shells and yolk sac of one day old chicks from various countries.

Meningitis, osteitis were observed in experimentally infected SPF chickens by intravenous challenge with 20% mortality but no airsacculitis as seen in the field (Goovaerts *et al.*, 1998).

Nagaraja *et al.* (1998) noted that the ORT was able to survive in the ovary and oviduct of experimentally infected breeder hens without clinical signs.

Mild respiratory symptoms and growth retardation, increased mortality were reported by Van Empel *et al.* (1999). Postmortem investigations showed a foamy, yellow-white "yoghurt like" exudate predominately in the abdominal airsacs and sometimes accompanied by a purulent pneumonia often unilateral and / or exudate in the trachea.

In layers, ORT has been associated with outbreaks of respiratory disease, decreased egg production and mortality in South Africa and Egypt. Many chickens were found dead without antemortem signs, postmortem findings were tracheitis, airsacculitis, pericarditis, ruptured ovarian follicles and egg yolk peritonitis (Stephanie *et al.*, 2000)

Treatment can be difficult because acquired resistance against the regular antibiotics is common in ORT isolates and an inactivated vaccine for broiler breeders has been developed (Van Veen, 2000)

Van Veen *et al.* (2001) observed that ORT was able to induce lesions after aerosol challenge without a previous priming with virus and thus ORT was proven to be a primary pathogen.

Van veen *et al.* (2004) observed in an experiment in SPF broiler chickens that the respiratory tract lesions at postmortem (P.M) examination were positive for ORT by bacteriological and immunohistological examination.

## **MATERIALS and METHODS**

### **Samples:**

A total of 80 dead balady chicken embryos and 140 samples from freshly dead layers were collected (35 samples from joints, 35 samples from oviducts, 35 samples from brain and 35 samples from airsacs, lungs and trachea) collected from different farms at Assiut Governorate and were subjected for bacteriological examination.

### **Media used:**

Sheep blood agar, trypticase soy agar, brain heart infusion broth, trypticase soy broth, triple sugar iron agar, glucose phosphate broth, gelatin agar media, koser's citrate medium, christensen's media, nitrate broth, sugar fermentation media and pepton water (2%).

### **Reagents and solutions :**

Kovac's reagent, methyl red reagent, nitrate reagent, mercuric chloride solution,  $\alpha$  - Naphthol 5%, 40% potassium hydroxide in absolute ethyl alcohol, 1% solution of tetramethyl -P- phenylens diamine dihydrochloride and 30% hydrogen peroxide.

### **Stain used: Gram's stain.**

### **Experimental birds:**

Twenty five, 30 days old balady chickens obtained from the faculty of Agriculture Assiut University poultry farm were used in our experiment.

### **Antibiotic sensitivity discs:**

Include: Enrofloxacin (5 $\mu$ g). Ampicillin (10 $\mu$ g), Chloramphenicol (30  $\mu$ g), Amoxycillin (10 $\mu$ g), Gentamycin (10 $\mu$ g), Trimethoprim (5 $\mu$ g), Neomycin (30  $\mu$ g), Oxytetracycline (30 $\mu$ g), Danofloxacin (5 $\mu$ g), Streptomycin (10 $\mu$ g), Spectinomycin (25 $\mu$ g), Tetracycline (30 $\mu$ g) and lincomycin (2 $\mu$ g).

## **Methods:**

### **I – Isolation**

- a- Samples from dead chicken embryos were taken from lung, trachea, liver and yolk sac and cultured into brain heart infusion broth supplemented with gentamycin as 10µg/ml media according to Back et al. (1996), and incubated at 37°C for 24 – 48 hr., under 7.5 – 10% CO<sub>2</sub> tension by using gas bags in candle jar according to Vandamme *et al.* (1994) Traverse *et al.* (1996) and Rojs *et al.* (2000). Then a loopfull from the cultured broth was streaked onto sheep blood agar plates supplemented with 10µg gentamycin /ml media to inhibit growth of enterobacteriaceae. Plates were incubated at 37°C for 48 hr. under 7.5 – 10 % CO<sub>2</sub>.
- b- Dead laying chickens were subjected to p.m. examination. Swabs taken from joint, oviduct, brain, airsac, lung and trachea were cultured as mentioned before.

### **II – Identification of the isolated organism**

The suspected colonies were examined for their morphology (shape – colour – size- odour) and films from suspected colonies were stained by Gram's stain.

- Biochemical reactions: The most important biochemical reactions of ORT in comparison to these of *Kingella* spp. and the gram – negative rods related to the family Pasteurellaceae potentially pathogenic for fowl and also be differentiated from other pathogenic gram – negative rods in the families Enterobacteriaceae and Neisseriaceae were done according to Van Empel *et al.* (1997).

### **Pathogenicity test:**

Twenty five, 30-days-old healthy chickens were used and 5 birds from them were tested before experiment and proved to be free from ORT organism. The other birds were divided as follow:

- 1<sup>st</sup> group: ten, 30 days old chickens inoculated intravenous with 1ml of a whole culture of brain heart infusion broth adjusted to a count of approximately 10<sup>9</sup> colony forming units/ml according to Saeb *et al.* (2002).
- 2<sup>nd</sup> group: five, 30 days old chickens inoculated intra airsac with 1ml of inoculum of ORT according to Saeb *et al.* (2002)
- 3<sup>rd</sup> group,: five – 30 days old chickens were left as control. Reisolation of ORT from experimental birds were done.

**In vitro sensitivity test:**

The determination of sensitivity of the isolated organism against different antibiotic discs was done according to the technique of Bauer – Kirby *et al.* (1966).

**RESULTS**

Postmortem examination of naturally infected freshly dead layers revealed airsacculitis, pneumonia, tracheitis and inflammation of the oviduct.

Bacteriological examination showed that the suspected ORT colonies were pin point size, greyish white and sometimes with a reddish glow appearance and a distinct odour, similar to the odour of butyric acid.

Gram's stain revealed gram negative, pleomorphic plump short rods.

Biochemical reactions revealed that the isolated organism was oxidase positive, ferments glucose, lactose, galactose and fructose but does not ferment sucrose and maltose and for other tests it was negative for catalase, methyl red, voges proskauer, gelatin liquefaction, citrate utilization, indole, nitrate reduction and triple sugar iron.

According to the cellular and colonial, morphology of the organism as well as the biochemical reactions, the frequency of the isolated ORT was 40% from dead chicken embryos while in dead layers it was 34% from the samples of (lung, trachea, airsac) and was 8.5% from the brain, 17% from joint and was 22.8% from the oviduct samples.

**Pathogenicity test:**

Chickens in group I inoculated intravenously showed in the 1<sup>st</sup> week postinoculation (PI) decrease in the body weight, depression, ruffling feather (Fig. 1) eye affection, difficult in breathing and nasal exudate (Fig. 2). In addition three birds showed nervous signs before death (30% mortality) in the 3<sup>rd</sup> and 4<sup>th</sup> day P.I (Fig. 3). In the 2<sup>nd</sup> week PI, some birds revealed joint affection and the bird could not stand (Fig. 4). Postmortem examination showed airsacculitis perihepatitis, pericarditis (Fig. 5), Pneumonia and oedema of the lung (Fig. 6) while the other birds showed yellow caseous material in the abdominal air sacs (Fig. 7) and others showed congestion of the brain (Fig. 8).

Chickens in group II inoculated intraairsacs also showed signs similar to those in group I but without deaths and revealed the same

gross lesions as in group I but appeared later than those observed in group I.

Neither signs or lesions or deaths were recorded in bird of group III.

Reisolation of ORT from lung, airsac, joint and brain in experimentally infected birds was successful.

**Sensitivity test:**

The effect of the different antibiotics on the isolated ORT are illustrated in Table I.

**Table I:** Illustrates the results of the in vitro sensitivity test:

Antibiotic discs	Conc.	Sensitivity of ORT isolates
Amoxycillin	10 µg	+++
Tetracycline	30 µg	+++
Enrofloxacin	5 µg	+++
Chloramphenicol	30 µg	++
Ampicillin	10 µg	++
Streptomycin	10 µg	++
Oxytetracycline	30 µg	++
Neomycin	30 µg	+
Danofloxacin	5 µg	+
Lincomycin	2 µg	+
Spectinomycin	25 µg	+
Trimethoprim	5 µg	-
Gentamycin	10 µg	-

+++ Sensitive (zone size of inhibition is above 26 mm).

++ moderate sensitive (zone size of inhibition is equal or less than 26 mm).

+ weak sensitive (zone size of inhibition is above 18 mm).

- resistant (zone size of inhibition is less than 18 mm).



**Fig. 1:** Experimentally infected chicken showing depression and ruffling feathers.



**Fig. 2:** Eye affection and difficult in breathing in experimentally infected chicken.





**Fig. 3:** Experimentally infected chicken showing nervous signs before death.



**Fig. 4:** Experimentally infected chicken showing joint affection and the bird could not stand




**Fig. 5:** Pericarditis and perihepatitis of experimentally infected chicken.



**Fig. 6:** Pneumonia, congestion and oedema of the lung of experimentally infected chicken.



**Fig. 7:** Yellow caseous material in the abdominal air sacs of experimentally infected chicken.



**Fig. 8:** Congestion of the brain of experimentally infected chicken.

## DISCUSSION

Respiratory infections are the most serious group of diseases affecting poultry and are accompanied by heavy economic losses due to increased mortality, increased medication costs, increased condemnation rates, drop in egg production, reduction of eggshell quality and decreased hatchability. ORT has been identified as newly emerging respiratory bacterial pathogen that has caused significant economic losses to the poultry industry with a reported incidence that ranged from 30 – 35% (Travers *et al.*, 1996 and Malik *et al.*, 2003).

In this study, postmortem examination of freshly dead layers showed peritonitis, tracheitis, airsacculitis, pericarditis and ruptured of ovarian follicles, this finding is similar to that reported by Stephanie *et al.* (2000) and Soriano *et al.* (2002).

Bacteriological examination revealed that ORT was recovered from airsac, lung and trachea as that reported by Van Empel *et al.* (1999), Saeb *et al.* (2002) and Soriano *et al.* (2002). Also we could isolate the organism from the joint, ovary and brain, this finding is in agreement with the result obtained by Van Empel *et al.* (1997).

Also we were able to recover ORT from the yolk sacs of newly hatched chicks and this result is similar to that obtained by Stephanie *et al.* (2000).

Experimental infection of the isolated organism to 30 days old chickens either intravenously or intraairsac revealed decrease of the weight gain, depression, ruffling feather, eye affection, difficult in breathing and nasal exudate. Nervous signs, meningitis and arthritis were also reported in some birds. These findings are similar to those observed by Travers *et al.* (1996), Goovaerts *et al.* (1998) and Saeb *et al.* (2002).

The gross lesions showed airsacculitis, perihepatitis, pericarditis, pneumonia and oedema of the lung, this result goes hand in hand with that observed by Travers (1996), Ryll *et al.* (1997), Sprenger *et al.* (1998) and Saeb *et al.* (2002).

In our experiment, we showed 30% mortality in the infected birds, this result is somewhat similar to that reported by Goovaerts *et al.* (1998) who recorded 20% mortality rate in their experiment.

The obtained results in our experiment and those that reported by other authors (Leorat *et al.*, 1994 and Travers *et al.*, 1996) differ with the data mentioned by Van Empel *et al.* (1996) who said that ORT culture

was only capable of reproducing the clinical features of the disease if the birds had been previously infected with respiratory viruses.

The most prominent gross lesions seen in some birds was foamy white, yoghurt-like' exudate in the airsacs this result is similar to that observed by Van Empel and Hafez (1999).

We were successful in reisolation ORT from experimentally infected birds and this result is in agreement with those observed by Van Beek *et al.* (1994), Nagaraja *et al.* (1998) and Soriano *et al.* (2002).

In vitro sensitivity test revealed that amoxicillin, tetracycline and enrofloxacin were the most effective drugs against the isolated organism, our result is relatively similar to that observed by Hafez (1996) who found that amoxicillin, tetracycline and chloramphenicol were effective to ORT isolates, but Saeb *et al.* (2002) found that tetracycline was the only drug which was sensitive to ORT isolates Amal (2002) recorded that amoxicillin, chloramphenicol and ampicillin were the most effective drugs.

Our conclusion in this study proved that ORT can be transmitted vertically through eggs as well as horizontally and ORT can be regarded as a primary pathogen in chickens to cause respiratory symptoms.

## REFERENCES

- Amal Ahmed Mohamed (2002):* Preliminary studies on *Ornithobacterium rhinotracheale* infection in poultry in upper Egypt M.V.Sc. Thesis Fac. Vet. Med. Assiut Univ.
- Back, A.G.; Halovorson, D. and Nagaraja, K. (1996):* Experimental studies of *Ornithobacterium rhinotracheale* (ORT) infection Proceedings of the 46<sup>th</sup> western poultry Disease conference, sacramento. PP. 7-8.
- Bauer, A.W.; Kirby, W.M.; Sherris, J.C. and Turch, M. (1966):* Antibiotic susceptibility testing by a standardized single disk method. *Am. J. Clin. Pathol.* 45: 493 – 496.
- Charlton, B.R.; Channing- Santiago, S.E.; Bickford, A.A.; Cardona, C.J.; Chin, R.P.; Cooper, G.L.; Droual, R.; Jeffrey, J.S. and Walker, R.L. (1993):* Preliminary characterization of a pleomorphic – negative rod associated with avian respiratory disease. *Journal of Veterinary Diagnostic Investigations*, 5: 47 – 51.
- Goovaerts, D.; Vrijenhoek, M. and Van Empel, P. (1998):* Immuno – histochemical and bacteriological investigation of the pathogenesis of *Ornithobacterium rhinotracheale* infection in South Africa in chickens with osteitis and encephalitis

- syndrome. In: proc. 16<sup>th</sup> Meeting of the European society of veterinary pathology, Lillehammer, Norway. P. 81
- Hafez, H.M. (1996):* Current status on the role of *Ornithobacterium rhinotracheale* (ORT) in respiratory disease complexes in poultry. Arch. Geflugelk. 60, 208 – 211.
- Hafez, H.M.; Hruse, W.; Empele, J. and Sting, R. (1993):* Eine Atemwegsinfektion bei Mastputen durch *Pasteurella* ähnliche Erreger: Klinik Diagnostik und therapie. Proc. of international conference on poultry diseases. Potsdam Germany. PP 105 – 112.
- Leorat, J.; Dudouyt, J.; Dore, C. and Gardin, Y. (1994):* *Ornithobacterium rhinotracheale*: un nouvelle raison de tousser. Filières Avicoles 559: 69 – 70.
- Malik, Y.S.; Olsen, K.; Kumar, K. and Goyal, S.M. (2003):* In vitro antibiotic resistance profiles of *Ornithobacterium rhinotracheale* strains from Minnesota turkeys during 1996 – 2002. Avian Diseases Jul-sep: 47 (3) 588 –93.
- Nagaraja, K.; Back, A.; Sorenger, S.; Rajshekara, G. and Halvorson, D. (1998):* Tissue distribution post-infection and antimicrobial sensitivity of *Ornithobacterium rhinotracheale*. In Proc. of the 47<sup>th</sup> western poultry Disease conference, sacramento, p.p. 57 – 60.
- Rojs – Zorman, O.; Zdovc, I.; Bencina, D. and Mrzel, I. (2000):* Infection of turkeys with *Ornithobacterium rhinotracheale* and *Mycoplasma synoviae*. Avian Diseases. 44: 1017 – 1022.
- Ryll, M.; Hinz, K-H.; Behr, H-P.; Kruse, W.; Lohren, U. and Sudbeck, M. (1997):* Zu pathogenität und verbreitung der *Ornithobacterium rhinotracheale* – infection bei der pute. In proc. of the 52<sup>nd</sup> Meeting of the fachgruppe “Geflugelkrankheiten” der Deutsche Veterinärmedizinische Gesellschaft Hannover pp. 39 – 60.
- Saeb, N.; El-Sukhon Asad Musa and Majed Al-Attar (2002):* Studies on the bacterial etiology of airsacculitis of broilers in Northern and Middle Jordan with special reference to *Escherichia coli*, *Ornithobacterium rhinotracheale* and *Bordetella avium* Avian Diseases. 46: 605 – 612.
- Soriano, V.E.; Longinos, M.G.; Navarrete, P.G. and Fernandez, R.P. (2002):* Identification and characterization of *Ornithobacterium rhinotracheale* isolates from Mexico. Avian Diseases, 46: 686 – 690.

- Sprenger, J.; Back, A.; Shaw, P.; Nagaraja, K.; Roepke, D. and Halvorson, D. (1998):* Ornithobacterium rhinotracheale infection in turkeys: experimental reproduction of the disease. *Avian Diseases*, 42: 154 – 161.
- Stephanie, J.; Sprenger David, A.; Halvorson, K.V.; Nagaraja, Radivoje spasojevic Richard S. Dutton, and Daniel P. Shaw (2000):* Ornithobacterium rhinotracheale infection in commercial laying type chickens. *Avian Diseases*, 44: 725 – 729.
- Travers, A. (1996):* Concomitant Ornithobacterium rhinotracheale and Newcastle disease infection in broilers in South Africa. *Avian Diseases*, 40: 488 – 490.
- Travers, A.; Coetzee, L. and Gummow, G. (1996):* Pathogenicity differences between South Africa isolates of Ornithobacterium rhinotracheale. *Onderstepoort Journal of Veterinary Research*, 63: 197 – 207.
- Van Beek, P.; Van Empel, P.C.M.; van den Bosch, G.; Storm, P.; Bongers, J. and Dupreez, J. (1994):* Ademhalings problemen, groeivertraging en gewrichtsontsteking bij Kalkoenen en Vleeskuikens dooreen Pasteurella – achtige bacterie : Ornithobacterium rhinotracheale of “taxon 28”. *Tijdschrift voor Diergeneeskunde*, 119, 99 – 101.
- Vandamme, P.; Segers, P.; Vancaneyt, M.; Van Hover, K.; Mutters, R.; Homme, J.; Dewirst, F.; paster, B.; Kersters, K.; Falsen, F.; Devrize, L.; Bisgaard, M.; Hinz, K.H. and Mannheim, W. (1994):* Description of Ornithobacterium rhinotracheale een. nov. sp. nov., isolated from the avian respiratory tract. *International Journal of systematic Bacteriology*, 44; 24 – 37.
- Van Empel, P.C.M. and Hafez, H.M. (1999):* Ornithobacterium rhinotracheale: a review. *Avian pathology* 28: 217 – 227.
- Van Empel, P.C.M.; Van den Bosch, H.; Goovaerts, D. and storm, P. (1996):* Experimental infection in turkeys and chickens with Ornithobacterium rhinotracheale. *Avian Diseases*, 40: 858 – 861.
- Van Empel, P.C.M.; Van den Bosch, H.; Loeffen, P. and Storm, P. (1997):* Identification and serotyping of Ornithobacterium rhinotracheale. *Journal of clinical Microbiology*, 35: 418–421.
- Van Empel, P.C.M.; Vrijenhoek, M.; Goovaerts, D. and van den Bosch, H. (1999):* Immuno-histochemical and serological investigation of experimental Ornithobacterium rhinotracheale infection in chickens. *Avian Pathology* 28: 187 – 193.

- Van Veen, L. (2000):* Ornithobacterium rhinotracheale infection in poultry: a review. Tijdschr Diergeneeskd, 15; 125 (4): 113 – 116.
- Van Veen, L.; Van Empel, P. and Fabri, T. (2001):* Ornithobacterium rhinotracheale a primary pathogen in broilers. Avian Diseases, : 896 – 900.
- Van Veen, L.; Vrijenhoek, M. and Van Empel, P. (2004):* Studies of the transmission routes of Ornithobacterium rhinotracheale and immunoprophylaxis to prevent infection in young meat turkeys. Avian Diseases, 48 (2): 233-7.