

EFFECT OF ACRYLAMIDE, AMYGDALIN, CAPSICUM, FURFURAL AND VANILLIN ON SOME FOOD BORNE PATHOGENIC BACTERIA

Youssef, Kh. M. and A.A. Shatta

Food Technology Dept., Fac of Agriculture, Suez Canal Univ. 41522, Ismailia, Egypt.

E-mail: me505073@yahoo.com

ABSTRACT

The growth of eleven food spoilage bacterial strains; four Gram negative (*Enterobacter (Ent.) aerogenes*, *Escherichia (E.) coli*, *Pseudomons (Ps.) aeruginosa* and *Ps. fluorescens*) and seven Gram-positive strains (*Bacillus (B.) cereus*, *B. firmus*, *B. pumilus*, *B. subtilis*, *Micrococcus (M.) luteus*, *M. varians* and *Staphylococcus (S.) aureus*) was studied in liquid media in the presence of acrylamide, amygdalin, capsicum, furfural and vanillin. These compounds are naturally occurred in plant products or formed or added during food processing.

The inhibitory effect of acrylamide, amygdalin, capsicum, furfural and vanillin is concentration and strain dependent. The inhibitory action increased as the concentration increased. Furthermore, the presence of amygdalin, vanillin, furfural as well as capsicum inhibited Gram-negative bacterial strains, namely *Ps. aeruginosa* and *Ps. fluorescens*, while Gram-positive, namely *B. pumilus*, *B. subtilis*, *M. luteus*, and *M. varians* were inhibited by furfural (750 ppm) and vanillin (3000 ppm).

These findings indicate that vanillin and capsicum could be useful as preservatives for minimally processed products.

Keywords: Pathogenic bacteria, Gram-negative strains, Gram-positive strains, acrylamide, amygdalin, capsicum, furfural, vanillin, growth, inhibition.

INTRODUCTION

The increase of international trade of food commodities has raised the risk of dispersion of pathogenic bacteria from the production sites to far places of consumption. High quality and safety of food are of major concern to the nowadays consumer, who prefers fresher, additive-free and minimally processed products. Hence, the use of natural antimicrobial agents for the preservation of foods could guarantee both high quality and safety. Natural antimicrobial compounds seem important not only to the preservation of food, but also to the control of human diseases of microbial origin (Gould, 1996; Baratta *et al.*, 1998 and Quattara *et al.*, 2000).

Earlier Kim and Ryeon (1979) reported the antibacterial effect of capsaicin from Korean hot peppers on *B. cereus*, *B. subtilis* and *Sarcina lutea*. Later on Serruti and Alzamora (1996) showed that vanillin (one of the capsaicin analogues) inhibits the growth of yeast. The trend of using natural antimicrobial compounds has led to growing interest in investigating a series of such compounds from the viewpoint of their antimicrobial activity (Alzamora and Guerrero, 2003).

Acrylamide occurs in plant material as potatoes, carrots, radish, lettuce, Chinese cabbage, parsley, onions, spinach and paddy rice (Arikawa and Shiga, 1980). Acrylamide does not occur naturally but it is found in

various fried, deep-fried and oven-baked foods. It concerns foods, which are regularly consumed throughout the year, like chips (French fries), crisps and bread, also biscuits, crackers and breakfast cereals (Tareke *et al.*, 2002). So far no acrylamide has been found in foods that are boiled (Rosen and Hellenäs, 2002). Acrylamide has been shown to be neurotoxic in humans and laboratory animals. It has also been shown to induce tumors in experimental animals, and has been classified as 'probably carcinogenic for humans' (IARC, 1994; EC, 2002; FAO/WHO, 2002 and Vatterm and Kalidas, 2003).

Amygdalin is a natural substance, known as cyanogenic glycosides; it was first isolated from bitter almond. It is found in the seeds of apricot, peach, plum, cherry, apple, sprouted alfalfa, mung, wheat, millet, lentil, watercress, lima beans, black eyed peas, clover, sorghum, broccoli, cabbage, flax seed, linseed...etc. (Moss, 1996 and Rosenberg, 1990). Herbert (1979) reviewed that intestinal bacteria contain enzymes (β -glucosidases) that activate the release of cyanide after amygdalin had been ingested. When amygdalin interacts with the enzyme β -glucosidase, or undergoes hydrolysis (breakdown upon reaction with water) in the absence of enzymes, hydrogen cyanide, benzaldehyde and glucose are produced (Newmark *et al.*, 1981 and Rauws *et al.*, 1982).

Red pepper (*Capsicum annum*, *Solanaceae*), an important flavoring and coloring spice, is used as a natural flavoring. Its pungency is attributed to capsaicinoid, which is composed of 80 % to 90 % capsaicin and dehydrocapsaicin (Pulselove *et al.*, 1988 and Perucka & Oleszek, 2000). *Capsicum annum* extract has an inhibitory effect on *B. cereus* and *S. aureus* (Dorantes *et al.*, 2000 a). Also, the same authors proposed the use of this extract as a natural antibacterial agent in a food such as raw beef, which is often contaminated (Dorantes *et al.*, 2000 b). Also, Careaga *et al.* (2003) reported that the *Capsicum annum* (bell pepper extract) exhibited an inhibitory effect against *S. typhimurium* and *Ps. aeruginosa*.

Furfural occurs naturally in many fruits, tea, coffee and cocoa and mostly is formed during the processing and domestic preparation of a broad range of foods (Monti *et al.*, 2000 and Murata *et al.*, 2002). It is also carried over into food from its use as an extraction solvent or as a component of flavor mixtures. Furfural and its derivatives from consumption of foods in which it occurs naturally is approximately 0.3 (Stofberg and Grundschober, 1987) to 0 - 0.5 mg /kg /day (IARC, 1994). The highest concentrations of furfural in food have been reported in cocoa (55 - 255 ppm), wholegrain-bread (26 ppm), heated skim milk (230 ppm) and coffee (90 - 881 ppm) (Adams *et al.*, 1997).

Vanillin, a plant phenol, is the predominant phytochemical that occurs in vanilla beans and is generally regarded as safe (GRAS). It is used widely as a food flavoring agent in confectioneries, chocolates, butter, toppings, icings, ice cream, beverages, biscuits, desserts, etc. (Beuchat & Golden, 1989; Hocking, 1997; Spillman *et al.*, 1997 and Ramachandra Roa & Ravishankar, 2000). Its antioxidant and antimutagenic activity had been observed in both bacteria and mammalian cells (Tamai *et al.*, 1992; Ohta, 1993; Prince & Gunson, 1994 and Santosh Kumar *et al.*, 2000).

As information on the sensitivity of pathogenic and spoilage bacteria to some additives or compounds formed in foods are scanty, the aim of the present study was to test the effect of acrylamide, amygdalin, capsicum, furfural and vanillin on the behavior of some food borne pathogenic bacteria *in Vitro*.

MATERIALS AND METHODS

MATERIALS

Additives

The compounds examined are acrylamide at 500, 1000 and 2000 ppm (SERVA, Heidelberg, Germany), amygdalin at 3.5, 50 and 100 ppm (Fluka Chemie, AG, CH-9470 Buchs, Switzerland), capsicum at 50, 100 and 200 ppm (Concentrated Capsicum, FLPT038-10, Ransom, William Ransom & Sonpic Hitchin, Herts, SG5, ILY, England), furfural at 250, 500 and 750 ppm (Merck, Darmstad, Germnay) and vanillin at 750, 1500 and 3000 ppm (Zhonghua Chemicals, Zhejiang, China). The minimal concentrations of these additives are naturally occurred or formed or added during food processing. Solutions of the additives were sterilized by filtration through a sterile 0.20 µm cellulose nitrate filter (Sartorius, AG. 37070 Goettingen, Germany) and then added at the concentrations mentioned above to the growth media.

Microorganisms and culture media

The following 11 Gram-positive and Gram-negative bacteria were used for antimicrobial activity studies. Gram-negative bacteria: *Enterobacter (Ent.) aerogenes* ATCC 15050, *Escherichia (E.) coli* ATCC 15130, *Pseudomonas (Ps.) aeruginosa* DSM 50071 and *Ps. fluorescens* DSM 50090. Gram-positive bacteria: *Bacillus (B.) cereus* DSM 31, *Bacillus firmus* ATCC 14575, *Bacillus pumilus* ATCC 14884, *Bacillus subtilis* DSM 10, *Micrococcus (M.) luteus* ATCC 15307, *M. varians* ATCC 15306 and *Staphylococcus (S.) aureus* ATCC 6538.

These strains were obtained from the American Type Culture Collection, Rockville, Maryland, USA (ATCC) and the German Collection of Micro-organisms, Braunschweig, Germany (DSM). All these strains were checked up and stored on Brain Heart Infusion (BHI) slants at 4 °C then subcultured twice in Brain Heart Infusion Broth (LAB M, Topley House, 52 Wash Lane, Bury, Lancashire, BL9 6AU, UK) and incubated at 22 °C (for *Ps. aeruginosa* and *Ps. fluorescens*) and 37 °C (for the rest of strains) for 24 h before use.

METHODS:

Determination of bacterial growth and inhibition activity

Flasks of BHI broth containing various concentrations of additives and control (without additives) were inoculated with apriori prepared cultures at 1% level (initial counts, 10^6 - 10^7 cfu ml⁻¹) and incubated at 22 and 37 °C. Triplicate flasks were treated for each additive at each concentration. The growth of each culture was monitored in two ways, by measuring its absorbance at 600 nm (OD₆₀₀) by a Spectronic 20D (Milton Roy Company, USA) at intervals for a total period of 72 hrs, and by plating on BHI agar (1.2% w/v) at 0, 3, 6, 9, 12, 24, 48, 72 hrs suitably diluted aliquots of the culture (viable counts). All experiments were repeated thrice.

Growth analysis

The growth percentage of 12 h culture equals
$$\frac{(OD_t - OD_{t_0})_{test}}{(OD_t - OD_{t_0})_{control}} \times 100,$$

Where: OD is the optical density at 600 nm, t is time after 12 h, t_0 is the initial time 0 h, *test* makes reference to the culture grown with additive(s) and *control* makes reference to the culture grown without additives (Nazer *et al.*, 2005). This variable indicates how much the growth is reduced in the presence of additives. A time of 12 h is chosen for the best discrimination of growth curves.

The inhibition percentage of the examined additives was calculated as follows:

$$\text{Inhibition \%} = [\text{Log } N_2 - \text{Log } N_1 / \text{Log } N_2] \times 100,$$

Where: Log N_1 : Log cfu ml⁻¹ of the sample at the last hour (72nd hr).

Log N_2 : Log cfu ml⁻¹ of control without additives at the last hour (72nd hr)

Statistical analysis

The results are presented as means \pm standard deviation from three replicates of each experiment. A P-value ≤ 0.01 is used to denote significant differences among mean values determined by analysis of variance (ANOVA) (CoStat program ver. 3.03, 1986).

RESULTS AND DISCUSSION

The inhibitory effect of acrylamide, amygdalin, capsicum, furfural and vanillin on the growth of eleven food borne pathogenic bacteria was studied and the results are given in Tables (1-2) and Figures (1-2)

Acrylamide

Acrylamide showed slowest decrease of the growth percentage of Gram-negative (except *Ps. aeruginosa*) and Gram-positive bacteria. In case of *B. pumilus* it gave the highest growth percentage at different concentrations. In other words acrylamide seemed to be a promoter of the growth of this bacterium (Figs. 1 and 2).

As shown in Tables (1 and 2), acrylamide has no effect on *Ent. aerogenes*, *E. coli*, *Ps. fluorescens*, *B. firmus* and *B. pumilus*, whereas inhibits the growth of *Ps. aeruginosa* (by 60.08% at 2000 ppm). Gram-positive bacteria, namely *B. cereus*, *B. subtilis*, *M. luteus*, *M. varians* and *S. aureus* are significantly affected by acrylamide. The inhibition percentages are 20.09, 32.12, 45.00, 49.06 and 42.43, respectively.

Amygdalin

Amygdalin leads a slowest decrease of the growth percentage of Gram-negative strains. The growth percentage of Gram-positive bacteria gives highest values at different concentrations. Amygdalin exhibited itself as a growth promoter for Gram-positive bacteria (Figs. 1 and 2).

Thus, from Table (1), it is seen that Gram-negative bacteria are significantly inhibited by amygdalin. *Ps. aeruginosa* and *Ps. fluorescens* are sensitive among the Gram negative or positive bacteria, at 3.5, 50 and 100 ppm. The inhibition percentages are 94.73, 96.51 and 96.68; 97.20, 97.38 and 98.48, respectively.

These results may be due to the assimilation of amygdalin by the bacteria, and the resulting product (HCN) in return affected them (Conn, 1978).

For some bacterial strains, these results may indicate that the bacteria do not ferment or assimilate amygdalin or hydrolyze it and release a free HCN (very toxic). Another assumption is that the bacteria secrete rhodanase, which possesses a disulphide bridge that can be broken down by the thiosulphate ion, with the release of non-toxic Na_2SO_4 and SCN (Rosenberg, 1990).

Capsicum

The same trend of amygdalin is observed. The growth percentage is higher for Gram-positive bacteria (Fig. 2).

The antibacterial activity of capsicum is considerably lower towards Gram-positive strains than towards Gram-negative, i.e. capsicum has no effect on the Gram-positive bacteria, whereas the growth of Gram-negative (*Ps. aeruginosa* and *Ps. fluorescens*) is significantly inhibited. The inhibition percentages are 95.27, 97.23 & 97.57; 98.11, 98.11 and 98.45% at 50, 100 and 200 ppm, respectively.

These results are in agreement with those of Dorantes *et al.* (2000 a) who reported that *Ps. aeruginosa* was sensible to *Capsicum annuum* extract. Also, Careaga *et al.* (2003) reported that the *Capsicum annuum* (bell pepper extract) exhibited an inhibitory effect against *S. typhimurium* and *Ps. aeruginosa*. The authors added that the minimum inhibitory concentration of the extract to prevent the growth of *Ps. aeruginosa* was 0.3 ml of the extract 100 g⁻¹ of meat as, bacteriostatic effect, while a concentration of 3 ml of the extract 100 g⁻¹ of meat was bactericidal. On the contrary, Dorantes *et al.*, (2000 b) mentioned that *Capsicum annuum* extract had an inhibitory effect on *B. cereus* and *S. aureus* using an agar diffusion method

Furfural

The growth percentage after 12 h of Gram-positive bacteria has no linear relation with furfural concentration (Figs. 1 and 2). There is almost no inhibition below some threshold concentration, then the growth percentage quickly decreases as the concentration of furfural increases. For instances, at 250 ppm of furfural the growth percentage is 91.23 while at 500 ppm, 17.81 for *Ps. aeruginosa*. Furthermore, at 750 ppm the growth percentage is 1.84. Also, this attitude holds true for *Ps. fluorescens*, 80.74% and 54.15% at 250 and 500 ppm, whereas at 750 ppm the corresponding percentage is 20.40 (Fig. 1).

For Gram-negative bacteria Table (1) shows that furfural has an inhibitory effect on *Ps. aeruginosa* and *Ps. fluorescens*. The inhibition percentages are 96.40 and 68.96 at 750 ppm, respectively. However, the inhibitory effect is concentration-dependent. *B. cereus*, *B. firmus* and *Ent.*

Thus, from Table (1), it is seen that Gram-negative bacteria are significantly inhibited by amygdalin. *Ps. aeruginosa* and *Ps. fluorescens* are sensitive among the Gram negative or positive bacteria, at 3.5, 50 and 100 ppm. The inhibition percentages are 94.73, 96.51 and 96.68; 97.20, 97.38 and 98.48, respectively.

These results may be due to the assimilation of amygdalin by the bacteria, and the resulting product (HCN) in return affected them (Conn, 1978).

For some bacterial strains, these results may indicate that the bacteria do not ferment or assimilate amygdalin or hydrolyze it and release a free HCN (very toxic). Another assumption is that the bacteria secrete rhodanase, which possesses a disulphide bridge that can be broken down by the thiosulphate ion, with the release of non-toxic Na_2SO_4 and SCN (Rosenberg, 1990).

Capsicum

The same trend of amygdalin is observed. The growth percentage is higher for Gram-positive bacteria (Fig. 2).

The antibacterial activity of capsicum is considerably lower towards Gram-positive strains than towards Gram-negative, i.e. capsicum has no effect on the Gram-positive bacteria, whereas the growth of Gram-negative (*Ps. aeruginosa* and *Ps. fluorescens*) is significantly inhibited. The inhibition percentages are 95.27, 97.23 & 97.57; 98.11, 98.11 and 98.45% at 50, 100 and 200 ppm, respectively.

These results are in agreement with those of Dorantes *et al.* (2000 a) who reported that *Ps. aeruginosa* was sensible to *Capsicum annum* extract. Also, Careaga *et al.* (2003) reported that the *Capsicum annum* (bell pepper extract) exhibited an inhibitory effect against *S. typhimurium* and *Ps. aeruginosa*. The authors added that the minimum inhibitory concentration of the extract to prevent the growth of *Ps. aeruginosa* was 0.3 ml of the extract 100 g^{-1} of meat as, bacteriostatic effect, while a concentration of 3 ml of the extract 100 g^{-1} of meat was bactericidal. On the contrary, Dorantes *et al.*, (2000 b) mentioned that *Capsicum annum* extract had an inhibitory effect on *B. cereus* and *S. aureus* using an agar diffusion method

Furfural

The growth percentage after 12 h of Gram-positive bacteria has no linear relation with furfural concentration (Figs. 1 and 2). There is almost no inhibition below some threshold concentration, then the growth percentage quickly decreases as the concentration of furfural increases. For instances, at 250 ppm of furfural the growth percentage is 91.23 while at 500 ppm, 17.81 for *Ps. aeruginosa*. Furthermore, at 750 ppm the growth percentage is 1.84. Also, this attitude holds true for *Ps. fluorescens*, 80.74% and 54.15% at 250 and 500 ppm, whereas at 750 ppm the corresponding percentage is 20.40 (Fig. 1).

For Gram-negative bacteria Table (1) shows that furfural has an inhibitory effect on *Ps. aeruginosa* and *Ps. fluorescens*. The inhibition percentages are 96.40 and 68.96 at 750 ppm, respectively. However, the inhibitory effect is concentration-dependent. *B. cereus*, *B. firmus* and *Ent.*

aerogenes are not affected while the growth of *B. pumilus*, *B. subtilis*, *M. luteus*, *M. varians* and *S. aureus* is significantly inhibited. The inhibition percentages are 87.82, 69.77, 67.50, 59.06 and 43.64, respectively at 750 ppm.

Table (1): The effect of acrylamide, amygdalin, capsicum, furfural and vanillin on the growth of some Gram-negative food borne pathogenic bacteria

Item tested	Inhibition percent			
	Ent. <i>aerogenes</i>	<i>E. coli</i>	<i>Ps. aeruginosa</i>	<i>Ps. fluorescens</i>
Acrylamide,				
500 ppm	0.00 ± 0.01 ^a	0.00 ± 0.01 ^a	32.36 ± 4.97 ^b	0.00 ± 0.01 ^b
1000 ppm	0.00 ± 0.01 ^a	0.00 ± 0.01 ^a	43.46 ± 4.76 ^b	0.00 ± 0.01 ^b
2000 ppm	0.00 ± 0.01 ^a	0.00 ± 0.01 ^a	60.08 ± 3.38 ^a	5.56 ± 0.93 ^a
Amygdalin				
3.5 ppm	12.83 ± 0.33 ^b	14.77 ± 1.90 ^b	94.73 ± 0.35 ^a	97.20 ± 0.61 ^a
50 ppm	19.74 ± 1.98 ^a	23.81 ± 1.91 ^a	96.51 ± 0.96 ^a	97.38 ± 0.37 ^a
100 ppm	19.74 ± 1.98 ^a	24.24 ± 1.14 ^a	96.68 ± 0.72 ^a	98.48 ± 0.25 ^a
Capsicum				
50 ppm	16.12 ± 2.96 ^a	11.02 ± 0.42 ^b	95.27 ± 0.41 ^b	98.11 ± 0.55 ^a
100 ppm	17.11 ± 1.98 ^a	12.50 ± 3.41 ^b	97.23 ± 0.93 ^a	98.11 ± 0.55 ^a
200 ppm	17.76 ± 0.01 ^a	25.57 ± 3.22 ^a	97.57 ± 0.45 ^a	98.45 ± 0.65 ^a
Furfural				
250 ppm	0.00 ± 0.01 ^b	0.00 ± 0.01 ^b	0.00 ± 0.01 ^c	4.88 ± 0.01 ^c
500 ppm	0.00 ± 0.01 ^b	0.00 ± 0.01 ^b	22.95 ± 3.77 ^b	24.09 ± 0.92 ^b
750 ppm	3.95 ± 1.14 ^a	23.67 ± 2.47 ^a	96.10 ± 0.21 ^a	68.96 ± 0.92 ^a
Vanillin				
750 ppm	0.00 ± 0.01 ^b	0.00 ± 0.01 ^b	85.10 ± 0.04 ^c	66.34 ± 0.52 ^c
1500 ppm	0.00 ± 0.01 ^b	0.00 ± 0.01 ^b	98.32 ± 0.10 ^b	70.67 ± 1.53 ^b
3000 ppm	29.93 ± 0.33 ^a	28.79 ± 0.38 ^a	98.87 ± 0.03 ^a	85.40 ± 0.27 ^a

Within each column and for each extract, means having the same superscripts are not significantly different at $p \leq 0.01$

Vanillin

Figs. 1 and 2 show that the growth percentage decreases as the vanillin concentration increases. However, this relation is not linear. The growth percentage of *B. firmus* at 750 ppm is 81.33%, while it falls to 26.35% at 3000 ppm. The same trend is observed in the case of *S. aureus*. In the case of *B. subtilis*, the growth percentage of the culture in the presence 750 ppm is almost equal to 74.35% and falls to 9.57% at 3000 ppm of vanillin.

From Tables (1 and 2), vanillin at low concentrations (750 ppm and 1500 ppm) is ineffective against *Ent. aerogenes*, *E. coli*, *M. luteus* and *S. aureus*. The inhibitory effect is against *Ps. aeruginosa* and *Ps. fluorescens*. The inhibition percent increases with the increase of vanillin concentration, being 85.10, 98.32 and 98.87 and 66.34, 70.67 and 85.40%, (at 750, 1500 and 3000 ppm) respectively. Notably, the inhibition percentages for *B. cereus*, *B. firmus*, *B. pumilus*, *B. subtilis*, *M. luteus*, *M. varians* and *S. aureus* are 59.64, 74.03, 87.39, 91.54, 22.02, 82.27 and 53.92 at 3000 ppm of vanillin, respectively. These results are in agreement with Fitzgerald *et al.*

(2004) who found the action of vanillin to be bacteriostatic and sometimes bactericidal (Friedman *et al.*, 2002; Ultee, *et al.*, 2002). Moreover, Cerrutti *et al.*, (1997) reported that vanillin was used as an antimicrobial agent in fruit purées, where it inhibited yeast, molds and bacteria at 2000 – 3000 ppm. Fitzgerald *et al.*, (2004) found that the inhibitory activity of vanillin resides primarily in its ability to detrimentally affect the integrity of the cytoplasmic membrane, with the resultant loss of ion gradient, pH homeostasis and inhibition of respiratory activity.

Table (2): The effect of acrylamide, amygdaline, capsicum, furfural and vanillin on the growth of some Gram-positive food borne pathogenic bacteria

Item	Inhibition percent						
	<i>B. cereus</i>	<i>B. firmus</i>	<i>B. pumilus</i>	<i>B. subtilis</i>	<i>M. luteus</i>	<i>M. varians</i>	<i>S. aureus</i>
Acrylamide							
500 ppm	12.85 ± 1.22 ^a	0.00 ± 0.01 ^a	0.00 ± 0.01 ^a	28.46 ± 1.92 ^a	32.14 ± 1.19 ^a	39.17 ± 3.96 ^b	31.35 ± 2.70 ^b
1000 ppm	12.85 ± 1.17 ^a	0.00 ± 0.01 ^a	0.00 ± 0.01 ^a	30.38 ± 3.85 ^a	35.86 ± 0.01 ^a	45.63 ± 1.25 ^{a,b}	33.24 ± 1.35 ^b
2000 ppm	20.09 ± 3.74 ^a	0.00 ± 0.01 ^a	0.00 ± 0.01 ^a	32.12 ± 0.58 ^a	45.00 ± 5.78 ^a	49.06 ± -0.52 ^a	42.43 ± 0.28 ^a
Amygdalin							
3.5 ppm	0.00 ± 0.01 ^c	0.00 ± 0.01 ^a	0.00 ± 0.01 ^a	0.00 ± 0.01 ^a	0.00 ± 0.01 ^b	7.56 ± 3.91 ^a	3.03 ± 0.01 ^a
50 ppm	5.51 ± 1.30 ^b	0.00 ± 0.01 ^a	0.00 ± 0.01 ^a	0.00 ± 0.01 ^a	13.10 ± 1.20 ^a	9.90 ± 3.13 ^a	7.06 ± 0.01 ^a
100 ppm	10.05 ± 2.11 ^a	0.00 ± 0.01 ^a	0.00 ± 0.01 ^a	0.00 ± 0.01 ^a	14.29 ± 1.19 ^a	15.63 ± 2.09 ^a	9.41 ± 1.18 ^a
Capsicum							
50 ppm	0.00 ± 0.01 ^b	0.00 ± 0.01 ^a	0.00 ± 0.01 ^b	0.00 ± 0.01 ^a	0.00 ± 0.01 ^b	0.00 ± 0.01 ^c	0.00 ± 0.01 ^a
100 ppm	0.00 ± 0.01 ^b	0.00 ± 0.01 ^a	0.00 ± 0.01 ^b	0.00 ± 0.01 ^a	0.00 ± 0.01 ^b	0.00 ± 0.01 ^c	0.00 ± 0.01 ^a
200 ppm	4.21 ± 1.87 ^a	0.00 ± 0.01 ^a	11.76 ± 0.14 ^a	0.00 ± 0.01 ^a	11.31 ± 1.79 ^a	13.39 ± 1.31 ^a	0.00 ± 0.01 ^a
Furfural							
250 ppm	0.00 ± 0.01 ^b	0.00 ± 0.01 ^b	58.34 ± 1.24 ^c	10.00 ± 0.77 ^c	37.05 ± 0.75 ^b	52.40 ± 5.31 ^a	0.00 ± 0.01 ^b
500 ppm	0.00 ± 0.01 ^b	0.00 ± 0.01 ^b	69.06 ± 2.83 ^b	21.54 ± 0.39 ^b	37.65 ± 0.45 ^b	56.04 ± 0.21 ^a	0.00 ± 0.01 ^b
750 ppm	4.05 ± 0.45 ^a	5.70 ± 1.32 ^a	87.82 ± 0.01 ^a	69.77 ± 5.31 ^a	67.50 ± 3.15 ^a	59.06 ± 1.56 ^a	43.64 ± 0.01 ^a
Vanillin							
750 ppm	36.40 ± 0.01 ^c	21.43 ± 1.08 ^c	69.06 ± 0.80 ^b	35.38 ± 0.39 ^c	0.00 ± 0.01 ^b	58.85 ± 2.61 ^b	23.35 ± 0.01 ^b
1500 ppm	44.59 ± 0.64 ^b	54.22 ± 0.82 ^b	69.21 ± 1.09 ^b	62.69 ± 0.23 ^b	0.00 ± 0.01 ^b	59.43 ± 1.72 ^b	23.36 ± 1.20 ^b
3000 ppm	59.64 ± 0.72 ^a	74.03 ± 0.19 ^a	87.39 ± 0.94 ^a	91.54 ± 0.16 ^a	22.02 ± 0.60 ^a	82.27 ± 2.46 ^a	53.92 ± 0.14 ^a

Within each column and for each extract, means having the same superscripts are not significantly different at $p \leq 0.01$




Fig (1): The effect of acrylamide, amygdalin, capsicum, furfural and vanillin on the growth percentage at 12 h of some Gram-negative food borne pathogenic bacteria.




Fig (2): The effect of acrylamide, amygdalin, capsicum, furfural and vanillin on the growth percentage at 12 h of some Gram-positive food borne pathogenic bacteria.

In conclusion, capsicum and vanillin could be used as preservative agents for minimally processed foods.

REFERENCES

- Adams, T. B.; Doull, J.; Goodman, J. I.; Munro, I. C.; Newberne, P.; Portoghese, P. S.; Smith, R. L.; Wagner, B. M.; Weil, C. S.; Woods, L. A. and Ford, R. A. (1997). The FEMA GRAS assessment of furfural used as a flavour ingredient. *Food and Chemical Toxicology*, 35, 739-751
- Alzamora, S. M. and Guerrero, S. (2003). Plant antimicrobials combined with conventional preservatives for fruit products. In S. Roller (Ed.), *Natural antimicrobials for the minimal processing of foods* (pp. 235– 249). Boca Raton, FL: CRC Press LLC.
- Arikawa, A. and Shiga, M. (1980). Determination of trace acrylamide in the crops by gas chromatography. *Bunseki Kagaku*, 29(7), 3339; *Chem. Abstr.*: 93, 202742.
- Baratta, T. M.; Dorman, D. J. H.; Deans, G. S.; Figueiredo, C. A.; Barroso, G. J. and Ruberto, G. (1998). Antimicrobial and antioxidant properties of some commercial oils. *Flavour and Fragrance J.*, 13, 235-244.
- Beuchat, L. R. and Golden, D. A. (1989). Antimicrobial occurring naturally in foods. *Food Technol.*, 43, 142-143.
- Careaga, M.; Fernandez, E.; Dorantes, L.; Mota, L.; Jaramillo, M. E. And Hernandez-Sanchez, H. (2003). Antibacterial activity of Capsicum extract against *S. typhimurium* and *Pseudomonas aeruginosa* inoculated in raw beef meat. *Int. J. Food Microbiol.*, 83, 331-335
- Cerrutti, P.; Alzamora, S. M. and Vindales, S. L. (1997). Vanillin as an antimicrobial producing shelf-stable strawberry puree. *J. Food Sci.*, 62, 608–610.
- Conn, E. E. (1978). Cyanogenesis, the production of hydrogen cyanide by plants. In R. F. Keeler; K. R. Van Krampen and L. F. James, (Eds.), *Effects of poisonous plants on livestock* (pp. 301-310). Academic Press, New York.
- CoStat Program (1986). CoStat 3.03, CoHort Software, Berkeley CA 94701.
- Dorantes, L.; Acero-Ortega, C.; Hernández-Sánchez, H., and López-Malo, A. (2000a). Effect of capsicum extracts on the growth of some microorganisms important in products. June 10-4; Dallas, Tex.: Inst. of Food Technologists Annual Meeting Book of Abstracts. Chicago, Illinois: Inst. of Food Technologists. USA, pp. 145-146.
- Dorantes, L.; Colmenero, R.; Hernández, H.; Mota, L.; Jaramillo, M. E., Fernández, E. and Solano, C. (2000b). Inhibition of growth of some foodborne pathogenic bacteria by *Capsicum annum* extracts. *Int. J. Food Micro.*, 51, 125-128.
- European Commission (EC) (2002). Risk assessment of acrylamide (CAS No. 79-06-1, EINECS No. 201-173-7). Summary Risk Assessment Report. European Commission, Joint Research Center, European Chemicals. Bureau Ispra, Italy.

- FAO/WHO (2002). Health implications of acrylamide in food. Report of a Joint FAO/WHO consultation (Geneva: WHO), pp. 8-9.
- Fitzgerald, D. J.; Stratford, M.; Gasson, M. J.; Ueckert, J.; Bos, A. and Narbad, A. (2004). Mode of antimicrobial action of vanillin against *E. coli*, *Lactobacillus plantarum* and *Listeria innocua*. J. Appl. Microbiol., 97, 104-113.
- Friedman, M., Henika, P. R., and Mandrell, R. E. (2002). Bactericidal activities of plant essential oils and some of their isolated constituents against *Campylobacter jejuni*, *Escherichia coli*, *Listeria monocytogenes*, and *Salmonella enterica*. J. Food Prot., 65, 1545-1560.
- Gould, G. (1996). Industry perspectives on the use of natural antimicrobials and inhibitors for food applications. J. Food Prot., Supplement, 59, 82-86.
- Herbert, V. (1979). Laetrile: the cult of cyanide. Promoting poison for profit. American J. Clin. Nutr., 32, 1121-1158.
- Hocking, M. B. (1997). Vanillin: synthetic flavouring from spent sulfite liquor. J. Chem. Ed., 74, 1055-1059.
- IARC Monographs on the Evaluation of Carcinogenic Risks to Humans. Some Industrial Chemicals (1994). International Agency for Research on Cancer, Lyon, 60, 389-433.
- Kim, K. and Ryeon, K. (1979). A study on content and antibacterial effects of capsaicin from Korean hot pepper. Report of the National Institute of Health, Korea. 16, 241-251.
- Monti, S. M.; Borrelli, R. C.; Ritieni, A. and Fogliano, V. (2000). A comparison of color formation and maillard reaction products of a lactose-lysine and lactose N-alpha-acetyl lysine model system. J. Agric. Food Chem., 48, 1041-1046.
- Moss, R. W. (1996). The Cancer Industry: The Classic Exposé on the Cancer Establishment. Brooklyn, NY: First Equinox Press. pp. 131-185.
- Murata, M.; Shinoda, Y. and Homma, S. (2002). Browning of model orange juice solution and changes in the components. Int. Congress Series, 1245, 459-460
- Nazer, A. I.; Kobilinsky, A.; Tholozan, J. L. and Dubois-Brissonnet, F. (2005). Combinations of food antimicrobials at low levels to inhibit the growth of *Salmonella* sv. *typhimurium*: a synergistic effect?. Food Micro., 22, 391-398.
- Newmark, J.; Brady, R. O. and Grimley, P. M. (1981). Amygdalin and prunasin beta-glucosidases: distribution in gram-free rat and in human tumor tissue. Proceedings of the National Academy of Sciences USA. 78, 6513-6516.
- Ohta, T. (1993). Modification of genotoxicity by naturally occurring flavorings and their derivatives. Crit. Rev. Toxicol., 23, 127-146.
- Perucka, I. and Oleszek, W. (2000). Extraction and determination of capsaicinoids in fruit of hot pepper *Capsicum annum* L. by spectrophotometry and high-performance liquid chromatography. Food Chem., 71, 287-291.

- Prince, R. C. and Gunson, D. E. (1994). Just plain vanilla? Trends Biochem. Sci., 19, 521.
- Pulseglove, J. W.; Brown, E. G.; Green, C. I. and Robbins, S. R. J. (1988). Spices. Vol. 1. New York: Longman Scientific & Technical., p. 331-439.
- Quattara, B.; Simard, R. E.; Piette, G.; Begin, A. and Holley, R. A. (2000). Diffusion of acetic and propionic acids from chitosan-based antimicrobial packaging films. J. Food Sci., 65, 768-773.
- Ramachandra Roa, S. and Ravishankar, G. A. (2000). Vanilla flavor: production by conventional and biotechnological routes. J. Sci. Food and Agric., 80, 289-304.
- Rauws, A. G.; Olling, M. and Timmerman, A. (1982). The pharmacokinetics of prunasin, a metabolite of amygdalin. J. Toxicol: Clinical Toxicology., 19, 851-856.
- Rosen, J. and Hellenäs, K. E. (2002). Analysis of acrylamide in cooked foods by liquid chromatography tandem mass spectrometry. Analyst, 27, 880-882.
- Rosenberg, H. (1990). Cancer chemo (toxico) therapy revisited and alternative ways of healing. Ph.D. Thesis. Anglo-American Institute of Drugless (Naturapathy).
- Santosh Kumar, S.; Gosh, A.; Devasagayam, T. P. A. and Chauhan, P. S. (2000). Effect of vanillin on methylene blue plus light-induced single-strand breaks in plasmid pBR322 DNA. Mutat. Res., 469, 207-214.
- Serruti, P. and Alzamora, S. M. (1996). Inhibitory effects of vanillin on some food spoilage yeast in laboratory media and fruit purees. Int. J. Food Microbiol., 29, 379-386.
- Spillman, P. J.; Pollnitz, A. P.; Liacopoulos, D.; Skouroumounis, G. K. and Sefton, M. A. (1997). Accumulation of vanillin during barrel-ageing of white, red, and model wines. J. Agric. Food Chem., 45, 2584-2589.
- Stofberg, J. and Grundschober, F. (1987). Consumption ratio and food predominance of flavoring materials. Perfumer and Flavorist., 12, 27.
- Tamai, K.; Tezuka, H. and Kuroda, Y. (1992). Different modifications by vanillin in cytotoxicity and genetic changes induced by EMS and H₂O₂ in cultured Chinese hamster cells. Mutat. Res., 268, 231-237.
- Tareke, E.; Rydberg, P.; Karlsson, P.; Ericksson, S. and Tornqvist, M. (2002). Analysis of acrylamide, a carcinogen formed in heated foodstuffs. J. Agric. Food Chem., 50, 4998-5006.
- Ultee, A., Bennik, M. H. J., and Moezelaar, R. (2002). The phenolic hydroxyl group of carvacrol is essential for action against the foodborne pathogen *Bacillus cereus*. Applied and Environmental Microbiology, 68, 1561-1568.
- Vattem, D. and Kalidas, S. (2003). Acrylamide in food: a model for mechanism of formation and its reduction. Innovative Food Science and Emerging Technologies, 4, 331-338.

تأثير الاكريلاميد والاميجدالين و الكابسيكم و الفرفيورال والفانيلين على بعض البكتيريا الممرضة في الغذاء

خالد يوسف و عادل شطا

قسم الصناعات الغذائية- كلية الزراعة- جامعة قناة السويس ٤١٥٢٢ الإسماعيلية- ج. م. ع.

درس تأثير كل من الاكريلاميد والاميجدالين و الكابسيكم و الفرفيورال والفانيلين على نشاط إحدى عشرة سلالة بكتيرية ممرضة في الغذاء، أربع منها سالبة لجرام وهي (*Enterobacter (Ent.) aerogenes, Escherichia (E.) coli, Pseudomonas (Ps.) aeruginosa and Ps. fluorescens*) وسبع سلالات موجبة لجرام وهي (*Bacillus (B.) cereus, B. firmus, B. pumilus, B. subtilis, Micrococcus (M.) luteus, M. varians and Staphylococcus (S.) aureus*) وذلك في مرق Brain Heart Infusion يحتوي على تركيزات مختلفة من المركبات السابقة الذكر. ومن المعروف أن هذه المركبات تتواجد طبيعيا في بعض المنتجات النباتية أو تتكون أو تضاف أثناء التصنيع الغذائي.

أعتمد تأثير المواد المضافة على التركيز المستخدم وكذا السلالة المختبرة. ثبت كل من الاميجدالين والفانيلين والفرفيورال والكابسيكم نمو السلالات البكتيرية السالبة لجرام (*Ps. aeruginosa and Ps. fluorescens*). وكانت سلالة *Ps. fluorescens* من أكثر السلالات حساسية تجاه المركبات المختبرة. أما السلالات الموجبة لجرام ومنها *B. pumilus, B. subtilis, M. luteus, M. varians*, فتبطلت عند تركيز ٧٥٠ جزء في المليون فرفيورال و ٣٠٠٠ جزء في المليون فانيلين.

في ضوء النتائج يمكن التوصية باستخدام الفانيلين و الكابسيكم كمواد حافظة في الأغذية وخاصة في الأغذية محنونة المعاملة.