

## **DETECTION OF SOME HEAVY METAL RESIDUES IN SOME CAMEL'S MEAT PRODUCTS**

**JEHAN, R. DAOUD\*; MONA, A. A. EL-KAHKY\*\* and TAREK, H. ALLAM\*\*\***

\* Animal Health Research Institute, Dokki, Egypt.

\*\* Tanta Provincial Laboratory, Animal Health Research Institute.

\*\*\* Zagazig Provincial Laboratory, Animal Health Research Institute.

**Received:** 16.8.2005.

**Accepted:** 25.8.2005.

### **SUMMARY**

Lead, cadmium and copper residues were determined in each of 15 samples of camel's luncheon and frozen sausage, collected from different supermarkets in Cairo, Giza and Zagazig Cities. The residues of these heavy metals were detected by using Atomic Absorption Spectrophotometer. The obtained results revealed that the mean values  $\pm$  S.E. of lead, cadmium and copper (p.p.m) in camel's luncheon were  $0.138 \pm 0.008$ ,  $0.055 \pm 0.004$  and  $3.662 \pm 0.221$ ; respectively, while in frozen sausage, such residues were  $0.133 \pm 0.008$ ,  $0.058 \pm 0.004$  and  $3.789 \pm 0.189$  p.p.m; respectively. The present results were compared to the permissible limits of FAO/WHO (1972), WHO (1972) and Egyptian Organization for Standardization and Quality Control "E.O.S.Q.C." (1993). Public health importance and the hazardous toxic

effects of these heavy metals as well as the suggestive recommendations to minimize the pollution with heavy metals were discussed.

---

### **INTRODUCTION**

Heavy metals and their salts constitute the most widely distributed group of highly toxic and long retained pollutants. They are commonly found in effluents discharging from metal processing factories. Mining products, chemical and sewage sludge effluents have high levels of heavy metals that might cause potential pollution to fresh water resources (Dean and Suess, 1985; Khalaf-Allah, 1998). Heavy metals in soil and water may enter the food chain through the biologic cycle which include bioconcentration by plants and animals (El-Shorbagy, 2004).

Lead is a toxic substance which accumulates in the body from old house paints. Lead arsenate contains large quantities of lead and acts as a possible source of lead for both man and animal. Chronic lead poisoning is particularly characterized by anemia, liver dysfunction, muscular pain, lead nephropathy and neuropathy. Its potential carcinogenic nature has also been shown by Zawurska and Medras (1988) and Goldfrank et al. (1990).

Cadmium is also a toxic element for human being. It is virtually absent from the human body at birth. It is accumulated with age in body tissues and causes renal failure (Gracey and Collins, 1992). Cadmium had a significant role in the incidence of some diseases as diabetes mellitus (Merali and Singhal, 1977), chronic renal failure (Friberg, 1984), human hypertension and anemia in all species (Watanabe and Murayama, 1974 and Nishiyama et al., 1986), reproductive toxicity and bone defects (Noroderg, 1995). Air pollution with cadmium from industrial sources (manufacturing of plastics, solder alloys, nickel cadmium batteries, photo cells and rubber tires) may be transmitted to man through contaminated vegetables used as food stuffs or through food of animal origin (Carstensen and Poulsen, 1974). The International Agency for Research on Cancer (IARC) in (1993) classified cadmium and cadmium compound as class "1" human carcinogens in (1993).

Uptake through food and digestion is regarded as the most significant source of cadmium for the general population (WHO, 1989).

Copper is widely distributed in nature and also present in food. It is important in formation of erythrocytes, development of bone, C.N.S and connective tissues. Acute exposure to copper causes hypotension and haemolytic anemia but chronic exposure to copper causes jaundice in human (Gossel and Bricker, 1990).

The purpose of this study was carried out to determine the levels of lead, cadmium and copper residues in either of camel's luncheon and frozen sausage which collected from different supermarkets located in Cairo, Giza and Zagazig Cities.

## MATERIALS AND METHODS

A total of 30 samples, 15 each of camel luncheon and frozen sausage were collected from different supermarkets in Cairo, Giza and Zagazig Cities. The weight of each sample ranged from 100-150 g. The samples were kept in polyethylene bags and then frozen stored at - 20°C until examined. The collected samples were prepared and digested according to the technique recommended by Khan et al. (1995) as follows: One gram of each sample was macerated in screw capped tube by sharp scalpel. Ten ml of digestion mixture consists of 7 parts of ultrapure concentrated nitric acid (HNO<sub>3</sub>) and 3 parts of perchloric acid (HClO<sub>4</sub>) were added.

ed to the sample. The tubes were tightly closed and the contents were vigorously shaken and allowed to stand overnight at room temperature, then heated until evaporated (at 95°C). The residues were redissolved in 10 ml of 1 N HNO<sub>3</sub>.

The solutions were filtered through Whatman no. 41 filter paper and analyzed for the presence of lead, cadmium and copper residues by using Atomic Absorption Spectrophotometer (UNI-CAM 696) under the following conditions:

Conditions \ Metal	Lead	Cadmium	Copper
Method	Normal segmented curve fit	Normal segmented curve fit	Normal segmented curve fit
Measurement time	4.0 second	4.0 second	4.0 second
Lamb current/m. am	15	8	5
Wave length (nm)	217	228.8	324.8
Technique	Flame	Flame	Flame
Flame Type	Air /Acetylene	Air /Acetylene	Air /Acetylene
Air/I	30	30	30
Acetylene/I	20	20	20
Fuel Flow (L/Min)	1.1	1.2	1.1

**N.B:** The estimation of such heavy metals in each examined sample was in p.p.m on the basis of wet weight sample.

#### Statistical analysis:

The obtained results were statistically analyzed according to the method of Selvin (1996).

### RESULTS AND DISCUSSION

Heavy metals make up one of the most important groups of pollutants, so it is necessary to monitor the levels of heavy metals residues which may

be avoidably present in luncheon and frozen sausage of camels meat.

**Lead (Pb):** The obtained results in table (1) and (2) revealed that the mean values of lead in luncheon and frozen sausage of camels samples were  $0.138 \pm 0.008$  and  $0.133 \pm 0.008$  p.p.m; respectively.

**Table (1): Heavy metal residues (p.p.m) in camel's uncheon.**

<b>Metal</b>	<b>Min.</b>	<b>Max.</b>	<b><math>\bar{X}</math></b>	<b>S.D</b>	<b><math>\pm</math> S.E.</b>
Lead (Pb)	0.089	0.182	0.138	0.032	0.008
Cadmium (Cd)	0.035	0.089	0.055	0.017	0.004
Copper (Cu)	2.461	5.971	3.662	0.857	0.221

N.B: No. of examined samples = 15.

p.p.m: Part per million.

Min. Minimum value.

Max.: Maximum value.

$\bar{X}$ : Mean values

S. D : Standard Deviation.

$\pm$  S. E. : Standard Error.

The obtained findings of lead residues in table (1) and (2) were within the permissible limits which intended by WHO (1972), Casarett and Doull (1975) and E.O.S.Q.C. (1993) in table (3). The present data of lead in table (1) and (2) were agreed with those reported by Amodio-Cocchieri and Fiore (1987) in sheep muscles and Shehata et al. (1999); El-Sharkawy (2000) in camel's meat. On the other hand high lead findings were recorded by Youssef et al. (1994) in buffalo muscles; Daoud et al. (1998) in cattle and sheep muscles; Lopez-alanse et al. (2000) in cattle muscles and

El-S horbagy (2004) in muscles of camels.. Low lead values in camels were detected by Diab (1995). Low lead levels in the present data might be attributed to the collection of meat from slaughtered camels subjected to low environmental pollution with lead. On the other hand, the presence of high levels of lead by some authors may be due to collection of samples from animals subjected to high environmental pollution and to the accumulative effect of such metal in tissues. This held the view reported by Doganoc (1996).

**Table (2): Heavy metal residues (p.p.m) in frozen sausage of camels.**

Metal	Min.	Max.	$\bar{X}$	S.D	$\pm$ S.E.
Lead (Pb)	0.075	0.185	0.133	0.032	0.008
Cadmium (Cd)	0.038	0.091	0.058	0.015	0.004
Copper (Cu)	2.187	5.192	3.789	0.733	0.189

N.B: No. of examined samples = 15.

p.p.m: Part per million.

Min. Minimum value.

Max.: Maximum value.

$\bar{X}$ : Mean values

S. D : Standard Deviation.

$\pm$  S. E. : Standard Error.

**Cadmium (Cd):** The results illustrated in table (1) and table (2) showed that the mean values of cadmium residues in either camel luncheon and frozen sausage were  $0.055 \pm 0.004$  and  $0.058 \pm 0.004$  p.p.m; respectively. The obtained data were within the limits of FAO/WHO (1972); Casarett and Doull (1975) and E.O.S.Q.C. (1993) in table (3). High levels of cadmium were recorded by Daoud et al. (1998) in muscles of cattle and sheep; Sallam and Morshdy (2000) and El-Shorbagy (2004) in muscles of camels. The high values of cadmium may be attributed to the grazing of animals in polluted pasture with cadmium from industrial sources. The actual extent of absorption depends on a number of dietary factors such as the intake of protein, calcium, vitamin D and other trace metals. Following absorption,

cadmium is transported and bound to certain proteins of the plasma and red blood cells to other sites throughout the body. However, the metabolism of the cadmium was antagonized with the copper and iron leading to anemia. This agrees with that reported by Underwood (1977). Low values in muscles of camels were detected by Diab (1995).

**Copper (Cu):** The results recorded in table (1) and (2) showed that the mean values of copper residues were  $3.662 \pm 0.221$  and  $3.789 \pm 0.189$  p.p.m; respectively. The obtained data were within the permissible limits of Casarett and Doull (1975) and E.O.S.Q.C. (1993) in table (3). High values of copper in camels meat were detected by Diab (1995). On the other hand low values were

recorded by Daoud et al. (1998) in muscles of cattle and sheep.

The variation of such metal concentrations between the presented results and those reported by many other authors was referred to the differences of the degree of pollution among the different localities in which the animals were raised.

Therefore, the preventive measures intended for

minimizing the pollution of raw meat with such metals are of significant concern, including:

- 1- Minimizing the use of phosphates and sludge for land fertilization as possible.
- 2- Periodical examination should be done for meat, meat products and their content for heavy metals should be evaluated according to the international guide lines as a fruitful advise to delay environmental contamination.

**Table (3): Recommended levels of heavy metals in food.**

References	Heavy metals		
	Lead (Pb)	Cadmium (Cd)	Copper (Cu)
Casarett and Doull (1975)	Human daily intake 0.3 mg	Human daily intake 0.018 - 0.20 µg	Human daily intake 3.2 mg
E.O.S.Q.C. (1993)	Human weekly intake 0.05 mg/Kg body weight. 0.5 mg/Kg in frozen sausage and luncheon of camels	Human weekly intake 0.0067 - 0.0081 mg/Kg body weight	Daily intake 0.05 - 0.5 mg/Kg body weight 15 mg/Kg in frozen sausage
FAO/WHO (1972)	_____	Not exceed 0.04 - 0.05 mg/Kg in food	_____
WHO (1972)	Human weekly intake 3 mg/person or 0.05 mg/Kg body weight	_____	_____

E.O.S.Q.C.: Egyptian Organization for Standardization and Quality Control.

FAO: Food and Agriculture Organization.

WHO: World Health Organization.

## REFERENCES

- Amodio-Cocchieri, R. and Fiore, P. (1987): Lead and cadmium concentrations in livestock breed in Campania, Italy. *Bull. Environ. Contam. Toxicol.*, 39(3): 460-454.
- Carstensen, J. and Poulsen, E. (1974): Public health aspects of environmental pollution with mercury and cadmium in Scandinavia. Problems of the contamination of man and his environment by mercury and cadmium. Published by the Commission of the European Communities, Luxembourg.
- Casarett, L. J. and Doull, J. (1975): Toxicology "The Basic Science of Poisons" Chapter 18, PP: 458, Macmillan Publishing Co., Inc., New York.
- Daoud, J. R.; Kamel, A. A. and Mostafa, A. R. (1998): Determination of lead, cadmium and copper residues in muscles, liver and kidneys of slaughtered cattle and sheep in El- Sharkia Governorate. *Vet. Med. J. Giza*, 46 (4A): 339-348.
- Dean, R. B. and Suess, M. J. (1985): The risk to health of chemicals in sewage sludge applied to land. *Waste Management Res.*, 3: 251-278.
- Diab, O. M. A. (1995): Chemical constituents of beef, buffaloes and camel's meat. Ph. D., Vet. Thesis, Cairo University, Egypt.
- Doganoc, D. Z. (1996): Lead and cadmium concentrations in meat, liver and kidney of Slovenian cattle and pigs from 1989 to 1993. *Food Add. & Contam.*, 13 (2): 237-241.
- Egyptian Organization for Standardization and Quality Control "E. O. S. Q. C. " (1993): Maximum residue limits for heavy metals in food. Ministry of Industry. No. 2360/1993. PP: 5. Cairo, Egypt.
- El-Sharkawy, R. E. (2000): Some heavy metal residues in camel carcasses. Ph. D., Vet. Thesis. Zagazig University, Egypt.
- El-Shorbagy, I. M. (2004): Lead and cadmium residues in meat retailed at Sharkia markets. *Sc. Vet. Med. J.*, 1: 71-82. (4th Int. Sci. Conf., Mansoura).
- FAO/WHO (1972): 16 Joint Expert Committee on Food Additives, evaluation of mercury, lead, cadmium and the food additives amaranth, diethyl pyrocarbonate and acyle galleic. WHO Food Additives Ser. 1972. No. 4.
- Enberg, L. (1984): Cadmium and the kidney. *Environ. Health Persp.*, 54:1-11.
- Goldfrank, L. R.; Folmenbaum, N. E.; Lewin, N. A.; Weisman, R. S. and Howland, M. A. (1990): Goldfrank's Toxicology Emergencies. 4th Ed. Prentice-Hall International Inc. New Jersey, USA.
- Gossel, T. A. and Bricker, J. D. (1990): "Principles of Clinical Toxicology" 2nd Ed. P: 182-183. Raven Press Ltd., New York.
- Gracey, J. F. and Collins, D. S. (1992): Meat Hygiene, 9th Ed., P: 219. Bailliere Tindall, London.
- International Agency For Research On Cancer "IARC" (1993): Beryllium, Cadmium, Mercury and Exposure in the Glass Manufacturing Industry. IARC. Monographs on the Evaluation of Carcinogenic Risks to Humans, Vol. 58 (Eyon: World Health Organization).
- Khalaf-Allah, S. S. (1998): Monitoring of heavy metal pollution in river Nile around Helwan, Egypt. *J. Egypt. Vet. Med. Assoc.*, 58(4): 571-589.
- Khan, A. T.; Diffay, B. C.; Datiri, B. C.; Forester, D. M.; Thompson, S. J. and Mielke, H. W. (1995): Heavy metals in livers and kidneys of goats in Alabama. *Bull. Environ. Contam. Toxicol.*, 55: 568-573.

- Lopez-alonse, M.; Bendito, J. L.; Miranda, M.; Castillo, C.; Hernandez, J. and Shore, R. F. (2000): Toxic and trace elements in liver, kidney and meat from cattle slaughtered in Galicia (NW Spain). *Food Add. Contam.*, 17(6): 447-457.
- Merali, Z. and Singhal, R. L. (1977): Long term effects of orally administrated cadmium on neonatal rats. *Proc. Of the 1st Intern. Congress on Toxicol. Held March, 30-April in Toronto.*
- Nishiyama, S.; Nakamura, K. and Konish, Y. (1986): Blood pressure and urinary sodium and potassium excretion in cadmium treated male rats. *Environ. Res.*, 40: 357-364.
- Noroderg, G. (1995): Human cadmium exposure in the general environment and related health risks. Ph. D. Thesis, Umea University.
- Sallam, Kh. I. and Morshdy, A. M. (2000): Selenium and some heavy metal residues in camel carcasses in relation to human health. *Zag. Vet. J.*, 28: 120-131.
- Selvin, W. (1996): *Statistical Analysis of Epidemiologic Data.* 2nd Oxford and I.B.H. Publishing Company Pomby.
- Shehata, R. E.; Atabany, A. I. and Shaaban, F. E. (1999): Lead and cadmium residues in camels (*Camelidae Dromedarius*) tissues in Sharkia Province. *Zag. Vet. J.*, 27:38-42.
- Underwood, E. J. (1977): *Trace elements in human and animal nutrition*, 4th Ed. Academic Press, New York.
- Watanabe, H. and Murayama, H. (1974): Studies on the changes of renal tubular function of inhabitants in cadmium pollution areas. *Kanko Hoken Report*, 31:12-17.
- World Health Organization "W.H.O." (1972): *Health hazards of the human environment.* WHO report, Geneva, PP: 61.
- Youssef, H.; Fathi, S.; Nagwa, M.; El-Sawi and Baulis, W. (1994): Some trace elements in tissues of camel, cattle and buffalo slaughtered in Assuit Upper Egypt, 6th Sci. Cong., 20-22 Nov. *Vet. Med. Assuit, Egypt.*
- W. H. O. (1989): Evaluation of certain food additives and contaminants. 33rd Report of the Joint FAO/WHO Expert Contaminants on Food Additives. WHO Technical Report Series, 776.
- Zawurska, B. and Medras, K. (1988): Tumoren und Steorungen des pophyrin Stoff Wechsels bei Ratten. *Mitchronischer Experimentaller, Bleiintoxikation (1) Morphologische Studien. Zblt f Allgem. Patholog und Patholo. Anatom.* 54 3: 1.



## تقرير بقايا بعض المعادن الثقيلة في بعض منتجات لحوم الجمال

جيهان رجب محمد داود\* ، منى عبد اللطيف أحمد الكحكي\*\*،

طارق حسن علام\*\*\*

\* معهد بحوث صحة الحيوان بالدقى - جمهورية مصر العربية

\*\* معمل بيطرى طنطا معهد بحوث صحة الحيوان

\*\*\* معمل بيطرى الزقازيق معهد بحوث صحة الحيوان

تم تقدير بقايا عنصر الرصاص والكاديوم والنحاس في ١٥ عينة من لانشون اللحم الجملى وفي ١٥ عينة من سجق اللحم الجملى المجمد الذين تم جمعهم من السوبر ماركت المختلفة المتواجدة في مدينة القاهرة والجيزة والزقازيق. وقد تم تقدير بقايا هذه المعادن باستخدام جهاز إمتصاص الطيف الذرى.

وقد أظهرت النتائج أن القيم المتوسطة لعنصر الرصاص والكاديوم والنحاس في عينات لانشون اللحم الجملى كانت  $0.128 \pm 0.0008$  ،  $0.005 \pm 0.0004$  و  $3.662 \pm 0.221$  جزء فى المليون على التوالي بينما فى سجق اللحم الجملى المجمد كانت  $0.133 \pm 0.008$  ،  $0.008 \pm 0.0004$  و  $3.789 \pm 0.189$  جزء فى المليون على التوالي وقد تم مقارنة النتائج المواصفات القياسية لمنظمة الأغذية والزراعة/منظمة الصحة العالمية (١٩٧٢) ، منظمة الصحة العالمية (١٩٧٢) والهيئة المصرية العامة لتوحيد القياسى وجودة الإنتاج (١٩٩٢) وقد تم مناقشة الأهمية الصحية والتأثيرات السامة لهذه المعادن الثقيلة وكيفية الحد من التلوث لهذه المعادن التى تضر صحة المستهلكين.