

16S rDNA analysis for characterization of *Pseudomonas* sp. strain MF30 isolated from *Rumex acetocella* roots in northern Sweden

(Received: 15.09.2006; Accepted: 28.09.2006)

Idress H. Attitalla*; **Ali A. Bataw****; **Jolanta J. Borowicz***** and **Sture Brishammar*****

*Omar Al-Mukhtar University, Faculty of Science, Botany Department, Box 919, El-Beida, Libya

**Omar Al-Mukhtar University, Faculty of Science, Zoology Department, Box 919, El-Beida, Libya

***Maselaboratoratorerna AB, Box 148, Uppsala, Sweden

Corresponding author I. Idressattitalla2004@yahoo.com

ABSTRACT

A bacterial strain obtained from the northern part of Sweden previously classified as *Pseudomonas veronii* based on biochemical and physiological tests. In this study, phylogenetic tree was constructed using a nearly complete sequence within the 16S rDNA gene. The strain of *Pseudomonas* sp. subdivided into two rather distinctly related groups, neither of which is very close to the group within the *Pseudomonas fluorescens* cluster. Although the phylogenetic analysis is not conclusive, it is consistent with other observations, especially the capacities of this strain as a biocontrol agent. Taken all together, the results suggest that the MF30 strain should be classified as another *Pseudomonas* species, either *Pseudomonas antarctica* or *P. meridina*.

Key words: Phylogenetic analysis, 16S rDNA gene, *Pseudomonas* species, *P. veronii*, *P. antarctica*, *P. meridina*.

INTRODUCTION

Bacteria belonging to the fluorescent pseudomonads, known for the diversity of their metabolites (Leisinger and Margaff, 1979), contain species that are recognized as human and as animal pathogens (Nakazawa and Abe, 1996) and some are important phytopathogens (Jaunet *et al.*, 1995). Most species are however, saprophytic, and as a group, they are ubiquitous in water (Verhille *et al.*, 1997), soil (Lemanceau *et al.*, 1995), plant spermosphere (Fukui *et al.*, 1994), phyllosphere (Rainey *et al.*, 1994) and rhizosphere (Lambert *et al.*, 1990). In recent years certain species, for example *Pseudomonas chloroaphis*, *P. fluorescens*, *P. veronii*, *P. putida*, have

received attention even though they possess abilities to influence plant growth and development through different mechanisms (Weller, 1988; O'Sullivan and O'Gara, 1992). They are now recognised as being antagonistic to several opportunistic soil-borne fungi (Weller, 1988; Keel *et al.*, 1996) and to seed-borne fungi (Hökeberg *et al.*, 1997). Also, some strains are responsible for significant plant promoting effects (DeFreitas and Germida, 1991; Kropp *et al.*, 1996).

Taxonomically, this fluorescent *Pseudomonas* spp. belong to the rRNA group I of the gamma subclass of *Proteobacteria* (Palleroni, 1993; Kersters *et al.*, 1996). The species within this group have been differentiated on the basis of their DNA/Dna homologies (Johnson and Palleroni, 1989; Gardan *et al.*,

1992). This method of differentiating bacterial species has for several years been the most reliable, since traditional classification of bacteria based on phenotypic features does not always correlate with molecular taxonomy (Woese, 1987). Other methods based on cataloguing 16S rRNA have been shown to be reliable and effective (Woese *et al.*, 1984; Woese, 1987). Indeed, using the 16-rDNA approach, Woese *et al.* (1990) were able to show definitively that life can be divided into three major lines of descent presently named Archea, Bacteria and Eukarya.

For bacteria belonging to the genus *Pseudomonas*, data collected on 16S rDNA genes make taxonomical studies possible, including those in the rhizosphere. As a result of such studies, many organisms classified as species of this genus have been reclassified (Kerstens *et al.*, 1996). However, classification of the fluorescent pseudomonads has yet to be clarified completely, in part due to the many new species being described (Elomari *et al.*, 1996). This paper a case in point. In earlier study, Attiatalla *et al.* (2001) identified a bacterial strain (MF30) based on the characteristics revealed by API NE 20 test (BioMerieux, Marcy l'Etoile, Marcy, France), the bacterial strain MF30 was assigned to the genus *Pseudomonas*, also partial sequences of 16S rDNA showed 99.6% similarity to *Pseudomonas veronii* and high similarity to other members of RNA group I of the *Pseudomonas*. Strain testing was done in Deutsche Sammlung von Mikro-organismen aus Zellkulturen GmbH, Braunschweig, Germany (DSMZ), which also found that the profile of the cellular fatty acids was typical for that RNA-group, and although physiological tests were unable to distinguish between the species in this group, yet they were all closely related to biovars of *Pseudomonas fluorescens*. This strain has been shown to produce biologically active

metabolites namely massetolides type A, a low molecular weight lipopeptide (Gerard *et al.*, 1997). This compound, as well as several derivatives (massetolides B-H), were recently isolated from two *Pseudomonas* spp. isolates of a marine algae and a marine tube worm origin (Gerard, 1997). To date, there only two reports of *in vitro* anti-tuberculosis activity expressed by massetolides (El-Sayed *et al.*, 2000).

In the present paper, we successfully applied restriction fragment length polymorphism and sequencing of the 16S rDNA for clarifying the classification of the strain MF30. This study indicates that it differs from the *P. veronii*. However, since 16S rDNA data precluded the prospect of conclusive results from phylogenetic analysis, one can rule out the possibility that the strain can represent a specific group or biotype of *P. veronii*. Using sequencing data from nearly complete sequence of the 16S rDNA gene together with their biological characters, we describe the strain MF30. We proposed that this bacterium should be reclassified as a new *Pseudomonas* species, either *P. antarctica* or *P. meridina*.

MATERIALS AND METHODS

Bacterial strain and cultivation

Using the procedure described by Hökeberg *et al.*, (1997), the bacterial strain MF30 was isolated from root of a wild plant (*Rumex acetocella*) in the mountain area of northern Sweden: The strain MF30 from the dicotyledonous plant. The strain was maintained as freeze-dried or as deep-frozen (-70°C) stock culture. The bacterium was grown in 50% strength Tryptic Soy Broth (15g/k, TSB, Difco) it shaking (120-rev min⁻¹) at 20°C for 48 hours.

Secretion of extracellular proteases

Proteolytic activity was estimated in 48 hours old cell free supernatant from bacterial culture grown in 50% normal strength TSB (15g/l) (Tryptic Soy Broth, Difco). Bacterial culture was incubated on rotary shaker (120-rev min⁻¹). Cells were removed by centrifugation for 20 min at 16.000 g and supernatant was then filter sterilized. The radial diffusion assay (Dow *et al.*, 1990) in milk agar plates was used. Fifty microliters of supernatant was applied to a well cut (6 mm in diameter) in the middle of skim milk agar plates, which were then incubated at 37°C. Clear zones (mm) around the well were measured after 24 hours. The test was repeated twice with two Petri-dishes per treatment.

Phenotypic characterization

The strain was also characterized phenotypically with regard to production of fluorescent colonies on King'B medium, and for motility in a swarming assay (Bondesson and Haas, 2002; Thorsson, 2002).

Fluorescence

Siderophore production was detected as fluorescence under UV light (wave length = 320 nm) when cells were spread on KB medium (King *et al.*, 1954). Plates were incubated at 25°C for 2 days prior to visual examination of fluorescence on UV a transilluminator.

Swarming

To test swarming ability, 1 µl of bacterial suspension (a small loop of fresh bacterial cells suspended in 100 µl high purity water) of the strain was inoculated onto liquid medium (LM plates containing 10g/l Bacto-trypton, 6g/l Bacto-yeast extract, .5 g/l K₂HPO₄, 0.5 g/l NaCl, 0.4g/l MgSO₄·7H₂O).

DNA isolation and PCR amplification

The bacterium was grown in 50% normal strength TSB on a rotary shaker (120-rev min⁻¹) at 20°C for 48 hours. The RapidPrepTM genomic DNA Isolation Kit for Cells and Tissues (Pharmacia Biotech) was used to isolate DNA as advised by the manufacturer. DNA amplification was conducted on pure 2 to 3 µl DNA sample with about 150 ng of DNA per 1 µl of sample in a perkin Elmer (Nowalk, CT) thermocycler. The fD1 and rD1 primers amplifying the 16S rDNA gene in most of bacteria was used for DNA amplification (Weisburg *et al.*, 1991). PCR reactions were conducted in a final volume of 100 µl with 0.2 µM of each primer, 2.0 mM MgCl₂, 200µM dNTPs and 2.5 units of AmpliTag DNA polymerase mixed in the PCR buffer (Perkin Elmer). DNA was amplified over 35 cycles of denaturation for 1 min at 94°C annealing at 55°C for 1.5 min and extension at 72°C for 2 min. After the last cycle, DNA was extended at 72°C for 10 min. Amplification was confirmed by analyzing 5µl of PCR reaction mixture on 1% agarose gel (Promega).

DNA sequencing

The PCR-product was purified using QIAquick PCR Purification Kit (Qiagen), and sequenced using automatic ABI 310 DNA Sequencer, Big Dye Terminator Cycle Sequencing Ready Reaction Kit, Perkin Elmer. The sequencing was performed in both direction using previously described primers (Lane *et al.*, 1985; Lane, 1991). Sequencing data was analyzed by two different computer alignment programs, DNASTar (DNASTAR, Inc., USA) and Sequence Navigator (Perkin, Corp., USA).

Determination of phylogenetic relationships

The BLAST database (Altschul *et al.*, 1997) of National Center for Biotechnology

Information was used to compare resolved sequence of the MF30 strain with known 16S rDNA sequences. Determination of phylogenetic relationships was analyzed by the program Phylogenetic Analysis Using Parsimony (PAUP) version 4.0b1 for Macintosh (Swofford, 1993). The robustness of the internal branches of the trees was estimated by bootstrap analyses using 1000 replications in a heuristic search with random stepwise addition (3 replications) (Vinnere *et al.*, 2002). Bootstrap majority-rule (> 50%) consensus trees were obtained.

DISCUSSION

Restriction fragment analysis of PCR-amplified 16S rDNA gene was used to classify

Table (1). The phenotype characterization of the strain MF30.

Bacterium	Fluorescence	Swarming	Protease
MF30 strain	+	+	++

The available gene databases on 16S rRNA genes presently contain a number of sequences, which allow reconstruction of phylogenetic relationships and precise positioning of even unknown species. In case of genus *Pseudomonas* (sensu stricto), nearly-complete sequences have been determined for the PCR amplified 16S rRNA genes of as many as 21 species (Moore *et al.*, 1996). In this paper, we have further investigated the taxonomic position of the bacterial strain (MF30), which we propose here represent a new species either *Pseudomonas antarctica* or *P. meridin*, upon sequencing most of the 16S rDNA gene, we discovered that the strain MF30 shares the same identical sequence, which is nearly the complete 16S rRNA gene. The identity of this sequence to the closest

an important bacterial strain. Attitalla *et al.*, (2001) attempted to identify the MF30 strain on the basis of morphological and biochemical characters; the strain was close to *P. veronii*. Considering the fact that the estimated number of culturable bacteria is probably only 0.1 to 1% of those present in nature (Bintrim *et al.*, 1997), many totally new bacterial strains are still being recovered (Borowicz, 1998). The traditional taxonomic methods based on morphological physiological, and biochemical characters are now accompanied by DNA-based methods like DNA-DNA hybridization, RFLP and sequencing of 16S rRNA genes (Grimont *et al.*, 1996; Hartung, 1998).

pseudomonads strains is 98-99%. Therefore, the previous identification of the strain MF30 by API system (Attitalla *et al.*, 2001), and the identification of this strain by fatty acid analysis as well as species of *P. veronii* was not confirmed.

It is interesting to note that in the phylogenetic tree constructed in this study, the strain MF30 is subdivided into 2 rather distantly related groups, neither of which is very close to the group within which the strains *P. antarctica* or *P. meridin* cluster (Fig. 1). The strain MF30 is clearly shown to be an extracellular protease producer (Table 1). The proteolytic activity on skim milk agar medium was visually determined, and by measuring the proteolytic activity zone of clearing in mm, the zone was 15.0 mm.




Fig. (1): The BLAST database (Altschul et al., 1997) of National Center for Biotechnology Information (NCBI) was used to compare resolved sequence of the MF30 strain with known 16S rDNA sequences. The Phylogenetic tree was reconstructed for the strain MF30.

The finding that protease production by the strain MF30 was inversely related to its growth rate in the tested medium (data not shown) further confirms this hypothesis. The protease-producing bacterium was also capable of secreting protease(s) when grown in non-protein medium such as mineral medium (MMB) supplemented with glucose as a single carbon source. This suggests that at least some protease(s) are constitutively produced (Whooley *et al.*, 1983).

Based on phylogenetic study, the analysis of the 16S rDNA gene sequences data for the strain MF30, *P. antarctica*, and *P. meridina* share the same sequence, but differ from *P. veronii*. The strain studied was probably too closely related to support a meaningful parsimony analysis and construction of a phylogenetic tree. The genetic relationships between the strain MF30 and known members of other species of *Pseudomonas* genus were estimated by parsimony analysis (Swofford, 1993) using heuristic search with TBR branch swapping (100 replicates). The bootstrap analyses were run with TBR MULPARS and 1000 replicates. Nine equally parsimonious trees, which showed few differences in topology analysis is shown in Fig. (1). These findings support further taxonomic analysis of the isolates by sequencing of the full 16S rRNA gene by DNA/DNA hybridization and/or by PCR analysis of other genes, preferably from non-coding DNA region.

ACKNOWLEDGEMENTS

We gratefully acknowledge Professor Siv Andersson and Dr. Paul Johnson (Uppsala University, Sweden) for assistance in interpreting the data and for critical reading and refining of the manuscript.

REFERENCES

- Altschul, F; Madden, T.L.; Schäffer, A.A.; Zhang, J.; Zhang, Z; Miller, W and Lipman, D.J. (1997). Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Research*, 25: 3389–3402.
- Attiatalla, I.H; Johansson; Maria, P.Brishammar, S, and Gerhardson, B. (2001). *Pseudomonas* sp. strain MF30 suppresses Fusarium wilt of tomato *in vivo*. *Phytopathologia Mediterranea*, 40: 234–239.
- Bintrim, S.B; Donohue, T.J; Handelsman, J; Roberts, G.P. and Goodman, R.M. (1997). Molecular phylogeny of *Archea* from soil. *Proceedings of National Academy of Science*, 94: 277–282.
- Bondesson, C. and Haas, D. (2000). Mechanism regulation and ecological role of bacterial cyanide biosynthesis. *Arch. Microbiology*, 173: 170–177.
- De Freitas, J.R. and Germida, J.J. (1991). *Pseudomonas cepacia* and *Pseudomonas putida* as winter wheat inoculants for biocontrol of *Rhizoctonia solani*. *Canadian Journal of Microbiology*, 37: 780–784.
- Dow, J.W; Clarke, B.R; Milligan, D.E; Tang, J.L. and Daniels, M.J. (1990). Extracellular proteases from *Xanthomonas campestris* pv. *campestris*, the black rot pathogen. *Applied Environmental Microbiology*, 56: 2994–2998.
- Elomari, M; Coroler, L; Hoste, B; Gillis, M; Izard, D. and Leclerc, H. (1996). DNA relatedness among *Pseudomonas* strains isolated from natural mineral waters and proposal of *Pseudomonas veronii* sp. nov. *International Journal of Systematic Bacteriology*, 46: 1138–1144.
- El Sayed, K.A; Bartyzel, P.; Shen, X., Perry, T.L; Zjawiony, J.K. and Hamann, M.T. (2000). Marine natural products as

- antituberculosis agents. *Tetrahedron*, 56: 949–953.
- Fukui, R; Schroth, M.N; Hendson, M, Hancock, J.G. and Firestone, M.K. (1994).** Growth patterns and metabolic activity of pseudomonads in sugar beet spermospheres: Relationship to pericarp colonization by *Pythium ultimum*. *Phytopathology*, 84: 1331–1337.
- Gardan, L.; Bollet, C, Abu Ghorrah, M. Grimont, F. and Grimont, P.A.D. (1992).** DNA relatedness among the pathovar strains of *Pseudomonas syringae* supsp. *Savastanoi* Janse (1982) and proposal of *Pseudomonas savastanoi* sp. nov. *International Journal of Systematic Bacteriology*, 42: 606–612.
- Gerard, J. Lloyd, R; Barsby, T; Haden, P, Kelly, M.T. and Andersen, R.J. (1997).** Massetolide A-H, antimycobacterial cyclic depsipeptides produced by two pseudomonads isolated from marine habitats. *Journal of Natural Products*, 60: 223–229.
- Grimont, P.A.D; Vancanneyt, M; Lefèvre, M; Vandemeulebroecke, K. Vauterin, L, Brosch, R, Kersters, K and Grimont, F. (1996)** Ability of biologic and Biotype-100 systems to reveal the taxonomic diversity of the pseudomonads. *Systematic Applied Microbiology*, 19: 510–527.
- Hartung, J. S. (1998)** Molecular probes and assays useful to identify plant pathogenic fungi, bacteria, and marked biocontrol agents. In: Boland, GJ and Kuykendall, LD (eds) *Plant microbe interactions and biological control*. Marcel Dekker, Inc. New York, Basel and Hongkong. 393–413.
- Hkeberg, M; Gerhardson, B. and Johnson, L. (1997).** Biological control of cereal seed-borne diseases by seed bacterization with greenhouse-selected bacteria. *European Journal of Plant Pathology*, 103: 25–33.
- Jaunet, T. Laguerre, G; Lemanceau, P, Frutos, R and Notteghem, J.L. (1995).** Diversity of *Pseudomonas fucovaginae* and other fluorescent pseudomonads isolated from diseased rice. *Phytopathology*, 85: 1534–1541.
- Johnson, J.L. and Palleroni, N.J. (1989).** Deoxyribonucleic acid similarities among *Pseudomonas* species. *International Journal of Systematic Bacteriology*, 39: 230235.
- Keel, C. Weller, D.M; Natsch, A; Défago, G; Cook, R.J. and Tomashow, L.S. (1996).** Conservation of the 2,4-diacetylphloroglucinol biosynthesis locus among fluorescent *Pseudomonas* strains from diverse geographic locations. *Applied Environmental Microbiology*, 62: 552–563.
- Kersters, K; Ludwig, W; VanCanneyt, M; De Vos, P; Gillis, M and Schleifer, K-H. (1996).** Recent changes in the classification of the pseudomonads: an overview. *Systematic Applied Microbiology*, 19: 465–477.
- King, E.O. Ward, M.K. and Raney, D.E. (1954).** Two simple media for the demonstration of Pyocyanin and fluorescin. *J. Lab. Clin. Med.* 44: 301–307.
- Kropp, B.R; Thomas, E; Pounder, J.I. and Anderson, A.J. (1996).** Increased emergence of spring wheat after inoculation with *Pseudomonas chlorophis* isolate 2E3 under field and laboratory conditions. *Biological Fertility of Soil*, 23: 200–206.
- Lane, D.J; Pace, B; Olsen, G.J; Stahl, D.A; Sogin, M.L. and Pace, N.R. (1985).** Rapid determination of 16S ribosomal RNA sequences for phylogenetic analysis. *Proceedings of National Academy of Science*, 82: 6955–6959.
- Lane, D.J. (1991).** 16/23S rRNA sequencing: In Stackebrandt, E and Goodfellow, M (eds) *Nucleic acid techniques in bacterial systematics*. John Wiley and Sons, Chichester, New York, Brisbane, Toronto and Singapore. 115–175.
- Lambert, B; Meire, P; Joos, H. ;Lens, P. and Swings, J. (1990).** Fast-growing, aerobic, heterotrophic bacteria from the

- rhizosphere of young sugar beet plants. *Applied Environmental Microbiology*, 56: 11: 3375–3381.
- Lemanceau, P. Corberand, T; Gardan, L.; Latour, X; Laguerre, G; Boeufgras, J-M and Alabouvette, C. (1995).** Effect of two plant species flax (*Linum usitatissimum* L.) and two tomato species (*Lycopersicon esculentum* Mill.) on the diversity of soilborne opulations of florescent pseudomonaads. *Applied Environmental Microbiology*, 61: 1004–1012.
- Leisinger, T. and Margraff, R. (1979).** Secondary metabolites of the fluorescent pseudomonads. *Microbiology Review*, 43: 422–442.
- Moore, ERB, Mau; M; Arnscheidt, A; Bttger, EC; Hutson, R.A; Collins, MD, van de Peer, Y, de Wachter, R and Timmis, K.N. (1996).** The determination and comparison of the 16S rRNA gene sequences of species of the genus *Pseudomonas* (sensu stricto) and estimation of the natural intrageneric relationships. *Systematic Applied Microbiology*, 19: 478–492.
- Nakazawa, T. and Abe, M. (1996).** Pathogenesis of *Burkholderia cepacia* and export of protease by general secretory pathway involving disulfied bond formation in the periplasm. In: Nakazwa, T., Furukawa, K Haas, D. and Silver, S. (eds). *Molecular Biology of Pseudomonads*. ASM Press, Washington, D.C. 462–471.
- O'Sullivan, D.J. and O'Gara, F. (1992).** Traits of fluorescent *Pseudomonas* spp. Involved in suppression of plant root pathogens. *Microbiology Review*, 56: 662–676.
- Palleroni, N.J. (1993).** *Pseudomonas* classification: A new case history in the taxonomy of Gram-negative bacteria. *Antonie van Leeuwenhoek*, 64: 231251.
- Pecknold, P.C. and Grogan, R.G. (1973).** Deoxyribonucleic acid homology groups among phytopathogenic *Pseudomonas* species. *International Journal of Systematic Bacteriology*, 23: 111121.
- Rainey, P.B; Bailey, M.J. and Yhomson, I.P. (1994).** Phenotypic and genotypic diversity of florescent pseudomonads isolated from field-grown sugar beet. *Microbiology*, 140: 2315–2331.
- Swofford, D.L. (1993).** PAUP: Phylogenetic analysis using parsimony, Version 3.1.1. Illinois Natural History Survey, Champaign, IL.
- Thorsson, L. 2002.** Characterisation of four regulatory mutants of *Pseudomonas chlororaphis* Vb10, a biocontrol agent. Final Thesis. Mlardalens Hgskola.
- Verhille, S; Elomari, M; Coroler, L.; Izard, D and Leclerc, H. (1997).** Phenotypically based taxonomy of fluorecent *Pseudomonas* strains isolated from four natural mineral waters. *Systematic Applied Microbiology*, 20: 137–149.
- Vinnere, O.; Fatehi, J.; Wright, S. A. I.; and Gerhardson, B. (2002).** The causal agent of anthracnose of *Rhododendron* in Sweden and Latvia. *Mycological Research*, 106: 60–69.
- Weisburg, W.G. ; Barns, S.M; Pelletier, D.A. and Lane, DJ. (1991).** 16S ribosomal DNA amplification for phylogenetic study. *Journal of Bacteriology*, 173: 697–703.
- Weller, D.M., (1988).** Biological control of soilborne plant pathogens in rhizosphere with bacteria. *Annual Review of Phytopathology*, 26: 379–407.
- Whooley, M.A.; O'Callaghan, J.A. and Mcloughlin, A.J. 1983.** Effects of substrate on the regulation of exoprotease production by *Pseudomonas aeruginosa* ATCC10145. *Journal of General Microbiology* 129, 4: 981–988.

Woese, C.R; Blanz, P. and Hahn, C.M. (1984). What isn't pseudomonads: the importance of nomenclature in bacterial classification. Systematic Applied Microbiology. 5, 179-195.

Woese, C.R. (1987). Bacterial evolution. Microbial Review 6, 60-69.

Woese, C.R, ; Kandler, O. and Wheelis, M.L. (1990). Towards a natural system of organisms: Proposal for the domains Archea, bacteri and Eucarya. Proceeding Natural Academy of Science, 7, 4578-4579.

الملخص العربي

تحليل الـ 16S rDNA لتوصيف السلالة MF30 لبكتريا السيدوموناس المعزولة من جذور نبات *Rumex acetocella* في شمال السويد

إدريس حمد عطية الله *، وعلى عبد القادر بطاوي، **جولانتا بروكز، **ستورا بريسهمر ***
 جامعة عمر المختار-كلية العلوم-قسم علم النبات-ص.ب. 919-البيضاء-الجمهورية العظمى*
 جامعة عمر المختار-كلية العلوم-قسم علم الحيوان-ص.ب. 919-البيضاء-الجمهورية العظمى**
 معمل ماسا AB-ص.ب. 148-أيسالا-المملكة السويدية***

السلالة البكتيرية MF30 والمعزولة من شمال المملكة السويدية والتي في السابق صنفت كسلالة *Pseudomonas veronii*، وذلك إستناداً إلى الإختبارات البيوكيميائية والفسولوجية. في الدراسة الحالية، شجرة التطور الوراثية والتي تم إستخلاصها بإستخدام تتابع (قريب من التتابع الكامل) لجين 16S rDNA. وهذه السلالة *Pseudomonas* قُسمت إلى مجموعتين محددين، أي من كليهما قريب من مجموع *Pseudomonas fluorescens*، وبالرغم من أن تحليل التطور الوراثي ليس شاملاً، لكنه أظهر بأن يتفق مع أو يدعم المشاهدات الأخرى، خصوصاً لسعة هذه السلالة كعامل في المكافحة الإحيائية. بأخذ كل هذه الإعتبارات جملة، أظهرت الاستنتاجات أو توصلت إلى أن سلالة MF30 يجب أن تصنف كنوع آخر من جنس *Pseudomonas*، إما أن يكون *Pseudomonas antarctica* أو *Pseudomonas merdina*.