

Dept. of Animal Prod.
Fac. of Agric. Al-Azhar University, Nasr City, Cairo, Egypt

**ALTERATIONS IN CONCENTRATIONS OF INHIBIN
IN RELATION TO FSH AND STEROIDS HORMONE
DURING ESTROUS CYCLE IN EGYPTIAN
BUFFALOES**
(With One Figure)

By

H.M. BADR

(Received at 13/9/2007)

**التغير في تركيزات الأنهيبين وعلاقته بالحائفة الجريبية وهرمونات المبيض
أثناء دورة الشياح في الجاموس المصري**

حسين محمد بدر

أجريت هذه الدراسة في محطة بحوث الإنتاج الحيواني بمسطررد التابعة لقسم الإنتاج الحيواني بكلية الزراعة جامعة الأزهر بالقاهرة. استخدم في هذه الدراسة عدد سبع جاموسات أظهرت الشياح المصاحب بالتبويض وتم معرفة ذلك عن طريق الجس المستقيمي وكشف الشياح ثلاث مرات يومياً. استهدفت الدراسة متابعة التغيرات في بعض الهرمونات التي اشتملت على هرمونات الأنهيبين والحائفة الجريبية وهرمون الاستروجين وهرمون البروجسترون. تراوحت مستويات هرمون الأنهيبين بين ٤٣٩ إلى ٥٨٣ بيكوجرام/ملل أثناء المراحل المختلفة لدورة الشياح، وتقد كان أعلى تركيز لهرمون الأنهيبين يوم الشياح بالنسبة، الحائفة الجريبية فقد كان أعلى تركيز لهذا الهرمون يوم الشياح ٢٥,٤٨ نانوجرام/ملل ثم انخفض وتذبذب مستواه باتجاه نهاية دورة الشياح حيث بدأ في الزيادة مرة أخرى على نهاية دورة الشياح. كان أعلى تركيز لهرمون الاستروجين ١٨,٥٤ بيكوجرام/ملل يوم الشياح بينما كان أعلى تركيز لهرمون البروجسترون ٤,٣٩ نانوجرام/ملل اليوم الثاني عشر من دورة الشياح. يمكن أن نستخلص من هذه الدراسة أن كل من هرمون الأنهيبين والاستروجين من الممكن أن يكون لهما تأثير رجعي على تنظيم إفراز الحائفة الجريبية في الجاموس المصري.

SUMMARY

Seven Egyptian buffaloes were used to study the plasma changes of inhibin, FSH, estradiol-17 β and progesterone concentrations during the estrous cycle. These hormones were determined at estrous and at two interval until day twenty two of estrous cycle. Plasma inhibin estimated by enzyme-linked immunoasorbent assay (ELISA). Both FSH, estradiol-

17 β and progesterone were estimated by radioimmunoassay (RIA). In order to elucidate the feedback system controlling mentioned hormones during estrous cycle, variations in the levels of hormones and their inter-relationships were studied. Plasma inhibin levels ranged between 439.28 and 584 pg/ml during various stages of estrous cycle, peak of inhibin level was at day of estrous by an average of 584. pg/ml. Follicle stimulating hormone levels were higher at day of estrous (25.48 ng/ml), decreased and fluctuated towards end of estrous cycle and then increased at day twenty two of estrous cycle. Peak of estradiol-17 β was 18.54 pg/ml at day of estrous, while the peak of progesterone was 4.39 ng/ml at twelfth day from the start of estrous. It could be concluded that each of inhibin and estradiol-17 β seems to have a feedback regulatory effect on FSH secretion in buffaloes.

Key words: *Inhibin, FSH, estradiol-17 β , progesterone, estrous cycle, buffalo*

INTRODUCTION

Inhibin is secreted by granulosa cells in response to FSH and its major action is the negative feedback control of pituitary FSH secretion (Ying, 1988). In mammalian species, ovulation is associated with the morphological, chemical and physiological changes in the ovulatory follicle. All these changes occur in response to the preovulatory LH and FSH (Dhali, *et al.* 2007).

Inhibin is a glycoprotein with two subunits α and β that belongs to the forming factor- β superfamily (Padilla-Rivas and Holtz 2006). The ovarian follicles have been identified as the major source of inhibin in sheep (Mann, *et al.* 1989), in goats (Medan, *et al.* 2005) in cattle (Rodgers, *et al.* 1989) and buffaloes (Palta, *et al.* 1996). This hormone have been studied in buffaloes by only a few authors over the past few years. Palta, *et al.* (1997) observed an increase of inhibin concentrations during the follicular phase with a peak occurring on day zero (estrous day defined by the lowest progesterone value). Inhibin has been implicated in the feedback regulation of FSH in heifers (Turzillo and Fortune 1990), in cows (Kaneko, *et al.* 1995) and in buffaloes (Singh, *et al.* 2001).

In cows, plasma FSH surge was significantly higher around estrous (Dhali, *et al.* 2007). Medan, *et al.* (2005) reported that inhibin A was negatively correlated with FSH, while it was positively correlated

with estradiol-17 β in goats. However, few information are available on the inhibin levels during estrous cycle in buffaloes. Therefore, the present study was undertaken to investigate the inhibin levels during different stages of estrous cycle in relation to FSH, estradiol-17 β and progesterone in Egyptian buffaloes.

MATERIALS and METHODS

Seven Egyptian buffaloes, aged 3-4 years, from the experimental farm of the Faculty of Agriculture Al-Azhar University, were used. Animals were fed according to their live body weight and production, on concentrate mixture containing 17% crude protein, 2.5% fat and 15% crude fiber. In addition to rice straw or wheat straw and berssem hay, Egyptian clover, (*Trifolium alexandrinum*) which was offered at 20 to 25 Kg/animal/day.

Blood samples were collected from the jugular vein of buffaloes at day of estrous and at two days intervals thereafter for 22 days. Buffaloes were observed for estrous sings three times daily, at morning, mid day and afternoon. A separate fertile bull in addition to rectal palpation were used for of determination estrous day. Plasma was separated and stored at- 20⁰C till determination of the inhibin, follicle stimulating hormone (FSH), estradiol-17 β and progesterone.

Inhibin level was determined using enzyme-linked immunoasorbent assay (ELISA) kit, manufactured by Biosource Europe S. A, Rue de l'industrie, 8 B-1400 Nivelles-Belgium, as described by Buckler (1992).

The mature form of inhibin has a molecular weight of 32.000 daltons, and consists of one alpha-chain (approx. 18 KDa) and one beta-chain (14 KDa) linked by disulphide bridges (Ying 1988). Plasma FSH, estradiol-17 β and progesterone were estimated by direct radioimmunoassay (RIA) using coat A-count kit. (DPC-Diagnostic, Corporation, Los Angeles CA 90045, U.S.A.) According to Razdan, *et al.* (1982), Xing, *et al.* (1993) the and Abraham (1981) for three the hormones receptively. Statistical analysis was carried out using Duncan's multiple range. Correlation coefficient among various hormones in the different phases was evaluated according to GL M procedure of SAS (1988).

RESULTS

The mean concentrations of inhibin, FSH, estradiol-17 β and progesterone from day of estrous to day twenty two of estrous cycle are shown in Fig. (1). The highest concentrations of plasma inhibin (584 pg/ml) and of FSH (25.48 ng/ml) were recorded at the day of estrous. A general decrease of both inhibin and FSH concentrations was observed to reach the lowest level (439.28 pg/ml and 13.25 ng/ml) at days 8 and 12 of estrous cycle, respectively. The concentrations of inhibin and FSH were slightly increased to reach 448.14 pg/ml and 18.11 ng/ml, respectively, at the end of estrous cycle. The differences of inhibin and FSH concentrations during the estrous cycle were significant ($P < 0.01$).

As shown in Fig (1) the general pattern of secretion of steroids showed that the highest estradiol-17 β level was 18.54 pg/ml at day of estrous. At the same time the level of progesterone was low 0.64 ng/ml. Estradiol-17 β has drastically dropped (12.07 pg/ml) at day two of the estrous cycle, continual to drop progressively to reach a nadir on day 18 (2.2pg/ml) then fluctuated and increased again 8.17 pg/ml towards the end of the cycle. The statistical analysis of estradiol-17 β concentrations among days of the estrous cycle was significant ($P < 0.01$). During estrous, the concentrations of inhibin/FSH and inhibin/estradiol-17 β were positively correlated (0.88 and 0.40 respectively). It seems that there are a synergistic interaction between inhibin and estradiol-17 β in the control of FSH secretion in buffaloes. The plasma progesterone profiles confirmed that all experimental animals have normal estrous cycle and ovulated following the onset of estrous. Progesterone concentrations reached a high level (4.39 ng/ml) at twelfth day of the estrous cycle. A gradual decrease of progesterone concentration was observed (0.52 ng/ml) at the end of estrous cycle.




Fig. 1: Plasma concentrations of inhibin, FSH, estradiol-17 β and progesterone during the estrous cycle.

DISCUSSION

The inhibin levels reported in this study are in coincidence with the levels reported by Singh, *et al.* (2001) and close to those reported by Palta, *et al.* (1997). The same trend was observed by Palta, *et al.* (1996) who found an increase of inhibin concentrations during the follicular phase with a peak occurring on day zero (estrous day). Kaneko, *et al.* (1992) reported a maximum inhibin concentration in cows two hours

after the LH peak. Vanmontfort, *et al.* (1998) found that inhibin concentration showed no significant variation during the estrous cycle in sheep, however, highest levels were recorded during the follicular phase, which then decreased during the luteal phase to basal levels.

In the present study the inhibin levels declined continuously after the day of estrous and were at their lowest value until twelfth day of the cycle which is corresponded to the absence of follicular growth on the ovary at this time in buffaloes. Singh, *et al.* (2001) demonstrated that the emergence of follicular waves seems closely tied to the elevation of plasma FSH. They postulated that species possessing large-size follicles, follicles do not develop during the luteal phase and have higher levels of feedback regulators such as inhibin.

A consistent fall of inhibin 1-4 h after LH peak, reaching a nadir at the time of the second surge of FSH has been reported (Findlay, *et al.* 1990). Amounts of inhibin in the follicular fluid and inhibin α subunit mRNA in granulosa cells are greater in healthy follicles compared to atretic follicles at several stages of development (Handerson, *et al.* 1984 and Ireland and Ireland 1994).

The present study showed that inhibin concentrations slowly increased again from the eighteenth day until the end of estrous cycle, suggesting that this increase was due to development of follicle on the ovary for new the cycle. Singh, *et al.* (2001) reported that the rise of inhibin and setradiol-17 β could be attributed to the growing crop of follicles on the ovary, which seems to have started around the eighth day of the cycle in buffaloes.

In general, the level of FSH reported in this study are in agreement with those noticed by Singh *et al.* (2001) and Dhali, *et al.* (2007). The later authors observed that the concentration of FSH was higher during the first three days of estrous cycle. The FSH level decreased to the basal level on day four of the estrous cycle. The results of the present study were lower than those found by Razdan, *et al.* (1982) on the day of estrous. However, Janakiraman, *et al.* (1980) have reported even higher levels during the whole estrous cycle.

The estradiol-17 β levels found in this study were at the same order as observed by Barkawi (1984), El-Terbany (1998) and Singh, *et al.* (2001) and lower than reported by Kumar, *et al.* (1991) on the day of estrous. Estradiol-17 β levels were found to be higher than those reported by Avenell, *et al.* (1985) on the day of estrous. The results showed that plasma progesterone concentrations are similar to those found by Barkawi (1984), Panchal, *et al.* (1992), Sarvaiya, *et al.* (1993) and Singh, *et al.* (2001).

Medan, *et al.* (2005) indicated that growing follicles secreted inhibin and estradiol-17 β which in turn suppressed FSH secretion in goats. Kaneko, *et al.* (1995), showed that passive immunization against inhibin during the follicular phase of the cycle in resulted in a transient increase of FSH, in cows thereby showing that inhibin has a definite role in the control of FSH secretion. Bleach, *et al.* (2001) showed that circulating inhibin A concentrations were positively correlated with levels of estradiol-17 β , whereas circulating concentrations of FSH were negatively correlated with inhibin A levels.

Fluctuation of the circulating FSH levels is involved in the recruitment of the follicles (Medan, *et al.* 2005). Also, they showed that inverse relationship between inhibin and FSH confirms the hypothesis that inhibin inhibits FSH secretion. Medan, *et al.* (2003) found an inverse relationship between FSH and inhibin A during estrous cycle in goats that confirms the hypothesis that inhibin A contributes to the inhibition of FSH secretion.

The present investigation showed high progesterone levels during mid of estrous cycle which indicates fully developed corpus luteum. There is no positive correlation between inhibin and progesterone (-0.39). From the obtained results, it could be confirmed that corpus luteum is the source of inhibin in buffaloes. Similar observations were reported by Singh, *et al.* (2001) in buffaloes and Kaneko, *et al.* (1992) in cattle.

It could be concluded that each inhibin and estradiol-17 seems to have a feedback regulatory effect on FSH secretion in Egyptian buffaloes.

REFERENCES

- Abraham, G.E. (1981):* The application of natural steroid radioimmunoassay to gynecologic endocrinology. In: Abraham, G.E., editor Radioassay Systems in Clinical Endocrinology. Basel: Marcel Dekker, 475-529.
- Avenell, J.A.; Saepudin, Y. and Fletcher, I.C. (1985):* Concentration of luteinizing hormone, estradiol-17 β and progesterone in the peripheral plasma of Swamp buffalo cows around the time of estrous. *J. Reprod. Fert.* 74: 419-424.
- Barkawi A.M. (1984):* Pre-partum and post partum hormonal pattern and reproductive activity in Egyptian buffaloes. Ph.D Thesis, Fac. Agric., Cairo Univ., Egypt.

- Bleach E.C.; Glencross R.G.; Feist S.A.; Groome N.P. and Knight P.G (2001):* Plasma inhibin A in heifers: relationship with follicle dynamics, gonadotropins, and steroids during the estrous cycle and after treatment with bovine follicular fluid. *Biol. Reprod.* 64: 743-752.
- Buckler, H.M. (1992):* Immunoreactive inhibin levels during ovarian stimulation may predict granulosa cell maturity. *Clin. Endocr.* 37: 1-6.
- Dhali, A.; Mishra, D.P.; Mech, A.; Prakash, B. and Rajkhowa, C. (2007):* Evidence of multiple preovulatory LH and FSH surges in mithun (*Bos frontalis*). *J. of cell and Anim. Biol.* 1: 1-6.
- El-Terbany, H.M.B. (1998):* Study of some reproductive aspects for post-partum period in Egyptian buffaloes. Ph.D Thesis, Fac. Agric. Al-Azhar Univ., Cairo, Egypt.
- Findlay, J.K.; Clarke, J. and Robertson, D.M. (1990):* Inhibin concentrations in ovarian and jugular venous plasma and relationship of inhibin with follicle stimulation hormone and luteinizing hormone during the ovine oestrous cycle. *Enocri.* 126: 528-535.
- Handerson K.M; Franchimont P.; Charlet-Renard, C. and McNatty, K.P. (1984):* Effect of follicular atresia on inhibin production by bovine granulosa cells in vitro and inhibin concentration in the follicular fluid. *J. Reprod. Fertil.* 72: 1-8.
- Ireland J.L. and Ireland J.J. (1994):* Changes in expression of inhibin/activin alpha A and beta B subunit messenger ribonucleic acids following increases in size and during different stages of differentiation or atresia of non-ovulatory follicles in cows. *Biol. Reprod.* 50: 492-501.
- Janakiraman, K.; Desai, M.C.; Amin, D.R.; Shett, A.R.; Moodbidri, S.B. and Wadadekar, K.B. (1980):* Serum gonadotrophin levels in buffaloes in relation to phases of estrous cycle and breeding periods. *Indian J. Anim. Sci.* 50: 601-606.
- Kaneko, H.; Nakamishi, Y.; Akagi, S.; Arai, K.; Taya, K.; Watanabe, G.; Sasamoto, S. and Hasegaw, Y. (1995):* Immunoneutralisation of inhibin and estradiol during the follicular phase of the estrous cycle in cows. *Biol. Reprod.* 53: 931-939.
- ko, H.; Watanbe, G.; Taya, K. and Sasamoto, S. (1992):* Changes in peripheral levels of bioactive and immunpreactive inhibin, estradiol-17 β , progesterone, luteinizing hormone and follicle stimulating hormone associated with follicular development in

- cows induced to superovulation with equine chorionic gonadotrophin. *Biol. Reprod.* 47: 76-82.
- Kumar P.; Jindal, R. and Rattan, P.J.S. (1991):* Plasma hormonal Profiles during estrous cycle of Murrah heifers. *Indian J. Anim. Sci.* 61: 382-385.
- Mann, G.E.; McNeilly, A.S. and Baird, D.T. (1989):* Source of ovarian inhibin secretion during the oestrous cycle of the sheep. *J. Endocrinol.* 123: 181-188.
- Medan M.S.; Watanabe, G.; Sasaki, K.; Groome, N.P.; Sharawy, S. and Taya, K. (2005):* Follicular and hormonal dynamics during the estrous cycle in goats. *J. Reprod. Develop.* 51: 455-463.
- Medan, M.S.; Watanabe, G.; SasaKi, K.; Sharawoy, S.; Groome, N.P. and Taya, K. (2003):* Ovarian dynamics and their associations with peripheral concentrations of gonadotropins, ovarian steroids and inhibin during the estrous cycle in goats. *Biol. Reprod.* 69: 57-63.
- Padilla-Rivas, G. and Holtz, W. (2006):* LH secretion in Boer goats actively immunized against recombinant ovine inhibin α subunit. *Vet. Mex.*, 37: 51-58.
- Palta, P.; Mondal, S.; Prakash, B.S. and Madan, M.L. (1997):* Peripheral inhibin levels in relation to climatic variations and stage of estrous cycle in buffalo (*Bubalus bubalis*). *Theriogenology* 47: 989-995.
- Palta, P.; Prakash, B.S. and Madan, M.L. (1996):* Peripheral inhibin levels during the oestrous cycle in buffalo (*Bubalus bubalis*). *Theriogenology*, 46: 993-998.
- Panchal, M.T.; Dhami, A.J.; Mehta, V.M. and Kodagali, S.B. (1992):* Plasma progesterone profile of fertile and infertile estrous cycle in repeat breeding buffaloes. *Indian Vet. J.* 69: 766-797.
- Razdan, M.N.; Kaker, M.L. and Galhotra, M.M. (1982):* Serum FSh levels during oestrous and a 4-week period following mating in Murrah buffaloes (*Bubalus bubalis*). *Theriogenology* 17: 157-181.
- Rodgers, R.J.; Stuchberry, S.J. and Findlay, J.K. (1989):* Inhibin mRNAs in ovine and bovine ovarian follicles and corpora lutea throughout the estrous cycle and gestation. *Mol. Cell Endocrinol.* 62: 95-101.
- S.A.S. Institute, Inc. (1988):* User's Guide Statistics, 1988. Ed., SAS, Inst. Cary NC., U.S.A.

- Sarvaiya, N.P.; Patel A.V. and Pathak M.M. (1993):* Blood levels of ovarian steroids during cyclic stages and anestrus condition of pubertal Surti buffalo heifers. *Indian J. Anim. Sci.* 63: 296-299.
- Singh, B.; Dixit, V.D.; Singh, P.; Georgie, G.C. and Dixit, V.P. (2001):* Plasma inhibin levels in relation to steroids and gonadotrophins during oestrous cycle in buffalo. *Reprod. Dom. Anim.* 36: 163-167.
- Turzillo, A.M. and Fortune, J.E. (1990):* Suppression of secondary FSH surge with bovine follicular fluid is associated with delayed ovarian follicular development in heifers. *J. Reprod. Fertile.* 89: 463-653.
- Vanmontfort, D.; Peeters, R.; Buys, N.; Rombauts, L. Yerhoeven, G.; Charlet-Renard, C.; Franchimont, P. and Decuyper, E. (1998):* Circulating levels of inhibin and FSH during the oestrous cycle in 4 genotypes of sheep with different reproductive performances. *Small Ruminant Research*: 29: 213-224.
- Xing, S.; Cekan, S.Z. and Diczfalusy, U.L. (1983):* Validation of radioimmunoassay for estradiol-17 β (by isotope dilution-mass spectrometry and by a test of radiochemical purity. *Clin. Chim. Acta*, 135: 189-201.
- Ying, S.Y. (1988):* Inhibins, activins, and follistatins: gonadal proteins modulating the secretion of follicle stimulating hormone. *Endocrine Reviews* 9: 267-293.