

Prevalence of Salmonella and E.coli O157 in some foods

N. S. Abd El-Atty, A. M. S.Meshref

*Food Hygiene and Control Department, Faculty of Veterinary Medicine, Beni-Suef University,
Beni-Suef 62511, Egypt.*

A total of 200 raw food samples including milk, kareish cheese, fresh sausage and hawawshy (spiced minced meat) (Fifty of each) were randomly collected from farmer's houses, butcher's shops and retail markets in Beni-Suef Governorate. All were screened for the presence of *E.coli* O157 and *Salmonella*. *E.coli* O157 could be detected in 1 (2%) and 1 (2%) of kareish cheese and sausage samples, respectively, while it could not be detected in any of milk or hawawshy samples. *Salmonella* were detected in 2 (4%), 2 (4%) and 1 (2%) of kareish cheese, sausage and hawawshy samples, respectively, while they could not be recovered from the examined milk samples. The isolated serotypes from kareish cheese samples were *S.menden* and *S.allerton*, while two strains of *S.III arizonae* were isolated from sausage samples, but *S.anatum* was recovered from hawawshy samples. The public health significance of isolated strains as well as suggested control measures was discussed.

Salmonella and *E.coli* O157 are two of the most common agents of food borne illness in humans (Buzby *et al.*, 1996). *E.coli* O157 and *Salmonella* have been isolated from beef and dairy cattle at all stages of production, and although their shedding are intermittent and can be difficult to detect, these bacteria appear to be fairly widespread throughout the bovine population (Fedorka-Cray *et al.*, 1998; Hancock *et al.*, 1998; Elder *et al.*, 2000).

E.coli O157 was first recognized as food borne pathogen in 1982 (Riley *et al.*, 1983). Since that, it has become a pathogen of major concern in both food and dairy industries, because of its ability to cause severe illness, in particular, hemorrhagic colitis, hemolytic uremic syndrome and thrombotic thrombocytopenic purpura (Abdul-Raouf *et al.*, 1993).

Several outbreaks of *E.coli* O157 have been epidemiologically attributed to consumption of contaminated fresh and dry sausage and other meat products (Chinen *et al.*, 2001). Other foodstuffs including unpasteurized milk and dairy products manufactured from raw milk, however, have been implicated in many outbreaks, (Maher *et al.*, 2001).

Foodborne Salmonellosis continues to be a significant public health problem. Various meat products have been associated with outbreaks caused by *Salmonella*, *Shigella* and *E.coli* O157 (CDC, 2002). During slaughter, pathogenic

bacteria may contaminate the carcasses and subsequently be distributed via cut meat or meat materials intended for further processing into meat products (Borch and Arinder, 2002). In addition, human Salmonellosis has been recognized due to consumption of raw or improperly pasteurized milk and milk products (McEwen *et al.*, 1988). Although the heating kills the pathogens, the heated food may become recontaminated by food handlers or from cross contamination with vegetables salad during processing, marketing and in food service operations.

Therefore, this study was carried out to determine the prevalence of *E.coli* O157 and *Salmonella* in raw milk, kareish cheese, fresh sausage and hawawshy collected from farmer's houses, butcher's shops and retail markets in Beni-Suef Governorate as well as discussing the public health significance of the isolated microorganisms and suggestive control and preventive measures.

Materials and methods

Collection of samples. A total of 200 raw food samples including cow's milk, kareish cheese, fresh sausage and hawawshy (spiced minced meat) (50 samples of each) were randomly collected from farmer's houses, butcher's shops and retail markets in Beni-Suef governorate. All samples were identified and rapidly delivered to the laboratory in an insulated ice-box to be examined for presence of such pathogens.

Experimental techniques.

Table (1): Incidence of *E.coli* O157 in examined food samples

Samples	No. of samples	Positive samples	
		No.	%
Raw milk	50	0	0
Kareish cheese	50	1	2
Sausage	50	1	2
Hawawshy	50	0	0
Total	200	2	1

Table (2): Incidence of *Salmonella* in examined food samples

Samples	No. of samples	Positive samples		Serotype
		No.	%	
Raw milk	50	0	0	0
Kareish cheese	50	2	4	S.menden & S.allerton
Sausage	50	2	4	S.III arizonae (2 strains)
Hawawshy	50	1	2	S.anatum
Total	200	5	2.5	

Isolation of *Escherichia Coli* O157 (De-Boer and Heuvelink, 2000). Twenty five grams of each hard sample were separately homogenized with 225 ml of modified Tryptone soy broth supplemented by novobiocin (20 mg/ l) for 2 min. using sterile homogenizer(Universal Laboratory, Poland). Milk sample (25ml) was directly added to modified Tryptone soy broth supplemented by novobiocin. The inoculated broth was incubated at 37 °C for 24 h. A loopful from the incubated broth was streaked onto Tellurite- Cefixime Sorbitol Mac Conkey agar plate and incubated at 37 °C for 24h. Sorbitol negative colonies (colourless) were picked up and purified then examined biochemically and serologically.

Isolation of *Salmonella*. The applied technique was recommended by (Flowers *et al.*, 1992) as follows: Twenty five grams of each hard sample were homogenized in 225 ml of buffered peptone water (BPW), under aseptic conditions for 2 min. by using sterile homogenizer, while 25 ml of milk sample were thoroughly mixed with 225 ml of BPW. All samples were incubated at 35 °C for 24 ± 2 h. One hundred microliters from the pre-enriched sample was transferred to 10 ml of Rappaport Vassiliadis (RV) enrichment broth and incubated at 43 °C for 24h. Loopfuls from enriched RV broth were separately streaked onto each of xylose lysine desoxycholate (XLD) agar and *Salmonella*-*Shigella* (SS) agar plates and incubated at 37 °C for 24h. Two or three of typical or suspected colonies (colourless with black center on SS and

red colonies with black center on XLD) were selected from each selective medium and streaked onto nutrient agar slope which incubated at 37 °C for 24h., for further biochemical and serological identification.

Serological identification of *E.coli* O157 and *Salmonella*. Serological identification of the strains was carried out in the Clinical Microbiology Department, Central Health Laboratories of Ministry of Health and Population, Egypt.

Results and Discussion

Because the reservoir of *E.coli* is the intestinal tract of both man and animals, the presence of such organism in foods and water is used as indicator of fecal pollution either directly or indirectly (ICMSF 1978).

According to the data summarized in Table (1), *E.coli* O157 was detected in one sample (2%) of each kareish cheese and sausage samples. However, it could not be detected in raw milk and hawawshy samples.

Similar results of cheese samples were reported by Aksu *et al.*, 1999; Moustafa 2004, while Abd El-Hady, *et al.*, 1995; Gonul, 1997; El-Kosi, 2001, reported higher values. On the other hand Ansay and Kaspar (1997); Aman, *et al.*, (1998); Svoboda, *et al.*,(1998) ; Ibrahim and Sobeih, (2006) failed to isolate *E.coli* O157 from cheese samples.

The presence of *E.coli* O157 in kareish cheese could be attributed to the fact that it is usually made from raw skim milk, in addition to the primitive way of processing, handling and selling.

The lack of isolation of *E.coli O157* from raw milk is not surprising, since also Hancock *et al.*, (1994); Ansay and Kaspar, (1997); Aman *et al.*, (1998); Heuvelink *et al.*, (1998); Coia *et al.*, (2001) did not find *E.coli O157* in raw milk. On contrary, Padhye and Doyle (1991); Abd El-Hady *et al.*, (1995); Abdul-Raouf *et al.*, (1996); Abd El-Khalek, *et al.*, (2001); Sayed and Hussein (2003); Reuben, *et al.*, (2003); Amer and Soliman (2004); Hammad (2004); Moustafa, (2004); Picoozzi, *et al.*, (2005) could isolate the *E.coli O157* from milk at various percentages.

The failure in detection of *E.coli O157* in milk and hawawshy is mainly returned to isolation of *E.coli O157* is often difficult as it is present sporadically at very low levels among very high levels of competitor organisms (Siriken, *et al.*, 2006).

Nearly similar results of *E.coli O157* in sausage were reported by (Magwira, *et al.*, 2005). High figures were reported by (Chapman *et al.*, 2000; Chinen *et al.*, 2001). Low figures were reported by (Vernozy-Rozand, *et al.*, 1997; Dontorou *et al.*, 2003). On the other hand (Saleh, 2001; Kassem and Sabry 2003; Abd El-Rahman 2006; Soliman and El-Tabiy 2006) failed to isolate *E.coli O157* from sausage samples .

The presence of *E.coli O157* in sausage could be attributed to the contamination from faeces of infected animals as well as reflect the unsatisfactory hygienic measures during manufacturing and handling. This held the view reported by Suthienkul, *et al.*, (1990); Soliman and El-Tabiy (2006).

E.coli O157 is considered to be a typical food borne pathogen and responsible for several outbreaks. The majority of outbreaks have been commonly associated with the consumption of a variety of raw foods of animal origin such as undercooked ground beef, fresh and dry sausage, raw milk and milk products, vegetables salad and untreated water (Martin *et al.*, 1986; Tuttle and Gomez 1990; Griffin and Tauxe 1991; Chinen *et al.*, 2001; Licene *et al.*, 2001). Manifestation of illness caused by *E.coli O157* include non bloody diarrhea, hemorrhagic colitis, cystitis hemolytic uremic syndrome and thrombotic thrombocytopenic purpura due to production of shiga like toxin (verocytotoxin) (Wells *et al.*, 1991 ; Abdul-Raouf *et al.*, 1993).

It is evident from table (2) that the incidence of isolated *Salmonella* organism from kareish cheese , sausage and hawawshy samples were 2 (4%), 2 (4%) and 1 (2%), respectively.

Many studies have been conducted to determine the prevalence of *Salmonella* in fresh sausage. Similar result was recorded by Sadek, (1963). High prevalence were recorded by Weissman, and Carpenter,(1969); Abd El-Aziz, (1979); Tolba, (1986); Abd El-Aziz, (1987); Ahmed, (1988); Khalafalla, (1988); El-Mossalami, *et al.*, (1989); Fathi, *et al.*, (1994); Helmy, *et al.*, (1995); Abraham, *et al.*, (1998); Escartin, *et al.*, (1999) and Soliman, *et al.*, (2002). Low incidence was reported by Fathi and Thabet, (2001). On the other hand Lotfi, *et al.*, (1988); Mousa, *et al.*, (1993) ; Abd El-Atty (2005) failed to isolate *Salmonella* from sausage samples.

The presence of *Salmonella* in sausage may be attributed to the fact that this product is made from raw meat in addition to natural casing is often used in the manufacture which may be important source of *Salmonella* especially if proper hygienic measures is neglected (Escartin *et al.*, 1999).

No *Salmonella* could be detected in the examined milk samples (table, 2). These result was in acceptance to the findings reported by Richter, *et al.*, (2000); Eleftheriadou, *et al.*, (2002), while McEwen, *et al.*, (1988); Saudi and Moawad (1990); Jayarao and Henning, (2001); Abd El-All, *et al.*, (2005) could isolate *Salmonella* from raw milk with different values..

There is no much available data about the prevalence of *Salmonella* in kareish cheese, however, El-Kosi, (2001) reported low (3.33%) incidence of *Salmonella* in kareish cheese. On the other hand El-Bassiony (1977); El-Kholy (1989); Bahout and Moustafa (2006) failed to recover *Salmonella* from kareish cheese samples.

Kareish cheese is a popular Egyptian food due to its high nutritive value and cheap price, manufactured from raw skim milk and consumed fresh giving rise to concern that this product may be a threat to consumer health and safety as a result of presence of some pathogenic microorganisms especially *Salmonella* due to using raw milk in the production accompanied by improper sanitary practices during manufacturing, handling and selling.

This study detected that the prevalence of *Salmonella* in hawawshy was 2 % in contrast to those reported by El-Mossalami, (2006), who has reported that the hawawshy was free from *Salmonella*.

Concerning *Salmonella* serotypes, *S.menden* and *S.allerton* were isolated from two kareish

cheese samples, while *S.III arizonae* (2 strains) were isolated from sausage samples, but *S.anatum* was detected in hawawshy samples. It is generally accepted that the presence of any serotype of *Salmonellae* in a food should be regarded as a potential hazard for man (Fathi *et al.*, 1994).

In recent years, several outbreaks of Salmonellosis in humans have been recorded due to consumption of raw or improperly pasteurized milk or milk products, meat and meat products (McEwen *et al.*, 1988; Escartin *et al.*, 1999).

From the present data it can be concluded that the kareish cheese, sausage and hawawshy represents a potential hazard for consumers, due to the potential presence of *E.coli O157* and *Salmonella* as well as there is neglected sanitary measures adopted during manufacturing, handling and distribution of such fresh foods. Consequently, food manufacturers and specialists should design comprehensive programs as good manufacturing practices (GMP) and implementation of HACCP system to ensure the freedom of such foods from these pathogens. In addition, effective heat treatment for foods, provision of information to food handlers and consumers as well as application of strict hygienic measures during manufacturing, storage and selling of these products to improve its quality and safeguard the consumers against infections of such organisms. Due to limitation of data about the occurrence of *E.coli O157* and *Salmonella* in kareish cheese and hawawshy, further studies should be directed in this kind of foods.

References

- Abd El All, A. A.; Mohamed, K. H. F. and Dardir, H. A. (2005)** : Evaluation of a rapid improved PCR assay for detection of Salmonella spp. in raw milk and some dairy products. J. Egypt. Vet. Med. Associ., 65:1, 225-235.
- Abd El-Atty, N. S. (2005)**: Studies for improving the quality of locally manufactured sausage. Ph.D., Thesis, Fac. Vet. Med, Beni-Suef University, Egypt.
- Abd El-Aziz , A. A. (1979)**: Studies on hygienic quality of manufactured fresh sausage. M.V.Sc., Thesis, Alexandria University, Egypt.
- Abd El-Aziz , A. T. (1987)**: Microbiological load of some meat products as influenced by the hygienic status of the producing plant. M.V.Sc., Thesis, Fac. Vet. Med., Cairo University, Egypt.
- Abd El-Hady, H. M.; Halawa, M. A. and El-Shenawy, S. H. (1995)**: Surveillance of enterohemorrhagic *Escherichia coli* (*E.coli* O157 :H7) in milks and kareish cheese. Assiut Vet. Med. J., 33:66, 110-113.
- Abd El-Khalek, A.; El Gaml, A. and El Sherbini, M. (2001)**: Prevalence of *Escherichia coli* O157 in milk and faeces of dairy farm animals in Dakahlia province. 1st Cong. Food Hygiene and Human Health, 6-8 Feb. 2001, Fac. Vet. Med., Assiut, Egypt.
- Abd El-Rahman A.M.S. (2006)**: *Escherichia coli* O157 :H7 in formed meat products. M.V.Sc., Thesis, Fac. Vet. Med., Cairo University, Egypt.
- Abdul-Raouf, U.M.; Ammar, M.S. and Beuchat, L.R. (1996)**: Isolation of *E.coli* O157 :H7 from some Egyptian foods. Int. J. Food Microbiol., 29: 423-426.
- Abdul-Raouf, U.M.; Beuchat, L.R. and Ammar, M.S. (1993)**: Survival and growth of *Escherichia coli* O157 :H7 in ground, roasted beef as affected by pH, acidulants and temperature. Appl. Environ. Microbiol., 59:8, 2364-2368.
- Abraham, A ; Papa, A.; Soutos, N.; Ambrosiadiss, I. and Antoniaadis, A. (1998)**: Antibiotic resistance of Salmonella spp. and Listeria spp. isolated from traditionally made fresh sausage in Greece. J. Food Prot. 61, 1378
- Ahmed, S. (1988)** : Salmonellae in locally manufactured meat products. M.V.Sc., Thesis, Fac. Vet. Med., Cairo University, Egypt.
- Aksu, H., Arun, Ö. Ö.; Aydin, A., and Ugur, M. (1999)**: *Escherichia coli* O157 :H7 nin hayvansal kokenli gida maddelerinde varligi. Pendik Veteriner Mikrobiyoloji Dergisi, 30:2:77-81.
- Aman, I.M.; Knappstein, K. and Hann, G. (1998)**: Examination of verotoxin producing *Escherichia coli* in some Egyptian dairy products with special reference to serotype O157 :H7 . Michwissenschaft, 53:12, 676-679.
- Amer, A.A. and Soliman N. F. (2004)**: Prevalence of enterohemorrhagic *E.coli* O157:H7 in raw milk and effect of some chemical preservatives on its viability. Assiut Vet. Med. J., 50:102.
- Ansary, S.E. and Kaspar, C.W. (1997)**: Survey of retail cheeses, dairy processing environments and raw milk for *Escherichia coli* O157 :H7 . Lett. Appl. Microbiol., 25: 131-134.
- Bahout, A.A. and Moustafa, A.H. (2006)**: Occurrence of some microorganisms in relation to public health in kareish cheese . Assiut Vet. Med. J., 52:111.
- Borch, E. and Arinder, P. (2002)**: Bacteriological safety issues in red meat and ready to eat meat products, as well as control measures. Meat Sci., 62:3, 381-390.
- Buzby, J.C.; Roberts, T.; Jordan Lin, C.T. and MacDonald, J.M. (1996)**: Bacterial food-borne disease: medical costs and productivity losses. Washington, D.C., US department of agriculture, Economic research service. Report 741.
- CDC. (2002)**: US foodborne disease outbreaks, searchable data 1990-1995. (Online) from <http://www2a.cdc.gov/ncid-od/food-borne/outbreaksreport.asp>.
- Chapman, P.A.; Siddons, C.A.; Cerdan Malo, A.T. and Harkin, M.A. (2000)**: A one year study of *Escherichia coli* O157 in raw beef and lamb products. Epidemiol. Infect., 124, 207-213.
- Chinen, I.; Tanaro, J.D.; Milliwebsky, E.; Lound, L.H.; Chillemi, G.; Ledri, S.; Baschkier, A.; Scarpin, M.; Manfredi, E. and Rivas, M. (2001)**: Isolation and characterization of *Escherichia coli* O157 :H7 from retail meats in Argentina. J. Food Prot. 64: 1346-1351.
- Coia, J.E.; Johnston, Y.; Steers, N.J. and Hanson, M.F. (2001)**: A survey of the prevalence of *Escherichia coli* O157 in raw meats, raw cows milk and milk cheeses in South East Scotland. Int. J. Food Microbiol., 66: 63-69.
- De Boer, E. and Heuvelink, A.E. (2000)**: Methods for detection and isolation of shiga toxin producing *Escherichia coli*. J. Appl. Microbiol., 88: 133-143.
- Dontorou, C.; Papadopoulou, C.; Filioussis, G.; Economou, V.; Aspostolou, I.; Zakkas, G.; Saamouram, A.; Kansouzidou, A. and Levidiotou, A. (2003)**: Isolation

- of *Escherichia coli* O157:H7 from foods in Greece. Int. J. Food Microbiol., 82: 273-279.
- El-Bassiony, T. A. (1977)** : Incidence of pathogenic microorganisms and viability of some food poisoning bacteria in kareish cheese during storage. Ph.D., Thesis, Fac. Vet. Med., Assiut University, Egypt.
- Elder, R. O.; Keen, J. E.; Siragusa, G. R.; Barkocky-Gallagher, G. A.; Koohmaraie, M. and Laegreid, W. W. (2000)**: Correlation of enterohemorrhagic *E.coli* O157 prevalence in feces, hides and carcasses of beef cattle during processing. Proceedings of the National Academy of Science, 97:2999-3003.
- Eleftheriadou, M.; Varnava- Tello, A.; Metta-Loizidou, M.; Nikolaou, A.S. and Akkelidou, D. (2002)** : The microbiological profile of foods in the Republic of Cyprus: 1991-2000. Food Microbiol., 19:463-471.
- El-Kholy, A. M. (1989)**: Enterobacteriaceae in Egyptian soft cheese and their public health significance. Assiut Vet. Med. J., 21:42, 70-75.
- El-Kosi, O. H. R. (2001)**: Occurrence of some enteric pathogens and their indicators in some Egyptian raw milk products. Assiut Vet. Med. J., 45:89, 48-61.
- El-Mossalami E. I. K. (2003)** : Risk assessment of ready prepared meat products. Ph.D. Thesis, Fac. Vet. Med., Cairo University, Egypt.
- El-Mossalami, M. E.; Safwat, A. A. S.; Abd El-Rahim, L. and El-Sawah, H. (1989)**: Salmonellae in locally produced meat products. J. Egypt. Vet. Med. Assoc., 49:1-2, 99-108.
- Escartin, E. F.; Castillo, A.; Hinojosa-Puga, A. and Saldana-Lozano, J. (1999)**: Prevalence of Salmonella in chorizo and its survival under different storage temperature. Food Microbiol., 16:479-486.
- Fathi, S.; El-Khateib, T.; Moustafa, S. and Hassanein, K. (1994)**: Salmonellae and enteropathogenic *Escherichia coli* in some locally manufactured meat products. Assiut Vet. Med. J., 31:62, 190-199.
- Fathi, S. and Thabet, E. I. R. (2001)**: Incidence of Salmonella and *E.coli* in packed meat products sold in Assiut City. Assiut Vet. Med. J., 46:91, 187-199
- Fedorka-Cray, P. J.; Dargatz, D. A.; Thomas, L. A and Gray, J. T. (1998)**: Survey of Salmonella serotypes in feedlot cattle. J. Food Protect., 61:525-530.
- Flowers, R. S.; Andrews, W.; Donnelly, C. W. and Koening, E. (1992)**: Pathogens in milk and milk products. P. 106. In, Standard methods for the examination of dairy products. APHA, R.T. Marshall (ed.), Washington, D.C.
- Gonul, A. (1997)**: Cig sut ve peynir orneklerinde enterohemorajik *E.coli* :O157:H7 rastlanma sikligi. Kukem Dergisi, 20:2:69-73.
- Griffin, P. M. and Tauxe, V. (1991)**: The epidemiology of infections caused by *Escherichia coli* O157:H7 and other *E.coli* associated hemolytic uremic syndrome. Epidemiol. Rev., 13:60-98.
- Hammad, A. M. M. (2004)**: Microbiological studies on raw milk and some dairy products. M.V.Sc., Thesis, Fac. Vet. Med., Minufiya University.
- Hancock, D. D.; Besser, T. E.; Kinsel, M. L.; Tarr, P. I.; Rice, D. H. and Paros, M.G. (1994)**: The prevalence of *Escherichia coli* O157:H7 in dairy and beef cattle in Washington state. Epidemiol. and Infect., 113: 199-207.
- Hancock, D. D.; Besser, T. E. and Rice, D. H. (1998)** : Ecology of *E.coli* O157: H₇ in cattle and impact of management practices. In *E.coli* O157:H7 and other shiga toxin producing *E.coli* strains ed. Kaper, J.B and O'Brien, A.D. pp. 85-91, Washington D.C.: American society of microbiology press.
- Helmy, N.A. (1995)**: Salmonellae in meat and poultry products using immunofluorescent technique. M.V.Sc., Thesis, Fac. Vet. Med. Cairo University.
- Heuvelink, A.E.; Bleumink, B.; Van den Biggelaar F.L.; Te Giffel, M.C.; Beumer, R.R. and De Boer, E., (1998)**: Occurrence and survival of verocytotoxin-producing *Escherichia coli* O157 in raw cow's milk in the Netherlands. J. Food Prot., 61: 1597-1601.
- Ibrahim E.M.A. and Sobeih A.M.K. (2006)**: Diarrhoeagenic *E.coli* in kareish cheese manufactured by different methods with special reference to *Escherichia coli* O157:H7. Minufiya Vet. J., 4:1, 169-178.
- ICMSF (1978)**: Microorganisms in foods, their significance and methods of enumeration. 2nd ed. Uni. of Toronto Press, Toronto.
- Jayarao, B.M. and Henning, D.R. (2001)**: Prevalence of foodborne pathogens in bulk tank milk. J. Dairy Sci., 84: 2157-2162.
- Kassem G.M., and Sabry M.A. (2003)**: Incidence of enteropathogenic *E.coli* in meat products with special reference to *Escherichia coli* O157. J. Egypt. Vet. Med. Assoc., 63:5, 61-71.
- Khalafalla, F.A. (1988)**: Sanitary status of meat, meat products and fish in Beni-Suef Governorate. Ph.D.. Thesis, Fac. Vet. Med., Cairo University, Egypt.
- Licence, K.; Oates, K.R.; Syngé, B.A. and Reid, T.M.S. (2001)** : An outbreak of *E.coli* O157:H7 infections with evidence of spread from animals to man through contamination of private water supply. Epidemiol. Infect., 126:135-138.
- Lotfi, A.Y.; Youssef, H.; EL-Khatieb, T.; Seddik, I.; El-Timmawy, A. and Ali, G. (1988)** : Enterobacteriaceae in meat products in Upper Egypt. Assiut Vet. Med. J., 17:77.
- Magwira, C.A.; Gashe, B.A. and Collison, E.K., (2005)**: Prevalence and antibiotic resistance profiles of *Escherichia coli* O157:H7 in beef products from retail outlets in Gaborone, Botswana. J. Food Prot., 68:2, 403-406.
- Maher, M.M.; Jordan, K.N.; Upton, M.E. and Coffey, A. (2001)**: Growth and survival of *E.coli* O157:H7 during the manufacture and ripening of a smear ripened cheese produced from raw milk. J. Appl. Microbiol., 90: 201-207.
- Martin, M.L.; Shipman, L.D.; Wells, J.G.; Potter, M.E.; Hedberg, K.; Wachsmuth, I.K.; Tauxe, R.V.; Davis, J.P.; Arnolai, J. and Tilleli, J. (1986)**: Isolation of *Escherichia coli* O157:H7 from dairy cattle associated with two cases of hemolytic uremic syndrome. Lancet, 1:2, 8414: 1043.
- McEwen, S.A.; McClure, L.H. and Martin, S.W. (1988)**: Farm inspection scores and milk quality criteria as indices of Salmonella in bulk milk. J. Food Prot. 51:12, 958-962.
- Mousa, M.M.; Awad, H.A.; Yassien, N.M. and Gouda, H.I. (1993)**: Microbial quality of some meat products. Vet. Med. J (Giza), 41:3, 59-62.
- Moustafa, M.F.H. (2004)**: Studies on *Escherichia coli* O157:H7 in milk and some milk products. Ph.D., Thesis, Fac. Vet. Med., Assiut University, Egypt.
- Padhye, N.V. and Doyle, M.P. (1991)**: Rapid procedure for detecting enterohemorrhagic *Escherichia coli* O157:H7 in food. Appl. Environ. Microbiol., 57: 1693-2698.
- Picozzi, C.; Foschino, R.; Heuvelink, A. and Beumer, R. (2005)**: Phenotypic and genotypic characterization of sorbitol negative or slow fermenting (Suspected O157) *Escherichia coli* isolated from milk samples in Lombardy region. Letters Applied Microbiol., 40: 491-496.
- Reuben, A.; Treminio, H. and Chaves, C. (2003)**: Presence of *Escherichia coli* O157:H7, *L. monocytogenes* and *Salmonella* spp. In food from animal origin in Costa

Rica. Arch Latinoam Nutr., 53:4, 389-392.

Richter, J.; Becker, H. and Martlbauer, E. (2000): Improvement in Salmonella detection in milk and some dairy products: comparison between the ISO method and the Oxoid sprint Salmonella test. Letters in Appl. Microbiol., 31: 443-448.

Riley, L. W.; Remis, R. S.; Helgerson, S. D.; McGee, H. B.; Wells, J. G.; Davis, B. R.; Hebert, R. J.; Olcott, E. S.; Johnson, L. M.; Hargrett, N. T.; Blake, P. A. and Cohen, M. L. (1983): Hemorrhagic colitis associated with a rare *Escherichia coli* serotype. N. Engl. J. Med., 308: 681-685.

Sadek, M. I. (1963): Studies on locally manufactured fresh sausage. M.V.Sc., Thesis, Fac. Vet. Med., Cairo University.

Saleh, S. K. (2001): Prevalence of enterohemorrhagic *E.coli* in some meat products. J. Egypt. Vet. Med. Assoc., 61:4, 173-178.

Saudi, A. M. M. and Moawad, A. A. (1990): Incidence of Enterobacteriaceae in market milk in Cairo and suburbs. Assiut Vet. Med. J., 24:47, 134139

Sayed A. S. M. and Hussein A. A. A. (2003): Occurrence of *E.coli* O157:H7 in apparently healthy dairy cattle and retail milk. Assiut Vet. Med. J., 49:97.

Siriken, B.; Pamuk, S.; Ozakin, C.; Gedikoglu, S. and Eyigor, M. (2006): A note on the incidences of *Salmonella* spp., *Listeria* spp. and *Escherichia coli* O157: H7 serotypes in Turkish sausage (Soudjouck). Meat Science, 72: 177-181.

Soliman, M. R.; Abd El-Monem, K. H. and Saad, S. M. (2002): Microbiological quality of ready to eat meat products and fishes in urban and rural areas. J. Egypt. Vet. Med. Assoc., 62:39

Soliman Z. I. and El-Tabiy A. A. (2006): A study on the occurrence of *E.coli* in some beef products with special

reference to *E.coli* O157 :H7. Assiut Vet. Med. J., 52:110, 75-87.

Suthienkul, O.; Brown, J. E.; Seriwatana, J and Escheverria, P. (1990): Shiga-like toxin producing *E.coli* in retail meats and cattle in Thailand. Appl. Environ. Microbiol., 56:1135-1139.

Svoboda, P.; Gautsch, S., and Luthi, T. M. (1998): Investigation on the occurrence of verotoxin producing *Escherichia coli* in ready to eat and non ready to food. Mitteilungen aus dem gebiete der lebensmitteluntersuhung und hygiene, 89:6, 758-774.

Tolba, K. S. (1986): Antibiotic resistant microorganisms in some meat products. M.V.Sc., Thesis, Fac. Vet. Med., Cairo University, Egypt.

Tuttle, J. and Gomez, T. (1990): Lessons from a large outbreak of *Escherichia coli* O157 :H7 infections insights into the infectious dose and methods of wide spread contamination of hamburger patties. Epidemiol. Infect., 122: 185-192.

Vernozy-Rozand, C.; Mazuy C.; Ray- Gueniot, S.; Boutrand-Loi,S.; Meyrand, A. and Richard, Y. (1997) : Detection of *Escherichia coli* O157 in French food using an immunomagnetic separation and the VIDAS *Escherichia coli* O157 . Letter in Appl. Microbiol., 25: 422-446.

Weissman, M. A. and Carpenter, J. A.(1969): Incidence of salmonellae in meat and meat products. J. Appl. Microbiol. 17, 899.

Wells, J. G.; Shipman, L. D.; Greene, K. D.; Sowers, E. G.; Green, J. H.; Cameron, D. N.; Downes, F. P. Martin, M.L. et al., (1991): Isolation of *Escherichia coli* O157 :H7 and other shiga like toxin producing *E.coli* from dairy cattle. J. Clin. Microbiol., 29: 985-989.

مدى تواجد ميكروب السالمونيلا والإيشيريشيا كولاي O157 في بعض الاغذية

اجريت هذه الدراسة على ٢٠٠ عينة من الاغذية الخام لتقرير ما بها من السالمونيلا والإيشيريشيا كولاي O157 . شملت الدراسة فحص كلا من اللبن الخام والجبن القريش والسجق الطازج والحواشى بواقع ٥٠ عينة من كل منتج. امكن عزل ميكروب الإيشيريشيا كولاي O157 من عينة واحدة من كلا من الجبن القريش والسجق الطازج بنسبة ٢ % لكل منهما بينما لم يتم عزلها من اللبن الخام والحواشى. اما ميكروب السالمونيلا فقد تم عزله من عينتين من الجبن القريش و عينتين من السجق الطازج و عينة حواشى بنسبة ٤ % ، ٢ % و ٢ % على التوالي بينما لم يتم عزله من اللبن الخام. تم عزل عترتى السالمونيلا مندن والسالمونيلا الرتاون من عينتى الجبن القريش وعتره السالمونيلا III أريزونا من عينتى السجق وعتره السالمونيلا أنيتم من عينة الحواشى. هذا وقد نوقشت الاهمية الصحية والاقتصادية للنتائج واقترح مايجب اتباعة من اجراءات لضمان حماية صحة المستهلك.