

## SEQUENTIAL USAGE OF BACTERIAL SUPERNATANTS IN MICROPROPAGATION OF DATE PALM

HALA M.A. FARRAG<sup>1</sup>, WEDAD E.E. EWEDA<sup>2</sup>, A.A. GUIRGIS<sup>3</sup>, A. M. IBRAHIM<sup>3</sup>,  
A.A. EL-BANNA<sup>1</sup> AND I.A. IBRAHIM<sup>3</sup>

1 Central Laboratory For Date Palm Researches and Development, ARC, Giza

2 Department of Agricultural Microbiology, Faculty of Agriculture, Ain Shams University.

3 Genetic Engineering and Biotechnology Research Institute (GEBRI), Menofya University.

(Manuscript received 28 Nov. 2007)

### Abstract

The present investigation was aimed to use phytohormones produced by some bacterial strains like *Azospirillum brasilense*, *Azotobacter chroococcum*, *Bacillus megaterium* and *Klebsiella pneumoniae* instead artificial hormones as a protocol in date palm micropropagation through somatic embryogenesis induced from shoot tip or leaf primordial explants, which were isolated from offshoot apex. In order to test the possibility to use supernatant of the strains, which are auxin over producers and/or cytokinin over producers in culture medium composition according to development stages for maximizing yield and to study the effect on each stage, i.e., callus initiation, embryo formation, shooting and rooting stages. In general, results showed that *Klebsiella* and *Azotobacter* have an auxin effect but *Azospirillum* and *Bacillus* were more related to cytokinin nature.

### INTRODUCTION

Several soil microorganisms are capable of producing physiologically active quantities of plant growth regulators (**PGRs**), which may have a pronounced effect on plant growth and development. A diverse array of bacteria, including species of *Azospirillum*, *Azotobacter*, *Bacillus* and *Klebsiella* have mechanisms which these rhizobacteria enhance plant growth through producing substances such as auxins, GA3 and cytokinins ( phytohormones) and mineralize organic phosphorus and enhance minerals uptake (Grayston *et al.* 1990 and Joo *et al.* 2004). So that, using these natural substances *in vitro* in culturing economic plants become necessary to overcome increasing artificial substances and that is of variable dangerous on healthy.

It is known that, the date palm is on one of the most important fruit crop in the Middle East and Arabian lands (FAO, 1984 and Moursy and Saker, 1996). Date palm is a monocot and dioecious fruit tree, which are difficult to study due to their long life. Therefore, *in vitro* techniques appear to be promising tools to study palm growth and development compared to field and greenhouse experimentation (Tisserat, 1983).

The concentration and type of hormones especially, auxins and cytokinins are considered the most critical factors in stimulating callus production and subsequent embryogenesis (Reynolds and Murashige, 1979 and Tisserat, 1979).

Generally, cytokinins are required for inducing cell division, callus formation, cell extension, shoot proliferation and shoot morphogenesis (Pan and Van Staden, 1998) Naphthaleneacetic acid (NAA) and kinetin at levels from 1-10 mg/L induced callus growth (Khan *et al.*, 1983). An embryo can be defined as the earliest recognizable multicellular stage of an individual that occurs before it has developed the structures or organs characteristic of a given species (Trigiano and Gray, 2000). Somatic embryogenesis is a process by which somatic (non-gametic) cells undergo differentiation to form a bipolar structure containing both root and shoot axes. These somatic embryos are similar to zygotic embryos and can mature and germinate (Smith, 2000).

Abdel-Hamid *et al.* (2001) found that in date palm micropropagation, the highest rate of shoot multiplication induced from shoot tip explants was obtained through callus cultured on medium containing 3 mg/L 2iP.

## MATERIALS AND METHODS

This work was carried out in the Central Laboratory for Research and Development of Date Palm, Agricultural Research Center (ARC), Ministry of Agriculture, Giza, Egypt and Plant Biotechnology Department, Genetic Engineering and Biotechnology Research Institute (GEBRI), Minufiya University during the period 2003-2006.

### **Source of bacterial strains:**

Representative bacterial strains used throughout this study were *Klebsiella pneumoniae* and *Azospirillum brasilense*, which were isolated through a previous study carried out at the Environmental Studies and Research Unit (ESRU) Department of Agricultural Microbiology, Faculty of Agriculture, Cairo University, Giza by Farrag (2000). The other two strains *Bacillus megaterium* and *Azotobacter chroococcum* were kindly obtained from the Culture Collection of Microbiological Resource Center, Faculty of Agriculture, Ain Shams University, Cairo, Egypt.

### **Purification and maintenance of the isolates:**

The selected strains were purified by streaking on agar plates containing the proper selective medium for each bacterium type as the following:

- I. Nitrogen deficient medium (NFD) for *Klebsiella* (Dixon *et al.*, 1980)
- II. Semi-solid malate medium for *Azospirillum* (Dobereiner and Day 1976)
- III. Nitrogen – deficient medium of Ashby for *Azotobacter* (Hegazi and Niemela, 1976)

#### IV. *Bacillus* medium (Allen, 1959).

The previous selected strains were surface inoculated with single strain and incubated at  $28 \pm 2^\circ\text{C}$  for 48hr to 7 days (5-7 times or more) until pure colonies were obtained.

#### **Plant material:**

*Phoenix dactylifera* cv. Sewi offshoots of 2-3 years old, 6-9 Kg in weight, about 1-1.5 m in length and 20-30 cm in diameter were selected from date palm that were grown in El- Wahaat El- Baharia in Egypt. The selected offshoots were carefully separated from the mother trees.

#### **Isolation and sterilization of explants:**

The selected young offshoots were immediately transferred to the laboratory. Two types of explants were separated from the offshoots, i.e., shoot tip and leaf primordial. For excising both types of explants, the green leaves of the offshoots were carefully removed by using sharp knife. Leaves separation were carried out starting from the base and acropetaly continued upwards until the white soft leaves, surrounding the shoot tip appeared. However, cork and other remaining tissues of the offshoot were discarded.

The plant materials including the shoot tip region (10 – 12 cm in length and 5 cm in diameter) were sterilized as ( Abdel Sattar, 2005).

The shoot tip regions including the shoot apex plus the associated primordial leaves were used. Shoot apical meristem plus few primordial leaves (4-5 leaves) were excised together, and would be considered as shoot tip explant. The shoot tip explants was about 1 cm in length and 0.5-0.8 cm in diameter. The shoot tip was divided longitudinally with a sharp cutter into 4 equal quarters. The leaf primordial explants were taken from the rest primordial leaves surrounding the meristem tip.

#### **Culture medium:**

The MS basal medium containing macro and microelements as well as vitamins ( Murashige and Skoog , 1962) was used in this study.

The MS basal medium was supplemented with 200 mg/L glutamine 100 mg/L myo- inositol, 3% sucrose and 0.6% agar. Medium was adjusted to  $\text{pH } 5.8 \pm 0.1$ .

The medium was distributed into the culture jars (250 mL), each jar contains 25 ml of MS medium. Jars were capped with polypropylene closure and autoclaved at  $121^\circ\text{C}$  and  $1.1 \text{ kg/cm}^2$  for 20 minutes. The jars were transferred to the culture cabinet to get cool. The different explants were cultured into MS medium supplemented with bacterial supernatants to obtain cultured medium containing all components as natural medium to study its effect on the yield of each studied stage.

**Callus initiation stage:**

Twenty-five offshoots were considered as a source of explants. The control medium was MS culture medium 10 mg/L, 2,4-D 3 mg/L 2iP, 30 g/L sucrose and all B. Ss (bacterial supernatants) were added to the MS medium without growth regulators at 150 mL/L concentration as recommended by Farrag (2000) in treatments A, a, B, b, C, c and D, d as the following:

A) *Azotobacter chroococcum* + tryptophane (as a producer for 41 µg/mL GA3 → 6.2 mL/150 mL),

a) *Azoto.* Cultured in free trypto. Medium (non significant production),

B) *Klebsiella pneumoniae* + tryptophane (IAA over producer 47 µg/mL → 7 mL/150 mL, also GA3 66 µg/mL → 10 mL/150 mL),

b) *Kleb.* Cultured in free trypto. Medium (non significant production),

C) *Azospirillum brasilense* + tryptophane ( cytokinin over producer 5.4 µg/mL → 0.81 mL/150 mL ),

c) *Azosp.* Cultured in free trypto. medium (non significant production),

D) *Bacillus megaterium* + tryptophane (non significant production),

d) *B. megaterium* Cultured in free trypto. medium ( cytokinin producer 5 µg/mL → 0.75 mL/150 mL),

Those bacterial strains ascertained in the considered treatments were chosen according to their ability to produce phytohormones from previous studies of Farrag *et. al.* (2007). According to Zaid and Tisserat (1983) cultures were transferred to fresh media every 6 weeks, and incubated in complete darkness for 24 weeks.

**The data recorded as follows:**

1. Callus formation degree.
2. Browning degree.
3. Swelling degree at the end of each subculture (4 subculture).

The compact callus, which formed from the previous culture were placed on media containing 10 mg/L 2,4-D and 3 mg/L 2iP to produced friable callus (control). Data were calculated visually as scores (Pottino, 1981) from 1- 4 as the following:

Negative results (-)	=1
Below average results (+)	=2
Average results (++)	= 3
Good results (+++)	= 4

**Embryogenic Callus Formation:**

Friable callus was transferred to the following media:

MS + free growth regulators (control) was (0).

MS + 150 mL/L B.Ss. A, a, B, b, C, c and D, d.

For embryogenesis, each treatment consisted of 3 replicates and each replicate contains about 1x1 cm friable callus. Cultures were incubated in total darkness for 3 months. Data were recorded as growth vigor and number of embryo.

#### **Shoot formation stage:**

Individual embryos were separated from previous stage and cultured on the same media. The data of growth and development were recorded as:

1. Number of shoots / plant.
2. Shoot length (cm).
3. Growth vigor.

#### **Root formation stage:**

Individual shoots were used as explants in this stage and they were cultured in MS supplemented with NAA 0.1 mg/L control and the same treatments. All culture media of each treatment were distributed into culture tubes. The culture tubes were generally placed in light condition for 16 h illumination of 3000 lux light intensity (white fluorescent tubes) at  $27 \pm 2^\circ\text{C}$ .

The data of this stage were recorded as:

1. Number of roots/ plant.
2. Root length (cm).
3. Growth vigor.

#### **Statistical analysis:**

Experiments were arranged in factorial complete design and data were statistically analyzed according to method described by Snedecor and Cochran (1972).

## **RESULTS AND DISCUSSION**

#### **callus initiation:**

Data presented in Table (1) showed callus formation, browning and swelling degree as affected by adding bacterial supernatants. Results revealed that the highest score value of callus formation degree was observed with *Klebsiella* supernatant in the presence of tryptophan (B) that is ( $\tau, \delta$ ) followed by *Azotobacter* supernatant(A), *Azospirillum* supernatant(C) with tryptophan and control (0), which recorded the same score (3). These treatments were the best in the swelling (3.25) except for the control (2.25). On the other hand, treatment (C) was the lowest in the browning degree (0.5), in the opposite, the highest value of browning degree was (2.5) in (A) and (D) treatments but the lowest values of callus degree was in *Azospirillum* supernatant free tryptophan (c) and *Bacillus* supernatant free tryptophan (d) which their score were (1) . In this respect, Chung and Tzeng (2004) showed that the production of IAA

in culture media was solely dependent on the presence of tryptophan. Moreover, One *et al.* (2005) stated that varying the concentration of tryptophan in batch experiments has an effect on both growth of *Azospirillum* and IAA synthesis. This results are in agreement with the obtained results which, in the present study, which gave positive effect by using bacterial supernatants from medium containing tryptophan as precursor for production of auxin.

Table 1. Effect of bacterial supernatant on callus formation of date palm (*Phoenix dactylifera cv.Sewi*) after 18 weeks from culturing (three subculture).

Treatments	Callus Formation	Browning	Swelling
<i>Azotobacter chroococcum</i> (A)	3.0	2.5	3.25
<i>Azoto</i> . free tryptophan (a)	2.0	1.5	2.0
<i>Klebsiella pneumoniae</i> (B)	3.5	1.0	3.25
<i>Klebsiella</i> free tryptophan (b)	1.5	1.0	1.25
<i>Azospirillum brasilense</i> (C)	3.0	0.5	3.0
<i>Azosp.</i> free tryptophan (c)	1.0	1.0	2.0
<i>Bacillus megaterium</i> (D)	2.5	2.5	3.25
<i>Bacillus</i> free tryptophan (d)	1.0	2.0	2.25
Control	3.0	2.0	2.75
L.S.D. at 5%	1.2	0.12	0.93

Moreover, Reuveni and Kipnis (1974) showed that auxins were found to be the most critical component in the media for callus production and Tisserat (1982) mentioned that (MS) medium supplemented with high level of auxin resulted in high percentage of callus. In this concern, Brackpool (1988) reported that in date palms micropropagation, high concentrations of the artificial auxins naphthaleneacetic acid (NAA) and 2,4-dichlorophenoxyacetic acid (2,4-D) are needed to induce callus formation. While (Trigiano and Gray 2000) reported that auxins play a role in many development processes, including cell elongation, swelling of tissue, apical dominance, adventitious root formation and somatic embryogenesis.

#### **Embryogenesis:**

The effect of bacterial supernatants on callus development is shown in Table (2). Results revealed that the greatest number of individual embryos was (30) in MS medium supplemented with *Azospirillum* supernatant with tryptophan (C) followed by (26) formed by using *Bacillus* supernatant with tryptophan (D) and (25) in *Bacillus* supernatant from medium free tryptophan (d) this results may due to the presence of cytokinin in this supernatants. In addition, the best growth vigor was due to (C) and

(d) which recorded a score (4) that may be owed to presence other substances like gibberellins and auxins. These results are in harmony with Gonzalez - Lopez *et al.*, (1986) who observed that the culture supernatant of *A. chroococcum* & *A. vinelandii* contained at least three gibberellins-like substances. Mahmoud *et al.* (1984) investigated the ability of different isolates belonging to genera *Bacillus* and *Azotobacter* to synthesize plant growth substances. The supernatant of the bacterial cultures showed the presence of a compound, which correspond to indole acetic acid together with unidentified derivatives. These organisms also secreted gibberellins.

Table 2 . Effect of bacterial supernatant added to MS medium on callus development and differentiation of date palm(*Phoenix dactylifera* cv.Sewi) after 18 weeks from culturing

Treatments	Growth vigor	Number of embryo
<i>Azotobacter chroococcum</i> (A)	3.0	15
<i>Azoto.</i> free tryptophan (a)	2.5	10
<i>Klebsiella pneumoniae</i> (B)	3.5	20
<i>Klebsiella</i> free tryptophan (b)	1.5	6.0
<i>Azospirillum brasilense</i> (C)	4.0	30
<i>Azosp.</i> free tryptophan (c)	3.0	17
<i>Bacillus megaterium</i> (D)	3.5	26
<i>Bacillus</i> free tryptophan (d)	4.0	25
Control	3.0	18
L.S.D. at 5%	1.01	4.4

At the same time , Omar (1988) investigated that ovular callus was noticed after 6 weeks on a medium containing 10 mg/L 2,4-D and 2 mg /L kinetin. While, Sharon *et al.* (1998) mentioned that white friable callus was initiated from leaf primordia of date palm cv. Yakubi cultured on modified MS medium supplemented with 50 mg/L 2,4-D, 1 mg/L kinetin and 0.5 mg/L 2iP and Ibrahim (1999) showed that, embryogenic callus was obtained from shoot tips and leaf primordial explants cultured on modified MS medium containing either 10 mg/l NOA, 5 mg/L 2,4-D, 1 mg/L kinetin and 1mg/L 2iP or 100 mg/L 2,4-D, 3 mg/L kinetin and 3 mg/L 2iP.

### **Shoot formation:**

Results in Table (3) showed that best growth vigor was in *Klebsiella* supernatant in the presence of tryptophan (B) that is (4) and the highest shoot was (7 cm) in the same treatment followed by *Azotobacter* supernatant with tryptophan (A), which gave growth vigor (3.5) and shoot length (6. 5 cm). The largest number of shoot/ plant was (5) due to culturing in MS medium containing *Azospirillum*

supernatant with tryptophan (C) these results are in agreement with those found by Kapulink *et al.* (1985) who found that *Azospirillum* has the ability to produce plant growth promoting substances such as auxins and cytokinins. El- Khawas *et al.* (1996) found that *Azospirillum* and other groups of PGPR were produce and release a broad spectra of plant growth regulators (auxins, several gibberellins and cytokinins). Garcia de Salamone *et al.* (2001) stated that five plant growth promoting rhizobacteria (PGPR) strains produced the cytokinin dihydrozeatin riboside (DHZR) in pure culture. In addition to, Saker *et al.* (1998) demonstrated that in *in vitro* tissue culture of date palm, only 2iP containing media was found to be satisfactory for shoot induction after a phase of callus formation. The highest percentage of shoot proliferation occurred when embryos were cultured on MS medium supplemented with 3 mg/L 2iP, 0.1 mg/L NAA and 3 g/L charcoal.

Table 3. Effect of bacterial supernatant added to MS medium on shoot formation of date palm (*Phoenix dactylifera cv.Sewi*) after 18 weeks from culturing

Treatments	Growth vigor	Shoot number	Shoot length
<i>Azotobacter chroococcum</i> (A)	3.5	1.0	6.5
<i>Azoto.</i> free tryptophan (a)	1.0	2.0	4.5
<i>Klebsiella pneumoniae</i> (B)	4.0	2.0	7.0
<i>Klebsiella</i> free tryptophan (b)	2.0	1.0	3.5
<i>Azospirillum brasilense</i> (C)	2.5	5.0	5.5
<i>Azosp.</i> free tryptophan (c)	1.5	2.0	4.0
<i>Bacillus megaterium</i> (D)	2.5	3.0	6.0
<i>Bacillus</i> free tryptophan (d)	3.0	4.0	4.0
Control	3.0	3.0	4.5
L.S.D. at 5%	0.95	1.35	1.68

Moreover Bekheet and Saker (1998) reported that in date palm micropropagation, the presence of cytokinins such as BA and 2iP in the culture media was considered as an important factor for bud differentiation. Shoot induction was confined to media containing 4 mg/L BA, 4 mg/L 2iP and 0.5 mg/L NAA. Also, Zaid (2003) showed that in date palm micropropagation the culture medium supplemented with 0.5 mg/l kinetine was more effective than 0.5 mg/l 2iP on stimulating the produced shoots number and shoot length.



**Rooting formation:**

Results in Table (4) revealed that, significant differences were observed among treatments, while *Klebsiella* supernatant in the presence of tryptophan (B) gave the best growth vigor value (4), the greatest root number (6)/ plant and the tallest root (6.5cm) followed by *Azotobacter* supernatant with tryptophan (A), which recorded growth vigor value (3.5), root number (4)/ plant and root length (5cm). These results and all the previous ones are in the same line with Hafeez *et al.* (2004) who conducted different experiments to determine the growth promotion activity of various rhizobacteria. The results showed that the root growth was depending on the ability of *Azotobacter* strains for IAA production.

Table 4. Effect of bacterial supernatant added to MSmedium on root formation of date palm (*Phoenix dactylifera cv.Sewi*) after 18 weeks form culturing

Treatments	Growth vigor	Root number	Root length
<i>Azotobacter chroococcum</i> (A)	3.5	4.0	5.0
<i>Azoto.</i> free tryptophan (a)	1.5	2.0	3.5
<i>Klebsiella pneumoniae</i> (B)	4.0	6.0	6.5
<i>Klebsiella</i> free tryptophan (b)	2.5	1.0	2.0
<i>Azospirillum brasilense</i> (C)	3.5	3.0	4.0
<i>Azosp.</i> free tryptophan (c)	3.0	1.0	2.5
<i>Bacillus megaterium</i> (D)	1.0	3.0	4.5
<i>Bacillus</i> free tryptophan (d)	3.0	1.0	2.0
Control	3.0	3.0	4.0
L.S.D. at 5%	0.57	1.16	0.84

Recently, Abd El- Kader(2007) cited that an isolate belongs to genus *Klebsiella* was IAA over producers. While, (Thorpe, 1981) reported that Auxin alone and\ or with a very low concentration of cytokinin is important for the induction of root primordial in plant tissue culture. Bekheet and Saker (1998) reported that in date palm tissue culture, rooting of proliferated shoots was achieved upon supplementation of MS culture medium with 1 mg/L NAA. Moreover, Gadalla (2003) demonstrated that in mass propagation of dry cultivars of date palm, the highest significant value of rooting percentage was observed when MS culture medium was supplemented with 3 mg/L of either NAA or IBA.

## REFERENCES

1. Abd-El-Hamid, M.A., M. H. A. Abou-Bakr, I. A. Ibrahim and M. A. Abd-El-Baky. 2001. Some aspects of *in vitro* micropropagation of date palm *Phoenix dactylifera* L.. J. Agric. Sci. Mansoura Univ., 26 (9): 5449 - 5466.
2. Abd El-Kader, E.M. 2007. Microbial secretions and their applications in agriculture. Ph. D. Thesis, Microbiol., Dept., Fac. of Agric., Cairo Univ., Egypt.
3. Abdel Sattar, M. M. 2005. Advanced physiological studies on date palm. Ph.D. Thesis, Plant Botany Dep., Fac. of Agric., Cairo Univ., Egypt.
4. Allen, O. M. 1959. Experiments in soil bacteriology. 1<sup>st</sup> Ed Burgss publishing Co. Minneapolis, Minnesota. USA
5. Bekheet, S. A. and M. M. Saker. 1998. *In vitro* propagation of Egyptian date palm: Direct and indirect shoot proliferation from shoot tip explants of *Phoenix dactylifera* L. Zaghlool. The First International Conference on Date Palms, In Egypt, 150- 155.
6. Brackpool, A. 1988. Commercial propagation of date palm *in vitro*. Plant Today, May / June, 82 - 85.
7. Chung, K. R. and D. D. Tzeng. 2004. Biosynthesis of IAA by the gall- inducing fungus *Ustilago esculenta*. J. Biol. Sci., 4(6): 744- 450.
8. Dixon, R.A., R. R. Eady, G. Espin, S. Hill, M. Laccarino, D. Khan and M. Merrich. 1980. Analysis of regulation of *Klebsiella pneumoniae* nitrogen fixation (*nif.*) gene cluster with gene. Nature. 286:128- 132.
9. Dobereiner, J. and J.M. Day. 1976. Associative symbioses in tropical grasses: Characterization of microorganisms and dinitrogen fixing sites. *In* Proceeding of the First International Symposium on Nitrogen Fixation. Vol.2. Edited by Newton, W.E. and C.J. Nyman, Washington State, University press, Pullman, USA., pp 518 - 538.
10. El- Khawas, H.M., I.A. Ibrahim, H.M. Anwar and N.A. Hegazi. 1996. *In vitro* propagation using phytohormones produced by associative diazotrophs. In : Malik K.(ed.) Abstracts, 7<sup>th</sup> International Symposium on Nitrogen Fixation with Non-Legumes, Faisalabad, Pakistan, October 16-21, printed at: Sheikh Ghulam Ali(PVT) Ltd., Lahore, Pakistan, P 67.
11. Farrag, Hala M. A. 2000. Plant hormones produced by microorganisms and tissue culture applications. M.Sc. Thesis, Microbiology Dept., Fac. of Agric., Cairo Univ., Egypt.
12. Farrag, Hala M.A., Eweda, E. E. Wedad, A.A. Guirgis, A. M. Ibrahim, A.A.El-Banna and I.A. Ibrahim 2007. Chemical and Molecular studies on some phytohormones producers bacteria(In press).
13. FAO 1984. *FAO Production year Book*. Food and Agriculture Organization of the United Nation, Rome, Italy.

14. Gadalla, E. G. 2003. Propagation of dry varieties of date palm. Ph.D. Thesis. Faculty of Agriculture, Cairo University. Egypt, pp. 109.
15. Garcia de Salamone, I. E. Hynes and L.M. Nelson. 2001. Cytokinin production by plant growth promoting rhizobacteria and selected mutants. *Can. J. Microbiol.*, 47:404- 411.
16. Gonzalez –Lopez, J., V.Salmeron, M.V.Martinze- Toledo, F. Ballesteros and A. Ramos- Gormenzana. 1986. Production of auxins, gibberellins and cytokinins by *Azotobacter vinelandii* ATCC 12837 in chemically defined media and dialyzed soil media. *Soil Biol. Biochem.*, 18: 119 – 120.
17. Grayston, S.J., J.H. Stephens and L.M. Nelson. 1990. Field and green house studies on growth promoting of spring wheat inoculation with co- existent rhizobacteria. Second Int. Workshop on PGPR, Interlaken, Switzerland.88- 96. (c.f.El- Hadidy,2003).
18. Hafeez, F. Y., M. E. Safdar, A. U. Chaudhry and K. A. Malik. 2004. Rhizobial inoculation improves seedling emergence, nutrient uptake and growth cotton. *Australian J. Experi. Agric.*, 44(6): 617- 622.
19. Hegazi , N.A. and S.I. Niemela. 1976. Note on the estimation of *Azotobacter* densities by membrane – filter technique. *J. Appl. Bact.*, 41: 311 – 314.
20. Ibrahim, A. I. 1999. Somaclonal variation during micropropagation of date palm *via* embryogenesis. The First International Conference, in Egypt, on Plant Tissue Culture and its Application. pp. 189 – 199.
21. Joo, G.J., Y.M. Kim, I.J. Lee, K.S. Song and I.K. Rhee. 2004. Growth promoting of red pepper plug seedling and the production of gibberellins by *Bacillus cereus*, *Bacillus macroides* and *Bacillus pumilus*. *Biotechnology Letters*. 26: 487- 491.
22. Kapulink, Y.R., R. Gafny and Y. Okon. 1985. Effect of *Azospirillum* spp. Inoculation on root development and NO<sub>3</sub> uptake in wheat in hydroponic systems. *Can. J.Bot.*, 63:627- 631.
23. Khan, M. A., M. S. Khalil and M. S. Al-Kahtani. 1983. *In vitro* culture of different tissue of date palm *Phoenix dactylifera* L. offshoot. 1<sup>st</sup> Symp. on date palm, King Faisal University, Al-Hassa, Saudi Arabia, pp.152–157.
24. Mahmoud, S.A.Z., E.M. Ramadan, F.M. Thabat and T. Khater. 1984. Production of plant growth promoting substances by rhizosphere microorganisms. *Zbl. Mikrobiol.*, 139: 227- 232.
25. Moursy, H. and M. Saker. 1996. Date palm problems and the need for biotechnology. Abst. Of the 5<sup>th</sup> Int. Conf. Desert. Development Texas Tech. Univ. USA, August 12- 17, p.44.
26. Murashige, T. and F. A. Skoog. 1962. A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Plant Physiol.*, 15: 473-479.

27. One, O., J. Van.Impe, E. Prinsen, J. Vanderleyden. 2005. Growth and IAA biosynthesis of *Azospirillum brasilense* Sp245 is environmentally controlled. FEMS Microbiology Letters. 246(1): 125- 132.
28. Omar, M. S. 1988. Callus initiation, asexual embryogenesis and plant regeneration in *Phoenix dactylifera* L. Date Palm Journal. 6: 265-275.
29. Pan, M. J. and J. Van Staden. 1998. The use of charcoal in *in vitro* culture. Plant Growth Regul., 26: 155-163.
30. Pottino, B. G. 1981. Methods in plant tissue culture. Dept. of Hort., Agric. College, Maryland Univ., College park, Maryland, USA., pp.8-29.
31. Reuveni, O. and H. L. Kipins. 1974. Studies of the *in vitro* culture of date palm (*Phoenix dactylifera* L.) tissue and organs. Pamphlet, No.145, pp.3-39.
32. Reynolds, J. F. and T. Murashige. 1979. Asexual embryogenesis in callus culture of plants. *In vitro*. 15: 383-387.
33. Saker, M. M., H. A. Moursy and S. A. Bekheet. 1998. *In vitro* propagation of Egyptian date palm: morphogenic responses of immature embryos. Bulletin of Faculty of Agriculture, Cairo University, 49 (2): 203-214.
34. Sharon, M., P. C. Shankar and M. Sharon. 1998. Somatic embryogenesis and plant regeneration from leaf primordial of *Phoenix dactylifera*. Indian. Journal Experimental Biology, 36 (5): 426-529.
35. Smith, R. H. 2000. Plant tissue culture techniques and experiments. Academic Press. New York,U.S.A.,pp. 15-28.
36. Snedecor, G. W. and Cochoran. 1972. Statistical Method 6<sup>th</sup>. The Iowa State Univ. Press, Ames., Iowa, U.S.A., pp.593.
37. Thorpe, T. A. 1981. Plant tissue culture methods and application in agriculture. Academic Press, New York, U.S.A., pp. 45-60.
38. Tisserat, B. 1979. Propagation of date palm *Phoenix dactylifera* L. *in vitro*. Journal Experimental Botany. 30: 1275 – 83.
39. Tisserat, B. 1982. Factors involved in the production of plantlets from date palm callus cultures. Euphytica. 31 (1): 201 -214.
40. Tisserat, B. 1983. Development of new tissue culture technology to aid in cultivation and crop improvement of date palms. Proc. First Symposium on Date palm, Saudi Arabia. Pp. 126- 140.
41. Trigiano, R. M. and D. J. Gray. 2000. Plant tissue culture concepts and laboratory exercises. CRC Press. New York, U.S.A., pp. 67.
42. Zaid, A. and B. Tisserat. 1983. *In vitro* shoot tip differentiation in *Phoenix dactylifera* L. Date palm J., 2 (2):163-182.
43. Zaid, Z. E. 2003. Comparative studies on the production of date palm cultivars via tissue culture technique. Ph.D. Thesis, Pomology Dept., Fac. of Agric., Cairo Univ., Egypt.

## تعاقب استخدام الراشحات البكتيرية في زراعة انسجه نخيل البلح

هاله محمد أنور فراج<sup>١</sup> ، وداد التهامي عويضة<sup>٢</sup> ، عادل أسخرون جرجس<sup>٣</sup> ، عبد المنعم عبد الودود ألبنا<sup>١</sup> ، عاطف محمد إبراهيم<sup>٢</sup> ، إبراهيم عبد المقصود إبراهيم<sup>٣</sup>

- ١ - المعمل المركزى للأبحاث و تطوير نخيل البلح - مركز البحوث الزراعية
- ٢ - قسم الميكروبيولوجيا الزراعية ، كلية الزراعة ، جامعة عين شمس
- ٣ - معهد الهندسة الوراثية والتكنولوجيا الحيوية - جامعه المنوفيه

يهدف هذا البحث إلى استخدام الهرمونات المنتجة طبيعيا بواسطة بعض السلالات البكتيرية مثل *Azospirillum brasilense*, *Azotobacter chroococcum*, *Bacillus megaterium* and *Klebsiella pneumoniae* بدلا من منظمات النمو المصنعه في زراعه انسجه نخيل البلح عن طريق تكوين الأجنة الجسديه.

بصفه عامه أظهرت النتائج أن: *Azotobacter chroococcum* و *Klebsiella pneumoniae* لها دور اوكسينى قوى بينما *Azospirillum brasilense* و *Bacillus megaterium* تنتمي في تأثيرها أكثر إلى السيتوكينين.

وقد تم استخدام الراشح المأخوذ من مزارع تلك السلالات ذات الكفاءة العاليه في إنتاج الأوكسين أو السيتوكينين ضمن مكونات بيئه الزراعة المستخدمه لتكوين الأجنة الجسديه من القمه الناميصة أو بادئات الأوراق المفصولة من فسانل نخيل البلح .