

## EFFICIENCY OF SOME SOIL MICROORGANISMS IN DEGRADATION OF DIAZINON PESTICIDE

### ABSTRACT

A pot experiment was carried out to study the efficiency of *Bacillus polymyxa*, *Pseudomonas fluorescens* and/or *Streptomyces aureofaciens* on biodegradation of the pesticide diazinon in sterilized and non-sterilized soil, either non-cultivated or cultivated. The obtained results showed that disappearance rate of diazinon was higher in the non-sterilized soil than in the sterilized one. Inoculation of soil with mixture of *Bacillus polymyxa* + *Streptomyces aureofaciens* and *Pseudomonas fluorescens* + *Streptomyces aureofaciens* exhibited higher dissipation rate of diazinon in either sterilized or non-sterilized soil, as compared with the soil inoculated with each microorganism individually. When the soil was cultivated with tomato plants and inoculation was made, biodegradation of diazinon increased, compared to the uncultivated and inoculated soil. The metabolites produced from biodegradation of diazinon by the tested microorganisms were 1,3-dimethyl-2-nitrobenzene, diethylphosphate, 2-isopropyl-6-methyl-4-hydroxypyrimidine (IMHP) and 2-(1-hydroxy-1-methyl)-6-methyl-4 (IH)-pyrimidine (HMMP).

**Key words:** Diazinon, biodegradation, *B. polymyxa*, *P. fluorescens* and *S. aureofaciens*.

### INTRODUCTION

Organophosphorus pesticides are more toxic to mammals than organochlorine ones. The persistence of pesticides has been of a considerable benefit to the user. Also, organophosphorus pesticide compounds introduce serious problems particularly with improper usage by farmers.

Although, there are many factors affect the persistence of pesticides in soil, such as temperature, pH, water content and others. But, soil microorganisms appeared to play a major role in the degradation of pesticides.

Primarily, biodegradation of pesticides under aerobic conditions is a major fate process for diazinon associated with water and soil. Also, diazinon can be biodegraded under anaerobic condition (Environmental Protection Agency, 1990). Diazinon has a relatively short half-life in soil ranging from 70 hrs to 12 weeks depending on pH, temperature as well as on the presence of specific lytic microorganisms (Ferrando *et al.* 1992 and Scheunert *et al.* 1993).

Also, Aislabie & Jones (1997) and Megharaj *et al.* (2003) reported that pesticides fate in the environment is mainly affected by microbial activity.


Li *et al.* (2002) reported that the degrading compounds from diazinon had identified as 2-isopropyl-6-methyl-4-hydroxypyrimidine (IMHP); IMHP is a major degradation product when low concentration of diazinon was studied in soil and water. Although (IMHP) is found to be potentially leacheable, it is less toxic than diazinon. Moreover, diazinon-O-analog (or diazoxon), 1,3-dimethyl-2-nitrobenzene, O,O,O,O-tetraethyl dithiopyrophosphate can also be produced as hydrolytic degradation. This research aimed at studying the efficiency of *Bacillus polymyxa*,

*Pseudomonas fluorescens* and *Streptomyces aureofaciens* on biodegradation of diazinon in sterilized and non-sterilized soils, both non-cultivated and cultivated.

## MATERIALS AND METHODS

### The pesticide used.

Organophosphorus pesticide, which is widely used in controlling many pests in Egyptian agricultural, was used, namely Diazinon (Dimpylate), "0,0-Diethyl-0-(2-isopropyl-4-methyl-6-pyrimidinyl) phosphorothioate".



Structural formula of diazinon

### Efficiency of the tested bacterial strains on biodegradation of diazinon in soil.

A pot experiment was carried out to study the efficiency of *Bacillus polymyxa*, *Pseudomonas fluorescens* and/or *Streptomyces aureofaciens* on biodegradation of diazinon in non-cultivated or cultivated, sterilized or non-sterilized soils. Textural class of the investigated soil was clayey loam. The collected soil was sieved and divided into two parts, one of them was sterilized by autoclaving at 121°C for 3 hrs. Whereas, the second part of soil was left without sterilization. A 30 cm diameter pots were filled with the sterilized or the non-sterilized soils (3 kg/pot). Each soil part was divided into three portions. The first and second potted portions were left without cultivation. The third portion was amended with NPK fertilizers at recommended dose and immediately planted with tomato seedlings c.v. super strain B. After 15 days of planting, each portion was divided into two sub-portion. One of them was left as a control (without pesticide application), while, the other was mixed with diazinon to give a final concentration (50 ppm). After 15 days of pesticide application, the following treatments were carried out:

- No bacterial inoculation.
- Inoculation with *B. polymyxa* (20 ml/pot,  $930 \times 10^6$  c.f.u. / ml) 48 hrs old culture.
- Inoculation with *P. fluorescens* (20 ml/pot,  $93 \times 10^6$  c.f.u./ ml) 48 hrs old culture.
- Inoculation with *S. aureofaciens* (20 ml/pot,  $7 \times 10^6$  spores/ml) 7 days old culture.
- Inoculation with *B. polymyxa* and *S. aureofaciens* (20 ml/pot) from each of them.
- Inoculation with *P. fluorescens* and *S. aureofaciens* (20 ml/pot) from each of them.

Soil samples were investigated five times throughout the experiment, i.e. initially and at 3, 7, 15, 35 days after inoculation with the tested microorganisms for assessment of the following:

- a) Dehydrogenase activity.
- b) Nitrogenase activity.
- c) Phosphatase activity.
- d) Persistent amount of the pesticide and its metabolites.

Dehydrogenase activity was assayed according to **Thalmann (1967)**. Phosphatase activity was determined by the method recommended by **Drobnikova (1961)**. Nitrogenase activity was measured using the technique adopted by **Dilworth (1970)**.

### Extraction of diazinon and its degradation products.

The method of **Lichtenstein et al (1967)** for extraction of diazinon and its metabolites from soil was used with some modifications. Gas Liquid Chromatography and Gas Chromatography/Mass Spectrometer were employed for detection of diazinon and its metabolites.

## RESULTS AND DISCUSSION

### Periodical changes of dehydrogenase activity in non-sterilized soil non-cultivated and cultivated with tomato.

Data illustrated in Table (1) showed that the untreated soil with diazinon gave higher values of DHA activity as compared with the soil treated with the pesticide. This result was expected since pesticides application to the soil inhibits the activities of different soil microorganisms. This result is in agreement with that obtained by **Balinove et al. (1997)** and **Lan et. al. (2006)** who reported that pesticides application to the soil had adverse effect on microbial populations and consequently the microbial enzyme activities were decreased.

Data in Table (1) also emphasized that soil treated with diazinon and uninoculated with tested microorganisms (control) gave lower values of DHA activity, compared to soil treated with diazinon and inoculated.

**Table (1): Periodical changes of dehydrogenase activity in non-sterilized soil non-cultivated and cultivated with tomato.**

Treatments Incubation periods(days)			Dehydrogenase activity (µg TPF / g dry soil / day)													
			Soil untreated with diazinon		Soil treated with diazinon*											
Planting	Pesticide addition	Inoculation	Non- cultivated soil	Cultivated soil	Non-cultivated soil						Cultivated soil					
					Control	A	B	C	A+C	B+C	Control	A	B	C	A+C	B+C
30	15	0	33.7	37.5	23.6	23.6	23.6	23.6	23.6	23.6	23.5	23.5	23.5	23.5	23.5	
33	18	3	30.1	38.7	20.1	25.0	26.0	22.6	36.2	38.8	24.9	29.3	28.6	28.4	43.5	
37	22	7	42.9	45.6	21.9	27.5	20.7	23.3	29.4	32.8	12.3	31.0	31.2	31.2	32.1	
45	30	15	32.3	41.3	18.3	35.7	30.7	28.0	38.6	40.0	14.4	39.1	37.1	37.1	41.3	
65	50	35	28.9	30.2	22.1	25.0	19.7	23.0	26.3	31.8	19.0	30.5	23.1	23.1	28.2	

\* A: *Bacillus polymyxa*

B: *Pseudomonas fluorescens*

C: *Streptomyces aureofaciens*

In addition, the soil treated with diazinon and inoculated with the mixture of the tested bacteria showed higher DHA activity than the soil inoculated with each one individually. Higher values of DHA activity in case of the soil inoculated with the mixture of the strains is likely due to the synergistic effect between the strains. Similar result was observed by Aislabie & Jones (1997) who mentioned that some pesticides are readily degraded by microorganisms including members of genera *Alcaligenes*, *Flavobacterium*, *Pseudomonas*, *Streptomyces* and *Rhodococcus*.

Moreover, data revealed that the soil treated with diazinon, inoculated with bacteria and cultivated with tomato plants gave higher values of DHA activity as compared with the uncultivated soil. Higher values of DHA activity in the cultivated soil may be due to the beneficial effect of root exudates and root debris which represent nutritional substances for soil microorganisms.

#### Periodical changes of $N_2$ -ase activity in non-sterilized soil non-cultivated and cultivated with tomato.

Data illustrated in Table (2) showed that the untreated soil gave higher values of  $N_2$ -ase activity in rhizosphere soil compared to the soil treated with diazinon. This result is in harmony with that of Trabue *et al.* (2001) who mentioned that the microbial activity was adversely affected by the nematicides application.

Yueh & Hensley (1993) reported that diazinon significantly decreased  $C_2H_2$  reduction. The soil amended with diazinon and uninoculated with the biodegrading strains showed lower values of  $N_2$ -ase activity, which can be attributed to the inhibitive effect of diazinon pesticide on free living  $N_2$ -fixers originally occurred in soil. This result is in accordance with Doggette & Rhodes (1991) who found that the growth rate of nitrogen fixers was suppressed by diazinon application at a concentration of 40 ppm.

Table (2): Periodical changes of  $N_2$ -ase activity in non-sterilized soil non-cultivated and cultivated with tomato.

Treatments			Nitrogenase activity (nmoles C <sub>2</sub> H <sub>4</sub> / g dry soil / hr.)													
			Soil untreated with diazinon		Soil treated with diazinon*											
Planting	Pesticide addition	Inoculation	Non-cultivated soil	Cultivated soil	Non-cultivated soil						Cultivated soil					
					Control	A	B	C	A+C	B+C	Control	A	B	C	A+C	B+C
30	15	0	50.06	95.48	45.28	45.28	45.28	45.28	45.28	45.28	45.25	45.25	45.25	45.25	45.25	45.25
33	18	3	50.01	65.62	15.02	15.9	11.68	15.06	25.35	25.33	18.26	26.16	28.14	25.96	55.17	48.31
37	22	7	40.51	55.59	5.40	15.73	15.95	20.23	30.89	25.96	20.85	30.65	32.27	23.0	65.3	60.55
45	30	15	50.01	55.09	5.07	15.28	15.03	15.07	25.16	25.02	24.47	33.90	28.68	28.19	70.60	65.66
65	50	35	43.8	48.2	21.6	26.1	24.18	21.36	28.61	26.8	30.38	36.69	34.4	31.14	78.6	68.16

\*A: *Bacillus polymyxa*

B: *Pseudomonas fluorescens*

C: *Streptomyces aureofaciens*

On the contrary, the soil inoculated with the investigated microorganisms and treated with diazinon showed higher values of  $N_2$ -ase activity than the uninoculated one. The soil amended with diazinon and inoculated with the mixture of strains gave higher values of  $N_2$ -ase activity as compared with soil inoculated with the strains individually.

The higher values of N<sub>2</sub>-ase activity in rhizosphere soil of inoculation treatments can be attributed to the hydrolysis and biodegradation of diazinon either by normal habitat or inoculated microorganisms in soil. Microbial degradation appeared to be the major pathway for the degradation of diazinon in soil (Aislabie and Jones, 1997).

Generally, the obtained data emphasized that the N<sub>2</sub>-ase activity was higher in the cultivated soil treatments compared with the uncultivated ones. Such increase is likely due to the difference in multiplication rate of different N<sub>2</sub>-fixers in the cultivated soil as a result of qualitative and quantitative changes in the nature of root exudates during different determination periods.

#### Periodical changes of phosphatase activity in non-sterilized soil non-cultivated and cultivated with tomato.

Data presented in Table (3) showed that the untreated soil gave higher values of phosphatase activity compared to the soil treated with diazinon during the first 15 days of the experimental period. This result was expected since pesticides application to the soil inhibits various microbial activities.

Whereas, results showed that phosphatase activity was enhanced later on the 35<sup>th</sup> day sample in the soil treated with diazinon as compared with the untreated one. This result can be attributed to the hydrolysis and biodegradation of diazinon either by the tested strains or by the natural flora of the soil. Similar results were obtained by Li *et al.* (2002) who found that metabolites of diazinon were less toxic than the parent compound.

Data also emphasized that the soil treated with diazinon and uninoculated with any of the tested microorganisms (control) gave lowest values of phosphatase activity in the periods of 3, 7 days compared to the soil treated with diazinon and inoculated with each strain individually.

**Table (3): Periodical changes of phosphatase activity in non-sterilized soil non-cultivated and cultivated with tomato.**

Treatments  Incubation periods (days)			Phosphatase activity (µg inorganic phosphorus / g dry soil / 24 hrs.)													
			Soil untreated with diazinon	Soil treated with diazinon*												
Planting	Pesticide addition	Inoculation	Non-cultivated soil	Cultivated soil	Non-cultivated soil						Cultivated soil					
					Control	A	B	C	A+C	B+C	Control	A	B	C	A+C	B+C
30	15	0	16.6	19.2	12.8	12.8	12.8	12.8	12.8	12.8	11.9	11.9	11.9	11.9	11.9	11.9
33	18	3	21.3	23.1	11.0	16.0	13.4	14.8	17.5	19.8	20.0	18.1	26.7	16.1	34.3	31.7
37	22	7	23.4	28.2	9.2	19.7	14.9	16.8	23.9	36.4	16.0	25.2	22.5	18.2	33.1	37.0
45	30	15	19.4	21.3	14.7	15.0	14.7	11.9	22.9	22.6	18.9	18.6	16.4	19.69	23.4	22.7
65	50	35	14.6	27.9	24.7	23.7	26.1	24.4	24.9	25.1	26.7	26.2	28.2	29.8	32.4	30.2

\*A: *Bacillus polymyxa*

B: *Pseudomonas fluorescens*

C: *Streptomyces aureofaciens*

In addition, the soil treated with diazinon and inoculated with a mixture of the tested strains showed higher phosphatase activity than the soil inoculated with each strain individually. Similar results were observed by Aislabie & Jones (1997) and Lui *et al.* (2005).

Moreover, the obtained data revealed that the soil treated with diazinon and cultivated with tomato plants gave higher values of phosphatase activity, compared to the uncultivated one. The phosphatase activity in the inoculated soil showed higher values than in the control; and the inoculation with the bacterial mixture apparently enhanced phosphatase activity than the individual strains. This may be due to the beneficial effect of root exudates and root debris which represent nutritional substances for different soil microorganisms.

#### **Persistence rate of diazinon in sterilized and non-sterilized soil.**

Persistence rates of diazinon are presented in Table (4) as percentages of the initial concentrations. The obtained results showed that disappearance rate of diazinon was higher in the non-sterilized soil than in the sterilized one. Disappearance rate of diazinon in the non-sterilized and uninoculated soils was faster than in the sterilized uninoculated one, since 42.0 and 72.1 % of the added amounts were detected in the non-sterilized uninoculated and the sterilized uninoculated soil, respectively at the end of the experiment. This result revealed the importance of indigenous soil microorganisms in diazinon decomposition. This result is in agreement with that obtained by Seyfried (1994) who found that losses of diazinon from sterilized samples of soil were much slower than from non-sterilized one, indicating microbial participation in diazinon degradation in soils. Talebi & Walker (1993) and Kale *et al.* (1996) found a similar trend of results when they studied the degradation of pesticide (carbofuran) in sterilized soil compared to non-sterilized one.

Dissipation rate of diazinon was accelerated in soil inoculated with *Bacillus polymyxa*, *Pseudomonas fluorescens* and *Streptomyces aureofaciens* compared to the uninoculated soil. Soil inoculated with *Bacillus polymyxa* showed the highest dissipation rate of diazinon compared to other microorganisms when each strain was individually used. These results revealed that all tested microorganisms are capable to hydrolyze diazinon. These results are in harmony with those reported by Kale *et al.* (1996), Trabue *et al.* (2001) and Singh *et al.* (2005) who revealed that there were many soil microorganisms capable of pesticides degradation when they were used as soil application.

Also, Ferrando *et al.* (1992) and Scheunert *et al.* (1993) found that diazinon was degraded by several genera of microorganisms and the microbial degradation appeared to be the major pathway for the degradation of diazinon in soils. Also, Aislabie & Jones (1997), Miguel *et al.* (2002), Wander *et al.* (2003) and Vijay *et al.* (2006) reported that some pesticides are readily degraded by microorganisms, others have proven to be recalcitrant. A diverse group of bacteria including members of genera *Pseudomonas*, *Alcaligenes*, *Flavobacterium* and *Rhodococcus* were able to metabolize pesticides.

**Table (4): Persistence rate of diazinon (%) in non-sterilized and sterilized soil.**

Treatments  Days after Diazinon addition		Uninoculated soil	Soil inoculated with *				
			A	B	C	A + C	B + C
Inoculation	Non-sterilized soil						
0	-	100	100	100	100	100	100
15	0	64.5	64.5	64.5	64.5	64.5	64.5
18	3	57.3	39.4	32.1	28.3	24.3	31.9
22	7	51.8	29.7	18.5	27.9	24.2	22.7
30	15	48.4	22.2	16.7	13.5	9.5	16.2
50	35	42.0	7.3	9.3	11.5	0.0	4.0
		Sterilized soil					
0	-	100	100	100	100	100	100
15	0	88.6	88.6	88.6	88.6	88.6	88.6
18	3	82.3	56.2	56.0	56.3	54.3	53.8
22	7	76.3	48.2	51.3	41.7	41.9	40.3
30	15	73.0	29.9	30.1	30.2	32.5	22.0
50	35	72.1	14.7	20.7	18.3	10.0	8.0

\*A: *Bacillus polymyxa*B: *Pseudomonas fluorescens*C: *Streptomyces aureofaciens*

Inoculation of the soil with mixtures of *Bacillus polymyxa* + *Streptomyces aureofaciens* and *Pseudomonas fluorescens* + *Streptomyces aureofaciens* exhibited higher dissipation rate of diazinon either in sterilized or non-sterilized soils, as compared with the soil inoculated with each microorganism individually. This increase in dissipation rate in the case of soil inoculation with the mixture of strains was likely being due to their synergistic interaction to improve the degradation of diazinon.

#### **Persistence rate of diazinon in non-sterilized soil uncultivated and cultivated with tomato.**

Data in Table (5) showed that the uninoculated soil gave lower dissipation rate of diazinon, compared to the inoculated one. This result was observed in either the cultivated or uncultivated soils. When the soil was cultivated with tomato plants and bacterially inoculated, biodegradation of diazinon was increased as compared with the uncultivated and inoculated soil. Since, the persistence percentages of diazinon at the end of the experiment in the cultivated soil was lower than those recorded in the uncultivated one. This difference in disappearance rate of diazinon might be due to the root excretions of tomato plants which may synergist the proliferation of soil microorganisms either the indigenous or the introduced and therefore the decomposition rate of diazinon was increased in the cultivated soil. These results are in agreement with those obtained by **Balinove et al. (1997)** **Trabue et al. (2001)** and **Dimitrios et al. (2005)** who found that the persistence rate of pesticides in soil cultivated with tomato was lower than in uncultivated one.

Inoculation of soil with *B. polymyxa* individually showed the highest disappearance rate of diazinon compared to soil inoculated with either *P. fluorescens* or *Streptomyces aureofaciens*. From data presented in Table (5) it was clear that soil inoculation with the mixtures of strains exhibited highly disappearance rate of diazinon as compared with the soil inoculated with each microorganism apart.

Increasing of disappearance rate of diazinon from the soil inoculated with the mixtures of strains is likely due to the synergistic effect that occurred between the bacterial inocula and subsequently the biodegradation rate of diazinon was increased.

**Table (5): Persistence rate of diazinon in non-sterilized soil non-cultivated and cultivated with tomato.**

Treatments Days after			Uninoculated soil	Soil inoculated with*				
				A	B	C	A + C	B + C
Planting	Diazinon addition	Inoculation	Non-cultivated soil					
15	0	-	100	100	100	100	100	100
30	15	0	64.5	64.5	64.5	64.5	64.5	64.5
33	18	3	57.3	39.4	32.1	28.3	24.3	31.9
37	22	7	51.8	29.7	18.5	27.9	24.2	22.7
45	30	15	48.4	22.2	16.7	13.5	9.5	16.2
65	50	35	42.0	7.3	9.3	11.5	0.0	4.0
			Cultivated soil					
15	0	-	100	100	100	100	100	100
30	15	0	52.0	52.0	52.0	52.0	52.0	52.0
33	18	3	50.2	30.8	21.8	20.9	19.5	18.9
37	22	7	47.8	22.4	15.7	19.9	13.0	16.0
45	30	15	34.4	13.0	9.6	19.2	12.4	11.1
65	50	35	35.5	1.4	1.9	2.9	1.2	0.0

\*A: *Bacillus polymyxa* B: *Pseudomonas fluorescens* C: *Streptomyces aureofaciens*

Analysis of diazinon biodegradable products by the tested bacterial strains in sterilized and non-sterilized soil using GLC.

Analysis of soil samples was carried out by GLC after 30 days of diazinon treatment (15 days after inoculation). The obtained results are given in Table (6) and illustrated by Fig (1 a, b).

From data in Table (6) it is obvious that the non-sterilized soil treatments showed more metabolites from the biodegradation of diazinon by the investigated bacteria, compared to those produced from sterilized one. These results reveal the importance of indigenous (native) soil microorganisms in diazinon decomposition in the soil. This result is in harmony with that obtained by **Getzin & Shanks (1990)** who found that the degradation of pesticides was slower in sterilized soil and the half-life of pesticides was exceeded. **Arunachalam & Lakshamanan (1990)** and **Zaghloul et al. (2003)** who found that in sterilized soil about 75 % of the added carbofuran was recorded as residue after 60 days, whereas in non-sterilized soil about 75 % of the added carbofuran was metabolized.

Extract of the non-sterilized soil inoculated with either *Bacillus polymyxa* or *Pseudomonas fluorescens* cultures showed five metabolites having Rt 0.91, 1.7, 1.8, 1.9 and 2.3 minutes. Whereas, the extract of the abovementioned cultures in the sterilized soil showed four metabolites having Rt 0.91, 1.7, 1.8 and 1.9. Extract of the soil inoculated with *Streptomyces aureofaciens* showed lower metabolites than the soil inoculated with either *Bacillus polymyxa* or *Pseudomonas fluorescens*. Similar trend of results was observed in both the sterilized and non-sterilized soil.



**Table (6): Analysis of diazinon biodegradable products by the tested bacterial strains in non-sterilized and sterilized soil using GLC.**

Compounds	Uninoculated soil	Retention time (Rt min.)*				
		A	B	C	A + C	B + C
	Non-sterilized soil					
Diazinon	4.8	4.8	4.8	4.8	4.8	4.8
metabolites 1	0.91	0.91	0.91	0.91	-	0.91
2	-	1.7	1.7	1.7	-	-
3	1.8	1.8	1.8	-	-	1.8
4	-	1.9	1.9	1.9	1.9	-
5	-	2.3	2.3	-	-	-
	Sterilized soil					
Diazinon	4.8	4.8	4.8	4.8	4.8	4.8
metabolites 1	0.91	0.91	0.91	0.91	0.91	0.91
2	-	1.7	1.7	1.7	-	-
3	1.8	1.8	1.8	-	-	-
4	-	1.9	1.9	1.9	1.9	-
5	-	-	-	-	-	-

\*A: *Bacillus polymyxa* B: *Pseudomonas fluorescens* C: *Streptomyces aureofaciens*


Metabolites produced from diazinon decomposition can't be identified by GLC analysis because their authentic materials were not available.

Gas mass spectrometer was used to determine the values of M/e for each compound to be as reference to use these values for identification of the compounds produced from the biodegradation of diazinon by the investigated microorganisms. The limited data are presented in Table (7) and shown in Fig. (2).

It could be noticed that the metabolites produced from the biodegradation of diazinon by the investigated bacteria showed molecular weights equivalent to the molecular weights of diazinon and its dominant metabolites which were observed and identified by **Robert *et al.* (2000)** in their study. They mentioned that diazinon pesticide was metabolized by microorganisms to diazoxone, diethylphosphate, 2-isopropyl-6-methyl-4-hydroxypyrimidine (IMHP), 2-(1-hydroxy-1-methyl)-6-methyl-4 (IH)-pyrimidine (HMMP) and 1,3-dimethyl-2-nitrobenzene. These compounds are recorded in Table (7).


Therefore, the metabolites produced from the biodegradation of diazinon by the tested bacterial strains were 1,3-dimethyl-2-nitrobenzene, diethylphosphate, HMMP and IMHP.

The identification deduction of the abovementioned metabolites was based on the relation between retention time obtained from GLC analysis and molecular weight obtained from Gas/Mass analysis which was achieved to confirm the obtained results. These results are in agreement with those obtained by **Li *et al.* (2002)** in their studies on diazinon decomposition. They reported that 2-isopropyl-6-methyl-4-hydroxypyrimidine was a major degradation product.



**Fig 1 a. Gas liquid chromatography spectra of the non-sterilized soil extracts at 15 days of:**

- A) Soil treated with diazinon.
- B) Soil treated with diazinon and inoculated with *Bacillus polymyxa* (B).
- C) Soil treated with diazinon and inoculated with *Pseudomonas fluorescens* (P).
- D) Soil treated with diazinon and inoculated with *Streptomyces aureofaciens* (S).
- E) Soil treated with diazinon and inoculated with (B + S).
- F) Soil treated with diazinon and inoculated with (P + S).



**Fig 1 b. Gas liquid chromatography spectra of the sterilized soil extracts at 15 days of:**


- A) Soil treated with diazinon.
- B) Soil treated with diazinon and inoculated with *Bacillus polymyxa* (B).
- C) Soil treated with diazinon and inoculated with *Pseudomonas fluorescens* (P).
- D) Soil treated with diazinon and inoculated with *Streptomyces aureofaciens* (S).
- E) Soil treated with diazinon and inoculated with (B + S).
- F) Soil treated with diazinon and inoculated with (P + S).

Dumas *et al.* (1990) stated that the organophosphorus hydrolase (OPH) is an organophosphotriester hydrolyzing enzyme that first discovered in the soil microorganisms *Pseudomonas diminuta* MG and *Flavobacterium* spp. The enzyme has broad substrate specificity and is able to hydrolyze a number of OPH pesticides such as paraoxon, parathion, coumaphos and diazinon. The reaction products, from hydrolysis by organophosphorus hydrolase (OPH) had generally reduced toxicity relative to the parent compound (Lai *et al.*, 1994 and Yakovlevsky *et al.*, 1997).

Summing up, from the obtained results it could be concluded that soil microorganisms play an important role in pesticides degradation.

**Table (7): Gas/Mass analysis of diazinon and related compounds in soil.**

Compounds	Molecular weight	M/e	Chemical formula
Diazinon	304	304	C <sub>12</sub> H <sub>21</sub> O <sub>3</sub> N <sub>2</sub> PS
Diethyl phosphate	154	154	C <sub>4</sub> H <sub>11</sub> O <sub>4</sub> P
2-(1-hydroxy-1-methyl)-6-methyl-4(1H)-pyrimidinone (HMMP)	168	167	C <sub>8</sub> H <sub>12</sub> O <sub>2</sub> N <sub>2</sub>
2-isopropyl-6-methyl-4-hydroxypyrimidine (IMHP)	152	152	C <sub>8</sub> H <sub>12</sub> ON <sub>2</sub>
1,3-dimethyl-2-nitrobenzene	151	151	C <sub>8</sub> H <sub>9</sub> O <sub>2</sub> N



**Fig (2): Mass spectrum of diazinon and its metabolites in soil.**

### Analysis of diazinon biodegradable products by the tested bacterial strains in cultivated and non-cultivated soil using GLC.

Data presented in Table (8) clearly show that extracts of the non-cultivated soil inoculated with either *Bacillus polymyxa* or *Pseudomonas fluorescens* showed five compounds having Rt 0.91, 1.7, 1.8, 1.9 and 2.3 minute. Whereas, inoculation of the cultivated soil showed in chromatographic analysis four compounds in case of *Bacillus polymyxa* having Rt 0.91, 1.7, 1.8 and 1.9 minutes and two compounds in case of *Pseudomonas fluorescens* having Rt 0.91 and 1.9 minutes.

Extract of the soil inoculated with *Streptomyces aureofaciens* showed lower metabolites than the soil inoculated with *Bacillus polymyxa*. Similar trend of results was observed in both the cultivated and uncultivated soil.

From the obtained results, it could be concluded that diazinon in soil decomposed or metabolized to five metabolites having Rt 0.91, 1.3, 1.7, 1.8 and 1.9 minutes.

Metabolites produced from diazinon decomposition was identified by the same method mentioned before in the sterilized and non-sterilized soil (Robert *et al.*, 2000).

Therefore, the metabolites products from the biodegradation of diazinon by various microorganisms used in this research were 1,3-dimethyl-2-nitrobenzene, diethylphosphate, HMMP and IMHP.

**Table (8): Analysis of diazinon biodegradable products by the tested bacterial strains in non-cultivated and cultivated soil using GLC.**

Compounds	Uninoculated soil	Soil inoculated with*				
		A	B	C	A + C	B + C
	Non-cultivated soil					
Diazinon	4.8	4.8	4.8	4.8	4.8	4.8
metabolites 1	0.91	0.91	0.91	0.91	0.91	0.91
2	-	-	-	-	-	-
3	-	1.7	1.7	1.7	-	-
4	1.8	1.8	1.8	-	-	1.8
5	-	1.9	1.9	1.9	1.9	-
6	-	2.3	2.3	-	-	-
	Cultivated soil					
Diazinon	4.8	4.8	4.8	4.8	4.8	4.8
metabolites 1	0.91	0.91	0.91	0.91	0.91	0.91
2	1.3	-	-	-	-	-
3	-	1.7	-	-	-	-
4	1.8	1.8	-	-	-	-
5	-	1.9	1.9	1.9	1.9	1.9
6	-	-	-	-	-	2.3

\*A: *Bacillus polymyxa* B: *Pseudomonas fluorescens* C: *Streptomyces aureofaciens*

Similar result was observed by Seyfried (1994) who reported that the main metabolite produced from biodegradation of diazinon was 2-isopropyl-6-methyl-4-hydroxypyrimidine (IMHP). Also, Michel *et al.* (1997) found that IMHP and diazoxon were the major degradation products. Whereas, Li *et al.* (2002) found that 1,3-dimethyl-2-nitrobenzene, compound was produced from the biodegradation of diazinon.

## REFERENCES

- Aislabie, J. and Jones, G. (1997). A review of bacterial degradation of pesticides . Australian Journal of Soil Research, Page 3127. Hamilton, New Zealand.
- Arunachalam, K. D. and Lakshmanan, M. (1990). Decomposition of <sup>14</sup>C-labelled carbofuran in a black tropical soil under laboratory conditions. Soil Biology and Biochemistry, 22 (3): 407-412.
- Balinove, I.; Balinova, A. and Naclikova, V. (1997). Behavior of some systemic soil applied N-methylcarbamate pesticides in greenhouse tomato plants. Bulgarian J. of Agricultural Science, 3(5): 551-555.
- Dilworth, M.J. (1970). The acetylene reduction method for measuring biological nitrogen fixation. Rhizobium News Later, 15 (7) : 155.
- Dimitrios G. K., Anastasia, F.; Urania, M.S. and Singh, B. K.(2005). Non-specific biodegradation of the organophosphorus pesticides, cadusafos and ethoprophos, by two bacterial isolates. Microbial Ecology, 53(3): 369-378.
- Doggette, S.M. and Rhodes, R.G. (1991). Effect of diazinon formulation on unialgal growth rates and phytoplankton diversity. Bull. Environ. Contam. Toxicol., 47, 36-42.
- Drobnikova, V. (1961). Factors influencing the determination of phosphatase in soil. Folia. Microbiol., 6, 260.
- Dumas, D. P.; Durst, H. D .; Landis, W. G.; Raushel, F.M. and Wild, J. R. (1990). Inactivation of organophosphorus nerve agents by the phosphotriesterase from *Pseudomonas diminuta*. Arch. Biochem. Biophys., 22 (7): 155-159.
- Environmental Protection Agency (EPA) (1990). Pesticide Fact Handbook. Noyes Data Corporation. Park Ridge, New Jersey. 1, 247-251.
- Ferrando, M.D.; Alarcon, V. and Fernandez-Casalderrey, A. (1992). Persistence of some pesticides in the aquatic environment. Bull. Environ. Contam. Toxicol., 48, 747-755.
- Getzin, L.W. and Shanks, C.H.J. (1990). Enhanced degradation of carbofuran in pacific north west soils. J. of Environ. Science and Health, part B, Pesticides Food Contamination and Agricultural Wastes, 25 (4): 433-446.
- Kale, S.P.; Kanthadai, R. and Raghu, K. (1996). Fate of C<sup>14</sup>-carbofuran in soil. Zeitschrift, Fur-Pflanzenernahrung und Bodenkunde, 159 (5): 519-523.
- Lai, K.; Dave, K. I. and Wild, J. R. (1994). Bimetallic binding motifs in organophosphorus hydrolase are important for catalysis and structural organization. J. Biol. Chem., 269, 16579-16584.
- Lan, W.S.; Gu, J.D.; Zhang, J.L. ; Shen, B.C. ; Jiang, H. ; Mulchandani, A. ; Chen, W. and Qiao, C.L.(2006). Coexpression of two detoxifying pesticide-degrading enzymes in a genetically engineered bacterium. International Biodeterioration & Biodegradation, 58(2): 70-76.
- Li, P.C.H.; Swanson, E.J. and Gobas F.A.P.C. (2002). Diazinon and its degradation products in agricultural water courses in British Columbia, Canada. Bull. Environ. Contam. Toxicol., 69, 59-65.
- Lichtenstein, E. P.; Fuhreman, T. W.; Scopes, N. E. A. and Skrentny, R. F. (1967). The method of extraction of diazinon and its metabolites J. Agric. Food Chem., 15, 864-869.

- Lui, A.; Huimin, Z. and Jeffrey, P.O. (2005).** Recent advances in the bioremediation of persistent organic pollutants via biomolecular engineering. *Enzyme and Microbial Technology*, 37(5): 487-496
- Megharaj, M.; Singh, N.; Kookana, R.S.; Naidu, R. and Sethunathan, N. (2003).** Hydrolysis of fenamiphos and its oxidation products by a soil bacterium in pure culture, soil and water. *Applied Microbiology and Biotechnology*, 61 (3): 252-256.
- Michel, J.F.C.; Reddy, C. A. and Forney L. J. (1997).** Biodegradation and bioremediation fate of C<sup>14</sup>-diazinon during the composting of yard trimmings. *J. Environ. Qual.*, 26, 200-205.
- Miguel A. S. and Eugenio V. (2002).** Enzymes involved in the detoxification of organophosphorus, carbamate and pyrethroid insecticides through hydrolysis. *Toxicology Letters*, 128(1-3): 215-228.
- Robert, A.; Yokley, N. S. and Max, W.C. (2000).** Determination of two oxy-pyrimidine metabolites of diazinon in urine by gas liquid chromatography/mass selective detection and liquid chromatography /electrospray-ionization mass spectrometry/mass spectrometry. *J. of AOAC International*. Vol.38 (5): 1229- 1238.
- Scheunert, I.; Mansour, M. and Doerfler, U. (1993).** Fate of pendimethalin, carbofuran and diazinon under abiotic and biotic conditions. *Science of the Total Environment*, 132, 361-369.
- Seyfried, B. (1994).** Degradation of C<sup>14</sup>-diazinon in one soil incubated under various experimental conditions. Project No. (351358). Basel, Switzerland, Novartis Crop Protection (Unpublished report).
- Singh, B. K.; Allan, W. and Denis, J. W. (2005).** Cross-enhancement of accelerated biodegradation of organophosphorus compounds in soils: Dependence on structural similarity of compounds. *Soil Biology & Biochemistry*, 37: 1675-1682.
- Talebi, K. and Walker, C.H. (1993).** A comparative study of carbofuran metabolism in treated and untreated soils. *Pestic. Sci.*, 39(1): 65-69.
- Thalmann, A. (1967).** Über die microbiello Aktivitat und ihr beziehung zu fruchtbarkeits merkmalen einiger acherboden unter besonderer berucksichtigung der dehydrogenase akativitat (TTC reduktion). Biss Gieben Ph.D.Thesis. W. Germany.
- Trabue, S.L.; Ogram, A.V. and Ou, L.T. (2001).** Dynamics of carbofuran degrading microbial communities in soil during three successive annual applications of carbofuran. *Soil Biology and Biochemistry*. 33(1): 75-81.
- Vijay A.K.; Gundi, B. and Reddy, B.R.(2006).** Degradation of monocrotophos in soils. *Chemosphere*, 62(3): 396-403
- Wander, W. S.; Annereinou D.; Gabriel J. M. Z.; Miranda P. A.; Paul C. M. N. and John R. P.(2003).** Competition of a parathion-hydrolyzing *Flavobacterium* with bacteria from ditch water in carbon-, nitrate- and phosphate-limited continuous cultures. *Microbial Ecology*, 43 (1): 45-53.
- Yakovlevsky, K.; Govarhan C.; Grimsley, J.; Wild, J. and Margolin, A. (1997).** Cross-linked crystals of organophosphorus hydrolase: an efficient catalyst for the breakdown of the O.P. compounds. *Proceedings of the 213<sup>th</sup> American Chemical Society National Meeting*. April 13-17. San Francisco. Paper BIOT-0012.

Yueh, L.Y. and Hensley, D.L. (1993). Pesticide effect on acetylene reduction and nodulation by soybean and lima bean. Journal of the American Society for Horticultural Science, 118 (1): 73-76.

Zaghloul, R. A.; Ehsan, A. Hanafy; Rahal, Gh. A. and Iobna, A. Moussa (2003). Bioremediation of polluted soil with carbamate pesticides by *Streptomyces violaceusniger* or/and *Azospirillum brasilense*. Eleventh Conference of Microbiology, Cairo, Egypt, Oct. 12-14, pp: 17.

### كفاءة بعض ميكروبات التربة في تحليل مبيد الديازينون

راشد عبد الفتاح زغلول<sup>١</sup> - أحمد غنيم رحال<sup>٢</sup> - إسمان أحمد حنفى<sup>١</sup> - طلعت محمد الحسينى<sup>١</sup> - هانى محمد عبدالرحمن<sup>١</sup>

١- كلية الزراعة - جامعة بنها

٢- معهد بحوث الأراضي والمياه والبيئة - مركز البحوث الزراعية

أقيمت تجربة بهدف دراسة كفاءة بعض ميكروبات التربة وهى

في *Pseudomonas fluorescens*, *Bacillus polymyxa*, *Streptomyces aureofaciens*

تحليل مبيد الديازينون فى التربة المعقمة وغير المعقمة وكذلك فى التربة المنزرعة بنباتات الطماطم. ولقد أوضحت نتائج هذه الدراسة ما يلى:-

كان معدل إختفاء مبيد الديازينون من التربة أعلى فى حالة التربة الغير معقمة مقارنة بمعدل إختفاؤه من التربة المعقمة. وكذلك أوضحت النتائج أن تلقيح التربة بمخلوط السلالات الميكروبية قد أدى إلى حدوث تحلل أسرع للمبيد مقارنة بالتلقيح الفردى بأى من السلالات. وعندما زرعت التربة بنباتات الطماطم وتم تلقيحها بأحد الميكروبات سالفة الذكر وجد أن مبيد الديازينون قد تحلل بمعدل أسرع مقارنة بالتربة غير المنزرعة.

و من أهم المركبات التى نتجت من تحلل مبيد الديازينون مركب ٣،١ داي ميثيل -٢- نيتروبنزين و داي إيثيل فوسفات و ٢- أيزوبروبيل -٦- ميثيل -٤- هيدروكسى بيريميدين و ٢- (١- هيدروكسى -١- ميثيل) -٦- ميثيل بيريميدين.

و إتضح من نتائج هذه الدراسة أن الأراضى المصرية غنية بالميكروبات ذات القدرة على تحليل مثل هذه المبيدات ولذلك يمكن إعداد لقاحات من هذه الميكروبات لتلقيح التربة بعد إضافة المبيدات و القيام بدورها فى القضاء على الافات و ذلك للتخلص من آثارها المتبقية وللحفاظ على التربة الزراعية من التلوث.