

Morphological and Molecular Genetical Studies on Restoring Ability for WA Type CMS Lines in Rice (*Oryza sativa* L.)

M. E. El-Denary¹, A. E. Draz², S. A. Dora³ and W. F. Ghidan²

1- Department of genetics, Faculty of Agriculture, Tanta University

2- Rice Research and Training Center, Sakha, Kafr El-Sheikh

3- Department of genetics, Faculty of Agriculture, Kafr El-Sheikh University

ABSTRACT

Three cytoplasmic male sterile lines wild abortive (WA); IR 58025 A, IR 69625 A and IR 70368 A were tested with nine different testers for some morphological, biochemical and molecular studies. A total of 27 F₁ crosses were made and harvested separately for each cross. The results appeared a wide range of variations was recorded for yield and related traits among the parental lines with their crosses. This means that the direction of heterosis was negative for some traits such as days to heading. On the other hand, positive heterosis was recorded for the majority of remained characters. With respect to general combining ability effects, the CMS line IR 69625 A was the best combiner for grain yield / plant. Also, for straw weight, IR 58025 A and IR 70368 A recorded significant and highly significant negative differences, respectively. But for specific combining ability effects, two crosses IR 58025 A x IRGA 370- 42- 1- 1F- C-1 and IR 69625 A x Giza 175 recorded significant and positive values for No. of tillers. In the peroxidase zymogram, results showed that, sub-patterns were observed among the different tested genotypes; these have specific bands which could be used to distinguish among the three CMS lines. While in respect of the esterase zymogram, a complex of sub-patterns was observed among the different tested genotypes. Total soluble proteins differences were detected in the CMS lines IR 58025 A and IR 69625 A and their hybrids. Generally, some bands were missing in the CMS lines,

while present in most of the hybrids, as well as, the testers. These bands might be derived from the testers. From DNA studies, a level of molecular polymorphism was detected. For example, band no. 7 was found in many testers and F_1 S in the two CMS lines; IR 58025 A and IR 70368 A, in spite of its missing in the CMS line itself. Therefore, further experiments could be applied for studying the correlation of this band with the restoring ability of CMS lines.

INTRODUCTION

Rice (*Oryza sativa*, L.) is considered as one of the most important cereal crops not only in Egypt, but also all over the world. In Egypt the yearly area cultivated annually with rice is almost more than one fifth of the total area.

Hybrid varieties developed by making use of WA CMS account for approximately 99% of the hybrid rice produced in China. These varieties have significantly outyielded conventional pure line cultivars. The WA CMS system has also been used extensively in hybrid rice breeding programs of other countries to produce hybrids in various genetic backgrounds (Virmani, 1994).

The wild abortive (WA) type of CMS source is used extensively for hybrid rice seed production, in China, as well as, in tropical rice-growing areas.

Many scientists in China have studied the relationship between isozymes and heterosis in rice. They found that the hybrid combination whose F_1 had dominant complementary bands for the esterase isoenzyme was closely associated with expression of heterosis Kato *et al.* (1994), Li *et al.*, (1996), Banumathy *et al.* (2003) and Chakraborty *et al.* (2002). The complementary enzyme bands may be used as one of the biochemical indicators for predicting heterosis.

Molecular marker studies have also been employed to determine the map positions of the fertility restorer genes. Zhang *et al.* (1996) surveyed DNA

polymorphism using randomly amplified polymorphic DNA (RAPD) analysis in a set of near-isogenic lines (NILs) developed by introgressing of the fertility restorer genes from IR 24 into the genetic background of Zhenshan 97 (Zhang *et al.*, 1997).

This study was undertaken to achieve the following objectives:

- i. Estimation of heterosis and combining ability for morphological, yield and its components in parental lines and their F₁ hybrids.
- ii. Study of the relationship between isozymes and total soluble protein and restoring ability in rice.
- iii. Using RAPD markers to identify the relationship between morphological and yield characters and molecular markers.

MATERIALS AND METHODS

This study was carried out at the experimental farm and Biotechnology laboratory of the Rice Research and Training Center (RRTC), Sakha, Kafr EL-Sheikh, Egypt and laboratories of Genetics Department, Faculty of Agriculture, Kafr EL-Sheikh University, during the two growing seasons 2002 and 2003. Three cytoplasmic male sterile lines; wild abortive (WA) IR 58025 A, IR 69625 A and IR 70368 A were tested with nine testers; Giza 175, Giza 178, Giza 181, Giza 182, IRGA 370-2-1 - 1F-C-1, WAB 326 - B-B-15, IR 69312-14-2-2-2, CT 9685-14-M-1-2-3-2P-M-1 and IET 12515.

Parents and hybrids were used to study the genetic analysis of yield and its components. These components were: plant weight, panicle length, grain yield per plant, 1000 grain weight and harvest index %. Some morphological traits such as days to flowering, plant height and number of tillers per plant were studied.

Biochemical analysis:

For isozymes detection, the method of Scandalios (1969) was used for the detection of esterase, and the method described by Tu *et al.* (1968) was used for the detection of peroxidase.

PCR based RAPD Analysis:

DNA isolation and purification was carried out using CTAB (Cetyl-tetramethyl ammonium bromide) method, (Murray and Thompson 1980).

DNA quantification:

DNA was quantified using gel quantification method, in which the samples were loaded on 0.8% agarose gel in 0.5 x TAE buffer and using diluted λ genomic DNA as standard. The intensity of individual sample was compared with a range of known amounts of λ DNA (25, 50, 75 and 100 ng). According to the sample intensity, samples were diluted, and again loaded on 0.8% gels till all of the samples finally reached around 50 ng/ μ l in a uniform manner.

The five 10-mer oligonucleotide random primers used in this study were as follows:

Primer	Sequence
RBL.4	5- AGC TTG CCC C- 3
RBL.5	5- GAA GGC GGT G- 3
RBL.6	5- GGG GTA CCG T- 3
RBL.7	5- CGT CAA GCC C- 3
RBL.8	5- GAG GCG GCT T- 3

RBL, Rice Biotechnology Laboratory (RRTC)

PCR amplification condition used in this part in 20 μ l PCR volume containing 1.0 μ l (50ng template DNA), 0.2 μ l dNTPs, (10 mM), 1.6 μ l Mg Cl₂ (25 mM), 2.0 μ l 10X buffer (10 mM tris, pH 8.0, 50mM KCl and 50 mM

ammonium sulphate), 4.0 μ l primer (15 pmole), 0.1 μ l taq polymerase (10u/ μ l). The volume was brought up to 20 μ l by autoclaved double distilled H₂O.

Statistical Analysis:

Twenty-seven hybrid combinations were produced from three lines and nine diverse testers. The data were subjected to analysis of variance for randomized complete block design as suggested by Panse and Sukhatme (1954) and analysis of variance for line x tester crossing according to Kempthorne (1957). All the present data were subjected to arcsine transformation before analysis.

Estimation of combining ability:

Assuming there are (L) lines and (t) testers and data are obtained by plot mean, two methods of analysis according to Kempthorne (1957) and Virmani *et al.* (1997) were used.

Estimation of heterosis:

$$\text{Heterosis over the mean parent \% } (\overline{MP}) = \frac{F_1 - \overline{MP}}{\overline{MP}} \times 100$$

Where: F₁ = mean value of the first generation and \overline{MP} = mid. parent value.

RESULTS AND DISCUSSION

The development of hybrid rice technology and adoptions of hybrid rice cultivars to Egyptian environments offer one approach to the problem of matching food supply to expected demand.

Mean performance:

A wide range of differences among the parental lines and their crosses are shown in Tables (1 and 2), and for each trait there were significant differences. The data in Table (1) showed that, the mean values of the testers

Table (1): Mean values for some studied characters for parents.

Testers	Days to heading (day)	Plant height (cm)	No. of tillers	Panicle length (cm)	Grain yield/plant (g)	Straw weight/plant (g)	Harvest index (%)	1000- grain weight (g)
Giza 175	95.00	95.00	19.00	22.02	23.72	47.40	41.36	17.67
Giza 178	100.00	96.76	16.33	20.27	15.08	33.05	35.96	19.87
Giza 181	112.00	97.67	20.00	24.80	38.04	59.73	40.80	23.83
Giza 182	92.00	70.00	14.67	22.40	62.09	67.70	59.21	25.17
IRGA 370-42-1-1F-C-1	100.00	97.13	20.33	19.53	25.89	65.15	27.12	22.83
WAB 326-B-B-15	94.00	109.87	10.67	20.60	28.12	54.84	35.46	24.13
CT9685-14-M-1-2-3-2F-M-1	102.00	103.47	19.00	22.93	41.46	59.63	42.08	27.00
IET 12515	104.00	103.07	18.33	20.07	21.36	51.95	32.96	22.23
IR 69312-14-2-2-2	99.00	92.47	10.33	21.93	19.40	24.72	66.90	26.90
Mean	99.78	96.15	16.52	21.62	30.57	51.57	42.43	23.29
Lines	Days to heading (day)	Plant height (cm)	No. of tillers	Panicle length (cm)	Grain yield/plant (g)	Straw weight/plant (g)	Harvest index (%)	1000- grain weight (g)
IR 58025 A	107	87.27	24.67	21.67	36.21	64.34	36.01	20.50
IR 69625 A	100	91.33	26.00	22.33	46.30	72.87	38.84	24.20
IR 70368 A	101	84.27	18.33	23.27	41.51	54.49	43.24	25.40
Mean	102.67	87.62	23.00	22.42	41.34	63.90	39.36	23.37
L.S.D at 5%	3.14	15.62	3.16	1.34	6.20	6.94	8.08	1.92
L.S.D at 1%	4.18	20.77	4.20	1.78	9.23	9.23	10.75	2.55

Table (2): Mean values for some studied characters for the crosses.

No	Crosses	Days to heading (day)	Plant height (cm)	No. of tillers	Panicle length (cm)	Grain yield /plant (g)	Straw weight/plant (g)	Harvest index	1000-grain weight (g)
1	IR58025A x Giza 175	94.03	102.47	26.00	23.60	44.07	61.00	38.99	19.63
2	IR58025A x Giza 178	95.33	107.93	18.67	21.13	52.59	70.61	57.97	19.43
3	IR58025A x Giza 181	107.67	108.93	18.00	23.73	59.09	103.46	38.33	18.00
4	IR58025A x Giza 182	95.67	95.40	21.67	22.73	48.03	57.93	60.94	23.87
5	IR58025A x IRGA 370-42-1-1F-C-1	97.33	91.27	37.67	25.00	33.47	58.40	39.24	25.27
6	IR58025A x WAB 326-B-B-1	90.33	122.80	33.00	25.80	26.26	84.50	23.65	29.23
7	IR58025A x CT9685-14-M-1-2-3-2P-M-1	101.00	103.07	19.33	22.33	22.35	53.95	37.47	18.10
8	IR58025A x IET 12515	101.03	96.33	21.33	23.93	27.70	54.15	36.63	25.23
9	IR58025A x IR 69312-14-2-2-2	101.00	96.07	30.00	25.00	13.61	121.67	10.10	8.00
10	IR 69625 A x Giza 175	94.67	101.33	36.33	22.47	89.59	126.95	39.53	23.00
11	IR 69625 A x Giza 178	100.33	108.53	25.67	24.80	47.41	66.03	51.11	24.27
12	IR 69625 A x Giza 181	94.33	96.07	22.33	21.60	42.02	48.26	50.17	24.93
13	IR 69625 A x Giza 182	94.67	100.40	28.00	22.40	57.96	63.12	55.13	27.37
14	IR 69625 A x IRGA 370-42-1-1F-C-1	94.33	97.60	25.00	21.87	57.38	67.12	42.36	22.73
15	IR 69625 A x WAB 326-B-B-15	93.03	133.73	19.33	22.00	24.23	126.03	15.79	27.10
16	IR 69625 A x CT9685-14-M-1-2-3-2P-M-1	95.33	100.33	20.33	23.07	44.97	83.56	30.18	25.10
17	IR 69625 A x IET 12515	99.33	105.80	18.67	21.73	20.34	149.22	12.28	20.13
18	IR 69625 A x IR 69312-14-2-2-2	99.33	99.13	24.33	22.33	12.73	129.16	8.80	11.77
19	IR 70368 A x Giza 175	94.03	98.20	16.67	23.53	49.70	53.39	54.50	23.13
20	IR 70368 A x Giza 178	101.33	104.73	24.67	23.47	50.26	56.84	50.06	25.10
21	IR 70368 A x Giza 181	103.03	96.27	17.67	22.40	45.56	50.44	49.60	26.22
22	IR 70368 A x Giza 182	91.33	100.13	15.67	22.80	41.72	55.08	44.47	26.93
23	IR 70368 A x IRGA 370-42-1-1F-C-1	93.67	98.20	18.33	23.80	25.93	53.63	48.68	25.63
24	IR 70368 A x WAB 326-B-B-15	89.03	130.60	22.00	22.73	45.54	74.82	41.79	28.47
25	IR 70368 A x CT9685-14-M-1-2-3-2P-M-1	97.67	103.80	19.00	22.00	44.34	65.70	50.29	26.53
26	IR 70368 A x IET 12515	102.03	88.73	24.00	23.00	11.55	79.00	13.00	10.50
27	IR 70368 A x IR 69312-14-2-2-2	101.67	93.00	24.67	23.67	12.76	122.83	10.14	8.03
	Mean.	97.13	102.99	23.27	23.11	38.93	79.15	37.45	21.99

were (99.78, 96.15, 30.57, 42.43 and 23.29) for days to heading, plant height, grain yield per plant, harvest index and 1000-grain weight. But the mean performances of their crosses for these traits were 97.13, 102.99, 38.93, 37.45 and 21.99, respectively (as shown in Table 2).

For the CMS lines, the results indicated that the IR 69625 A gave the desirable values for days to heading (100), No. of tillers /plant (26.00) and grain yield/plant (46.30). While, the IR 70368 A gave the desirable values (84.27, 23.27, 54.49, 43.24 and 25.40) for plant height, panicle length, straw weight, harvest index and 1000-grain weight respectively.

Analysis of variance:

Results in Table (3) showed that analysis of variance revealed highly significant differences among the 39 genotypes (27 hybrids, 9 testers and 3 CMS lines) tested for all the studied parameters. The parental lines and the hybrids showed highly significant differences for some traits. Parents v.s crosses mean squares indicated that the average heterosis were significant in all crosses for all studied traits.

Heterosis :

Evaluation based on the standard heterosis revealed that 15 hybrids recorded significantly negative standard heterosis for early heading date. These findings indicated that heterosis effects can be used to get earliness in rice hybrids. The highest estimates of the 15 hybrids were recorded for the crosses IR 70368 A x WAB 326-B-B15, IR 58025 A x WAB 326-B-B-15 and IR 70368 x Giza 182 which recorded -11.00%, -10.00% and -9.00%, respectively (Table 4). While 21 cross of the 27 crosses recorded significantly positive standard heterosis for grain yield/plant (g). The highest evaluates were IR 69625A x Giza 175, IR 58025A x Giza 181 and IR 69625A x Giza 182 which recorded 494.10, 291.84 and 284.42, respectively (Table 4 cont).

Table (3): Mean square values of lines, testers and their crosses for some studied characters.

Source of variance	d . f	Days to heading (day)	Plant height (cm)	No. of tillers	Panicle length (cm)	Grain yield / plant (g)	Straw weight/plant (g)	Harvest index	1000-grain weight (g)
Replication	2	0.718 ^{ns}	75.820 ^{ns}	2.872 ^{ns}	3.176 [*]	4.227 ^{ns}	44.104 ^{ns}	9.427 ^{ns}	0.923 ^{ns}
Genotype	38	72.541 ^{**}	382.926 ^{**}	104.305 ^{**}	5.768 ^{**}	853.652 ^{**}	2439.752 ^{**}	661.287 ^{**}	83.988 ^{**}
Parents	11	10.212 ^{**}	35.777 ^{ns}	7.465 [*]	0.761 ^{ns}	60.728 ^{**}	66.040 ^{**}	40.802 ^{ns}	2.697 [*]
Crosses	26	6.613 [*]	38.460 ^{ns}	10.974 ^{**}	0.439 ^{ns}	109.509 ^{**}	304.399 ^{**}	88.259 ^{**}	12.311 ^{**}
Parents V.S Crosses	1	2472.280 ^{**}	13157.690 ^{**}	3596.154 ^{**}	199.406 ^{**}	28923.336 ^{**}	84069.772 ^{**}	22385.363 ^{**}	2841.783 ^{**}
Lines (GCA)	2	3.346 ^{ns}	8.127 ^{ns}	20.200 ^{**}	1.338 ^{ns}	59.402 [*]	628.397 ^{**}	31.36 ^{ns}	3.766 ^{ns}
Testers (GCA)	8	14.077 ^{**}	108.032 ^{ns}	8.788 [*]	0.166 ^{ns}	238.410 ^{**}	477.989 ^{**}	227.887 ^{**}	29.911 ^{**}
Lines x Testers (SCA)	16	3.290 ^{ns}	7.466 ^{ns}	10.914 ^{**}	0.463 ^{ns}	51.335 ^{**}	177.104 ^{**}	25.557 ^{ns}	4.579 ^{**}
Error	76	3.727	91.465	3.731	0.670	14.447	18.011	24.424	1.388

^{*}, ^{**} : Significant at 5% and 1% , respectively.

^{ns} : Not significant .

Table (4): Estimates of standard heterosis (S.H), better parent ($B\bar{P}$) and mid - parent ($M\bar{P}$) for some studied characters.

No	Crosses	Days to heading			Plant height (cm)			No. of tillers			Panicle length		
		S.H%	B.P.%	M.P.%	S.H%	B.P.%	M.P.%	S.H%	B.P.%	M.P.%	S.H%	B.P.%	M.P.%
1	IR58025A x Giza 175	-6.90**	-1.05**	-6.93*	6.90**	17.42**	12.44**	59.22**	5.93**	19.07*	16.43*	6.93*	7.91*
2	IR58025A x Giza 178	-5.00**	-5.00**	-8.21**	11.66**	23.67**	17.35**	14.27**	-24.32**	-8.93**	4.29**	-2.49**	0.76**
3	IR58025A x Giza 181	8.00**	0.93**	-1.37**	12.69**	24.82**	17.80**	10.23**	27.04**	-19.41**	17.12**	-4.31**	2.13**
4	IR38025A x Giza 182	-4.00**	4.35**	-3.52**	-1.31**	36.29**	21.32**	32.64**	-12.16**	10.17**	12.19**	1.47**	3.15**
5	IR58025A x IRGA 370-42-1-IF-C-1	-3.00**	-3.00**	-6.28**	-5.59**	4.58**	-1.01**	130.62**	52.70**	67.42**	23.33**	15.37**	21.36**
6	IR58025A x WAB 326-B-B-15	-10.06**	-4.26**	-10.45**	27.03**	40.71**	24.58**	102.08**	33.77**	86.76**	27.28**	19.06**	22.07**
7	IR58025A x CT9685-14-M-1-2-3-2P-M-1	1.00**	-0.98**	-3.35**	6.62**	18.10**	8.57**	18.37**	-21.65**	-11.47**	15.15**	1.74**	4.62**
8	IR58025A x IET 12515	1.00**	-2.88**	-4.27**	-0.34**	10.38**	1.22**	30.62**	-13.54**	0.79**	18.11**	10.43**	14.66**
9	IR58025A x IR 69312-14-2-2-2	1.00**	2.02**	-1.94**	-0.62**	10.08**	6.90**	83.71**	21.61**	71.43**	23.33**	14.00**	14.68**
10	IR 69625 A x Giza 175	-5.00**	0.00**	-2.56**	4.83**	10.95**	8.76**	122.47**	39.73**	61.47**	10.85**	0.63**	1.22**
11	IR 69625 A x Giza 178	0.00**	0.00**	0.00**	12.28**	18.83**	15.46**	57.13**	-1.27**	21.29**	22.35**	11.06**	16.43**
12	IR 69625 A x Giza 181	-6.00**	-6.00**	-11.32**	-0.62**	5.19**	1.66**	36.47**	-1.41**	-2.91**	6.65**	-12.90**	-8.34**
13	IR 69625 A x Giza 182	-5.00**	3.26**	-1.04**	3.86**	9.93**	24.47**	71.46**	7.69**	37.69**	10.51**	0.00**	0.16**
14	IR 69625 A x IRGA 370-42-1-IF-C-1	-6.00**	-6.00**	-6.00**	0.96**	6.87**	5.58**	53.09**	-3.85**	7.92**	7.89**	-2.06**	4.49**
15	IR 69625 A x WAB 326-B-B-15	-7.00**	-1.06**	-4.12**	38.35**	46.42**	52.93**	18.37**	-25.65**	5.43**	8.53**	-1.48**	2.49**
16	IR 69625 A x CT9685-14-M-1-2-3-2P-M-1	-5.00**	-5.00**	-5.94**	3.30**	9.85**	3.01**	24.49**	-21.81**	-9.64**	13.81**	0.61**	1.94**
17	IR 69625 A x IET 12515	-1.00**	-1.00**	-2.94**	9.44**	15.84**	8.85**	14.27**	-28.19**	-15.77**	7.25**	-2.69**	2.50**
18	IR 69625 A x IR 69312-14-2-2-2	-1.69**	0.00**	-0.50**	2.56**	8.54**	7.87**	48.99**	-6.42**	33.94**	10.21**	0.00**	0.90**
19	IR 70368 A x Giza 175	-6.00**	-1.05**	-4.08**	1.58**	16.53**	9.56**	2.02**	-12.26**	-10.69**	16.13**	1.12**	3.79**
20	IR 70368 A x Giza 178	1.00**	1.00**	0.50**	8.35**	24.28**	15.76**	51.01**	34.59**	42.35**	15.79**	0.86**	7.81**
21	IR 70368 A x Giza 181	3.00**	1.98**	-3.29**	-0.41**	14.24**	5.83**	8.14**	-11.65**	-7.80**	10.51**	-9.68**	-6.80**
22	IR 70368 A x Giza 182	-9.00**	-1.09**	-5.70**	3.59**	43.04**	29.81**	-4.10**	-14.51**	-5.03**	12.48**	-2.02**	-0.15**
23	IR 70368 A x IRGA 370-42-1-IF-C-1	-5.00**	-6.00**	-6.47**	1.58**	16.53**	8.27**	12.25**	-9.84**	-5.17**	17.41**	2.28**	11.21**
24	IR 70368 A x WAB 326-B-B-15	-11.00**	-5.32**	-8.72**	35.10**	54.98**	34.54**	34.72**	26.02**	51.72**	12.19**	-2.32**	3.62**
25	IR 70368 A x CT9685-14-M-1-2-3-2P-M-1	-2.00**	-2.97**	-3.45**	7.38**	23.18**	10.58**	16.35**	0.00**	1.79**	8.53**	-5.46**	-4.76**
26	IR 70368 A x IET 12515	2.00**	0.99**	-0.49**	-8.20**	5.29**	-5.27**	46.97**	30.93**	30.93**	13.47**	-1.16**	6.14**
27	IR 70368 A x IR 69312-14-2-2-2	2.00**	3.03**	2.00**	-3.80**	10.36**	5.24**	51.01**	34.59**	72.16**	16.77**	1.72**	4.73**
	LSM (S.E.)	3.08	3.14	2.74	16.55	15.02	13.43	5.13	33.15	2.74	3.37	1.34	1.15
	LSD (5%)	4.10	4.18	3.61	20.63	18.37	17.98	4.12	4.28	3.03	3.72	1.78	1.54

*, **: Significant at 5% and 1%, respectively.
ns : Not significant.

Table (4): Cont. Estimates of standard heterosis (S.H), better parent (BP) and Mid - parent (MP) for some studied characters.

No	Crosses	Grain yield (g/plant)			Straw yield (g/plant)			H.C. test index					
		S.H	BP	MP	S.H	BP	MP	S.H	BP	MP	S.H	BP	MP
1	IR58025 A x Giza 175	192.24 ^{**}	21.71 ^{**}	47.05 ^{**}	84.57 ^{**}	28.69 ^{**}	9.18 ^{**}	8.43 ^{**}	-5.73 ^{**}	0.77 ^{**}	-1.16 ^{**}	-4.24 ^{**}	2.83 ^{**}
2	IR58025 A x Giza 178	248.74 ^{**}	45.24 ^{**}	15.03 ^{**}	113.65 ^{**}	113.65 ^{**}	44.99 ^{**}	61.21 ^{**}	60.98 ^{**}	61.07 ^{**}	-2.16 ^{**}	-5.22 ^{**}	-3.76 ^{**}
3	IR58025 A x Giza 181	291.64 ^{**}	55.34 ^{**}	59.14 ^{**}	213.07 ^{**}	73.21 ^{**}	66.76 ^{**}	6.39 ^{**}	-6.05 ^{**}	-0.21 ^{**}	-9.41 ^{**}	-24.46 ^{**}	-18.81 ^{**}
4	IR58025 A x Giza 182	218.50 ^{**}	-22.64 ^{**}	-2.28 ^{**}	75.31 ^{**}	-9.95 ^{**}	-12.25 ^{**}	69.47 ^{**}	2.92 ^{**}	28.00 ^{**}	20.13 ^{**}	-5.16 ^{**}	4.51 ^{**}
5	IR58025 A x IRGA 370-42-1-1F-C-1	121.95 ^{**}	-7.57 ^{**}	7.79 ^{**}	76.70 ^{**}	-9.23 ^{**}	-9.81 ^{**}	9.12 ^{**}	8.97 ^{**}	24.30 ^{**}	27.18 ^{**}	10.69 ^{**}	16.61 ^{**}
6	IR58025 A x WAB 326-B-B-15	74.20 ^{**}	-27.48 ^{**}	-18.37 ^{**}	155.67 ^{**}	54.08 ^{**}	41.80 ^{**}	-34.23 ^{**}	-34.32 ^{**}	-33.83 ^{**}	47.16 ^{**}	21.14 ^{**}	30.96 ^{**}
7	IR58025 A x CT9685-14-M-1-2-3-2P-M-1	48.21 ^{**}	46.09 ^{**}	-42.46 ^{**}	63.24 ^{**}	-9.53 ^{**}	-12.97 ^{**}	4.20 ^{**}	-10.96 ^{**}	-4.05 ^{**}	-8.91 ^{**}	-32.96 ^{**}	-24.11 ^{**}
8	IR58025 A x IET 12515	83.69 ^{**}	-23.50 ^{**}	-3.79 ^{**}	63.84 ^{**}	4.23 ^{**}	-6.88 ^{**}	1.86 ^{**}	1.72 ^{**}	6.20 ^{**}	26.98 ^{**}	13.50 ^{**}	18.01 ^{**}
9	IR58025 A x IR 69312-14-2-2-2	-9.75 ^{**}	-62.41 ^{**}	-51.06 ^{**}	268.41 ^{**}	392.56 ^{**}	173.43 ^{**}	-71.91 ^{**}	-84.90 ^{**}	-80.37 ^{**}	-59.74 ^{**}	-70.26 ^{**}	-66.24 ^{**}
10	IR 69625 A x Giza 175	464.16 ^{**}	93.50 ^{**}	155.90 ^{**}	284.11 ^{**}	167.83 ^{**}	111.09 ^{**}	9.39 ^{**}	-4.42 ^{**}	-1.42 ^{**}	15.75 ^{**}	-4.96 ^{**}	9.84 ^{**}
11	IR 69625 A x Giza 178	214.39 ^{**}	2.40 ^{**}	54.48 ^{**}	99.79 ^{**}	99.79 ^{**}	24.68 ^{**}	42.13 ^{**}	31.59 ^{**}	36.66 ^{**}	22.14 ^{**}	0.29 ^{**}	10.12 ^{**}
12	IR 69625 A x Giza 181	178.65 ^{**}	-9.24 ^{**}	-0.36 ^{**}	46.05 ^{**}	-19.20 ^{**}	-27.21 ^{**}	39.52 ^{**}	22.97 ^{**}	25.99 ^{**}	25.52 ^{**}	3.02 ^{**}	3.79 ^{**}
13	IR 69625 A x Giza 182	284.42 ^{**}	-6.65 ^{**}	6.94 ^{**}	90.98 ^{**}	-6.77 ^{**}	-10.20 ^{**}	53.31 ^{**}	-6.89 ^{**}	12.44 ^{**}	37.75 ^{**}	8.74 ^{**}	10.83 ^{**}
14	IR 69625 A x IRGA 370-42-1-1F-C-1	280.50 ^{**}	23.93 ^{**}	58.95 ^{**}	103.09 ^{**}	3.02 ^{**}	-2.74 ^{**}	17.80 ^{**}	9.66 ^{**}	28.44 ^{**}	14.44 ^{**}	-6.07 ^{**}	-3.36 ^{**}
15	IR 69625 A x WAB 326-B-B-15	60.68 ^{**}	-47.67 ^{**}	-34.88 ^{**}	281.36 ^{**}	129.81 ^{**}	97.35 ^{**}	-56.09 ^{**}	-59.35 ^{**}	-57.50 ^{**}	36.39 ^{**}	11.98 ^{**}	12.12 ^{**}
16	IR 69625 A x CT9685-14-M-1-2-3-2P-M-1	198.28 ^{**}	-2.87 ^{**}	2.48 ^{**}	152.83 ^{**}	40.13 ^{**}	26.13 ^{**}	-16.07 ^{**}	-29.28 ^{**}	-25.41 ^{**}	26.32 ^{**}	-7.04 ^{**}	-1.95 ^{**}
17	IR 69625 A x IET 12515	34.95 ^{**}	-56.07 ^{**}	-39.88 ^{**}	351.50 ^{**}	187.24 ^{**}	139.10 ^{**}	-65.85 ^{**}	-68.38 ^{**}	-65.79 ^{**}	1.36 ^{**}	-16.82 ^{**}	-13.31 ^{**}
18	IR 69625 A x IR 69312-14-2-2-2	-15.58 ^{**}	-72.51 ^{**}	-61.25 ^{**}	290.80 ^{**}	422.49 ^{**}	164.67 ^{**}	-75.53 ^{**}	-86.85 ^{**}	-83.36 ^{**}	-40.76 ^{**}	-56.24 ^{**}	-53.93 ^{**}
19	IR 70368 A x Giza 175	229.58 ^{**}	19.73 ^{**}	52.36 ^{**}	61.57 ^{**}	12.64 ^{**}	4.79 ^{**}	51.56 ^{**}	26.04 ^{**}	28.84 ^{**}	16.46 ^{**}	-8.94 ^{**}	7.38 ^{**}
20	IR 70368 A x Giza 178	233.29 ^{**}	21.08 ^{**}	77.60 ^{**}	71.98 ^{**}	71.98 ^{**}	29.86 ^{**}	39.21 ^{**}	15.77 ^{**}	26.41 ^{**}	26.32 ^{**}	-1.18 ^{**}	10.87 ^{**}
21	IR 70368 A x Giza 181	202.19 ^{**}	9.76 ^{**}	14.53 ^{**}	52.65 ^{**}	-7.43 ^{**}	-11.68 ^{**}	37.93 ^{**}	14.72 ^{**}	18.04 ^{**}	32.06 ^{**}	3.27 ^{**}	6.54 ^{**}
22	IR 70368 A x Giza 182	176.72 ^{**}	-32.81 ^{**}	-19.46 ^{**}	66.69 ^{**}	1.08 ^{**}	-9.35 ^{**}	23.67 ^{**}	-24.39 ^{**}	-13.20 ^{**}	35.58 ^{**}	6.02 ^{**}	6.48 ^{**}
23	IR 70368 A x IRGA 370-42-1-1F-C-1	72.02 ^{**}	-37.53 ^{**}	-23.06 ^{**}	62.27 ^{**}	-1.58 ^{**}	-10.35 ^{**}	35.37 ^{**}	12.58 ^{**}	38.37 ^{**}	29.04 ^{**}	0.91 ^{**}	6.26 ^{**}
24	IR 70368 A x WAB 326-B-B-15	202.06 ^{**}	9.71 ^{**}	30.79 ^{**}	126.41 ^{**}	37.31 ^{**}	36.86 ^{**}	16.21 ^{**}	-3.35 ^{**}	6.20 ^{**}	43.28 ^{**}	12.09 ^{**}	14.49 ^{**}
25	IR 70368 A x CT9685-14-M-1-2-3-2P-M-1	194.03 ^{**}	6.82 ^{**}	6.87 ^{**}	98.82 ^{**}	20.57 ^{**}	15.14 ^{**}	39.85 ^{**}	16.30 ^{**}	17.89 ^{**}	33.57 ^{**}	-1.74 ^{**}	1.26 ^{**}
26	IR 70368 A x IET 12515	-23.41 ^{**}	-72.18 ^{**}	-63.26 ^{**}	139.03 ^{**}	52.07 ^{**}	48.44 ^{**}	-63.85 ^{**}	-69.94 ^{**}	-65.88 ^{**}	-47.16 ^{**}	-58.66 ^{**}	-55.92 ^{**}
27	IR 70368 A x IR 69312-14-2-2-2	-15.32 ^{**}	-69.26 ^{**}	-58.11 ^{**}	271.65 ^{**}	396.89 ^{**}	210.10 ^{**}	-71.80 ^{**}	-84.94 ^{**}	-81.59 ^{**}	-59.54 ^{**}	-70.15 ^{**}	-69.29 ^{**}
	L.S.D at 5%	6.18	3.87	5.38	6.30	6.04	6.30	8.16	9.06	9.38	10.15	10.15	10.15
	L.S.D at 1%	8.20	4.19	7.16	9.15	9.23	7.98	10.86	10.75	10.75	11.75	11.75	11.75

*, ** : Significant at 5% and 1% , respectively.

ns : Not significant .

Patel *et al.* (1994) reported that, the heterosis effects were high for grain yield / plant and similar reports have been made by Yolanda and Das (1996) for days to flowering. The results obtained by Panwar *et al.* (1998), Ramesha *et al.* (1998), Vishwakarma *et al.* (1999), Jin *et al.* (2001), Bhave *et al.* (2002), Singh *et al.* (2002) and Banumathy *et al.* (2003) were in agreement for the other traits with those obtained in this investigation.

General combining ability effects (GCA):

Among three lines tested (Table 5 A), IR 69625 A was the best combiner for grain yield / plant. Also IR 58025 A and IR 70368 A recorded significant and highly significant negative differences, respectively for straw weight.

The results in (Table 5 B) suggested that, the variety WAB 326-B-B-15 was the best general combiner for days to heading and 1000-grain weight, Giza 175 was a good combiner for grain yield / plant, Giza 182 was a good combiner for straw weight, grain yield / plant, harvest index and 1000-grain weight and finally Giza 178 was a good combiner for grain yield / plant, straw weight and harvest index.

Specific combining ability effects (SCA):

The results summarized in (Table 6) suggested that the IR 58025 A x WAB 326-B-B-15, IR 69625 A x Giza 181 and IR 70368 A x WAB 326-B-B-15 were good combiners for days to heading based on the standard heterosis and also the selection of WAB 326-B-B-15 as a tester was good donor to give significant and negative heterosis. Three crosses exhibited significant and highly significant and positive standard heterosis over the standard parent, the

Table (5 A): General combining ability (GCA) effects of lines for some studied characters.

Lines	Days to heading (day)	Plant height (cm)	No. of tillers	Panicle length (cm)	Grain yield/plant (g)	Straw weight/plant (g)	Harvest index	1000-grain weight (g)
IR 58025 A	0.34 ^{ns}	-0.10 ^{ns}	0.60 ^{ns}	0.24 ^{ns}	-0.86 ^{ns}	-1.72 [*]	0.24 ^{ns}	-0.41 ^{ns}
IR 69625 A	-0.36 ^{ns}	0.59 ^{ns}	0.39 ^{ns}	-0.21 ^{ns}	1.71 ^{**}	5.45 ^{**}	-1.17 ^{ns}	0.31 ^{ns}
IR 70368 A	0.01 ^{ns}	-0.49 ^{ns}	-0.99 ^{**}	-0.02 ^{ns}	-0.86 ^{ns}	-3.72 ^{**}	0.95 ^{ns}	0.10 ^{ns}
L.S.D at 5%	0.74	3.68	0.74	0.32	1.46	1.64	1.90	0.46
L.S.D at 1%	0.98	4.89	0.98	0.43	1.94	2.18	2.53	0.61

Table (5 B): General combining ability (GCA) effects of testers for some studied characters.

Testers	Days to heading (day)	Plant height (cm)	No. of tillers	Panicle length (cm)	Grain yield/plant (g)	Straw weight/plant (g)	Harvest index	1000-grain weight (g)
Giza 175	-0.92 ^{ns}	-0.77 ^{ns}	1.02 ^{ns}	0.03 ^{ns}	7.39 ^{**}	0.44 ^{ns}	2.30 ^{ns}	-0.02 ^{ns}
Giza 178	0.53 ^{ns}	1.36 ^{ns}	-0.09 ^{ns}	0.01 ^{ns}	3.72 ^{**}	-4.88 ^{**}	5.20 ^{**}	0.31 ^{ns}
Giza 181	1.53 [*]	-0.86 ^{ns}	-1.32 [*]	-0.17 ^{ns}	3.32 [*]	-3.92 ^{**}	2.86 ^{ns}	0.35 ^{ns}
Giza 182	-1.03 ^{ns}	-1.45 ^{ns}	-0.50 ^{ns}	-0.15 ^{ns}	3.43 ^{**}	-6.81 ^{**}	5.36 ^{**}	1.36 ^{**}
IRGA 370-42-1-1F-C-1	-0.69 ^{ns}	-2.43 ^{ns}	1.24 ^{ns}	0.15 ^{ns}	0 ^{ns}	-6.47 ^{**}	2.00 ^{ns}	0.85 [*]
WAB 326-B-B-15	-2.14 ^{**}	8.68 ^{**}	0.50 ^{ns}	0.14 ^{ns}	-2.31 ^{ns}	5.33 ^{**}	-3.42 [*]	2.09 [*]
CT 9658-14-M-1-2-3-2P-M-1	0.31 ^{ns}	-0.20 ^{ns}	-1.24 ^{ns}	-0.10 ^{ns}	-0.57 ^{ns}	-3.80 ^{**}	0.62 ^{ns}	0.42 ^{ns}
IET 12515	1.20 ^{ns}	-2.01 ^{ns}	-0.65 ^{ns}	-0.07 ^{ns}	-6.36 ^{**}	4.99 ^{**}	-5.60 ^{**}	-1.12 ^{**}
IR 69312-14-2-2-2	1.20 ^{ns}	-2.31 ^{ns}	1.02 ^{ns}	0.19 ^{ns}	-8.64 ^{**}	15.15 ^{**}	-9.25 ^{**}	-4.24 ^{**}
L.S.D at 5%	1.28	6.38	1.28	0.54	2.54	2.82	3.30	0.78
L.S.D at 1%	1.70	8.49	1.70	0.72	3.38	3.75	4.39	1.04

*, **: Significant at 5% and 1%, respectively.
ns : Not significant.

Table (6): Specific combining ability (SCA) of hybrids for some studied characters.

No	Cross	Days to heading (day)	Plant height (cm)	No. of tillers	Panicle length (cm)	Grain yield (t/ha)	Straw yield (t/ha)	Harvest index %	1000-grain weight (g)
1	IR58025A x Giza 175	-0.44 ^{ns}	0.697 ^{ns}	-0.713 ^{ns}	-0.103 ^{ns}	-4.820	-4.767 ^{ns}	-2.023 ^{ns}	-0.357 ^{ns}
2	IR58025A x Giza 178	-1.563 ^{ns}	0.387 ^{ns}	-2.047 ^{ns}	-0.907 ^{ns}	1.690 ^{ns}	3.757 ^{ns}	1.403 ^{ns}	-0.753 ^{ns}
3	IR58025A x Giza 181	1.770 ^{ns}	2.940 ^{ns}	1.040 ^{ns}	0.140 ^{ns}	4.257 ^{ns}	13.747 ^{ns}	-2.803 ^{ns}	-1.270 ^{ns}
4	IR58025A x Giza 182	0.330 ^{ns}	-0.980 ^{ns}	-0.637 ^{ns}	-0.213 ^{ns}	0.460 ^{ns}	1.460 ^{ns}	2.233 ^{ns}	-0.323 ^{ns}
5	IR58025A x IRGA 370-42-1-1F-C-1	0.233 ^{ns}	-1.377 ^{ns}	2.957 ^{ns}	0.243 ^{ns}	-0.963 ^{ns}	1.277 ^{ns}	-1.640 ^{ns}	0.653 ^{ns}
6	IR58025A x WAB 326-B-B-1	-0.569 ^{ns}	-1.977 ^{ns}	2.140 ^{ns}	0.520 ^{ns}	-1.057 ^{ns}	-1.823 ^{ns}	-1.417 ^{ns}	0.733 ^{ns}
7	IR58025A x CT9685-14-M-1-2-3-2P-M-1	0.657 ^{ns}	0.327 ^{ns}	-0.677 ^{ns}	-0.063 ^{ns}	-4.100 ^{ns}	-2.877 ^{ns}	-0.850 ^{ns}	-1.367 ^{ns}
8	IR58025A x IET 12515	-0.233 ^{ns}	-0.110 ^{ns}	-0.600 ^{ns}	0.107 ^{ns}	3.473 ^{ns}	-11.600 ^{ns}	5.090 ^{ns}	2.610 ^{ns}
9	IR58025A x IR 69312-14-2-2-2	-0.233 ^{ns}	0.103 ^{ns}	0.620 ^{ns}	0.203 ^{ns}	1.057 ^{ns}	0.777 ^{ns}	-0.103 ^{ns}	-0.013 ^{ns}
10	IR 69625 A x Giza 175	0.587 ^{ns}	-0.373 ^{ns}	2.940 ^{ns}	-0.030 ^{ns}	7.783 ^{ns}	10.047 ^{ns}	-0.433 ^{ns}	0.047 ^{ns}
11	IR 69625 A x Giza 178	0.803 ^{ns}	-0.103 ^{ns}	0.497 ^{ns}	0.767 ^{ns}	-2.607 ^{ns}	-4.940 ^{ns}	-0.527 ^{ns}	0.140 ^{ns}
12	IR 69625 A x Giza 181	-2.197 ^{ns}	-2.037 ^{ns}	0.613 ^{ns}	-0.120 ^{ns}	-3.873 ^{ns}	-11.823 ^{ns}	2.553 ^{ns}	0.320 ^{ns}
13	IR 69625 A x Giza 182	0.797 ^{ns}	-0.003 ^{ns}	1.683 ^{ns}	0.127 ^{ns}	1.200 ^{ns}	-3.980 ^{ns}	1.707 ^{ns}	0.123 ^{ns}
14	IR 69625 A x IRGA 370-42-1-1F-C-1	0.023 ^{ns}	0.043 ^{ns}	-1.057 ^{ns}	-0.350 ^{ns}	4.437 ^{ns}	-3.987 ^{ns}	0.810 ^{ns}	-0.913 ^{ns}
15	IR 69625 A x WAB 326-B-B-15	1.140 ^{ns}	0.977 ^{ns}	-2.207 ^{ns}	-0.297 ^{ns}	-4.303 ^{ns}	4.850 ^{ns}	-2.627 ^{ns}	-0.697 ^{ns}
16	IR 69625 A x CT9685-14-M-1-2-3-2P-M-1	-0.643 ^{ns}	-1.277 ^{ns}	-0.133 ^{ns}	0.300 ^{ns}	0.870 ^{ns}	-0.177 ^{ns}	-1.870 ^{ns}	0.307 ^{ns}
17	IR 69625 A x IET 12515	-0.200 ^{ns}	2.357 ^{ns}	-1.277 ^{ns}	-0.177 ^{ns}	-1.550 ^{ns}	12.920 ^{ns}	-1.617 ^{ns}	0.190 ^{ns}
18	IR 69625 A x IR 69312-14-2-2-2	-0.200 ^{ns}	0.433 ^{ns}	-1.060 ^{ns}	-0.247 ^{ns}	-1.807 ^{ns}	-3.927 ^{ns}	0.873 ^{ns}	0.520 ^{ns}
19	IR 70368 A x Giza 175	-0.117 ^{ns}	-0.337 ^{ns}	-2.233 ^{ns}	0.133 ^{ns}	-2.943 ^{ns}	-5.303 ^{ns}	2.437 ^{ns}	0.300 ^{ns}
20	IR 70368 A x Giza 178	0.767 ^{ns}	-0.290 ^{ns}	1.543 ^{ns}	0.133 ^{ns}	0.913 ^{ns}	1.167 ^{ns}	-1.943 ^{ns}	0.627 ^{ns}
21	IR 70368 A x Giza 181	0.433 ^{ns}	-0.890 ^{ns}	0.440 ^{ns}	-0.043 ^{ns}	-0.253 ^{ns}	-1.927 ^{ns}	0.243 ^{ns}	0.963 ^{ns}
22	IR 70368 A x Giza 182	-0.907 ^{ns}	0.987 ^{ns}	-1.047 ^{ns}	0.070 ^{ns}	-1.643 ^{ns}	2.510 ^{ns}	-3.967 ^{ns}	0.187 ^{ns}
23	IR 70368 A x IRGA 370-42-1-1F-C-1	-0.347 ^{ns}	1.323 ^{ns}	-1.900 ^{ns}	0.103 ^{ns}	-3.477 ^{ns}	1.687 ^{ns}	0.797 ^{ns}	0.263 ^{ns}
24	IR 70368 A x WAB 326-B-B-15	-0.563 ^{ns}	1.013 ^{ns}	0.063 ^{ns}	-0.243 ^{ns}	5.370 ^{ns}	-3.050 ^{ns}	3.920 ^{ns}	-0.030 ^{ns}
25	IR 70368 A x CT9685-14-M-1-2-3-2P-M-1	-0.013 ^{ns}	0.960 ^{ns}	0.803 ^{ns}	-0.240 ^{ns}	3.230 ^{ns}	5.040 ^{ns}	2.713 ^{ns}	0.993 ^{ns}
26	IR 70368 A x IET 12515	0.430 ^{ns}	-2.253 ^{ns}	1.880 ^{ns}	0.057 ^{ns}	-1.910 ^{ns}	-1.317 ^{ns}	-3.497 ^{ns}	-2.810 ^{ns}
27	IR 70368 A x IR 69312-14-2-2-2	0.430 ^{ns}	-0.530 ^{ns}	0.423 ^{ns}	0.020 ^{ns}	0.773 ^{ns}	3.133 ^{ns}	-0.600 ^{ns}	-0.513 ^{ns}
	L.S.D at 5%	2.22	1.043	2.24	0.964	4.388	4.990	5.707	1.819
	L.S.D at 1%	3.47	1.637	3.34	1.25	6.870	6.518	7.589	2.819

*, **: Significant at 5% and 1%, respectively.
ns: Not significant.

crosses were IR 58025 A x Giza 178, IR 69625 A x Giza 182 and IR 70368 A x Giza 178 in harvest index trait.

Estimates of SCA effects for the 27 rice crosses are shown in Table (6) when considering SCA effects for selection of superior hybrids, the estimate of SCA effects for number of tillers indicated that, two crosses IR 58025 A x IRGA 370- 42- 1- 1F- C-1 and IR 69625 A x Giza 175 recorded significant and positive values ranging between 2.940 and 2.957. For grain yield / plant the crosses IR 69625 A x Giza 175 (7.783), IR 70368 A x WAB 326- B- B-15 (5.370) and IR 69625 A x IRGA 370- 42- 1- 1F- C-1 (4.437) recorded significant and highly significant positive values.

From these Data, the tester Giza 182 , Giza175 and WAB 326-B-B-15 were good combiners and could be used to produce F₁ hybrid seeds when crossed with IR 69625 A which was good stable under the Egyptian conditions.

Biochemical and Molecular Analysis:

1. Isozyme Analysis:

In the peroxidase zymogram, results showed that sub-patterns were observed among the different tested genotypes, these sub-patterns have specific bands which could be used to distinguish among the three CMS lines. Where as the IR 58025 A has the band No. 7, this band was absent in their crosses except the genotype IR 58025 A x IR 69312-14-2-2-2. On the other hand, bands No. 4, 10 and 11 were absent in the IR 58025 A (Figure 1 A). Bands No. 4, 7 and 10 were also absent in the IR 69625 A (Figure 1 B), in the IR 70368 A bands No. 10 and 11 were absent (Figure 1 C). While for the testers, all the testers were differed in the number (ranged from seven to ten) and activity of the bands (Figure 2).




Fig. (1) Isozyme diagrams:
 (A) of peroxidase for IR 58025 A (lane 1) and their nine F_1 hybrids (lane 2 - 10).
 (B) of peroxidase for IR 69625 A (lane 1) and their nine F_1 hybrids (lane 2 - 10).
 (C) of peroxidase for IR 70368 A (lane 1) and their nine F_1 hybrids (lane 2 - 10).





Fig. (2) Isozyme diagram of peroxidase for the nine testers.


With respect to the esterase zymogram, a complex of sub-patterns were observed among the different tested genotypes. Description of esterase isozyme patterns of the three CMS lines and the nine testers showed that IR 58025 A and IR 69625 A exhibited the same number of bands (eight bands) as shown in (Figure 3 A and B), the band No.7 was absent in both lines. But the IR 70368 A line showed nine bands with different activity (Figure 3 C). On the other hand, all testers were differed in number of bands (ranged from seven to nine) and the activity (Figure 4).

2. Total Soluble Protein Analysis:


Total soluble proteins were detected using Coomassi Brilliant blue-R250 for the CMS lines IR 58025 A and IR 69625 A and their hybrids as shown in (Table 7). The first CMS line exhibited a total number of 13 bands, most of them were present in the tested entries, while their intensity varied in most cases.

For the CMS line IR 69625 A and their hybrids, most of bands were present in the ten tested entries and their intensity varied from case to another, except band No. 2 which was absent in the crosses IR 69625 A x IRGA 370-42-1-1F-C-1 and IR 69625 A x IR 69312-14-2-2-2.


Generally some bands were missing in the CMS lines while present in most of the hybrids ,as well as, the testers. These bands might be derived from the testers. The bands were missing in both the CMS and the testers may present in the F₁ hybrids. Such observation as in the case of IR 58025 A x Giza 181 might have some relation with restorer genes, since Giza 181 is a good restorer. The reverse also was true in other cases, where the band was present in both CMS and tester while it was absent in their F₁' s. Thus, in general terms,



(A)



(B)



(C)




Fig. (3) Isozyme diagrams:

- (A) of esterase for IR 58025 A (lane 1) and their nine F_1 hybrids (lane 2 - 10).
- (B) of esterase for IR 69625 A (lane 1) and their nine F_1 hybrids (lane 2 - 10).
- (C) of esterase for IR 70368 A (lane 1) and their nine F_1 hybrids (lane 2 - 10).




Fig. (4) Isozyme diagram of esterase for nine testers.

Table (7): Description of total soluble protein of IR 70368 A with their F₁ hybrids and nine testers.

Genotype Bands No	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	
1	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++
2	.	++	++	++	++	++	++	++	++	++	++	++	++	++	++	++	++	++	++	++
3	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++
4	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++
5	++	++	++	++	++	++	++	++	++	++	++	++	++	++	++	++	++	++	++	++
6	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
7	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++
8	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++
9	.	.	+++	++	++	++	++	++	++	++	++	++	++	++	++	++	++	++	++	++
10	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++
11	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++
12	+++	++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++
13	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++	+++

++++: Very strong

++ : Weak

+++ : Strong

+ : Very weak

++ : Intermediate

. : Absent

1- IR 70368 A	11- Giza 175
2- IR 70368 A x Giza 175	12- Giza 178
3- IR 70368 A x Giza 178	13- Giza 181
4- IR 70368 A x Giza 181	14- Giza 182
5- IR 70368 A x Giza 182	15- IRGA 370-42-1-1F-C-1
6- IR 70368 A x IRGA 370-42-1-1F-C-1	16- WAB 326-B-B-15
7- IR 70368 A x WAB 326-B-B-15	17- CT9685-14-M-1-2-3-2P-M-1
8- IR 70368 A x CT9685-14-M-1-2-3-2P-M-1	18- IET 12515
9- IR 70368 A x IET 12515	19- IR 69312-14-2-2-2
10- IR 70368 A x IR 69312-14-2-2-2	

there was poor correlation between total soluble proteins and restoring ability despite the polymorphism detected.

3. DNA Marker Analysis:

Using RAPD marker RBL.4 for PCR and subsequent separation of PCR products using 1.5% Agarose gel electrophoresis, a total of 14 amplified fragments were detected for the CMS line IR 58025 A along with nine testers and their respective hybrids (Fig. 5 A). The amplified fragments varied in their intensity and mobility, 50 bp ladder DNA was used as a size marker as shown in the first lane. Some of the amplified fragments were monomorphic and appeared in most of the tested materials. Bands were numbered on the basis of their respective sizes.

The first four bands had fragment sizes larger than 1031 bp, the largest band in the 50 bp ladder used. The largest fragment with approximately size of 1420 bp was present in eight genotypes and absent in 11 cases. The second band with size of 1350 bp appeared in 12 genotypes and their hybrids with the CMS line IR 58025 A. Band No. five with a size of 925 bp appeared in ten cases and was absent in nine lanes. The size of the remained 12 fragments ranged between 1155bp and 217bp. They varied in their intensity and presence or absence in the CMS line IR 58025 A and their F₁s as shown in (figure 5 A).

The sixth band with size of 816 bp was present in all samples, while it was very faint in case of CMS line IR 58025 A.

As for CMS line IR 69625 A along with the nine testers and their respective hybrids (Figure 5 B), the first four bands also had fragment sizes larger than 1031 bp. The largest fragment with band size of 1420 bp was present in 13 genotypes and absent in six.

The CMS line IR 70368 A with nine testers and their hybrids are shown in (Figure 5 C), 14 amplified fragments were detected for the CMS line with three respective hybrids. The first four bands had fragments were larger than 1031 bp. The largest fragment with approximate size of 1420 bp was present in ten genotypes and absent in nine.

From DNA studies, a level of molecular polymorphism was detected. For example, band No. 7 in the two CMS lines; IR 58025 A and IR 70368 A was found in many testers and their F_1 'S, in spite of its missing in the CMS line itself (Figure 5 C). Figure 6 a, b and c are diagrams for (Figure 5 a, b and c) respectively. Therefore, further experiments could be applied for studying the correlation of this band with the restoring ability of CMS lines. Also, the same trend of polymorphism was detected in the case of total soluble proteins in all CMS lines, testers and their F_1 'S. Most of the testers are considered to be good restorers. In case of the rest of detected bands, a quite poor correlation between the detected polymorphism and restoring ability was evident. These results are in correspondence with those obtained by Jena and Pandey (1999) and Li *et al.*, (2000), who found few polymorphic DNA profiles between A (CMS) and R (restorer) lines.




Fig. (5): DNA polymorphism:

- (A) for CMS line IR 58025 A and their respective F_1 's detected by RAPD marker RBL.4.
 (B) for CMS line IR 69625 A and their respective F_1 's detected by RAPD marker RBL.4.
 (C) for CMS line IR 70368 A and their respective F_1 's detected by RAPD marker RBL.4.




Fig. (5 a)




Fig. (6 b)




Fig. (6 c)

REFERENCES

- Banumathy, S.; K. Thiyagarajan and P. Vaidyanathan (2003).** Study on magnitude of heterosis of rice hybrids for yield and its components. *Crop Research Hisar*. 25: 2, 287 – 293; 23 ref.
- Bhave, S.G.; B.L. Dhonukshe and V.W. Bendale (2002).** Heterosis in hybrid rice. *Journal of Soils and Crops*. 12: 2, 183 – 186; 11 ref.
- Chakraborty, S.; S. Bhattacharya and N. Mandal (2002).** Analysis of peroxidase activities in CMS, restorer and F₁ hybrids of rice (*Oryza sativa* L.) during reproductive stages. *Journal of Interacademia*. 6: 1, 13 –16; 12 ref.
- Jena, K.K. and S.K. Pandey (1999).** DNA markers for purification of A and B lines for hybrid rice improvement. *Hybrid Rice Newsletter*, Vol. 2, No. 1.
- Jin, D.M.; W.J. Wang; S.Y. Lan; Z.X. Xu and S.H. Yang (2001).** Heterosis and correlative analysis of yield components and panicle characters of rice (*Oryza sativa* L.) hybrid Pei-Ai 64_s / E32. *Journal of Huazhong Agricultural University*. 20: 6, 516 – 521; 10 ref.
- Kato, H.; K. Tanaka; H. Nakazmi; H. Araki; T. Yoshida and Y. Ogi (1994).** Heterosis of biomass among rice ecospecies and isozyme polymorphism and RFLP. National Agriculture Research Center, Tsukuba 305, Japan, *Breeding, Sci.* 44: 3, 271 – 277, 339 – 340; 20 ref.
- Kempthorne, O. (1957).** An introduction to genetic statics. John Wiley and Sons Inc., New York, 458-471 pp.
- Li, G.R.; Q.X. Liu and B.X. Li (1996).** A study of the difference in the zymograms of esterase isozymes in rice genotypes. *J. of Southwest. Agric. Uni. China*. 18:2, 152-154; 4 ref.
- Li, L.; J.B. Yang; X.F. Wang; T.H. Xiang and J.D. Wu(2000).** Identification of Xieyou combinations of hybrid rice and their three parents by using RAPD and microsatellite markers. *Chinese Journal of Rice Science*. 14: 4, 203 – 207; 12 ref.
- Murray, M.G. and W.F. Thompson (1980).** Rapid isolation of high molecular weight plant DNA. *Nucleic Acids Research* 8:4321-4325.

- Panse, V.G. and Sukhatme (1954).** Statistical methods for Agricultural Workers. Indian Council of Agric. Res., New Delhi, India, 227 p.
- Panwar, D.V.S.; R. Kumar; A. Singh and B.S. Mehla (1998).** Studying heterosis for grain yield and its components in hybrid rice. International Rice Research Notes. 23: 2, 15.
- Patel, S.R.; N.M. Deasi and M.U. Kukadia (1994).** Heterosis for yield and yield-contributing characters in upland rice. Gujarat Agricultural University Research Journal. 20: 1, 162 – 163; 10 ref.
- Ramesha, M.S.; B.C. Viraktamath; M.I. Ahmed; C.H.M. Vijayakumar and S. Singh (1998).** Yield system analysis in rice hybrids. International Rice Research Notes. 23: 2, 14 – 15.
- Scandalios, J.G. (1969).** Genetic control of multiplex molecular forms of enzymes in plant. Biochem. Gent. 3:37-79.
- Singh, R.V.; J.L. Dwivedi and R.K. Singh (2002).** Heterosis studies in rice hybrids involving WA sources of CMS line. Annals of Agricultural Research. 23: 4, 541 – 547; 12 ref.
- Tu, A.T.; J.A. Reinose and Y.Y. Hsiao (1968).** Peroxidase activity of hemeptides for horse heart cytochrome. C. Experintia, 24:219-221.
- Virmani, S.S. (1994).** Heterosis and hybrid rice breeding. Berlin (Germany): Springer – verlag. 189 p.
- Virmani, S.S.; B.C. Viraklamath; C.L. Casal; R.S. Toledo; M.T. Lopez and J.O. Manalo (1997).** Hybrid Rice Breeding Manual. IRRI Inter. Rice Research Institute, Philippines, pp. 151.
- Vishwakarma, D.N.; D.M. Maurya; G.P. Verma; S.R. Vishwakarma and R. Kumar (1999).** Heterosis for yields components in rice hybrids (*Oryza sativa* L.). Indian Journal of Agricultural Sciences. 69: 7, 530 – 532; 4 ref.
- Yolanda, J.L. and L.D.V. Das (1996).** Heterosis in hybrid rice. Madras Agricultural Journal. 83: 2, 115 – 117; 8 ref.
- Zhang, G.; T.S. Bharaj ; Y. Lu ; S.S. Virmani and N. Huang (1996).** Mapping of the Rf-3 nuclear fertility-restoring gene for WA cytoplasmic male sterility in rice using RAPD and RFLP markers. Theor. Appl. Genet. 94: 27-33.
- Zhang, G.; T.S. Bharaj; Y. Lu; S.S. Virmani and N. Huang (1997).** Mapping of the Rf-3 nuclear fertility – restoring gene for WA cytoplasmic male sterility in rice using RAPD and RFLP markers. Theoretical and Applied Genetics. 94: 1, 27 – 33, 30 ref.

الملخص العربي

دراسات مورفولوجية ووراثية جزئية على المقدره على إعادة الخصوبة لسلالات الأرز العقيمة ذكراً

مدحت عراقى الدينارى^١ - عبد السلام عبيد دراز^٢ - سعيد عبد السلام دره^٣ - ونيد فؤاد غيطان^١

١- قسم الوراثة - كلية الزراعة جامعة طنطا

٢- مركز البحوث والتدريب فى الارز - سخا - كفر الشيخ

٣- قسم الوراثة - كلية الزراعة جامعة كفر الشيخ

أستخدم فى هذه الدراسة ثلاث سلالات أرز عقيمة ذكراً (CMS) وهى IR 58025 A ، IR 69625 A و IR 70368 A وتم اختبارها لبعض الصفات المورفولوجية والبيوكيميائية والجزئية بواسطة تسعة أصناف مختلفة من الأرز. وقد استخدم نموذج line x tester بين الثلاث سلالات العقيمة ذكراً (lines) والتسعة أصناف (testers) وذلك للحصول على ٢٧ هجين جيل أول (F₁) وتم حصاد كل هجين منفصل عن الآخر.

أظهرت النتائج أن السلالات الأبوية و الهجن بها اختلافات معنوية واضحة. فكان اتجاه الخلط سلبى فى بعض الصفات مثل أيام الطرد، و كان الخلط ايجابى فى كثير من الصفات الأخرى. وتشير النتائج إلى أن السلالة IR 69625 A أعطت قيم مرغوبة لصفة عدد أيام التزهير وعدد الفروع فى النبات والمحصول فى النبات بينما أعطت السلالة IR 70368 A قيم مرغوبة فى صفة طول النبات ، طول السنبله ، وزن القش ، دليل الحصاد ووزن الـ ١٠٠٠ حبة وأخيراً أعطت السلالة IR58025 A قيم غير مرغوبة فى معظم الصفات المدروسة تحت الظروف المصرية. وبالنسبة للقدرة العامة على التألف أظهرت السلالة IR 69625 A أحسن قدرة تألف خاصة بالنسبة لصفة محصول الحبوب/للنبات، بينما سجلت السلالة IR 58025 A

والسلالة IR 70368 A اختلافات معنوية ومعنوية جدا على التوالي بالنسبة لصفة وزن القش. وبالنسبة لتأثيرات القدرة الخاصة على التألف سجل الهجين IR 58025 A x IRGA 370-42-1-1f-c1 والهجين IR 69625 A x Giza 175 قيما ايجابية معنوية بالنسبة لعدد الأشطاء.

أوضحت النتائج أن هناك إختلاف في عدد و شدة بعض الحزم بين التراكيب الوراثية المختلفة بالنسبة لنشاط إنزيم البيروكسيديز، بعض هذه الحزم كان يميز السلالات العقيمة ذكريا. كما أوضحت النتائج بالنسبة لنشاط إنزيم الأستيريز أن هناك اختلافات معقدة في الحزم بين التراكيب الوراثية المختلفة. كما تم استكشاف البروتينات الكلية الذاتية في السلالات العقيمة ذكريا IR 58025 A و IR 69625 A وهجتها . بشكل عام فإن تحليل البروتينات الذاتية الكلية اظهر وجود بعض الحزم في اغلب الهجن والأصناف تحت الدراسة في حين كانت غائبة في حالة السلالة العقيمة ذكريا و اغلب الظن أن هذه الحزم هي نفسها الموجودة في الإباء على إعتبار أنها غير موجودة في السلالة العقيمة ذكريا. ويمكن الربط بين مثل تلك الحزم وبين القدرة على إعادة الخصوبة. كما أظهرت الدراسة بمعلمات الـ DNA الجزيئية درجة عالية من الإختلاف بإستخدام المعلم الجزيئي RBL4 إذ لوحظت حزم محددة في الهجن وكذلك في الأصناف تحت الدراسة في حين كانت تلك الحزم غير موجودة في السلالة العقيمة ذكريا. إلا أن الإرتباط بين تلك الإختلافات والقدرة على إعادة الخصوبة كان ضعيفا إلى حد كبير ويرجع ذلك إلى إستخدام معلمات الـ RAPD وهي معلمات عادة ما تكون غير متخصصة للصفة محل الدراسة وهذا يحتاج مزيدا من الدراسة.

