

## FTIR ANALYSIS OF A CRYOPRECIPITATED SOY PROTEIN

R. E. Abdolgader

Department of Food Science and Technology Omar Almkhtar University,  
El-Baida - Libya

(Received: Nov., 7, 2006)

**ABSTRACT:** *The effects of heating on the NaOH-CP structure, in the presence of varying concentration of NaCl and reducing agents, were examined by Fourier transform infrared (FTIR) spectroscopy. FTIR spectroscopy revealed that at 25 °C the NaOH-CP protein showed four bands at 1633, 1650, 1667, and 1689 cm<sup>-1</sup>. At 85 °C, the temperature associated with gelation, the four bands decreased in the intensity with the simultaneous increase of a band at 1644 cm<sup>-1</sup> as result of the protein unfolding, furthermore two new bands at 1616 and 1685 cm<sup>-1</sup> appeared. The latter two bands are attributed to intermolecular  $\beta$ -sheet formed during gelation. No major changes were observed in the amide I bands upon heating to the solutions up to 100 °C in the presence of 0.5 or 1.0 M NaCl. In the presence of SDS as function of increasing temperature reveals that a 50 mM of SDS stabilizes the secondary structure against denaturation on the other hand 100 mM of SDS major changes were observed in the amide I above 50 °C.*

**Kew words:-** *Soybean proteins, FTIR, Secondary structure.*

---

## INTRODUCTION

Soy is a popular component of the Asian diet and more recently has found increasing acceptance in western diets due to the Food and Drug Administration claim of linking the consumption of soy protein with a lower of heart disease (FDA 1999). Additionally soy products are being recognized as having potential roles in the prevention and treatment of chronic diseases, most notably cancer (Fukutake *et al* 1996, Applelt and Reicks 1999), osteoporosis, and kidney disease and to decrease plasma triacylglycerol and cholesterol levels in humans (Urade *et al* 2003, Vittadini and Vodovotz 2003).

Structurally, soy protein is classified based on sedimentation characteristics; the conventional nomenclature for soy protein fractions is the 2S, 7S, 11S and 15S fractions (Yamauchi *et al* 1991), the major fractions are the 7S fraction (7S globulin or conglycinin) and 11S fraction (11S globulin or glycinin).

Very little information on the structural conformation of  $\beta$ -conglycinin can be found in the scientific literature.  $\beta$ -conglycinin is a compactly folded protein with high degree of unstructured regions. The  $\alpha$ -helix,  $\beta$ -structure and random coil contents of the secondary structure are 5, 35, and 60%, respectively (Lewis and Chen 1979). However, the secondary structure of

glycinin was predicted to be 25 %  $\alpha$ -helix structure, 25 %  $\beta$ -Sheet structure, 42 % turns and 8 % unordered forms on the basis of molecular modeling its amino acid sequence (Argos *et al* 1985). Recently, Abbott *et al* (1996) reported Fourier transform infrared (FTIR) spectra of glycinin as 24 %  $\alpha$ -helix structure, 30 %  $\beta$ -Sheet structure, 31 % turns, and 12 % unordered. Their data also indicated that glycinin has the same secondary structure in solution and in hydrate solids. While, Dev *et al* (1988) concluded that the secondary structure of 11S consisted mainly of  $\beta$ -sheet,  $\beta$ -turns and unordered structure with very little  $\alpha$ -helix structure.

The ordered structure of glycinin can be affected by several factors. Alkali treatment decreased ordered structure, especially the  $\beta$ -Sheet; adding ethanol to the alkali denatured glycinin increased the ordered structure (particularly  $\alpha$ -helix) and decreased the random coil contents (Ishino *et al* 1980). Acetylation of glycinin decreased the  $\beta$ -sheet and increased the random coil content at 90% acetylation (Yamauchi *et al* 1979). The observed similarity in the amounts of  $\alpha$ -helix and  $\beta$ -sheet secondary structures may indicate that their interior conformations are very similar or highly conserved (Marccone 1999). The present work was conducted to study the secondary structure of cryoprecipitated soy proteins and to investigate the effect of heating temperature, ionic strength, and reducing agents on the secondary structure of that protein.

## **MATERIALS AND METHODS:**

### **Sodium Hydroxide Extraction / Cryoprecipitation**

Proteins were extracted using the procedure of Davidson *et al* (1979) with some modifications. The defatted soybean meal (100g) was mixed with dilute NaOH (1L, 0.02 %, pH 10.7) and allowed to stand for 1 h at room temperature with intermittent stirring. The mixture was centrifuged (12000 x g) for 10 min. The extract was refrigerated (4 °C) for 18 h and the proteins which precipitated were recovered by centrifugation (12000 x g) for 10 min followed by lyophilization. This isolate was designated as sodium hydroxide cryoprecipitate (NaOH-CP)

### **Preparation of Protein Samples for FTIR Analysis**

FTIR analysis was used for determination of NaOH-CP secondary structure. NaOH-CP (12 % w / v) was prepared by dissolving the protein in deuterium oxide ( $D_2O$ ) or  $H_2O$ . To study the effect of ionic strength, the protein was dissolved in  $D_2O$  solutions containing 0.5 and 1.0 M NaCl. To study the effect of denaturing agents, NaOH-CP (12 % w / v) were prepared using 50 and 100 mM sodium dodecyl sulfate (SDS), and 0.1 and 0.5 % 2-mercaptoethanol (2-MeSH).

Infrared spectra were recorded with a 8210E Nicolet fourier transform infrared spectroscopy (FTIR) spectrometer equipped with a deuterated triglycine sulfate detector. The spectrometer was purged with dry air from

Balston dryer (Balston, Haverhill, MA, 018350723, USA). Samples (7  $\mu$ l) were held in an IR cell with a 25 $\mu$ m path length (for samples prepared in D<sub>2</sub>O) and 12 $\mu$ m (for samples prepared in H<sub>2</sub>O) between two CaF<sub>2</sub> windows. The IR cell was placed in a temperature-controlled cell holder. The temperature was increased in 5 °C increments and the cell allowed to equilibrate for 10 min before recording the spectrum, for the cooling cycle, the temperature was decreased in 10 °C increments and the cell allowed to equilibrate for 10 min before recording the spectrum. The second derivative spectrum was obtained as described by Cameron and Moffatt (1987).

## **RESULTS AND DISCUSSION**

### **FTIR of NaOH-CP**

Figure 1a shows a plot of stacked second derivative spectra of the amide I band in the infrared spectra of NaOH-CP dissolved in D<sub>2</sub>O. Figure 1b shows a plot of stacked second derivative spectra of the amide I band in the infrared spectra of NaOH-CP dissolved in H<sub>2</sub>O. The second derivative spectra were generated from the spectra of NaOH-CP after subtraction of the absorption due to water from each spectrum. At 25 °C the amide I band in the infrared spectrum of NaOH-CP in D<sub>2</sub>O (Figure 1a) showed four peaks at 1633, 1650, 1667, and 1689 cm<sup>-1</sup>. In H<sub>2</sub>O similar peaks were observed, except that the frequency of the band at 1654 cm<sup>-1</sup> in H<sub>2</sub>O shifted to 1650 cm<sup>-1</sup> in D<sub>2</sub>O, this shifting due to hydrogen-deuterium exchange. The band at (a) 1633 cm<sup>-1</sup> has been assigned to  $\beta$ -sheet structure (Susi and Byler 1988, Dong *et al* 1990, Herald and Smith 1992); at (b) 1650 cm<sup>-1</sup> has been assigned to  $\alpha$ -helix structure (Susi and Byler 1988, Mitchell *et al* 1988, Holloway and Mantsch 1989, Dong *et al* 1990); at (c) 1667 cm<sup>-1</sup> has been assigned to the turn structure (Herald and Smith 1992); and the band at (d) 1689 cm<sup>-1</sup> has been assigned to  $\beta$ -turn structure as reported by other researchers (Surezwiz and Mantsch 1988, Arrondo *et al* 1988, Susi and Byler 1988) or to the high frequency component band of an antiparallel  $\beta$ -sheet (Surewiz and Mantsch 1988). Abbott *et al* (1996) reported that based on FTIR spectroscopy, the secondary structure of glycinin was 33%  $\beta$ -sheet, 25%  $\alpha$ -helix, 31% turns and 12 % unordered structure, Dev *et al* (1988) had previously reported that the secondary structure of glycinin mainly of  $\beta$ -sheet structure,  $\beta$ -turns and unordered structure with very little  $\alpha$ -helix. Since both studies were carried out in H<sub>2</sub>O it was difficult to employ resolution enhancement techniques to ascertain the position of the amide I bands precisely leading to possible miss assignment of the unordered and  $\alpha$ -helical structures. It is evident that the secondary structure of NaOH-CP in D<sub>2</sub>O or H<sub>2</sub>O that the major secondary component is  $\beta$ -sheet with a smaller amount of  $\alpha$ -helix and turn structures.




Figure 1:- Stacked plot of the second derivative infrared spectra of NaOH-CP, a:D<sub>2</sub>O, b: H<sub>2</sub>O

### Effect of Temperature on Secondary Structure

Heating the NaOH-CP solution (12 % w / v in D<sub>2</sub>O) from 25-70 °C had no significant effect on the secondary structure (Figure 2). Above 70 °C the 1650 cm<sup>-1</sup> band attributed to α-helix decreases. Heating the protein above 85 °C (temperature associated with gelation) resulted in a substantial decrease in the intensities of the 1633, 1689 and 1667 cm<sup>-1</sup> bands and an increase in the intensity of a shoulder at 1644 cm<sup>-1</sup> attributed to random coil (Nagano *et al.* 1994) formed as a result of protein unfolding. The appearance of the 1644 cm<sup>-1</sup> coincides with the appearance of two additional bands at 1685 and 1616 cm<sup>-1</sup> attributed to intramolecular β-sheet formed as a result of re-association of unfolded peptides segments which lead to formation of a gel and aggregate structures (Clark *et al.* 1981, Ismail *et al.* 1992 and De Las Rivas 1997). These results are in contrast to those reported by Chen and Barber (1990) who observed that the amount of α-helix of glycinin was increased from 7.8 to 28 % as the temperature of protein solution increased from 25 to 90 °C employing FTIR spectroscopy. While Wang and Damodaran (1991) reported that at high protein concentration (10 % protein) thermal treatment results in extensive denaturation of the secondary structure in soy 11S (glycinin). They attributed that at high concentration, the propensity of

## Ftir analysis of a cryoprecipitated soy protein

intermolecular interactions apparently leads to aggregation and network formation and this prevents refolding and regaining of secondary structure. Nagano *et al.* (1994) investigated the secondary structure changes of glycinin during heat-induced gelation employing FTIR spectroscopy. They reported that bands at 1618 and 1680  $\text{cm}^{-1}$  attributed to  $\beta$ -sheet increased with the formation of heat-induced gel of glycinin attributed the bands at 1618 and 1680  $\text{cm}^{-1}$  to intermolecular hydrogen-bonded antiparallel  $\beta$ -sheet structure resulting from protein aggregation. Cooling cycle of the heated solution of NaOH-CP showed no changes in the secondary structure of the heated protein (Figure 3) this suggests that the denaturation temperature of NaOH-CP is irreversible.




Figure 2:- Stacked plot of the second derivative infrared spectra of NaOH-CP in D<sub>2</sub>O as function of increasing temperature




Figure 3:- Cooling cycle of NaOH-CP in D<sub>2</sub>O

### Effect of Ionic Strength on Secondary Structure

The FTIR spectra in the amide I absorption region of NaOH-CP in the presence of 0.5 and 1.0 M NaCl solutions recorded as a function of increasing temperature between 25-100 °C are shown in Figure 4 and Figure 5 respectively. No major changes were observed in the amide I bands upon heating to the solutions up to 100 °C in the presence of 0.5 M or 1.0 M NaCl. Using FTIR, Boye (1995) reported that increasing NaCl concentration up to 2 M lead to increase the thermal stability of  $\beta$ -lactoglobulin. A notable exception in the small drop in the relative intensity of the band 1650 cm<sup>-1</sup> ( $\alpha$ -helix) to the 1633 cm<sup>-1</sup> band. This suggests that the  $\alpha$ -helix component of glycinin is thermally labile. Elevating the heating temperature of NaOH-CP solution from 25 to 100 °C in the presence of 0.5 M and 1.0 M NaCl did not cause the appearance of the two bands at 1620 and 1684 cm<sup>-1</sup> attributed to protein aggregation. These results indicate that NaCl appear to stabilize the protein structure aggregate formation possibly through non-specific ion

## Ftir analysis of a cryoprecipitated soy protein

effects of NaCl on electrostatic interaction between charges groups on the protein (Léger and Arntfield 1993). Cooling cycle of NaOH-CP (Figure 6) in presence of 1 M NaCl showed the same secondary structure of heating cycle of this protein which indicates that higher heating temperature was required to disrupt the native protein structure at higher ionic strength.




Figure 4 :- Stacked plot of the second derivative infrared spectra of NaOH-CP in D<sub>2</sub>O as function of increasing temperature in the presence of 0.5 M NaCl





Figure 5 :- Stacked plot of the second derivative infrared spectra of NaOH-CP in D<sub>2</sub>O as function of increasing temperature in the presence of 1.0 M NaCl





**Figure 6:- Cooling cycle of NaOH-CP in D<sub>2</sub>O in the presence of 1.0 M NaCl**


### **Effect of Reducing Agents on Secondary Structure**

Examination of the FTIR spectra of NaOH-CP in the presence of SDS as a function of increasing temperature reveals that a 50 mM concentration of SDS stabilizes the secondary structure of NaOH-CP against denaturation or aggregation Figure 7. It is of interest to note that the  $\alpha$ -helical structure also remains intact. They suggested that SDS at low concentration could form a bridge between cationic groups of protein and ionic groups of SDS, which seem to stabilize the protein structure. The aggregation bands at 1616 and 1684  $\text{cm}^{-1}$  did not appear with increasing temperature in the presence of 50 mM SDS, indicating the absence of antiparallel  $\beta$ -sheet formation. These results are in a good agreement with Boye *et al.* (1996) who reported no aggregation bands were observed with bovine serum albumin in the presence of 50 mM of SDS.

In the presence of 100 mM SDS major changes in the amide I bands above 50 °C (Figure 8). A drop in the relative intensity of the 1633  $\text{cm}^{-1}$  to the 1650  $\text{cm}^{-1}$  band is observed. The bandwidth of amide I bands significantly increase above 70 °C along with the appearance of a weak band at 1616  $\text{cm}^{-1}$  attributed to aggregate formation. A broad band centered at approximately

1644  $\text{cm}^{-1}$  (unordered structure) is also observed above 70 °C. These results demonstrate that and the increase in SDS concentration favors the unfolding of the protein more than aggregate formation. Steinhardt (1975) reported that high concentrations of SDS favor a decrease in the enthalpy due to the non-specific binding of detergent to the protein facilitating its denaturation.


The spectra in Figure 9 and Figure 10 show the combined effects of 2-mercaptoethanol (2-MeSH) and heating on the secondary structure of NaOH-CP. NaOH-CP in the presence of 2-MeSH appear to lose its secondary structure form aggregates at approximately 80 °C in 0.1% 2-MeSH and 75°C in 0.5 % 2-MeSH.



**Figure 7 :- Stacked plot of the second derivative infrared spectra of NaOH-CP in D<sub>2</sub>O as function of increasing temperature in the presence of 50 mM SDS**

***Ftir analysis of a cryoprecipitated soy protein***

---



**Figure 8 :- Stacked plot of the second derivative infrared spectra of NaOH-CP in D<sub>2</sub>O as function of increasing temperature in the presence of 100 mM SDS**




Figure 9 :- Stacked plot of the second derivative infrared spectra of NaOH-CP in  $\text{D}_2\text{O}$  as function of increasing temperature in the presence of 0.1% 2-MeOH

## FTIR analysis of a cryoprecipitated soy protein




Figure 10 :- Stacked plot of the second derivative infrared spectra of NaOH-CP in D<sub>2</sub>O as function of increasing temperature in the presence of 0.5% 2-MeOH

### Conclusion:-

Heating temperature, ionic strength, and reducing agents were used to investigate the secondary structure of a cryoprecipitated soy proteins using FTIR spectroscopy. FTIR spectroscopy shows that glycinin has high amount of  $\beta$ -sheet structure, Heating temperature from 25 - 70 °C had no significant effect on the secondary structure while heating temperature above 80 °C had significant effect on the secondary structure. FTIR spectroscopy indicates that NaCl stabilize the structure of glycinin against denaturation while SDS and 2 MeOH appear to disrupt the secondary structure especially at temperature higher than 80 °C

**REFERENCES**

- Abbott, T. P., H. abetani, D. J. Sessa, W. J. Wolf (1996). Effects of bound water on FTIR spectra of glycinin. *J Agric Food Chem* 44 2220-2224.
- Applelt, L. C., M. M. Reicks,(1999). Soy induces phase II enzymes but does not inhibit dimethylbenz[a] anthracene-induced carcinogenesis in female rats. *J Nutr* 129: 1820-1826.
- Argos, P., S. V. L. Narayana, N. C. Nielsen (1985). Structural similarity between legumin and vicilin storage proteins from legumes. *EMBO J* 4 1111-1117.
- Arrondo, J. L. R., N. M. Young, H. H. Mantsch (1988). The solution structure of concanavalin A probed by FTIR spectroscopy. *Biochim Biophys Acta* 952 261-268.
- Boye, J. I. (1995). Thermodynamic and structural properties related to the gelation of whey proteins. Ph D. Thesis McGill University.
- Boye, J. I., I. Alli, A. A. Ismail (1996). Interactions involved in the gelation of bovine serum albumin. *J Agric Food Chem* 44 996-1004.
- Cameron, D. G., D. J. Moffatt (1987). A generalized approach to derivative spectroscopy. *Appl Spectrosc.*41 539-544.
- Chen, R., Y. Ker, C. Wu (1990). Temperature and shear rate affecting the viscosity and secondary structural changes of soy 11S globulin measured by a cone-plate viscometer and Fourier transform infrared spectroscopy. *Agric Biol Chem* 54 1165-1176.
- Clark, A. H., D. H. P. Saunderson, A. Suggett (1981). Infrared and Laser-ramab spectroscopic studies of thermal-induced globular protein gels. *Int J Peptide Protein Res.* 17 353-364.
- Davidson, R. M., R. E. Sand, R. E. Johnson (1979). Methods for processing soy protein and composition of matter. U.S. Patent 4, 172,828.
- De Las Rivas, J., J. Barber (1997). Structure and thermal stability of photosystem II reaction centers studied by infrared spectroscopy. *Biochem* 36 8897-8903.
- Dev, S. B., J. T. Keller, C. K. Rha (1988). Secondary structure of 11S globulin in aqueous solution investigated by FT-IR derivative spectroscopy. *Biochimica Biophysica Acta* 272-280.
- Dong, A., P. Huang, W. S. Caughey (1990). Protein secondary structure in water from second-derivative amide I infrared spectra. *Biochem.* 29 3303-3308.
- [FDA] Food and Drug Administration, (1999). Food labeling: health claims; soy protein and coronary heart disease. Final rule. 21 VER 101. Washington D.C. Dept. of Health and Human Services.
- Fukutake, M., M. Takahashi, K. Ishida, H. Kawamura, T. Sigimura, K. Wakabashi (1996). Quantification of genistein and genistin in soy beans and soy bean products. *Food Chem Toxic* 34:457-461.
- Herald, T. J., D. M. Smith (1992). Heat-induced changes in the secondary structure of hen egg S-ovalbumin. *J Agric Food Chem.* 40 1737-1740.
- Holloway, P., H. H. Mantsch (1989). Structure of cytochrome b<sub>5</sub> in solution by Fourier transform infrared spectroscopy. *Biochem* 28 931-935.

## ***Ftir analysis of a cryoprecipitated soy protein***

---

- Ishino K., S. Kudo (1980). Conformational transition of alkaline-denatured soybean 7S and 11S globulins by ethanol. *Agric Biol Chem* 44 537-543.
- Ismail, A. A., H. H. Mantsch, P. T. T. Wang (1992). Aggregation of chymotrypsin: Portrait by infrared spectroscopy<sup>1</sup>. *Biochimica Biophysica Acta* 1121 183-188.
- Lèger L. W., Arntfield, (1993). Thermal gelation of the 12S canola globulin. *JAOCs* 70 853-861.
- Lewis B. A., J. H. Chen (1979). Effect of conformation and structure changes induced by solvent and limited enzyme modification on the functionality of soy proteins. In: *Functionality and Protein Structure*, ed Pour-Ei A. Washington, D C, p.27-35.
- Marcone, M. F., (1999). Biochemical and biophysical properties of plant proteins: a current understanding with emphasis on 11S seed globulins. *Food Res Inter* 30 79-92.
- Mitchell, R. C., P. I. Haris, C. Fallowfield, D. J. Keeing, D. Chapman (1988). Fourier transform infrared spectroscopic studies on gastric H<sup>+</sup>/K<sup>+</sup> ATPase. *Biochem Biophys Acta* 941 31-38.
- Nagano T., T. Akasaka, K. Nishinan (1994). Dynamic viscoelastic properties of glycinin and  $\beta$ -conglycinin gels from soybeans. *Biopolymers* 34 1303-1309.
- Steinhardt, J. (1975). The nature of specific and non-specific interactions of detergent with proteins: complexing and unfolding. In: *Protein-Ligand Interactions*, eds H Sund and G Blauer, W de Gruyter New York and Berlin, p 412-426.
- Surewiz, W. K., H. H. Mantsch (1988). New insight into protein secondary structure from resolution-enhanced infrared spectra. *Biochim Biophys Acta* 952 115-130.
- Susi H., D. M. Byler (1988). Fourier transform infrared spectroscopy in protein conformation studies. In: *Methods for Protein Analysis*, ed Cherry J P. Barford R A. American Oil Chemists' Society: Champaign, IL, pp 235-255.
- Urade R., S. Okamoto, T. Yagi, T. Moriyama, T. Ogawa, M. Kito (2003). Functions of soy phosphatidylcholine in dough and bread supplemented with soy protein. *J Food Sci* 68 (4) : 1276-1282
- Vittadini, E., Y. Vodovotz (2003). Changes in the physicochemical properties of wheat- and soy-containing breads during storage as studied by thermal analyses *J Foo SCI* 68 (6): 2022-2026
- Wang, C., S. Damodaran (1991). Thermal gelation of globular Proteins: Influence of protein conformation on gel strength. *J Agric. Food Chem.* 39 433-438.
- Yamauchi, F., H. Ono, Y. Kamata, K. Shibasaki (1979). Acetylation of amino groups and its effect on the structure of soybean glycinin. *Agric Biol Chem.* 43 1309-1315.
- Yamauchi, F., T. Yamagishi, S. Iwabuchi (1991). Molecular understanding of heat-induced phenomena of soybean protein. *Food Rev Inter* 7 283-322.

## تحليل بروتينات فول الصويا المرسبة بالتبريد باستخدام مطياف متوسط

### المدى للاشعة تحت الحمراء

رمضان الصالحين عبدالقادر

قسم علوم وتقنية الأغذية - كلية الزراعة - جامعة عمر المختار - البيضاء - ليبيا

#### المخلص العربي

تأثير درجة الحرارة في وجود تركيزات مختلفة من كلوريد الصوديوم بالإضافة إلى العوامل المختزلة على بروتين فول الصويا المرسبة بالتبريد تمت دراستها باستخدام مطياف FTIR. بينت النتائج انه عند درجة حرارة ٢٥ م° فان بروتين فول الصويا قيد الدراسة اظهر أربع حزم عند  $1623, 1650, 1667$  و  $1689$   $cm^{-1}$ . عند درجة حرارة ٨٥ م° فان هذه الأربع حزم انخفضت في كثافتها مع زيادة في كثافة الحزمة  $1644$   $cm^{-1}$  والتي تظهر نتيجة لدنترة البروتين بالإضافة إلى زيادة كثافة هذه الحزمة فان هناك حزمتين تم ظهورهما عند  $1616$  و  $1685$   $cm^{-1}$ .

بينت الدراسة عدم حدوث أي تغيرات معنوية على التركيب الثانوي للبروتين الثنائي حتى درجة ١٠٠ درجة مئوية في وجود كلوريد الصوديوم بتركيز ٠,٥ و ١ مولر. أظهرت نتائج هذه الدراسة أن وجود SDS بتركيز ٥٠mM كان له تأثير على تثبيت البروتين ضد درجة الحرارة بينما تركيز ١٠٠mM من نفس المادة أحدثت تغيراً واضحاً على تركيب البروتين عند درجة حرارة أعلى من ٥٠ م°.