

IMPROVEMENT OF GROWTH AND YIELD OF BLACK CUMIN PLANT BY ACTIVE DRY YEAST AND SOME VITAMINS

Ismail, Sakina I. ¹, M. A. Khafagy² and M. R. Khater¹

1- Department of Medicinal and Aromatic Plants Research, Horticulture Research Institute, Agricultural Research Center, EGYPT

2- Department of Agricultural Botany, Faculty of Agriculture, Mansoura University, EGYPT.

ABSTRACT

Two pot experiments were carried out in the experimental farm of Agricultural Botany Department, Faculty of Agriculture, Mansoura University, Mansoura, Egypt, during the two growing seasons of 2004 and 2005 to investigate the effect of presoaking black cummin seeds in vitamins (ascorbic acid at 25 or 50 ppm, thiamine at 50 or 100 ppm and α tocopherol at 10 or 20 ppm) and active dry yeast at 1000 or 2000 ppm on black cummin plant growth and yield and its components.

The results indicated that presoaking seeds in vitamins and active dry yeast lead to an increase in vegetative growth in terms of plant height, number of leaves, number of lateral branches and fresh and dry weight per plant. Data also showed that α tocopherol at 10 and 20 ppm, thiamine at 50 ppm and yeast at 1000 ppm delayed significantly flowering and fruiting date. While, other treatments hastened flowering and fruiting in both seasons. Application of vitamins and active dry yeast caused a marked increase in number of capsules and seed yield per plant as well as volatile oil and fixed oil percentage in addition to oil content per plant. Yeast at 2000 ppm and α tocopherol at 20 ppm were the best treatments.

INTRODUCTION

Medicinal and aromatic plants occupy a prominent position in the Egyptian cultivation because of increasing interest demand of local industry and export.

Black cummin (*Nigella sativa* L.) belonging to family Ranunculaceae is an annual spicy herb native to the Mediterranean region and now cultivated in other parts of the world including Middle East, North Africa and Asia. Its seeds used for edible and medicinal purposes in many countries. In Egyptian folk medicine, *Nigella* seeds are used as carminative, diuretics and delayed menses and lactation, while its oil has protective action against histamine induced bronchospasm, cough and bronchial asthma (Soliman, 1978). Recent studies indicated that the *Nigella* oil has antibacterial, antifungal and antihelminthic effects (Salomi *et al.* 1992).

Organic and biodynamic agriculture considered as one of developmental techniques which produce no or less polluted yields. It has many angles, one of them application of biofertilization which had drawn the attention of research workers and had became in the last few decades a positive alternative to chemical fertilizers, biofertilization are reasonably more

safer to the environment and human compared to chemical fertilizers. In parallel to biofertilization and equal to its importance as well its sound able impact on the environment using natural and safety substituents, i.e. vitamins (B1 or C or E) and yeast as alternatives to using growth regulators in order to improve plant growth, flowering, fruit setting and yield.

Vitamins are organic compounds that are essential to the metabolism of living organisms. They are known as growth factors inflicting many physiological processes. They act as co-enzymes or constituents of enzymes cofactor. Vitamins have functions as growth regulators or hormone precursors; have antioxidative properties and probably also yet unknown modes of actions (Oertli, 1987). The various positive effects of applying active dry yeast was attributed to its own contents of different nutrients, high percentage of protein, larger amounts of vitamin B and natural plant growth regulators such as cytokinins. (Ahmed *et al.* 1997)

The beneficial effects of applying vitamins and yeast on growth and yield of medicinal and aromatic plants have been frequently reported, In early studies of Reda *et al.* (1977) they found that application of vitamins (thiamine and ascorbic acid) favoured the growth of roots, stems and leaves as well as fruits and rays of umbels of *Ammi visnaga* L. as indicated by increase in their dry weight. The most effective concentration of thiamine was 50 mg/L whether applied as soaking of the seeds or as foliar spray. Ascorbic acid was more effective when applied only as presowing treatment of the seeds especially at 50 and 100 mg/L. Thiamine treatment significantly increased the total yield of chromones and khellin as well as visnagin yield (mg/plant) in the fruits in both soaking and spraying applications, especially at 50 mg/L. The yield of different chromones in the fruits under the effect of ascorbic acid (50 and 100 mg/L soaking method) was about 3 folds that of corresponding control.

Ahmed *et al.* (1998) on roselle plant found that when active dry yeast were applied as foliar spray with three concentrations 0.0, 0.1 and 0.2 g/L at vegetative growth, flowering and fruiting stages improved significantly growth, yield of calyxes and active ingredients. They added that maximum values obtained when plants sprayed with 0.2 g/L. Ahmed and Ali (2001) found that active dry yeast at 0.0, 0.1 and 0.2 g/L significantly increased the vegetative growth of *Ambrosia maritima*, while damassin and ambrosin percentage significantly decreased by increasing the yeast concentration. Ali (2001) found that foliar spray with 4.5 g/L active dry yeast gave better results of *Calendula officinalis*.

Refaat and Balbaa (2001) obtained pronounced increment in lemongrass vegetative growth, yield and essential oil percentage due to applying thiamine. They also found that quantitative and qualitative changes in essential oil constitutes due to thiamine treatment. Naguib and Khalil (2002) reported that using yeast and thiamine on *Nigella sativa* L. plant had promising effects on vegetative growth, seed index and yield as well as fixed and essential oil yield. The treatment with the superiority was yeast at 2 g/L combined with 20 ppm thiamine in increasing *Nigella sativa* seed yield with good quality. Wahba (2005) mentioned that using yeast and riboflavin increased vegetative growth parameters and yield of *Oenothera biennis*.

Recently, Massoud (2006) confirmed that using yeast caused an increase in vegetative growth in terms of plant height, number of branches, herb fresh and dry weight per plant, herb yield and essential oil of sage (*Salvia officinalis* L.) plant.

It could be noticed that there is no enough literature concerning the effect of vitamins and yeast on the growth and yield of medicinal plants generally and particularly on black cumin. Thus this study aimed to give some spot lights on this topic and to investigate the effect of presoaking *Nigella sativa* seeds in different concentrations of active yeast, thiamine, ascorbic acid and α -tocopherol on plant growth and seed yield and its components.

MATERIALS AND METHODS

Two pot experiments were carried out in the farm at the Agricultural Botany Department, Faculty of Agriculture, Mansoura University during the period of 2004-2005, to investigate the effects of vitamins (ascorbic acid, thiamine and α -tocopherol) and yeast (active dry yeast) on black cumin plants growing under normal conditions.

The seeds of black cumin (*Nigella sativa* L.) used in the present study were secured from Medicinal and Aromatic Plants Research Department, Horticulture Research Institute, Agricultural Research Center, Giza, Egypt.

The pots used in this experiment were of plastic type 40 cm in diameter having drainage holes and were filled with 20 kg clean air dry soil. Homogenous lots of black cumin seeds were separately soaked for 12 hours in vitamins and yeast as follows: 1-Control (distilled water), 2- Ascorbic acid at 25 ppm, 3- Ascorbic acid at 50 ppm, 4- Thiamine at 50 ppm, 5- Thiamine at 100 ppm, 6- α -tocopherol at 10 ppm, 7- α -tocopherol at 20 ppm, 8- Yeast at 1000 ppm and 9- Yeast at 2000 ppm.

After soaking, thirty seeds were planted in each pot on 19th November in the two growing seasons. After 6 weeks from sowing, the plants were thinned to leave only 5 uniform plants per pot. Phosphorus fertilizer (calcium super phosphate 15.5% P_2O_5) was mixed with soil prior to sowing at the rate of 200 kg / feddan, while both nitrogen (ammonium sulphate 20.6% N) and potassium (potassium sulphate 48% K_2O) were added individually in two equal doses at the rate of 50 Kg/Feddan and 75Kg/Feddan, respectively. The first dose was added after thinning and the second half at the beginning of flowering stage. Irrigation was conducted whenever required throughout the experimental period.

After 90 days from sowing three plants from each treatment were taken randomly to study the following parameters. 1- Plant height, number of leaves / plant, number of lateral branches / plant, shoot system fresh weight (g) and shoot system dry weight (g). During growing season the number of days till flowering and fruiting were recorded while at the end of season the following yield parameters were taken, number of capsules / plant, seed yield / plant (g), weight of 1000 seed (g), oil yield per plant and oil percentage% (fixed and volatile).

Fixed oil % was carried out as described by A.O.A.C. (1990) and volatile oil was determined according to Guenther (1961).

The treatments were arranged in complete randomized block design and the obtained data were subjected to statistical analysis of variance according to Gomez and Gomez (1984) LSD value for comparison

RESULTS AND DISSCUSSION

1- Vegetative Growth

Data presented in Table (1) reveal that α -tocopherol at (10 or 20 ppm), thiamine at (50 and 100 ppm) and yeast at (1000 and 2000 ppm) increased significantly black cummin plant height in the two growing seasons. There are no significant differences between thiamine and yeast treatments especially at the high level. The highest values were obtained with the high level of them. In addition presowing seeds in active dry yeast at 2000 ppm proved to be more effective in increasing plant height. However, ascorbic acid at 25 ppm decreased plant height in the first season while, increased it in the second season.

Table (1): Effect of vitamins (Ascorbic acid, Thiamine and α – Tocopherol) and active dry yeast on black cummin plant vegetative growth characters during 2004 and 2005.

Treatments	Plant height cm		Number of leaves/plant		Number of lateral branches/plant		Plant fresh weight g		Plant dry weight g	
	2004	2005	2004	2005	2004	2005	2004	2005	2004	2005
Control	23.2	24.9	14.7	15.7	6.0	7.3	3.49	4.89	0.699	0.877
AsA 25 ppm	22.1	25.0	15.0	16.3	6.0	8.7	4.62	5.59	0.924	1.017
AsA 50 ppm	26.7	28.7	14.7	16.0	6.3	8.3	4.65	5.13	0.930	0.927
Thi. 50 ppm	27.1	29.5	15.3	16.7	7.3	9.3	4.39	5.76	0.877	1.053
Thi. 100 ppm	32.3	35.4	18.3	19.7	9.7	11.0	5.58	7.36	1.115	1.373
α -Toco. 10ppm	28.7	30.4	16.7	17.3	8.3	9.3	5.35	6.76	1.070	1.253
α -Toco. 20ppm	30.9	31.3	17.0	18.0	9.0	10.7	5.76	7.91	1.152	1.482
Y. 1000 ppm	27.6	30.4	16.0	17.0	9.0	10.3	4.50	6.26	0.899	1.153
Y. 2000 ppm	34.2	36.3	18.3	19.7	11.0	12.3	6.08	7.41	1.215	1.381
L.S.D. 5%	1.01	1.17	1.14	1.62	1.43	1.36	0.47	0.45	0.100	0.091

With regard to number of leaves per plant, data presented in Table (1) reveal that the presowing seeds in thiamine, α -tocopherol and yeast led to an increase in the number of black cummin leaves per plant, Moreover, both thiamine and yeast at high levels were more effective in this concern. While, ascorbic acid either at 25 or 50 ppm and thiamine at 50 ppm had no significant effect in this concern.

Regarding the effect of vitamins and active dry yeast on number of lateral branches per plant, data in the same table point out that yeast, α -tocopherol and thiamine increased significantly number of branches of black cummin plant but ascorbic acid caused a slight increase in this regard. Moreover, yeast application was more effective in this respect.

With respect to fresh and dry weights per plants, data recorded in Table (1) reveal that ascorbic acid, thiamine and α -tocopherol as well as

yeast treatments increased significantly both fresh and dry weight of black cummin plant in the two successive seasons. This increment was highly significant in yeast, α -tocopherol and thiamine at the high concentrations. It could be noticed that soaking black cummin seeds with vitamins (ascorbic acid, thiamine, and α -tocopherol) or yeast showed higher significant increases in all studied growth characters compared with control.

The promotive effects of ascorbic acid on the fresh and dry weights per plant could be attributed in part to its effect on many metabolic and physiological processes and/or increase the organic acids exerted from the roots into the soil and consequently increase the solubility of the most nutrients which slowly release into the rizosphere zone where it may be utilized by the plant (Negm *et al.* 1997)

Thus, it could be concluded that ascorbic acid in lower concentrations might probably acts as growth factor. Another approach for the role of ascorbic in intact plant growth was stated by Aberg, (1961) who attributed its effect to increasing the availability of iron and micronutrients in the plant. Tarraf *et al.* (1999) mentioned that AsA increased plant height and greatly increased the number of tillers/ plant of lemon grass plants. They added that ascorbic acid could be involved in the main metabolic processes especially with energy transfer coenzymes, carbohydrate metabolism and improved photosynthetic activity. Some investigators mentioned that ascorbic acid had a regulation effect upon oxidation reduction potential of cytoplasm (Aberg, 1961, Sana and Ota 1977). Our findings are in line with Reda *et al.* 1977 on *Ammi visnaga* L. and Saraswathamma and Jayachandra, 1981 on *Trigonella foenum* L.

Dealing with thiamine effects on growth, thiamine is connected with the role of thiamine pyrophosphate cocarboxylase, as a co enzyme in various types of decarboxylation involving pyruvic and α -ketoglutamic acid. Kodandaramaiah and Rao (1985) suggested that B. vitamins participate in plant growth and development indirectly by enhancing the endogenous levels of various growth factors such as cytokinins and gibberellins. Naguib and Khalil (2002) mentioned that thiamine has a promotive effect on vegetative growth of *Nigella sativa* L. they suggested that thiamine affect upon the meristem may partly be of an indirect nature and be mediated by the mature tissue through an altered supply of metabolite to the apex. Vitamin B1 has a function in intermediate carbohydrate metabolism (Robinson, 1973). These results are in accordance with those of ZhuKova (1977), Ramaiah *et al.* (1984), Oertli (1987), El-Ghamriny *et al.* (1999) and Youssef *et al.* (2005). The enhancing influence of α -tocopherol (Vitamin E) on *Nigella sativa* L. growth parameters which was observed in this study may be due to its physiological role in protecting membrane lipids from peroxidation, the vitamin E used up by this process is regenerated by vitamin C (Kunert and Ederer, 1985) and reducing oxidative stress, e.g. imposed by gaseous pollutants (SO_2 , O_3 , O_2 and OH), drought, chilling and herbicides (Fryer, 1992). Preliminary studies have shown that α -tocopherol can be absorbed by the plant tissue (Schmitz, 1997). The maximum of α -tocopherol uptake is achieved within 24 to 48 hours. Vitamin E results in higher membrane fluidity (Tanczos *et al.* 1982). These authors suggested that α -tocopherol is built into

the plant membranes, the fluidity of which is thus increased. Alpha tocopherol increased the water permeability of liposomes at low temperatures. Similar results were published by Mallet *et al.* (1994) who established a significant linear correlation between antioxidative capacity of lipophilic extracts and α -tocopherol content in leaves of 15 selected plant species. Finally, the improving effects of antioxidants on growth characters might be attributed to their positive action on enhancing cell divisions and protecting plant cells from free radicals that responsible for plant senescences (Raskin, 1992). The positive effect of α -tocopherol on plant growth is in harmony with many findings on earlier results from literature on some plant species i.e. Tanczos *et al.* (1982) on rice, Matakiaadis and Kintzios (2005) on cucumber plants.

Regarding the increasing effects with yeast, the positive effect of yeast extract on black cumin growth characters may be due to the fact that yeast extract is a natural source of cytokinins, vitamins, and most of the essential elements (Nagodawithana, 1991). In addition, the increase in the release of carbon dioxide through fermentation process effectively stimulates photosynthesis and accelerates the biosynthesis of carbohydrates. It increases synthesis of plant growth promoters especially GA_3 , IAA and cytokinins which lead to improving cell division and cell enlargement (Moor, 1979). Our results coincided with the results obtained by Ahmed *et al.* (1998) on roselle plants, who found that yeast treatments improved growth and yield of plants, Ahmed *et al.* (1998) on marjoram plant who showed that active dry yeast caused more branches, heavier herb and leaves and dry weight, Naguib 2002 on lemongrass, as well as Naguib and Khalil 2002 on black cumin plant.

2 Yield and Its Components

2-1 Flowering and fruiting date:

Data tabulated in Table (2) show that thiamine at 50 ppm and α -tocopherol at 10 and 20 ppm as well as yeast at 1000 ppm delayed significantly flowering and fruiting date. Yeast treatment at 1000 ppm delayed the appearance of the first flower by 6 days while α -tocopherol treatments delayed flowering by 5.4 and 5.7 days respectively followed by thiamine at 50 ppm which delayed flowering by 3.7 days in the first season. In addition, α -tocopherol at 10 ppm delayed the appearance of first flower by 4.6 days followed by thiamine at 50 ppm and yeast at 1000 ppm which delayed it by 4 days as well as α -tocopherol which delayed flowering by 3 days in the second season. On the contrary, ascorbic acid at 25 and 50 ppm and thiamine at 100 ppm as well as yeast at 2000 ppm hastened flowering in the two seasons. The rate of ascorbic acid at 25 ppm was more effective in this respect.

Table (2): Effect of vitamins (Ascorbic acid, Thiamine and α -Tocopherol) and active dry yeast on black cumin yield and its components during 2004 and 2005.

Treatments	Days till flowering		Days till fruiting		Number of capsules/plant		Seed yield/plant (g)		Weight of 1000 seeds		Volatile oil %		Fixed oil %		Oil content per plant	
	2004	2005	2004	2005	2004	2005	2004	2005	2004	2005	2004	2005	2004	2005	2004	2005
Control	110	115	114	118	4	5	1.68	1.96	6.27	6.44	1.30	1.32	30.1	30.3	0.528	0.620
AsA 25 ppm	109	113	112	116	5	5	1.96	2.24	6.00	6.27	1.31	1.34	31.9	31.5	0.651	0.736
AsA 50 ppm	109	114	114	117	5	6	2.10	2.38	6.31	6.43	1.35	1.38	31.6	31.8	0.692	0.790
Thi. 50 ppm	114	119	118	123	5	6	2.24	2.38	6.16	6.28	1.36	1.37	31.9	31.4	0.745	0.780
Thi. 100 ppm	109	113	113	117	7	7	2.80	3.08	6.39	6.54	1.37	1.37	32.4	32.3	0.946	1.037
α -Toco. 10ppm	116	119	119	122	6	6	2.66	2.66	6.31	6.45	1.32	1.42	32.6	31.8	0.902	0.884
α Toco. 20ppm	116	118	119	121	7	8	3.08	3.22	6.37	6.70	1.35	1.44	33.9	33.2	1.086	1.115
Y. 1000 ppm	116	119	120	122	5	7	1.96	3.08	6.31	6.63	1.40	1.52	32.8	34.2	0.670	1.100
Y. 2000 ppm	109	114	114	119	7	8	2.94	3.36	6.71	6.84	1.43	1.53	34.6	35.1	1.059	1.231
L.S.D. 5%	1.58	1.81	1.44	2.31	1.81	1.95	0.759	0.820	0.178	0.120	0.047	0.043	0.343	0.515	0.035	0.032

He mentioned that the superiority of vitamins and yeast on fixed and volatile oil may be due to the role of these vitamins as co-enzymes involved in specific biochemical reactions in the plant such as oxidative and non-oxidative decarboxylation. He stated also that the biochemical active pyrophosphates are the units which condense to form the many varied forms, which constitute of terpenes. In addition, Subba Rao, (1984) and Dewic (2000) mentioned that these results may be due to the stimulatory effect of yeast and vitamins which act as co-enzymes of photosynthesis and metabolism of carbohydrates and other metabolites in seeds.

REFERENCES

- A.O.A.C. (1990). Official Methods of Analysis. Association of Official Agriculture Chemists. P.O. Box 540, Benjamin, Franklin Station, Washington, D.C.
- Aberg, B. (1961). Vitamins as growth factors in higher plants. *Encyclopedia of Plant Physiology*. XIV: 418 – 448.
- Ahmed, E. T. (1998). Influence of concentration and time of spraying active dry yeast on growth and volatile oil content of marjoram plant. *J. Agric. Sci. Mansoura Univ.*, 23 (11). 5067-5081.
- Ahmed, F.F.; Ragab, M.A.; Ahmed, A.A. and Mansour, A.E.M. (1997). Improving the efficiency of spraying different nutrients for Red Roomy grapevines (*Vitis vinifera* L.) by using glycerol and active dry yeast. *Egypt. J. Horti.* 24 (1): 91 – 108.
- Ahmed, S.K. and Ali, A.F. (2001). Effect of salinity treatments and active dry yeast on growth and active ingredients of *Ambrosia maritima* L. . *Proc. The Fifth Arabian Horticulture Conference*, Ismailia, Egypt, 1:217-224.
- Ahmed, S.K.; El-Ghawas, E.O. and Aly, A.F. (1998). Effect of active dry yeast and organic manure on roselle plant. *Egyptian J.Agric. Res.* 76 (3). 1115 – 1142.
- Ali, A.F. (2001). Response of pot marigold plants (*Cleome officinalis* L.) to some rock phosphate source and yeast. *Proc. The Fifth Arabian Horticulture Conference*, Ismailia, Egypt, 1:31-42.
- Baszynski, T. (1967): *Naturwissenschaften* 54, 339. Cited from Oertli, J.J. (1987): Exogenous application of vitamins as regulators for growth and development of plants. *Z. Pflanzenernahr Bodenk*, 150: 375-391.
- Dewic, K.P.M. (2000). *Medicinal Natural Products, A Biosynthetic Approach*. 2nd Ed., pp. 306 – 356. John Wiley & Sons, N.Y.
- El-Ghamriny, E.A.; Arisha, E.A.H. and Nour, K.A. (1999). Studies on tomato flowering, fruit set, yield and quality in summer season. I. Spraying with thiamine, ascorbic acid and yeast.
- Fadi, M.; Reda, F.; Abdel-All, R.S. and El-Moursi, A. (1978). Physiological studies on *Ammi vesnaga* L. *Egypt. J. Physiol. Sci.* 5: 73 – 83.
- Faissal, F.A. and Hassan, A. M. (2004). Influence of some antioxidants on growth, vine nutritional status, yield and quality of berries in Banaty Grapevines. *Assiut. J. Agric. Sci.* 35 (4): 131 – 140.

- Fryer, M.J. (1992). The anti oxidative effects of thylakoid vitamin E (α -tocopherol). *Plant Cell and Environment*. 15: 381 – 392.
- Gomez, K.A. and Gomez, A.A. (1984). *Statistical Procedure for Agriculture Research*. 2nd Ed. Ghon Willy and Sons.
- Guenther, E. (1961). *The essential oils*. Vol. I, II and III. D. Van Naster and Comp. Inc., New York.
- Idso, S.B.; Idso, K.E. and Hooper, J.K. (1995). Effect of atmospheric CO₂ enrichment and foliar methanol application on net photosynthesis of sour orange trees (*Citrus aurantium*) leaves. *Amer. J. Botany*. 82 (1): 26 – 30.
- Kodandaramaiah, J. and Rao, P.G. (1985). Influence of B-vitamins on stomatal index, frequency and diurnal rhythms in stomatal opening in *Cyamopsis tetragonalba* (L.). *Taub. J. Biol. Res.* 5: 68 – 73.
- Kunert, K.J. and Ederer, M. (1985). *Physiol. Plant*. 65: 85 – 88. Cited from Oertli, J.J. (1987): Exogenous application of vitamins as regulators for growth and development of plants. *Z. Pflanzenernahr Bodenk*, 150: 375-391.
- Mallet, J.F.C.; Cerrti, E.; Ucciani, E.; Gamisans, J. and Gruber, M. (1994). Antioxidant activity of plant leaves in relation to their alpha-tocopherol content. *Food Chem*. 49: 61 – 65.
- Matakiadis, T. and Kintzios, S. (2005): The effect of ATP on cucumber (*Cucumis sativus* L.) regeneration from nodal explants: association with α tocopherol, H₂O₂ and size of culture vessel. *Plant Growth Regulation* 45:127-137.
- Michniewicz, M. and Kamienska, A. (1967). *Acta Soc. Botan. Polon*. 36, 67 – 72. Cited from Oertli, J.J. (1987): Exogenous application of vitamins as regulators for growth and development of plants. *Z. Pflanzenernahr Bodenk*, 150: 375-391.
- Moor, T.C. (1979). *Biochemistry and Physiology of Plant Hormones*. Pub. By Springer-Verlag, New York, USA.
- Nagodawithana, W.T. (1991). *Yeast Technology*. Universal Foods Corporation Milwaukee, Wisconsin Published by Van Nostrand Reinhold, New York. P. 273.
- Naguib, N.Y. (2002). Yield and quality of lemongrass plants (*Cymbopogon flexuosus* Stapf) as influenced by farm yard manure and foliar application of bread yeast. *Annals of Agric. Sci. Cairo*. 47 (3): 859 – 873.
- Naguib, N.Y. and Khalil, M.Y. (2002). Studies on the effect of dry yeast, thiamine and biotin on the growth and chemical constituents of black cummin (*Nigella sativa* L.). *Arab Univer, J. Agric. Sci.* 10 (3): 919 – 937.
- Negm, A.Y.; Zahran, F.A. and Rizk, N S. (1997): Foliar application of ascorbic acid, magnesium and nickel to lentil grown in newly reclaimed sandy soils. *Egypt. J. Agric. Res.*, 75(4):843-853.
- Oertli, J.J. (1987): Exogenous application of vitamins as regulators for growth and development of plants. *Z. Pflanzenernahr Bodenk*, 150: 375-391.
- Patil, B. and Lall, S.B. (1973). Effect of presowing treatment with L-ascorbic acid and gibberellic acid on growth and physiological constituents of wheat. *Botanique (Naqpur)*, 4, 5770 (Biol.Abst., 57 – 64).

- Ramaiah, J.K.; Venkataramaiah, C.; Rao, P.G. and Rao, K. N. (1984): Proc. Natl. Acad. Sci. India, Sect. B54, 1-5. Cited from Oertli, J.J. (1987): Exogenous application of vitamins as regulators for growth and development of plants. Z. Pflanzenernähr Bodenk, 150: 375-391.
- Raskin, I. (1992). Salicylate, a new plant hormone. Plant Physiology. 99: 799 – 803.
- Reda, F.; Fadl, W.; Abdel-Aal, R.S. and El-Moursi, A. (1977). Physiological studies on *Ammi visnaga* L. Lam. 5. The effect of Thiamine and ascorbic acid on growth and chromon yield. Egypt. J. Pharm. Sci. 18 (1): 19 – 27.
- Refaat, A. M. and Balbaa, L.K. (2001). Yield and quality of lemongrass plants (*Cymbopogon flexuosus stapf*) in relation to foliar application of some vitamins and micro-elements. Egypt Journal of Horticulture. 28 (1): 41 – 57.
- Robinson, F.A. (1973). Vitamins. In: Phytochemistry Vol. III: 195 – 220. Lawrence P. Miller (Ed.) Van-Nostrand, Reinhold Co., New York.
- Sahu, M.P.; Solanki, N.S. and Dashora, L.N. (1993). Effects of Thiourea, Thiamine and Ascorbic Acid on Growth and Yield of Maize (*Zea mays* L.). Journal of Agronomy and Crop Science. 171: 65 – 69.
- Salomi, N.; Nair, S.C.; Jayawarahanan, K.K. and Varghese, C.D. (1992): Antitumor principles from *Nigella sativa* seeds. Johns Hopkins, Al Mag., 63:33-36.
- Sana, J.S. and Ota, Y. (1977). Plant growth regulation activities of nicotinamide II. Effect of nicotinamide on growth of several crops. Japan. J. Crop Sci. 46: 8 – 12.
- Saraswathamma, D.N. and Jayachandra, N. (1981). Effect of presowing soaking with growth regulators on seedling growth in fenugreek (*Trigonella foenum-grecum* L.). Comparative Physiology and Ecology. 6, (2): 108 – 110.
- Schmitz, M.(1997). Bedeutung von vitaminen für die Abwehr von oxidativem Stre B bei Bohne und Apfel Dissertation, Rheinische Friedrich-wilhelms-Universität Bonn.
- Simkunas, R.; Mateikiende, I. and Bluzmanas, P. (1980): Mathm. Naturw. Reihe, 29, 85-86. Cited from Oertli, J.J. (1987): Exogenous application of vitamins as regulators for growth and development of plants. Z. Pflanzenernähr Bodenk, 150: 375-391.
- Soliman, M. A. (1981). A Pharmacognostical study of certain *Nigella* species growing in Egypt .M.Sc. Thesis, Cairo Univ., Egypt.
- Subba Rao, N.S. (1984). Biofertilizers in Agriculture. P. 189. Oxford. IBH Company, New Delhi.
- Tanczos, O.; Erdie, L.; Vigh, L.; Kuipers, B. and Keuper, J.C.(1982). Physiol. Plant. 55: 289 – 295. Cited from Oertli, J.J. (1987): Exogenous application of vitamins as regulators for growth and development of plants. Z. Pflanzenernähr Bodenk, 150: 375-391.
- Tarraf, S.A.; El-Din, K.M.G. and Balbaa, L.K. (1999). The response of regulative growth, essential oil of lemongrass (*Cymbopogon citrates* Hort.) to foliar application of ascorbic acid, nicotinamine and some micronutrients. Arab-Univer. J. Agric. Sci., 7 (1): 247 – 259.

- Wahba, H.E. (2002). Growth, yield and chemical composition of *Oenothera biennis* as affected by yeast, biotin and riboflavin foliar application. Arab Univ. J. Agric. Sci. 10 (3): 997 – 1017.
- Youssef, A.A.; Aziz, A.A. and Talaat, I.M. (2005). Influence of some antioxidants on growth, flower-heads and essential oil content of *Matricaria chamomilla* L. plants. Annals Agric. Sci., Moshtohor. 43 (2): 823 – 832.
- Zhukova, P. S. (1977). Effect of combined use of growth regulators, vitamins and fertilizers on growth and certain physiological and biochemical processes in vegetable crops. Plant growth Regul. Proc. Int. Symp. 2 nd, 736-743. Kudrev, T., Ivanova, I. and Karanov, E. (Eds), Banasofia, Bulgaria.

تحسين النمو و المحصول في نبات حبة البركة بالخميرة و بعض الفيتامينات

سكينة إبراهيم إسماعيل¹ , محمود عيد المنعم خفاجي² , مرتضى رضا خاطر¹

1- قسم بحوث النباتات الطبية و العطرية - معهد بحوث البساتين - مركز البحوث الزراعية - مصر

2- قسم النبات الزراعي - كلية الزراعة - جامعة المنصورة - مصر

اجريت تجرنتسى أصص فى المزرعة التجريبيه الخاصه بقسم النبات الزراعى ؛ كلية الزراعة ؛جامعة المنصوره خلال الموسم الشتوى 2004 و 2005 لدراسة تأثير نقع بذور نبات حبة البركة فى بعض الفيتامينات مثل حمض الاسكوربيك بتركيزات 25؛50جزء فى المليون ؛الثيامين بتركيزات 50؛100 جزء فى المليون والخميره الجافة النشطة بتركيزات 1000؛2000جزء فى المليون على النمو الخضرى و المحصول ومكوناته.

وقد أثبتت الدراسة أن معاملات النقع فى الفيتامينات والخميرة كانت فعالة فى زيادة صفات النمو الخضرى المتمثلة فى طول النبات و عدد الأوراق و عدد الفروع الحانيه للنبات و الوزن الطراز و الجاف للمجموع الخضرى و قد تفوقت معاملات الثيامين(100جزء فى المليون) و الالفاتوكوفيرول(20 جزء فى المليون) والخميرة(2000جزء فى المليون) على باقى المعاملات. كما أوضحت نتائج التجربة أن حمض الأسكوربيك (25؛50جزء فى المليون) و الثيامين(100جزء فى المليون) و الخميرة(2000جزء فى المليون) أدوا الى تشجيع الازهار و الاثمار بتقليل عدد الأيام اللازمة لتفتح أول زهرة و تكوين أول ثمرة بينما أدت باقى المعاملات الى تأخير كلا من الازهار و الاثمار .

أدى النقع فى أى من الفيتامينات و الخميرة الى زيادة ملحوظة فى عدد الكبسولات على النبات و من ثم زيادة المحصول البدرى للنبات و كذلك زيادة النسبة المئوية للزيت الثابت و الطيار و المحتوى الكلى للزيت و كانت المعاملة بالالفاتوكوفيرول (20جزء فى المليون) و الخميرة (2000جزء فى المليون) هما الأكفأ.