

DISTRIBUTION AND TRANSFORMATION OF PHOSPHORUS FORMS IN SOIL AFTER LONG-TERM APPLICATIONS OF P FERTILIZER AND CROP RESIDUES

Raghib, H.M.A.; Hala H. Gomah; M.F. Ghoneim and M.H. Raslan
Department of Soil and Water, Faculty of Agriculture, Assiut University.
Halagomaa71@hotmail.com

ABSTRACT

Two rates of crop residues (-CR and +CR) and six rates of P fertilizer (0, 100, 200, 300, 400, and 500 kg superphosphate fed^{-1}) were applied each growing season (corn and clover) in a long-term field experiment started at the Experimental Farm of Assiut University, Assiut, Egypt. The crop residues were added to the soil at the end of each growth season at rates of 5.45 and 3.71-ton fed^{-1} corn stover and Egyptian clover straw.

The response of soil P fractions to both CR and superphosphate fertilizer was small at the very beginning of the experiment, however it became more pronounced starting from the year 1999 up to the year 2003. Resin-P accounted for 0.81- 0.93%; and 1.30- 1.52% of total P in soil after 5 and 16 years of continuous application of CR and SP, respectively. At zero level of SP and no addition of CR the resin-P fraction was markedly decreased from 6.5 to 5.65.

The size of inorganic (P_i) and organic (P_o) phosphorus forms of the NaHCO_3 fractions were increased with the continuation of applying the P fertilization management for 16 years and ranged from 2.73 to 3.13 and from 1.84 to 2.19 for P_i and P_o forms, with and without crop residues, respectively. Five years after continuous application of SP and without application of CR, the relative concentrations of NaOH-P_i and NaOH-P_o increased from 3.69 to 4.44% and from 1.60 to 2.00% of total P, respectively. Also, increasing the application level of SP from zero (P_0) to 500 kg SP fed^{-1} (P_5) each growth season for five years increased the HCl-P and residual-P fractions linearly and significantly.

The path analysis showed that both resin- P_i and $\text{NaHCO}_3\text{-P}_i$ were mainly and directly affected by P fertilization. In soil that received P fertilization management with CR, resin-P was largely controlled by added- P_i , $\text{NaHCO}_3\text{-P}_i$ and HCl-P fractions.

Keywords: Long term, crop residues, P fertilizer, P fractions and P transformation.

INTRODUCTION

Phosphorus exists in soils in different fractions of great diversity in size and availability. The size of each fraction is controlled and changes with type of soil, plant, climate and agricultural practices, the most important of them are P fertilization practices, organic manuring, soil moisture contents, and time (Aulakh *et al.*, 2003). According to Stewart and Sharpley (1987) inorganic P forms are associated with amorphous and crystalline sesquioxides, and calcareous compounds. Organic P forms include relatively labile phospholipids and fulvic acids and more resistant humic acids. As mentioned by Stewart and Tiessen (1987), P forms are not discrete entities, as integrates and dynamic transformations between forms occur continuously to maintain equilibrium condition.

In their recent works, Rubaek and Sibbesen (1993, 1995) showed that P_o extracted by the resin revealed changes induced by drying and rewetting and biocidal treatments, and reflected seasonal variations in soil related to differences in P fertilization. Using the macroporous anion-exchange resin

extraction, Guggenberger *et al.* (1996) found that the soil under grass had the largest amount of resin P_o and diester-P, indicating a rapid turnover of P_o with building-up of a large potentially labile, microbial derived P_o pool. The amount of resin P_i appeared to be a primarily function of fertilization, and amounted to around 100 mg kg^{-1} in the fertilized soils irrespective of the soluble organic matter content and P source.

Application of fertilizer P and organic manure for many years resulted in accumulation of P_i and P_o fractions in the soil (McKenzie *et al.*, 1992_a and _b; Schmidt *et al.*, 1996). Reddy *et al.* (1996) and Singh *et al.* (2001_{a,b}) reported that the application of farmyard manure with fertilizer P enriched long-term P fertility of a Vertisol through enhancement in organic P content and a short-term P supply as recalcitrant P. In the most P-deficient soils under crop production, P fertilization over a long time increased both organic and inorganic P fractions (Richards *et al.*, 1995). These effects depend on the rates and forms of P fertilizers applied. The addition of P in the manure resulted in increasing the $\text{NaHCO}_3\text{-}P_i$ and decreasing the $\text{NaOH-}P_i$ (Mnkeni and MacKenzie, 1985; Xie *et al.*, 1991; Zhang and MacKenzie 1997). Soil $\text{NaOH-}P_i$ is held strongly by chemisorption to Fe and Al components of soil (Ryden *et al.*, 1977) and is considered as moderately labile P. The addition of organic material to that soil tended to decrease P sorption, possibly due to competition between phosphate ions and organic anions for retention sites in the soil (Hue, 1991; Iyamuremye *et al.*, 1996a). Zhang and MacKenzie (1997) reported that changes in soil P_o fractions displayed different patterns between fractions and between inorganic fertilizer P levels. Their data showed that $\text{NaHCO}_3\text{-}P_o$ decreased with added manure, probably due to accelerated mineralization caused by addition of organic C. On the other hand, the $\text{NaOH-}P_o$ fraction increased rapidly at first then slowly in relation to added fertilizer P.

Soil P transformations vary depending on soil type, climatic conditions, and management practices. In Mollisols, labile P was associated with inorganic bicarbonate and hydroxide extractable P, while in Ultisols, labile P was largely in the organic P forms (Tiessen *et al.*, 1984). Assessing the turn over rates of ^{33}P added in soybean residues to soils from three experimental sites in Michigan, Daroub (1994) found that the largest fraction of the applied ^{33}P was found in the inorganic labile form, which decreased with time accompanied by an increase in the NaOH fraction in all three soils tested.

MATERIALS AND METHODS

In 1988 a field experiment with corn- clover cropping system was started at the Experimental Farm of Assiut University, Assiut, Egypt. The soil of the experimental site is alluvial clay (37% clay and 33% silt) with pH 7.79, $\text{ECe } 2.96 \text{ ds/m}$, OM 1% and 5.12 ppm total Olsen P. Crop residues of corn (cv single cross 10) and the plant materials of the last cut of Egyptian clover were used as sources of organic materials, for long term, in P fertilization management. By the end of each growth season, the corn stover and the plant materials of the last cut of Egyptian clover were dried out in the open field, cut to small pieces and ploughed into the surface layer (0.00- 0.25m depth) of the experimental site. Two rates of crop residues (without,-CR and

with,+CR) and six rates of P fertilizer (0, 100, 200, 300, 400, and 500 kg super phosphate fed^{-1}) were applied each growth season. The crop residues were added to the soil at the end of each growth season at rates of 5.45 and 3.71 ton fed^{-1} corn stover and Egyptian clover straw. The treatments were arranged in a split-plot design with crop residues (CR) in the main plots and P fertilizer in the sub-plots of 3.0 m x 3.5 m each. Each treatment was replicated four times and laid out in the same plot each growth season, each year starting from 1988 up to 2003.

Fertilizer P (superphosphate 15.5% P_2O_5) was applied broadcasting and thoroughly mixed with the top 0.00 - 0.25 m surface soil layer immediately before sowing of each crop. All plots received the recommended agricultural practices and doses of N (105 kg N fed^{-1} added to corn plants) and K fertilizers (50 kg K_2O fed^{-1} for each of corn and clover) every year.

Total soluble salts were determined in the saturated soil-paste extract by measuring the electrical conductivity (Jackson, 1973). Organic matter content was determined using Walkely-Black method, following Baruah and Barthakur (1997). Soil pH was measured in 1:1 (soil: water) suspensions using a glass electrode (Jackson, 1973). Mechanical analysis was carried out using the pipette method (Piper, 1950). Olsen P extraction (Olsen *et al.*, 1954) with 0.5 M NaHCO_3 (pH 8.5, 1: 20 soil: extract ratio, 30 min shaking) was prepared and a portion was digested with potassium persulphate as described by Bowman (1989) and analyzed for total Olsen's extracted P (P_{Ot}).

The soil samples collected successively in years 1992, 1996, 1999 and 2003 after harvesting corn were thoroughly mixed, air-dried and ground to pass a 2-mm sieve and stored for chemical analysis and P fractionation. A modified version of Hedley *et al.* (1982) procedure as described by Zhang and MacKenzie (1997) was used to sequentially fractionate soil P (Fig. 1).

Duplicate soil samples of 1.00 gm soil passed through a 2-mm sieve were extracted sequentially as outlined in Fig. 1. Soil samples were placed, each, in a 50-ml centrifuge tube with a resin bag (containing 0.4 g of Dowex 1-X8 anion exchange resin in bicarbonate-saturated form) and 30-ml of distilled water was added and shaken for 16 h, then the resin bags were collected and the adsorbed P_i was desorbed using 20 ml of 0.5 M HCl solution as described by Nziguheba *et al.* (1998). After that, soil samples were extracted sequentially with 30 ml of each 0.5 M NaHCO_3 (pH 8.5), 0.1 M NaOH, and 1.0 M HCl by shaking the samples for 16 h, consecutively after shaking, the suspensions were centrifuged for 30 min at 6000 rpm and then filtered through 0.45 μ filters. Total P was determined by digestion of the whole soil sample (1.00g soil) with 5 ml concentrated H_2SO_4 and 5 ml H_2O_2 at 360 °C. The residual-P was calculated as difference between total P and sum of other fractions. A portion of the NaHCO_3 and NaOH extracts was acidified to precipitate the extracted organic matter and the supernatant was analyzed for P_i (Zheng *et al.*, 2002). In addition, another portion of NaHCO_3 and NaOH extracts was digested with potassium persulphate as described by Bowman (1989) and analyzed for P_i . The concentration of organic $\text{NaHCO}_3\text{-P}_o$ and NaOH-P_o were calculated as the difference between those of P_i and P_o . The P_i concentration in extracts was determined colorimetrically using phosphomolibdic-stannous chloride system as described by Jackson (1973).




Fig. 1. Modified procedure after Hedley et al. (1982) for P fractionation.

The data obtained of soil P fractions were subjected to standard analysis of variance and the means of treatments (crop residues and P fertilization rates) over years were tested for significant differences using least significant difference at $P = 0.05$. The CALIS statistical procedure of SAS statistical analysis program (SAS Institute, 2000) was used to derive the path coefficients between the P pools in the conceptual models. This statistical procedure also tests how well conceptual models represent graphic interpretations of P transformations in soil. Fit Goodness Index and Chi-square Test were used to assess model fit (SAS Institute, 2000; Zheng *et al.*, 2002).

RESULTS AND DISCUSSIONS


The changes in soil P fractions following the seasonal additions of crop residues (CR) and inorganic P fertilizer (SP) for 5 and 16 years are presented in Tables 1 and 2, respectively. Continuous application of CR and SP for 5 years increased the size of resin-P, NaHCO_3 -P and NaOH -P fractions; although the changes were small; they were significant. However, the more stable P fractions (HCl-P and residual fractions) showed no significant responses to continuous application of CR and SP during the first 5 years. After 16 years of continuous application of CR and SP, all P fractions were significantly increased. At low level of inorganic P fertilization, resin-P, NaHCO_3 -P, and HCl fractions (the inorganic P fractions) were mostly affected by the incorporation of CR into the 0.25 cm upper soil layer (Tables 1 and 2). However, with increasing the level of SP, these fractions were mostly affected by the inorganic P fertilizer (SP). This is in harmony with the data obtained by Saleque *et al.* (2004), they found that the concentrations of NaHCO_3 soluble P, NaOH extracted inorganic P (P_i) and acid P were affected under treatments with high proportions of organic fertilizers.

1-Resin-P:

The changes in resin-P as affected by continuous application of CR and SP over the laps of 16 years were significant. Resin-P accounted for 0.81- 0.93%; and 1.30- 1.52% of total P in soil after 5 and 16 years of continuous application of CR and SP, respectively.

Resin-P increased linearly and significantly ($p \leq 0.05$) with increasing the rate of SP application. The sharp increases were continuous with continuous application of SP from the year 1992 (Table 1) to the year 2003 (Table 2). These effects on resin-P were emotively enhanced by the incorporation of CR into the 0.25 m surface soil layer, and became more pronounced after 16 years of continuous application of CR and SP (Fig. 2).

The magnitude of the resin-P fraction linearly increased with long-term application of both CR and SP. These results confirm the earlier findings of Aulakh *et al.* (2003) whose results emphasized that the soil content of resin-P increased successively with each increment of applied P, and Motavalli and Miles (2002) observed that long-term application of inorganic P fertilizer or manure for 111 years significantly increased resin-P among different cropping systems compared with the unfertilized control and the native prairie site.



Po = 0.00, P3= 300, P5= 500 kg SP fed⁻¹ each growth season and CR= crop residues.

Fig. 2. Effects of long-term (16 years) application of P fertilizer (SP) and crop residues on resin-P fraction.

Application of CR to plots that received zero level of SP increased the concentration of resin-P from 6.5 to 6.89 mg P kg⁻¹ after 5 years of continuous application of CR (Table 1), and from 5.75 to 11.56 mg P kg⁻¹ after 16 years of continuous application of CR (Table 2). It seems that organic matter application to soil not only increases the more soluble P fractions, but also decreases the P binding affinity by soils and increases P desorption capacity (Sui and Thompson, 2000). These processes finally increase the magnitude of resin-P pool which is available to plant uptake. The competition between organic anions (resulting from the decomposition of CR) and phosphate anions for the same sites of adsorption increased the amount of phosphate anion desorbed into the soil solution (Nziguheba *et al.*, 1998).

Table 1. Effect of crop residues and P fertilization on P fractions in soil at the end of summer growth season, 1992 (continuous application for 5 years).

Treatments		Resin	NaHCO ₃		NaOH		HCl	Residual
			P _i	P _o	P _i	P _o		
mg P kg ⁻¹								
Without crop residues	P ₀	6.50	18.64	8.51	43.83	18.97	386	705
	P ₁	7.11	23.25	8.63	48.28	22.97	400	741
	P ₂	8.23	29.45	11.91	52.51	24.37	412	750
	P ₃	10.52	30.57	13.3	59.58	27.28	432	800
	P ₄	13.65	37.23	15.18	63.75	28.56	457	834
	P ₅	19.62	44.50	16.75	68.84	31.0	493	878
Mean		10.94	30.61	12.38	56.13	25.53	430	785
With crop residues	P ₀	6.89	20.45	9.31	45.94	22.92	390	692
	P ₁	8.23	24.75	10.14	51.88	24.72	409	734
	P ₂	10.9	33.75	12.5	55.97	28.22	416	768
	P ₃	12.5	36.19	15.0	68.84	30.41	441	804
	P ₄	17.11	48.00	16.22	74.06	33.98	473	835
	P ₅	21.35	55.13	17.0	81.88	39.1	488	886
Mean		12.83	36.38	13.36	63.10	29.89	436	787
LSD 0.05	CR	0.85	1.90	1.02	2.11	0.25	NS	NS
	P	0.93	1.39	1.16	1.81	1.50	20	6.15
	CR × P	1.45	2.56	NS	3.08	1.95	NS	10.50

P₀= 0.00, P₁= 100, P₂= 200, P₃= 300, P₄= 400, P₅= 500 kg SP fed⁻¹ for each growth season.
 P_i= Inorganic phosphorus, P_o= Organic phosphorus,
 CR= Crop residues, P= Phosphorus, NS= Non significant.

Table 2. Effect of crop residues and P fertilization on P fractions in soil at the end of summer growth season, 2003 (continuous application for 16 years).

Treatments		Resin	NaHCO ₃		NaOH		HCl	Residual
			P _i	P _o	P _i	P _o		
		mg P kg ⁻¹						
Without crop residues	P ₀	5.75	20.20	11	39.25	14.75	293	756
	P ₁	13.57	32.93	24.83	74.0	32.23	528	807
	P ₂	15.64	39.83	29.55	78.0	34.4	536	820
	P ₃	23.45	47.03	33.23	85.38	38.93	558	840
	P ₄	29.48	57.90	36.6	88.83	43.13	626	890
	P ₅	36.52	62.78	41.1	95.45	50.55	693	973
Mean		20.74	43.45	29.39	76.82	35.67	539	848
With crop residues	P ₀	11.56	32.85	27.24	75.75	27.45	438	764
	P ₁	16.75	43.65	31.93	88.5	38.43	513	827
	P ₂	21.45	46.40	34.2	96.7	43.35	528	860
	P ₃	26.77	56.05	35.0	102.6	56.5	558	912
	P ₄	37.74	64.43	44.9	102.8	65.31	648	917
	P ₅	42.15	78.53	51.63	105.4	73.5	734	1008
Mean		26.07	53.65	37.48	95.29	50.76	570	881
LSD 0.05	CR	1.73	1.67	0.78	2.39	1.69	15	5.15
	P	1.31	1.63	1.53	1.57	1.61	36	5.57
	CR× P	2.36	2.63	2.10	3.05	2.62	48	8.68

P₀= 0.00, P₁= 100, P₂= 200, P₃= 300, P₄= 400, P₅= 500 kg SP fed⁻¹ for each growth season. P_i= Inorganic phosphorus, P_o= Organic phosphorus, CR= Crop residues, P= Phosphorus.

2- Bicarbonate extractable P (NaHCO₃-P):

Soil NaHCO₃-P fraction includes the water-soluble and the freely exchangeable P, and represents a considerably readily and potentially available P for plant uptake (Aulakh et al., 2003; Bowman and Cole, 1978). Examination of the data presented in Table (1) reveals that the size of this

fraction was relatively small and ranged between 3.18 and 3.61% of total P, 26.86 - 28.80% of it was in the organic form, $\text{NaHCO}_3\text{-P}_o$. After 16 years of continuous application of P fertilization managements (Table 2), the relative size concentration of this fraction rose up and ranged between 4.57 and 5.32% of total P.

During the years of this experiment, $\text{NaHCO}_3\text{-P}$ forms linearly and significantly ($p \leq 0.05$) increased due to the increasing levels of SP, the incorporation of CR into the surface 0.25 m soil layer, and their interaction (Fig. 3). The sizes of these fractions were significantly increased with incorporating the CR into the P fertilization management, and became noticeably higher. The concentrations of $\text{NaHCO}_3\text{-P}_i$ and $\text{NaHCO}_3\text{-P}_o$ in plots that received the highest level of SP with CR for 5 years were 55.13 and 17.0 mg P kg^{-1} , respectively (Table 1).




Fig (3): Effect of long term (16 years) application of P fertilizer (SP) and crop residues on $\text{NaHCO}_3\text{-P}$ fractions.


Continuous incorporation of CR into the surface 0.25 m soil layer drastically improved the amount of NaHCO_3 extractable P. This indicates the transformation of crop residues P (inorganic and organic) to more labile forms (inorganic and organic) with time. However, the size of the $\text{NaHCO}_3\text{-P}_o$ was more affected especially in the absence of mineral P fertilization. In plots that received no SP, the changes in the concentration of $\text{NaHCO}_3\text{-P}_i$ over the lapse of 16 years due to the incorporation of CR were less than that of $\text{NaHCO}_3\text{-P}_o$ (Table 2). In plots with zero level of mineral P fertilizer, most of

NaHCO₃-P resulted from the decomposition of CR, which remained in the organic form, giving rise to the size of NaHCO₃-P_o pool. With high levels of SP, the accumulation of NaHCO₃-P exceeded the levels of available P that are recommended for most field crops (Cope and Rouse 1980). This may arise the potential of ground and surface water pollution.

The increase in NaHCO₃-P due to the application of CR was comparable to the increase that occurred due to SP, which further suggests that continuous application of CR may partially substitute for mineral P fertilizer. These results indicate that the size of labile NaHCO₃-P fraction can be conspicuously increased by the combination of SP fertilizer and CR. Using the crop residues for long time as a partial substitution for mineral P fertilizers could be an acceptable P fertilization management. Seasonal application of CR in combination with a low rate of mineral P fertilizer (not more than 100 kg SP fed⁻¹) could maintain the level of available labile P pools (Table 2, P_i= 43.6 and P_o= 31.9 mg P kg⁻¹) in soil at the recommended level for most field crops. This, in turn, leads to reducing the cost of input of the production system, and the potential pollution of either ground or surface water resources.

3- Sodium Hydroxide extractable P (NaOH-P):

Incorporation of CR into the surface 0.25 m soil layer, application of SP, and their interaction significantly ($p \leq 0.05$) increased the concentration of NaOH-P forms in all years of study (Fig. 4). The application of increased levels of SP linearly and significantly increased the concentration of both NaOH-P_i and NaOH-P_o. This was further magnified with incorporation of CR into soil. A similar increase in NaOH-P_i in soil due to organic matter addition was reported by Sharpley *et al.* (1984) and Iyamuremye *et al.* (1996_b).



P₀ = 0.00, P₃ = 300, P₅ = 500 kg SP fed⁻¹ each growth season and CR = crop residues.
Fig. 4. Effects of long-term (16 years) application of P fertilizer (SP) and crop residues on NaOH-P fractions.

The magnitudes of changes of NaOH-P forms were variable based upon the presence of CR or not, and the duration of applying the P fertilization management. At the beginning of the experiment (5 years after starting the experiment, Table 1) the changes in the relative size of NaOH-P_i and NaOH-P_o as a result of incorporation of CR were in the favor of NaOH-P_i form, which mean that most of P applied either as inorganic or organic form was transformed to the NaOH-P_i pool.

After 16 years of continuous application of SP and CR (Table 2), the situation was changed. The changes in relative size of NaOH-P_o form due to the application of CR and SP were more noticeable, which means that most of the P added to soil entered into the NaOH-P_o pool. Data obtained by Saleque *et al.* (2004) showed that the concentration of NaOH-P_i was lowest under treatments that contained no organic additions and a larger concentration of NaOH extracted P_i was observed under treatments with organic fertilizers.

Increasing the rate of SP application increased the rate of accumulation of both NaOH-P forms, although the rates of changes in the concentrations of NaOH-P_i were not altered much without the incorporation of CR. However, in the presence of CR the rate of changes in NaOH-P_o was considerably higher than that in the absence of CR (Fig. 4).


Comparing the concentrations of NaOH-P forms with the concentrations of resin-P and NaHCO₃-P forms in the zero SP plot (Tables 1 and 2), which almost remained constant over the time, proved that some transformations from NaOH-P_i and/or mineralization of NaOH-P_o to resin-P and NaHCO₃-P may have occurred during the lapse of experiment, 16 years. These results are similar to those reported by Zhang (1996).

4- Acid extractable P (HCl-P) and residual P:

The relative size of HCl-P and residual-P fractions varied from 25 to 35% and 48 to 66% of total P, respectively. At the very beginning of the experiment, continuous incorporation of CR into the surface 0.25 m surface soil layer had no significant effect on the HCl-P or residual-P (Table 1). However, the size of the HCl-P pool and residual-P fraction, after 16 years of continuous application of CR and SP, were significantly ($p \leq 0.05$) increased due to continuous application of CR, SP and their combination (Table 2 and Fig. 5).

The incorporation of CR into the surface 0.25 m soil layer stimulated the effects of increasing levels of SP on the concentrations of both HCl-P and residual-P fractions. HCl-P constitutes stable Ca-bound P (Williams *et al.*, 1980), while residual-P pool is likely composed of in the humus fraction (Stewart *et al.*, 1980) and insoluble inorganic P forms. It seems that, these two pools together constitute the storage pools of P in most of the Egyptian soils, and is expected to be transformed into the more soluble forms when the amount of P taken up by plants exceeds the amount of P fertilizers added to soil.

As a result of plant uptake, the readily- and moderately-labile P pools, which are in equilibrium with the HCl-P fraction, were depleted, and some of the less soluble HCl-P fraction eventually transformed to the other more soluble fractions, causing the recorded decreasing of HCl-P pool.



P₀ = 0.00, P₃ = 300, P₅ = 500 kg SP fed⁻¹ each growth season and CR = crop residues.

Fig. 5. Effects of long-term (16 years) application of P fertilizer (SP) and crop residues on HCl-P and residual P fractions.

Long-term effects of P fertilization managements on P transformation in soil:

In both P fertilization managements, added mineral P fertilization (P_i) was positively related with all P pools (Table 3 and 4). In soil received no CR (-CR) the correlations between added P_i and P_i pools (resin-P_i, NaHCO₃-P_i, and NaOH-P_i) were high, but the correlation between added P_i and NaHCO₃-P_i was the highest. This means that in the absence or with low input of organic matter, most of mineral P fertilizer has interred into the mineral forms. Using the crop residues in the P fertilization management, the correlations between added P₀ and the organic pools (NaHCO₃-P₀, NaOH-P₀) and NaOH-P_i were stronger and more associated (the r values were 0.83, 0.60 and 0.73 for NaHCO₃-P₀, NaOH-P₀ and NaOH-P_i, respectively). The presence of increased amount of P₀ in the P fertilization management resulted in building up the P₀ pools, which considered as a protected form against the precipitation and/or adsorption processes, and thus increased the availability of P in soil. In both P fertilization managements, resin-P was positively related with other P pools, but it was highly correlated with NaHCO₃-P_i. This may be because resin-P and NaHCO₃-P_i are a part of the same soil pool, which considered as a labile P pool.

In agreement with the results reported by Zheng *et al.* (2002), the residual-P was strongly associated with P_i added, but was less related to P_o added, which means that this pool consists mainly of inorganic forms of P.

Table 3. Pearson's correlation coefficients for soil P fractions as influenced by the addition of inorganic P fertilizer (coefficients significant at $p \leq 0.05$ are shown).

	Resin-P	NaHCO ₃ -P _i	NaHCO ₃ -P _o	NaOH-P _i	NaOH-P _o	HCL-P	Residual-P	Added P _i
Added P _i	0.94	0.98	0.91	0.90	0.93	0.89	0.94	1.00
Residual-P	0.93	0.95	0.83	0.86	0.87	0.86	1.00	
HCL-P	0.92	0.91	0.87	0.93	0.92	1.00		
NaOH-P _o	0.91	0.93	0.92	0.97	1.00			
NaOH-P _i	0.90	0.93	0.94	1.00				
NaHCO ₃ -P _o	0.89	0.90	1.00					
NaHCO ₃ -P _i	0.95	1.00						
Resin-P	1.00							

Added P_i= Inorganic P added

To investigate the effects of P fertilization managements on soil P pools and their transformations, the path relationships between different soil P pools as affected by P fertilization managements were constructed taking into account the two different P sources i.e. the mineral P fertilizer (added-P_i) and the organic P of CR (added-P_o). Resin-P is an estimate of the utmost soluble form of P in soil solution, while the residual-P is considered as the ultimate main storage form of P that mainly consisted of tightly bound-P_i and -P_o. The other P fractions are transformed mutually and continuously to resin-P and/or residual-P, the processes that are controlled by climatic factors, agronomic practices, and soil conditions. Thus the different P pools are conceptually grouped into soluble P (resin-P), labile and moderately labile P_i and P_o (NaHCO₃-P_i and -P_o), low labile P_i and P_o (NaOH-P_i and -P_o), acid-P, and residual-P. The impact of mineral P fertilizer (added-P_i) and the P of CR (added-P_o), as independent variables on resin-P, NaHCO₃-P_i, NaHCO₃-P_o, NaOH-P_i, NaOH-P_o, HCL-P and residual-P, as dependent variables were, therefore, explored using standardized multiple linear regression and path analysis using SAS PC version 8.2 (SAS institute, 2000). First, simple Pearson's correlations between P fractions were calculated (Table 3 and 4) and the standardized regression coefficients (β , path coefficient) of each relation was calculated.

The conceptual model for each P fertilization management was drawn illustrating the transformations among P pools in soil (Fig. 6 and 7). In the path diagrams, a straight single-headed arrow indicates a direct causal path. Indirect causal effects are connoted by a variable being linked to a given dependent variable via one or more intermediary variables. The path coefficients (numbers on lines) indicate the relative magnitudes of the direct effects of the explanatory variables linked to each dependent variable.

Table 4. Pearson's correlation coefficients for soil P fractions as influenced by the addition of inorganic P fertilizer and crop residues (coefficients significant at $p \leq 0.05$ are shown).

	Resin-P	NaHCO ₃ -P _i	NaHCO ₃ -P _o	NaOH-P _i	NaOH-P _o	HCl-P	Residual-P	Added P _i	Added P _o
Added P _o	0.49	0.44	0.83	0.73	0.60	0.52	0.44	0.46	1.00
Added P _i	0.98	0.95	0.83	0.86	0.96	0.94	0.94	1.00	
Residual-P	0.95	0.97	0.79	0.89	0.91	0.90	1.00		
HCl-P	0.95	0.92	0.83	0.84	0.92	1.00			
NaOH-P _o	0.95	0.92	0.88	0.89	1.00				
NaOH-P _i	0.87	0.88	0.90	1.00					
NaHCO ₃ -P _o	0.83	0.79	1.00						
NaHCO ₃ -P _i	0.97	1.00							
Resin-P	1.00								

Added P_i= Inorganic P added; Added P_o= Crop residues P added.

In the soil that received no CR (Fig. 6) 95.3% of the variability in resin-P was accounted for the variation in added-P_i and the size of NaHCO₃-P_i, NaHCO₃-P_o, and residual-P. The path analysis shows that both resin-P_i and NaHCO₃-P_i were mainly and directly affected by P_i fertilization with path coefficients (β) of 0.89 and 1.0, respectively, which mean that resin-P acts as a sink for P_i that came from added-P_i and directly transformed from residual-P. The added-P and residual-P consist the main and direct source to replenish the depletion in resin-P as a result of plant uptake. The NaOH-P_i form was significantly, negatively and directly affected by P_i addition ($\beta = -0.98$), but positively and indirectly responded to P_i fertilization through NaHCO₃-P_i ($\beta = 1.88 \times 0.92$) and through NaHCO₃-P_i and NaOH-P_o ($\beta = 1.88 \times 0.88 \times 1.10$). This means that added-P_i was transformed indirectly through the NaHCO₃-P_i pool into NaOH-P_i ($\beta = 1.88 \times 0.92$), and through NaHCO₃-P_i and NaOH-P_i pools into HCl-P ($\beta = 1.88 \times 0.92 \times 0.66$).

The direct effect of NaHCO₃-P_i on resin-P was not significant ($\beta =$ not significant), but resin-P was significantly and indirectly affected by NaHCO₃-P_i through residual-P ($\beta = 1.76 \times 0.40$). It appeared that resin-P and NaHCO₃-P_i pools worked as a sink for soluble P_i that came directly from added-P_i, and through the transformation of residual-P. The path coefficient between NaHCO₃-P_i and NaHCO₃-P_o was negative ($\beta = -1.06$). The negative sign of the path coefficient emphasized that the NaHCO₃-P_i was immobilized to replenish the decrease of NaHCO₃-P_o; especially in the absence of low input of inorganic P. Organic P pools (NaHCO₃-P_o and NaOH-P_o) have non-significant direct effect on resin-P.

Resin-P and NaOH-P_i have significant and direct effects on HCl-P ($\beta = 0.47$, and 0.66 , respectively). These results suggested that the HCl-P acted as sink for labile and moderately-labile P present in excess of plant removal, probably through the formation of Ca-phosphate. This means that a large portion of available P was transformed into secondary minerals, this result

support the theory that secondary P minerals are formed in the soil at the expense of more labile P_i forms. The HCl-P was considered as a secondary sink for residual fertilizer P in an Alfisol (Agbenin and Goladi, 1998) as well as a calcareous soil (Daroub *et al.*, 2000).

The $\text{NaHCO}_3\text{-P}_o$ was directly affected by NaOH-P_i ($\beta = 1.53$), and indirectly through NaOH-P_o and NaOH-P_i ($\beta = 0.88 \times 1.10 \times 1.53$), which means that a part of the inorganic P pool was transferred into the organic P pool. The biological immobilization of P_i through microbial activities and that associated with roots activities could be the direct causes for such kind of transformation. Therefore, path analysis clearly showed that under the condition of no and/or low addition of organic P sources mineral forms of P tended to transform into organic forms (P_o). Schmidt *et al.* (1996) and Simard *et al.* (1995) reported a larger amount of NaOH-P_o in agricultural soils that received a surplus of P_i addition concerning plant needs. They suggested that, with high P_i fertilizer rates in excess of plant removal, the excess P was supposed to reduce the phosphorelase activity and consequently reduce the mineralization of NaOH-P_o , with the final result of increasing the size of NaOH-P_o pool.

Residual-P was indirectly and significantly affected by added- P_i through $\text{NaHCO}_3\text{-P}_i$ ($\beta = 1.88 \times 1.76$). The other P pools have no significant direct effects on residual-P.

In soil that received the P fertilization management with CR (Fig. 7), resin-P was largely controlled by added- P_i , $\text{NaHCO}_3\text{-P}_i$ and HCl-P fractions ($\beta = 0.46, 0.28$ and 0.22 , respectively) and 97.4% of the variability in resin-P is accounted for by the variations in these variables. This means that in the presence of OM, both $\text{NaHCO}_3\text{-P}_i$ and HCl- P_i are easily transformed into resin-P and these P pools consist the main source for replenish the depletion in resin-P caused by plant uptake. The NaOH-P_i fraction has no significant effect on resin-P, while it has a high correlation with resin-P ($r = 0.87$). This was due to its indirect effect through $\text{NaHCO}_3\text{-P}_i$ ($\beta = 0.33 \times 0.28$). The $\text{NaHCO}_3\text{-P}_i$ was not significantly and directly affected by P_i fertilization, but it indirectly responded to P_i fertilization through residual-P ($\beta = 5.48 \times 0.85$).

The NaOH-P_i was positively related to residual-P ($\beta = 0.99$) and negatively related to resin-P ($\beta = -0.42$) fractions. In this P fertilization management the increase in residual-P and the decrease in resin-P were associated with the increase in NaOH-P_i . The NaOH-P_i was directly affected by added- P_o ($\beta = 0.47$). This may be due to the abilities of the NaOH extract solution to dissolve the organic compounds. This result concurs with Sui *et al.* (1999), whose results revealed that the order of the rate of increases in concentrations of P fractions with each increment of P applied with biosolids was $\text{NaOH-P}_i > \text{NaHCO}_3\text{-P}_i \approx \text{HCl-P} > \text{NaOH-P}_o > \text{H}_2\text{O} > \text{residual P} > \text{NaHCO}_3\text{-P}_o$.

The $\text{NaHCO}_3\text{-P}_o$ and NaOH-P_o fractions positively responded to added- P_o ($\beta = 0.46$ and 0.20 , respectively). The path coefficient showed that $\text{NaHCO}_3\text{-P}_o$ was more affected by the added- P_o .




Fig. 6. Path diagram for the relationship between P fractions in the 0.00- 0.25 m surface soil layer receiving continuous application of superphosphate (P fertilization management without CR). The percentage value (multiple correlation coefficient) indicates the total variation in the resin-P associated with the independent variables.

3247




Fig. 7. Path diagram for the relationship between P fractions in the 0.00- 0.25 m surface soil layer receiving continuous application of superphosphate and CR (P fertilization management with CR). The percentage value (multiple correlation coefficient) indicates the total variation in the resin-P associated with the independent variables.

The negative effects of $\text{NaHCO}_3\text{-P}_o$ and NaOH-P_o on the residual-P ($\beta = -1.9$ and -2.45 , respectively) partly support the idea that the crop residues application tended to promote the accumulation of $\text{NaHCO}_3\text{-P}_o$ and NaOH-P_o fractions in the soil. In the same context, the negative effect of $\text{NaHCO}_3\text{-P}_o$ on residual-P ($\beta = -1.9$) and the positively effect of NaOH-P_o on $\text{NaHCO}_3\text{-P}_o$ ($\beta = 0.37$), suggested that the addition of crop residues accelerates the mineralization of moderately labile P_o into labile P_o .

The positive effects of NaOH-P_o and $\text{NaHCO}_3\text{-P}_i$ on HCl-P ($\beta = 0.71$ and 0.68 , respectively) and the negative effect of HCl-P on residual-P ($\beta = -1.51$), suggest that the HCl-P was a reservoir of soil P as well as a source for plant P uptake, presumably through equilibration with resin-P ($\beta = 0.22$). The $\text{NaHCO}_3\text{-P}_o$ was directly affected by HCl-P ($\beta = 0.28$). Mineralization of CR resulted in increasing the solubility of secondary P_i forms, and releasing soluble and organic P of low molecular weight and hence may represent an easily accessible P source for microbes (Beck and Sanchez, 1994). The path analysis showed that the residual-P act as a sink for P_i and P_o fertilization ($\beta = 5.48$ and 1.75 , respectively), as well as long-term reservoir of P equilibrations with other P fractions.

REFERENCES

- Agbenin, J.O. and J.T. Goladi. 1998. Dynamics of P fractions in a savanna Alfisol under continuous cultivation. *Soil Use Manage.* 14:59-64.
- Aulakh, M.S.; B.S. Kabba; H.S. Baddesha; G.S. Bahl and M.P.S. Gill. 2003. Crop yields and phosphorous fertilizer transformations after 25 years of applications to a subtropical soil under groundnut-based cropping systems. *Field Crops Res.* 83:283-296.
- Baruah, T.C. and H.P. Barthakur. 1997. ("A Textbook of Soil Analysis"). Vikas Publishing House Pvt Ltd, New Delhi.
- Beck, M.A. and P.A. Sanchez. 1994. Soil phosphorus fraction dynamics during 18 years of cultivation on a Typic Paleudult. *Soil Sci. Soc. Am. J.* 58:1424-1431.
- Bowman, R.A. 1989. A sequential extraction procedure with concentrated sulfuric acid and dilute base for soil organic phosphorous. *Soil Sci. Soc. Am. J.* 53:362-366.
- Bowman, R.A. and C.V. Cole. 1978. An exploratory method for fractionation of organic phosphorous from grassland soils. *Soil Sci.* 125:95-101.
- Cope, J.T. and R.D. Rouse. 1980. Interpretation of soil test results. p. 35-54. In Walsh L.E. and J.D. Beaton (ed.) *Soil Testing and Plant Analysis*. SSSA Publication. SSSA, Madison, WI.
- Daroub, S. 1994. The effect of tillage on phosphorus transformations in soils. Ph.D. Dissertation, Michigan State University, East Lansing, Michigan, USA.
- Daroub, S.H., F.J. Pierce and B.G. Ellis. 2000. Phosphorus fractions and fate of Phosphorus-33 in soils under plowing and no-tillage. *Soil Sci. Soc. Am. J.* 64:170-176.

- Guggenberger, G., B.T. Christensen, G. Rubaek and W. Zech. 1996. Land-use and fertilization effects on P forms in two European soils: resin extraction and ^{31}P -NMR analysis. *European J. Soil Sci.* 47:605-614.
- Hedley, M.J., J.W.B. Stewart and B.S. Chauhan. 1982. Changes in inorganic and organic phosphorous fractions induced by cultivation practices and by laboratory incubations. *Soil Sci. Soc. Am. J.* 46:970-976.
- Hue, N.V. 1991. Effects of organic acids anions on P sorption and phytoavailability in soils with different mineralogies. *Soil Sci.* 152:463-471.
- Iyamuremye, F., R.D. Dick and J. Baham. 1996_a. Organic amendments and phosphorus dynamics: I. Phosphorus chemistry and sorption. *Soil Science* 161: 426-435.
- Iyamuremye, F., R.P. Dick, and J. Baham. 1996_b. Organic amendments and phosphorus dynamics: II. Distribution of soil phosphorus fractions. *Soil Sci.* 436-443.
- Jackson, M.L. 1973. ("Soil Chemical Analysis"). Prentice-Hall, Inc. Englewood Cliffs, N.J. New Delhi.
- McKenzie, R.H., J.W.B. Stewart, J.F. Domaar, and G.B. Schaalje. 1992_a. Long-term crop rotation and fertilizer effects on phosphorus transformations. I. In a Chernozemic soil. *Can. J. Soil Sci.* 72: 569-579.
- McKenzie, R.H., J.W.B. Stewart, J.F. Domaar and G.B. Schaalje. 1992_b. Long-term crop rotation and fertilizer effects on phosphorus transformations. II. In a Luvisolic soil. *Can. J. Soil Sci.* 72: 581-589.
- Mnkeni, P.N.S. and A.F. MacKenzie. 1985. Retention of ortho- and polyphosphates in some Quebec soils as affected by added organic residues and calcium carbonate. *Can. J. Soil Sci.* 65:575-585.
- Motavalli, P.P. and R.J. Miles. 2002. Inorganic and organic soil phosphorus fractions after long-term animal manure and fertilizer application. *Biol. Fertil. Soils.* 36: 35-42.
- Nziguheba, G., C.A. Palm, R.J. Buresh, and P.C. Smithson. 1998. Soil phosphorus fractions and adsorption as affected by organic and inorganic sources. *Plant and Soil.* 198:159-168.
- Olsen, S.R., C.V. Cole, F.S. Watanabe and L.A. Dean. 1954. Estimation of Available Phosphorus by Extraction with Sodium Bicarbonate. USDA Circ. 939. U.S. Gov. Print. Office, Washington, DC.
- Piper, C.J. 1950. ("Soils and Plant Analysis"). Inter. Sci. Publishers, Inc. New York.
- Reddy, K.S., A.S. Rao and P.N. Takkar. 1996. Transformation of fertilizer P in a Vertisol amended with farmyard manure. *Biol. Fertil. Soils.* 22: 279-282.
- Richards, J.E., T.E. Bates and S.C. Sheppard. 1995. Changes in the forms and distribution of soil P due to long-term corn production. *Can. J. Soil Sci.* 75: 311-318.
- Rubaek, G.H. and E. Sibbesen. 1993. Resin extraction of labile soil organic phosphorus. *J. Soil Sci.* 44:467-478.
- Rubaek, G.H. and E. Sibbesen. 1995. Soil phosphorus dynamics in a long-term field experiment at Askov. *Biology and Fertility of Soils.* 20:86-92.

- Ryden, J.C., J.R. McLaughlin and J.K. Syers. 1977. Mechanisms of phosphorus sorption of soils and hydrous ferric oxide gel. *J. Soil Sci.* 28:72-92.
- Saleque, M. A., U. A. Naher, A. Islam, A.B.M.B.U. Pathan, A.T.M.S. Hossain and C.A. Meisner. 2004. Inorganic and organic phosphorus fertilizer effects on the phosphorus fractionation in wetland rice soils. *Soil Sci. Soc. Am. J.* 68: 1635-1644.
- SAS Institute. 2000. SAS user's guide: Statistics, Ver 8.2; Cary, NC.
- Schmidt, J.P., S.W. Buol and E.J. Kamprath. 1996. Soil phosphorus dynamics during seventeen years of continuous cultivation: Fractionation analyses. *Soil Sci. Soc. Am. J.* 60:1168-1172.
- Sharpley A.N., S.J. Smith, J.W.B. Stewart, A.C. Mathers. 1984. Phosphorus forms in soil receiving cattle feedlot waste. *J Environ. Qual.* 13: 211-215.
- Simard, R.R., D. Cluis, G. Gangbazo and S. Beauchemin. 1995. Phosphorus status of forest and agricultural soils from a watershed of high animal density. *J. Environ. Qual.* 24: 1010-1017.
- Singh, M., A.K. Tripathi, K.S. Reddy and K.N. Singh. 2001_a. Soil phosphorus dynamics in a Vertisol as affected by cattle manure and nitrogen fertilization in soybean-wheat system. *J. Plant Nutr. Soil Sci.* 164: 691-696.
- Singh, M., Singh, V.P. and K.S. Reddy. 2001_b. Effect of integrated use of fertilizer nitrogen and farmyard manure or green manure on transformation of N, K and S and productivity of rice-wheat system on a Vertisol. *J. Indian Soc. Soil Sci.* 49: 430-435.
- Stewart, J.W.B. and A.N. Sharpley. 1987. Controls on dynamics of soil and fertilizer phosphorus and sulfur. pp. 101-121. In Follett, R.F., J.W.B. Stewart, and C.V. Cole (eds.), *Soil Fertility and Organic Matter as Critical Components of Production Systems*. Soil Sci. Soc. Am. Spec. Pub., 19, Soil Science Society of America, Madison, WI.
- Stewart, J.W.B. and H. Tiessen. 1987. Dynamics of soil organic phosphorus. *Biogeo. Chemistry.* 4: 41-60.
- Stewart, J.W.B., M.J. Hedley and B.S. Chauhan. 1980. The immobilization, mineralization and redistribution of phosphorus in soil. p. 276-306. In J. T. Harapiak (ed.) *Proc. Western Canada phosphorus symposium*. Alberta Soil Science Society, Edmonton, Canada.
- Sui, Y. and M.L. Thompson. 2000. Phosphorus sorption, desorption, and buffering capacity in a biosolids-amended Mollisol. *Soil Sci. Soc. Am. J.* 64: 164-169.
- Sui, Y., M.L. Thompson and C. Shang. 1999. Fractionation of phosphorus in a Mollisol amended with biosolids. *Soil Sci. Soc. Am. J.* 63: 1174-1180.
- Tiessen, H., J.W.B. Stewart and C.V. Cole. 1984. Pathway of phosphorus transformations in soils of differing pedogenesis. *Soil Sci. Soc. Am. J.* 48: 853-858.
- Williams, J. D.H., T. Mayer and J.O. Nriagu. 1980. Extractability of phosphorus from phosphate minerals common in soils and sediments. *Soil Sci. Soc. Am. J.* 35:250-255.

- Xie, R.J., J.W. Fyles. A.F. Mackenzie and I.P. O'Halloran. 1991. Lignosulphate retention in a clay soil: Causal modeling. *Soil Sci. Soc. Am. J.* 55:711-716.
- Zhang, T.Q. 1996. Chemical behavior of P in fertilized soils. Ph.D. diss. McGill Univ., Quebec, Canada.
- Zhang, T.Q. and A.F. MacKenzie. 1997. Changes of soil phosphorus fractions under long-term corn monoculture. *Soil Sci. Soc. Am. J.* 61:485-493.
- Zheng, Z., R.R. Simard, J. Lafond and L.E. Parent. 2002. Pathways of soil phosphorus transformations after 8 years of cultivation under contrasting cropping practices. *Soil Sci. Soc. Am. J.* 66: 999-1007

توزيع و تحولات الصور المختلفه للفوسفور فى التربه تحت تأثير الاضافات المستمره لبقايا المحاصيل و السماد الفوسفاتى
حسين محمد على راغب ، هاله حساتين جمعه، محمد فتحى غنيم و محمود رسلان
قسم الاراضى و المياه - كلية الزراعة - جامعة اسيوط

تم استخدام معدلين من بقايا المحاصيل (بدون وبإضافة بقايا المحاصيل) و ٦ مستويات من السماد الفوسفاتى، سوبر فوسفات، (٠، ١٠٠، ٢٠٠، ٣٠٠، ٤٠٠ و ٥٠٠ كجم سوبر فوسفات للفدان)، أضيفت لكل من المحصولين (الذرة والبرسيم)، تم خلط بقايا المحاصيل بالطبقة السطحية للتربة (٠،٢٥ - ٠،٢٥ م) عند انتهاء كل موسم نمو (٣،٧١ ، ٥،٤٥ طن للفدان من حطب الذرة و قش البرسيم ، على التوالى).

استجابة صور الفوسفور لإضافة كل من بقايا المحاصيل والسوبر فوسفات كانت قليلة إلى حد ما في بداية التجربة، على الرغم من أنها أصبحت أكثر ملاحظة بداية من سنة ١٩٩٩ إلى سنة ٢٠٠٣. التغيرات في Resin-P كانت معنوية نتيجة للإضافات المستمرة من بقايا المحاصيل والسوبر فوسفات لمدة ١٦ سنة. كانت نسبة ال Resin-P من الفوسفور الكلى في التربة من ٠،٨١-٠،٩٣ % و ١،٣-١،٥٢% بعد ٥ و ١٦ سنة من الإضافات المستمرة لبقايا المحاصيل والسوبر فوسفات، على التوالى. انخفض ال Resin-P من ٦،٥ إلى ٥،٦ جزء فى المليون عند استخدام المستوى صفر من السوبر فوسفات و بدون إضافة بقايا المحاصيل.

إزداد حجم الصورة العضوية والمعدنية للفوسفور المستخلص ب NaHCO_3 مع استخدام الإدارة للتسميد الفوسفاتى لمدة ١٦ سنة وتراوحت هذه القيم من ٢،٧٣ إلى ٣،١٣ ومن ١،٨٠ إلى ٢،١٩ للصورة المعدنية و العضوية، مع إضافة وعدم إضافة بقايا المحاصيل، على التوالى. أدت الإضافات المستمرة للسوبر فوسفات لمدة خمس سنوات بدون إضافة بقايا المحاصيل إلى زيادة التركيزات النسبية لل NaOH-P_i و NaOH-P_o من ٣،٦٩ إلى ٤،٤٤% ومن ١،٦٠ إلى ٢،٠٠% من الفوسفور الكلى، على التوالى. كما أن زيادة مستويات السوبر فوسفات المضافة من صفر (P_0) إلى ٥٠٠ كجم للفدان (P_5) لكل موسم نمو لمدة خمس سنوات أدت إلى زيادة كلا من HCl-P و residual-P زيادة معنوية و خطية.

أوضح استخدام تحليل المسار (path analysis) أن resin-P و $\text{NaHCO}_3\text{-P}_i$ تأثرت بشكل مباشر ومعنوي بالسماد الفوسفاتى. في إدارة التسميد الفوسفاتى مع إضافة بقايا المحاصيل، تأثر resin-P بشكل كبير بكل من السماد المعدني المضاف $\text{NaHCO}_3\text{-P}_i$ و HCl-P .