

## **EFFECT OF COOKING METHODS ON NUTRITIONAL QUALITY OF FABA BEAN**

**Saker, Dina M.\*; Akila S. Hamza\*; M. Abd-Alakhar\*\* ; M.A.A. Bekheit\*\* and El-Hashimy, Ferial S. A. \*\***

**\* Regional Center For Food and Feed, Agriculture Research center, Cairo, Egypt.**

**\*\*Food Science and technology, Faculty of Agriculture, Cairo university.**

### **ABSTRACT**

Faba bean products ,i.e., Medamnis, Bissara, Nabet soup, Falafel, green faba bean and cooked green faba bean were evaluated for its nutritional value . The obtained results revealed that green faba bean and cooked green faba bean contained the highest amount of tannins and phytic acid followed by dry faba bean , while Falafel and Bissara products showed the lowest contents. Green faba bean and dry faba bean showed the highest contents of trypsin inhibitor followed by cooked green faba bean, while other products showed the lowest contents. The results also indicated that Falafel, Bissara and dry faba bean showed the highest values of essential and non essential amino acids followed by Nabet soup, then Medamnis, while cooked and uncooked green faba beans showed the lowest values. Concerning protein utilization, the obtained results indicated that Bissara, Falafel and Nabet soup products showed the highest values of TD, BV and NPU, while cooked green faba bean , green faba bean and Medamnis products showed the lowest values . These obtained results revealed that the pretreatments practiced for preparation of these products and cooking method were found to have great effect on antinutritional factors and substantially on the nutritive value of the products.

### **INTRODUCTION**

Faba bean (*Vicia faba* L) is an important food in many countries such as China, Egypt, Italy, Brazil and Ethiopia. It is known by a variety of common names such as field bean, broad bean, horse bean, tick bean, windsor bean and others. It belongs to the family leguminosae (Fabaceae) (Adsule and Akpapunam, 1996).

Faba bean forms an important part of peoples diet in developing countries. It is consumed in the form of immature tender pods, green mature seed or as dry seeds after cooking. The method of cooking and consumption differ widely according to the geographical location. Some of the popular faba bean dishes in Egypt include Fool Medamnis, Taamia or Falafel and Fool Nabet (Askar, 1986).

Faba bean seeds have a relatively high protein content with low nutritional quality (Palmer and Thompson, 1975). Raw faba bean seeds were reported to contain antinutritional factors such as protease inhibitors, tannins, favism – inducing factors, haemagglutinins , flatus producing factors and phytates (Adsule and Akpapunam, 1996).

The present investigation was carried to study the antinutritional factors contents and the biological value of the different faba bean products which are widely consumed in Egypt.

## **MATERIALS AND METHODS**

### **Materials**

Dry and fresh faba beans (*Vicia faba*) Var. Giza-2- , the most common variety in Egypt, were obtained from the Legume Research Section, Agriculture Research Center , Ministry of Agriculture, Giza Egypt.

### **Methods**

#### **1-Preparation of faba bean products :**

Faba bean samples (the whole and dehulled seeds) were subjected to the following processes:

Soaking and germination processes were carried out as reported by Ghada Youssef (1999).

Cooked green faba bean was prepared by blanching the whole green faba bean horns in boiled water at the ratio of 1:1.5 W/V for 75 min with addition of 10 gm salt and 7 gm cumin in an aluminium pan.

Medammis product was prepared according to the method described by Ghada Youssef (1999).

Nabet soup, Falafel and Bissara products were prepared according to the method described by Bakr and Bayomy (1997).

Faba bean products were dried at 60°C for 18 hours in an electric air draught oven. The dried products were ground to pass through a 70 mesh sieve, packed into air- tight jars and kept at 4 °C. until analysis.

#### **2-Determination of amino acids contents :**

-Amino acid determination except for tyrosine and tryptophan was performed according to the method of the Official Journal of the European Communities 19-9-98.

-The system used for the analysis was high performance Amino Acid Analyzer, Beckman 7300 in Regional Center for Food and Feed, Agriculture Research Center, Cairo, Egypt.

#### **3- Determination of antinutritional factors:**

-Tannins contents were determined according to the method of Price *et al.*, (1978).

-Phytic acid contents were determined according to the method of Wheeler and Ferrel (1971).

-Trypsin inhibitor was determined according to the method of Kakada *et al.*, (1969).

#### **4- Biological evaluation:**

Biological evaluation of the processed products and green faba bean was carried out according to the procedures described by Eggum (1973) using adult male albino rats (90 animals) in 18 groups each of 5 rats. The experiment was spread over a period of 9 days. The rats were fed 150 mg N and 10 g dry matter per rat per day. The amount of dry matter was adjusted with N- Free diet supplied with vitamins and minerals. Nitrogen of urine and feces was determined according to the method described in A. O. A.C. (2002). True digestibility (T.D.), Biological value (B.V.) and net protein utilization (NPU) values were calculated according to the following equation:-

$$TD = \frac{N \text{ intake} - (\text{Faecal N} - \text{metabolic N})}{N \text{ intake}} * 100$$

$$BV = \frac{N \text{ intake} - (\text{Faecal N} - \text{metabolic N}) - (\text{urinary N} - \text{endogenous})}{N \text{ intake} - (\text{Faecal N} - \text{metabolic N})} * 100$$

$$NPU = \frac{T.D. * B.V.}{100}$$

where:

N=nitrogen

-Statistical analysis of the obtained results was carried out according to the method of Snedecor and Cochran (1980) using SAS program (1987) and IBM PC computer.

## RESULTS AND DISCUSSION

### 1-Amino acids contents of raw materials:

Amino acids content of raw materials under study were determined and the obtained results are presented in Tables (1 and 2) From these results it could be noticed that Falafel, Bissara and dry faba bean showed the highest values for essential amino acids contents followed by Nabet soup then Medamnis, while green faba bean and cooked green faba bean showed the lowest values. The same trend was also observed for nonessential amino acids. Youssef *et al.*, (1986) reported that severe heat processing, such as that used in the stewing process for Medamnis, drastically lowered some of the essential amino acids. This decline was explained by El-Mahdy(1974) on the basis of the effect of the prolonged heat treatment on the protein and their possible reactions with the reducing components or phenolic compounds of the beans. Youssef *et al.*, (1986) reported also that preparation of Bissara has improved the amino acid pattern, which can be attributed to advantages of the decortication process and addition of other ingredients to formulate Bissara. .Khalil (2001) found that germination of faba beans increased the total essential amino acid. Since tryptophane and tyrosine were not determined in our study, we compared our findings of essential amino acids, excluding the two mentioned amino acids, with FAO provisional pattern 1985. We found that the first limiting amino acid were methionine and cystine which were similar to (Marquardt and Campell, 1975) and El-Shemy (1996) who found that methionine and cystine were the first limiting amino acid in bean proteins. Supplementation of diets of faba bean with sulphur amino acids can improve its nutritional quality.

**Table 1: Essential amino acids contents of faba bean products (gm of amino acid per 16gm of protein nitrogen).**

Essential amino acids	dry faba bean	Medammis	Nabet soup	Falafel	Bissara	Green faba bean	Cooked green faba bean
Methionine	0.78	0.82	1.06	0.99	1.26	0.94	0.86
Threonine	3.50	3.56	3.52	4.33	4.29	2.49	2.40
Cystein	1.67	1.29	1.43	1.3	1.3	1.89	1.29
Valine	4.35	4.20	4.36	4.33	4.3	3.30	3.25
Isoleucine	3.77	3.51	3.37	4.39	4.11	2.76	2.53
Leucine	6.90	6.59	6.96	6.98	6.2	4.55	4.42
Phenylalanine	4.22	3.22	3.62	4.09	3.9	2.19	2.07
Lysine	6.08	5.98	6.26	6.47	6.22	4.35	4.03
Total determined essential amino acids	31.27	29.17	30.62	32.88	31.58	22.47	20.85

**Table 2: Non essential amino acids contents of faba bean products (gm of amino acid per 16 gm of protein nitrogen)**

Non essential amino acids	dry faba bean	Medammis	Nabet soup	Falafel	Bissara	Green faba bean	Cooked green faba bean
Aspartic	10.66	10.06	10.66	10.52	10.15	9.47	9.25
Serine	4.56	4.82	4.66	4.04	4.24	2.80	3.39
Glutamic	18.64	15.68	15.81	18.78	18.77	9.74	9.32
Proline	3.39	3.80	4.44	3.94	3.92	2.83	2.75
Glycine	4.47	4.02	3.66	4.5	4.3	2.73	2.70
Alanine	4.81	4.00	4.17	4.94	4.39	5.73	5.14
Histidine	2.13	2.38	2.20	2.42	2.44	1.69	1.60
Arginine	8.28	7.76	7.76	9.53	9.59	6.03	6.02
Total nonessential amino acids	56.94	52.32	53.56	58.65	57.80	41.02	40.17
total determined	88.21	81.49	84.18	91.53	89.38	63.49	61.02

## **2- Antinutritional factors of faba bean products**

Tannins, phytic acid and trypsin inhibitor, the major antinutritional factors of faba bean were determined and the obtained results are shown in Table (3) on dry weight basis.

**Table (3): Antinutritional factors contents of faba bean products**

Products	Total tannins (%)	Phytic acids (%)	Trypsin inhibitor Mg/100g
Dry faba bean	0.89	0.61	350
Medammis	0.55	0.35	50
Nabet Soup	0.48	0.36	42
Falafel	0.33	0.20	34
Bissara	0.33	0.25	40
Green faba bean	1.16	0.99	380
Cooked green faba bean	1.00	0.80	240

From the results presented in Table (3) it could be observed that green faba bean, cooked green faba bean and dry faba bean showed the highest contents for tannins and phytic acid, followed by Medammis and Nabet soup, while Falafel and Bissara showed the lowest contents for the same components. Concerning trypsin inhibitor, green faba bean and dry faba bean showed the highest contents followed by cooked green faba bean. However, Medammis, Nabet soup, Bissara and Falafel showed the lowest contents of

trypsin inhibitor, Thorn *et al.*, (1983) found that heat treatment reduced the activity of trypsin inhibitor in legumes by about 97%. Ologhobo and Fetuga (1984) reported that some processes, such as soaking, germination and cooking, were found to be most effective in decreasing phytate content and/or eliminate the phytic acid in legumes. Zienna (1989) found that up to 10% of the total tannins were decomposed by heat treatment during cooking of faba bean. Similar results were also reported by Vijayakumari *et al.*, (1996). They found that autoclaving of faba beans for 45 min lead to reduce tannins and phytic acid contents by about 71% and 44%, respectively. Soaking, germination and dehulling processes were also found to be effective in reducing trypsin inhibitor and phytic acid contents (Shekip *et al.*, (1989) and Sharma and Seghal, (1992).

**3-Biological evaluation of different faba bean products :**

Growing albino rats were fed on various faba bean products, i.e., Medammis, Falafel, Bissara, Nabet soup, cooked green faba bean and green faba bean in addition to casein diet for comparison. The nutritional quality of these different diets was evaluated and the true digestibility (TD), biological value (BV) and net protein utilization (NPU) values were calculated. The obtained results are shown in Table (4).

**Table (4) : Biological effect of different faba bean products**

Source of protein	True digestibility % (TD)	Biological value % (BV)	Net protein utilization % (NPU)
Medammis	74.16 <sup>m</sup> ± 3.78	61.56 <sup>bcde</sup> ± 7.35	45.76 <sup>ef</sup> ± 6.83
Falafel	90.33 <sup>abcd</sup> ± 2.52	72.50 <sup>ab</sup> ± 15.77	65.58 <sup>ab</sup> ± 14.55
Bissara	87.84 <sup>abc</sup> ± 0.90	65.82 <sup>bcde</sup> ± 15.01	57.80 <sup>bcde</sup> ± 12.98
Nabet Soup	84.54 <sup>a</sup> ± 5.55	64.50 <sup>bcde</sup> ± 13.08	54.50 <sup>bcde</sup> ± 7.63
Green faba bean	68.79 <sup>f</sup> ± 5.39	56.72 <sup>cde</sup> ± 11.40	39.02 <sup>f</sup> ± 10.15
Cooked green faba bean	72.12 <sup>g</sup> ± 2.01	53.85 <sup>cde</sup> ± 5.86	38.82 <sup>f</sup> ± 4.45
Casein	97.48 <sup>a</sup> ± 0.87	76.68 <sup>a</sup> ± 4.79	74.74 <sup>a</sup> ± 4.59
L.S.D. (0.05)	* 4.220	*** 11.663	** 9.851

Each value is the mean of 5 replicates; Means ± Standard error; Means in the same column with the same letter are not significantly different at P < 0.05; \*\*: Highly significant values; \*: Low significant values.

From the results presented in Table (4) it could be noticed that TD value of casein was 97.48% followed by Falafel (90.33%) then Bissara (87.84%) and Nabet soup (84.54%). Medammis, cooked green faba bean and green faba bean products showed the lowest TD Values i.e.74.16, 72.12 and 68.79%, respectively. The results of statistical analysis showed that all TD values of faba bean products were significantly different from casein diet . However no significant difference was observed between TD values of Falafel and Bissara products. The same trend was also found for TD values of Nabet soup and Bissara. Concerning BV , the obtained results indicated that Falafel , Bissara and Nabet soup were found to be not significantly different from that of casein diet while Medammis, green faba bean and cooked green faba bean showed significant differences in comparison with casein diet.

The results in the same Table indicated also that NPU of Falafel (65.58%)was not significantly different from that of casein (74.74%), while the

other studied products showed significant differences in compared to casein. However, no significant difference was observed for NPU values of Nabet soup, Falafel and Bissara. The same trend was observed for Medammis, green faba bean and cooked green faba bean. However, these results revealed that cooked green faba bean , green faba bean and Medammis products showed the lowest TD, BV and NPU values compared to other products .This might be due to its higher content of tannins (Table 3). It was reported that higher levels of tannins may have depressed the utilization of dietary protein by interacting with proteins to form indigestible compound, by inactivating proteolytic enzymes, by interference with the epithelial protective mucus of the intestine or by altering the absorption of digested nutrients OH and Hoff (1989). On the other hand , Bissara, Falafel and Nabet soup products showed the highest values of TD,BV and NPU . This could be related to the pretreatments practiced for preparing these products, such as soaking, germination and dehulling which lead to improve the nutritional value and caused a significant decrease in the antinutritional factors (Bakr, 1996).

From the obtained results it could be concluded that pretreatments practiced and cooking methods were found to have great effect on the nutritional quality of faba bean products and the products which were prepared from dehulled, soaked and germinated faba bean might be the best from the nutritional standpoint. The habit of consuming green faba bean should be limited.

## REFERENCES

- Adsule , R. N. and M. Akpapunam (1996). Faba bean (*Vicia faba* L.) in: Food and feed from legumes and oilseeds. Edited by E. Nwokolo J. Smartt. Published by Chapman and Hall. London- Glasgow- New York- Tokyo.
- A. O. A. C Association of Official Analytical Chemists (2002). Official methods of analysis association of official analytical. Chemists, 17<sup>th</sup> Edition , Revision 1,2002,Washington, USA.
- Askar, A.(1986).Faba beans (*vicia faba* (L.)) and their role in the human diet. Food Nutr. Bull., 8, 15-24.
- Bakr, A.A. (1996).Effect of Egyptian cooking methods of faba beans on its nutritive value, dietary protein utilization and iron deficiency anemia. 1. The role of main technological pretreatments.Plant Foods Hum Nutr 49(1):83-92
- Bakr, A.A. and M.F. Bayomy (1997).Effect of selected Egyptian cooking methods on faba bean nutritive value and dietary protein utilization 2: ability of faba bean products to support hemoglobin response in rats. Plant Foods Hum Nutr 50(1):81-91.
- Eggum, B.O.(1973). A study of certain factors influencing protein utilization in rats and Pigs. Kommission hos landhusholdningselskabets forlag Trykt frederikberg Bogtrykkeri.

- El-Mahdy, A.R.(1974). Evaluation of *Vicia faba* beans as a source of protein and the influence of processing thereon. PhD Thesis, Faculty of Agriculture university of Alexandria, Egypt.
- El- Shemy, H. A. (1996). Biochemical studies on some antinutritional factors. Ph.D. thesis, Fac. of Agric. Cairo Univ.
- Ghada Youssef, M.(1999). Effect of processing on antinutritional factors and biological value of major legumes. Ph.D. thesis, Fac. of Agric. Cairo Univ.
- Kakada, M.L.; N. Simons and I.E. liner (1969) . An evaluation of natural synthetic substrates for measuring the antitryptic activity of soy bean samples . *Cereal Chem.*, **46**: 518-526
- Khalil, M.M. (2001) .Effect of soaking, germination, autoclaving and cooking on chemical and biological value of guar compared with faba bean. *Nahrung* ;**45**(4):246-50
- Marquardt, R.R. and L.D. Campell (1975). Performance of chicks fed faba bean (*Vicia faba*) diets supplemented with methionine, sulphate and cystine. *Can. J. Anim. Sci.*, **55**, 213-18.
- Official journal of the European communities 19-9-98. Manual of high performance Amino Acid Analyzer, 7300 (Analyzer used). EZ Chrom ( soft ware used for data collection and processing.
- OH, H. J. and j.E. Hoff (1989). Effect of condensed grape tannins on in vitro digestion of proteins. *Nutr. Report Inter* **34**:51-58.
- C.F. Bakr, A.A. and Bayomy, M.F. *Plant Foods Hum Nutr* **50**(1) 81-91. (1997).
- Ologhobo, A. D. and B. L.Fetuga (1984).Distribution of phosphorus and phytate in some Nigerian varieties of legumes and some effects of processing. *J. of Food Sci.*, **49**, 199.
- Palmer, R. and R.Thompson (1975).A comparison of the protein nutritive value and composition of four cultivars of The nutritional evaluation of faba bean (*Vicia faba* L.). grown and harvested under controlled conditions . *J. Sci. Food Agric.*, **26** , 1577-83.
- Price, M. L.; S.V. Scoyoc and L.G. Butler (1978). A Critical Evaluation of the Vanillin Reaction as an Assay for Tannin in Sorghum Grain. *J.Agric .Food Chem.* **26** (5)1214-1217.
- SAS, (1987). SAS / STAT, guide for personal computers, version 6. SAS Institute, INC., Cary, NC.
- Sharma, A. and S. Seghal (1992). Effect of processing and cooking on the antinutritional factors of faba bean (*Vicia faba*). *Food Chem.*, **43**, 383.
- Shekib, L. A.; M. A. Abdel- Aal and M. M. Youssef (1989).The nutritive value of raw, decorticated, soaked and germinated and differently processed faba beans (*Vicia faba* L.). *Egypt. J. Food Sci.*, **17** (1-2), 47.
- Snedecor, G.W. and W.G. Cochran (1980). *Statistical methods*. 7<sup>th</sup> Ed. Published by the Iowa state university press, Ames ,Iowa, U.S.A.
- Thom, K. A.; A. M.Tinsley , C. W. Weber and J. W. Berry (1983). Antinutritional factors in legumes of Sonoran desert . *Ecology of Food and Nutr.*, **13**(4), 251.C.F. F.S.T.A., **17**(1), 1J142(1985).

- Vijayakumari, M. K.; P. Siddhuraju and K. Janardhanan (1996). Effect of different post-harvest treatments on antinutritional factors in seeds of the tribal pulse, *Mucuna pruriens* (L.) DC. *J Food Sci Nutr.* **47**(3):263-72.
- Wheeler E.L. and R.E. Ferrel (1971). A method for phytic Acid determination in wheat and wheat Fractions. *American Association of Cereal Chemists* **55**, 312-20.
- Youssef , M. M.; M.A. Hamza , M. H.Abdel Aal, L. A. Shekib, and A. A. El-Banna (1986). Amino acid composition and in vitro digestibility of some Egyptian foods made from faba bean (*Vicia faba* L.) . *Food Chem.*, **22**, 225.
- Ziena, H. M. S. (1989). Hard- to- cook phenomenon in relation to physical, chemical and biological properties of faba beans (*Vicia faba* L.). Ph.D. Thesis Fac. of Agric. Alex. Univ.

تأثير طرق الطبخ على الجودة الغذائية لمنتجات الفول البلدى  
دينا محمد صقر\* - عقيله صالح حمزه\* - محمود عبد الأخر\*\* -  
محمود على أحمد بخيت\*\* و فريال سيد أحمد الهاشمى\*\*  
\* المركز الأكليمي للأغذية والأعلاف - مركز البحوث الزراعيه - القاهره - مصر.  
\*\* قسم الصناعات الغذائية - كلية الزراعة - جامعة القاهرة.

تم دراسة لقيمة الغذائية لمنتجات الفول البلدى مثل الفول المدمس والبصارة وشورية الفول النابت والطعميه والفول الأخضر والفول الأخضر المطبوخ وتبين من النتائج أن الفول الأخضر والفول الأخضر المطبوخ أعلى قيمة في المحتوى من التانينات وحمض الفوتك يلي ذلك الفول الجاف بينما أظهرت النتائج أن منتجات الطعميه والبصارة أقل قيمة.  
كما لوحظ أن الفول الأخضر والفول الجاف أعلى قيمة في المحتوى من مثبطات التربسين ويلي ذلك الفول الأخضر المطبوخ . بينما لوحظ أن المنتجات الأخرى أظهرت قيمة أقل .  
المحتوى من الأحماض الأمينية الأساسية وغير الأساسية كان أعلى في الطعميه والبصارة والفول الجاف عنها في شوربة الفول النابت والفول المدمس بينما لوحظ أن الفول الأخضر المطبوخ وغير المطبوخ أظهرت قيمة أقل.  
لما فيما يخص الإستفادة القلبيه من البروتين فإن النتائج المتحصل عليها أظهرت أن منتجات البصارة والطعميه وشورية الفول النابت كانت الأعلى في القيمة الحيويه وكذلك في البروتين المستفاد والقابلية للهضم بالمقارنة بالفول الأخضر المطبوخ والفول الأخضر والفول المدمس التي أظهرت قيمة أقل لهذه المؤشرات.  
وتوضح من النتائج المتحصل عليها أن المعاملات الأولية التي تطبق أثناء إعداد هذه المنتجات وطرق الطبخ المختلفة كان لها تأثير كبير على مضادات التغذية وبالتالي على القيمة الغذائية لهذه المنتجات.