

BIOCHEMICAL STUDIES OF *Bacillus thuringiensis* AND *Spinosad* ON THE CARBOHYDRATES HYDROLYZING ENZYMES AND PHOSPHATASE ENZYMES OF COTTON LEAF WORM, *Spodoptera littoralis* WITH SPECIAL COMPARISON OF PYRETHROID COMPOUND

Salama. Z.A.; Ebied.A.A.; Ahmad.H. A. and El sheikh.T.A.
Faculty of Science Mansoura University

ABSTRACT

This investigation Show the effect of the three tested compounds on the activity of Haemolymph acid and alkaline phosphatase of 5th instar larvae of *S. littoralis* treated as 4th instar larvae with the LC₅₀ levels of *Spinosad*, *B. thuringiensis* and *Cypermethrin*. The results indicated that the acid phosphatase activity was insignificantly increased by about 2.55% more than the control in case of treatment with *B. thuringiensis*. On the other hand, *Spinosad* and *Cypermethrin* insignificantly decreased the activity of acid phosphatase. The application of the three tested materials to the 4th instar larvae of *S. littoralis* had changed the carbohydrate hydrolyzing enzymes as indicated that there was a large significant increase in amylase activity of 5th instar larvae of *S. littoralis* in case of *Spinosad* treatment; and treatment with *Cypermethrin* significantly decreased the amylase activity, whereas treatment with *B. thuringiensis* significantly decreased in compared to control. Invertase activity was significantly decreased in case of all tested compound treatments compared to control. Trehalase activity was significantly decreased, compared to control for *Spinosad* treatments. Whereas, *Cypermethrin* and *B. thuringiensis* treatment insignificantly decreased trehalase activity compared to control.

INTRODUCTION

The cotton leaf worm. *Spodoptera littoralis* which is considered as one of the major and economic pests not only in Egypt, but also in many parts of the world, infesting over 112 plant species, belonging to 44 families, including cotton *Gossypium hirsutum* (L.). The larval stage is known as a notoriously leaf eater accepting almost all herbaceous plants (Hill, 1975). The cotton leafworm larvae feed for about two weeks mostly on the leaves and occasionally on flowers and bolls. The use of insecticides for control of such pest proved to be the most accepted during the recent years. However, the practical application of different insecticides extensively has resulted in several problems such as development of resistance in field population of insects. There is many lepidopteran species that have been successfully controlled by microbial agents. This work was carried out to evaluate the control of *S. littoralis* larvae by the bacterium *Bacillus thuringiensis* var. *kurstaki*, the bacterium *spinosad* and the pyrethroid compound (*cypermethrin*).

MATERIALS AND METHODS

1- Tested Compounds:

In the present studies two entomopathogens and one pyrethroid compound were selected to test their effects against the 4th instar larvae of

Spodoptera littoralis.

These entomopathogens were,

1.1- *Bacillus thuringiensis* var. *kurstaki* Berliner

Produced by Valent Biosciences Corporation – USA

Commercial name *Diple 2x*

Common name *Bacillus thuringiensis* var. *kurstaki*

1.2- *Spinosad*

Commercial name Traceer

Common name *spinosad*

Chemical formula $C_{41}H_{65}NO_{10}$ (spinosyn A) + $C_{42}H_{67}NO_{10}$ (spinosyn D)

Spinosad (spinosyn A and spinosyn D) are a new chemical class of insecticides that are registered by the EPA to control a variety of insects. The active ingredient is derived from a naturally occurring soil dwelling bacterium called *Saccharopolyspora spinosa*, a rare actinomycete reportedly collected from soil in an abandoned rum distillery on a Caribbean Island in 1982 by a vacationing scientist. It has not been found in nature since that time, and was subsequently described as a new species. The bacteria produce compounds (metabolites) while in a fermentation broth. The first novel fermentation-derived compound was formulated in 1988. Spinosad has since been formulated into insecticides that combine the efficacy of a synthetic insecticide with the benefits of a biological pest control organism.

1.3- *Cypermethrin*

Commercial name Synthetic pyrethroid

Common name *cypermethrin*

Chemical formula $C_{22}H_{19}O_3NCl_2$

All of these tested compounds were obtained from the Plant Protection Research Institute, Ministry of Agriculture, Cairo.

2.1- Rearing of the Egyptian cotton leaf worm *Spodoptera littoralis* (Boisd.)

- The stock culture of susceptible Egyptian cotton leaf worm *Spodoptera littoralis* was reared on castor leaves (*Ricinus communis* L.) for several generations at laboratory conditions of 25 ± 2 °C and 70 ± 5% RH. Egg masses were placed on castor oil leaves in cylindrical glass jars. The jars were covered with muslin cloth and fastened with rubber band. First instar larvae hatched within 2-3 days. The newly hatched larvae were transferred into rearing jars bottomed with sheets of towel paper to absorb excess humidity. Castor bean leaves were provided daily to the larvae in sufficient amounts. The accumulated feces and debris were cleaned out daily. After pupation, pupae were collected and placed in wide clean jars until adult emergence. Then, the emerged adults were supplied with a piece of cotton wetted with 5-10% sugar solution and branches of Tafla (*Nerium oleander*) as suitable site for oviposition (El-Defrawi et al., 1964). Newly laid egg masses were collected daily and transferred into the rearing jars.

3- Preparation of samples for biochemical assay.

Haemolymph was collected from 3 pooled samples, each from 8-10 late 5th instar larvae fed as 4th instar for 24 hours on castor-oil leaves treated with the LC₅₀ values of each tested compound. One of the prolegs was removed and the Haemolymph was collected in cold tubes (on ice) previously coated

with crystals of phenylthiourea to prevent melanization. The samples were centrifuged at 2500 rpm for 5 minutes under cooling (4°C) to remove the blood cells. After centrifugation, the supernatant fluid was divided into small aliquots (0.5 ml) and stored at -20 °C until analysis.

Acid phosphatase activity was measured according to the method of Laufer and Schin (1971).

The methods used to determine the digestion of trehalose, starch and sucrose by trehalase, amylase and invertase enzymes respectively, were similar to those described by Ishaaya and Swirski (1976).

4- Susceptibility tests

A series of concentrations (in water) for each insecticide were prepared on the active ingredient (a.i) based on ppm by diluting the commercial formulation. Castor-bean leaves were dipped for 30 seconds in each concentration then left to dry for one hour. The 4th instars larvae of each tested strain were confined with treated leaves in glass jars covered with muslin for 24 hrs. Test also included a non treated leaves control in which leaves were dipped in water (as a check). Treated leaves were then removed and fresh untreated leaves provided for another day. Three replicates (each of 30 larvae) were tested for each concentration. Daily inspection was carried out for all treatments and mortality percentages were recorded after treatment. The average of mortality percentage was corrected using Abbott's formula (1925). The corrected mortality percentage of each compound was statistically computed according to Finney (1971). From which the corresponding concentration probit lines (Ic-p lines) were estimated in addition to determine 50 and 90% mortalities, slope values of tested compounds were also estimated.

RESULTS

1- Susceptibility test:

In the present studies Table(1) shows the susceptibility of the three tested compounds. The two bioinsecticides (*Bacillus thuringiensis*, *spinosad*) and pyrethroid compound (*cypermethrin*) used in the present work caused considerable toxic effects against the 4th larval instar of *Spodoptera littoralis* with cypermethrin and *Bacillus thuringiensis*. *Cypermethrin* was the most toxic to *S. littoralis* larvae but we should say that the toxicity of the bioinsecticide was significantly lower than that of the chemical insecticide. The LC₅₀ of *cypermethrin* was 1.675 followed by 7.59 and 28.868 ppm in case of treatment of the 4th instar larvae with *Bacillus thuringiensis* and *spinosad*, respectively.

2-Effects of LC₅₀ of *Bacillus thuringiensis*, *spinosad* and *cypermethrin* on carbohydrates hydrolyzing enzymes of 4th instar larvae of *Spodoptera littoralis*

2.1- Amylase enzyme

Amylase activity through the present study indicated that there was a large significant increase in amylase activity of 5th instar larvae of *S. littoralis* in case of *Spinosad* treatment; however treatment with *Cypermethrin* significantly decreased the amylase activity, whereas treatment with *B. thuringiensis* insignificantly decreased amylase activity in compared to the

untreated one as follow (1.92,1.15 and 1.24 with spinosad , cypermethrin and Bacillus thuringiensis respectively compared to 1.32 of the control as shown in Fig.(2) and Table (3)

2.2-Invertase enzyme

In this study all the tested compound had a remarked effect on Invertase activity in the Haemolymph oh 5th instar larvae of Spodoptera littoralis. Spinosad, cypermethrin and *B. thuringiensis* were significantly decreased the invertase activity compared to the untreated larvae as follow (72.37, 67.19 and 72.58% respectively compared to 100% for the untreated one.

2.3.Trehalase enzyme

The Change % in Trehalase enzyme decreased after the 4th day post- treatment in all treated larvae, but it was higher than the untreated larvae. *Spinosad* showed the highest level of enzyme activity followed by *cypermethrin* and *B.thuringiensis*, respectively. But the decrease was significant only with spinosad but non significant with larvae treated with *cypermethrin* and *B.thuringiensis*. Generally, change % in trehalase activity reached its maximum increase with *spinosad*.

3- Effects of LC₅₀ of *Bacillus thuringiensis*, spinosad and cypermethrin on Phosphatase in the Haemolymph of 4th instar larvae of *Spodoptera littoralis* Phosphatase activity:

Data obtained in table (2) and in Fig. (1) show the effect of the three tested compounds on the activity of Haemolymph acid and alkaline phosphatase of 5th instar larvae of *S. littoralis* treated as 4th instar larvae with the LC₅₀ levels of Spinosad, *B. thuringiensis* and Cypermethrin. The results indicated that the acid phosphatase activity was slight significantly increased by about 2.55% more than the control in case of treatment with *B. thuringiensis*. On the other hand, Spinosad and Cypermethrin insignificantly decreased the activity of acid phosphatase.

Table (1) Susceptibility of *Spodoptera littoralis* 4th instar larvae to Spinosad, Cypermethrin and *B. thuringiensis*

Treatments	LC ₅₀	95% Fiducial limits		Slope + SE	X ²
		Lower	Upper		
Spinosad	28.868	24.727	33.59	3.598 ± 0.576	2.026
Cypermethrin	1.675	1.441	1.99	3.509 ± 0.606	6.883
<i>B. thuringiensis</i>	7.59	6.332	9.29	2.583 ± 0.422	1.258

Table (2) Effects of LC₅₀ of *Bacillus thuringiensis*, spinosad and cypermethrin on Phosphatase in the Haemolymph in 5th instar larvae of treated 4th larval instar of *S.littoralis*

Treatments	Acid phosphatase activity	% change from the control	Alkaline phosphatase activity	%change from the control
Spinosad	4.30	99.53	6.13	70.37*
Cypermethrin	4.12	95.37	6.55	75.2*
<i>B. thuringiensis</i>	4.43	102.55	6.76	77.61*
Control	4.32	100	8.71	100

* =significant at P =0.05

But The tested compound caused a significant decrease in the activity of Haemolymph alkaline phosphatase of *S. littoralis* by 70.37, 75.2 and 77.61% with spinosad, Cypermethrin and *B. thuringiensis* respectively in respect to 100% in case of untreated larvae. Acid phosphatase activity predominated that of alkaline phosphatase in either non-treated or treated larvae Fig. (1).

Table (3): Effects of LC₅₀ of Spinosad, Bacillus thuringiensis and Cypermethrin on carbohydrates hydrolysing enzymes of 5th instar larvae of treated 4th larval instar of *S.littoralis*

Treatment	Amylase	% change from the control	Invertase	% change from the control	Trehalase	% change from the control
Spinosad	1.92	145.46*	3.22	72.37*	2.52	89.67*
Cypermethrin	1.15	87.12*	2.99	67.19*	2.65	94.3
<i>B. thuringiensis</i>	1.24	93.93	3.23	72.58*	2.68	95.37
Control	1.32	100	4.45	100	2.81	100

• =significant at P =0.05




Fig.(1): Effects of LC₅₀ of *Bacillus thuringiensis* , Spinosad and Cypermethrin on phosphatases enzymes in 5th instar larvae of *Spodoptera littoralis*.




Fig.(2): Effects of LC₅₀ of *Bacillus thuringiensis* , *Spinosad* and *Cypermethrin* on carbohydrates hydrolyzing enzymes in 5th instar larvae of *Spodoptera littoralis*.

DISCUSSION

1-Carbohydrate hydrolyzing enzymes

In the present study *spinosad* and *cypermethrin* caused decrease of Haemolymph amylase and invertase activities. These results agree with those obtained by El-Mageed *et al.* (2006) who observed that *Spinosad* decreased amylase and invertase activity in 4th instar larvae of *S. littoralis*.

1.1-Trehalase enzyme

The fat body is generally regarded as the principal site of trehalose biosynthesis in insects. Trehalase is activated for the production of glucose needed for chitin build-up in the newly synthesized cuticle; it is generally present in large amounts in the Haemolymph of most insects and it has the important function of energy supply to insect; and its activity might be an indicator of energy reserves resulting from availability of carbohydrate nutrient (Wyatt, 1967).

The Change % in Trehalase enzyme decreased after the 4th day post-treatment in all treated larvae, but it was higher than the untreated larvae. *Spinosad* showed the highest level of enzyme activity followed by *cypermethrin* and *B.thuringiensis*, respectively. But the decrease was significant only with *spinosad* but non significant with larvae treated with *cypermethrin* and *B.thuringiensis* generally, change % in trehalase activity reached its maximum increase with *spinosad*. Our results are in harmony with El-Ghar *et al.* (1995) who stated that *B. thuringiensis* at concentration of 200 p.p.m. reduced trehalase activity by 53% after 2 days of treatment. . Mordue and Blackwell (1993) reported that disrupted mid gut tissues would function abnormally at which the enzymes secretion and nutrient absorption would be disrupted. Rapid decrease of glucose concentration at the end of last instar larvae of *S. littoralis* was probably caused by high metabolic activity of the epidermis, which is known as a tissue with low trehalase, so it is unable

to utilize trehalose (Florkin and Jeauix, 1964). Therefore, as a result of treatment with spinosad, the activity of Haemolymph trehalase of treated insects in the present study lowers than that of non-treated ones

1.2- Amylase enzyme

Amylase activity through indicated that there was a large significant increase in amylase activity of 5th instar larvae of *S. littoralis* in case of *Spinosad* treatment; whoever treatment with *Cypermethrin* significantly decreased the amylase activity, whereas treatment with *B. thuringiensis* insignificantly decreased in compared to the untreated one. This is parallel to El-Ghar et al. (1995) found that *B. thuringiensis* (5 p.p.m.) caused a remarkable decrease in amylase activity at which maximum inhibition, about 77% was reduced 3 days after treatment also, at concentration 200 p.p.m. reduced amylase activity by 53% after 2 days of treatment.

1.3-Invertase enzyme

In this study all the tested compounds had a remarked effect on Invertase activity which significantly decreased in case of all tested compound treatments compared to control. Trehalase activity significantly decreased, compared to control similar to El-Ghar et al. (1995) found that *B. thuringiensis*, at from this result the great effect of this tested compounds appear with invertase enzymes.

2-Effect on phosphatases enzymes Acid and alkaline phosphatases:

Acid and alkaline phosphatases have been shown to be associated with insect development especially in relation to nutrition and egg maturation (Tsumuki and Kanehisa 1984). Acid phosphatase has received considerable attention in developmental studies because of its association with histolysis. This latter process is appreciable at the metamorphic moults of holometabolous species to which *S.littoralis* belongs. Show the effect of the three tested compounds on the activity of Haemolymph acid and alkaline phosphatase of 5th instar larvae of *S. littoralis* treated as 4th instar larvae with the LC₅₀ levels of *Spinosad*, *B. thuringiensis* and *Cypermethrin*. The results indicated that the acid phosphatase activity was insignificantly increased by about 2.55% more than the control in case of treatment with *B. thuringiensis*. On the other hand, *Spinosad* and *Cypermethrin* insignificantly decreased the activity of acid phosphatase agrees with the results obtained by Abdel Hafez et al. (1988) on *S. littoralis* larvae treated with diflubenzuron and triflumuron .

The present work showed insignificant decline in the activities of acid but indicated a significant decrease in alkaline phosphatase as a result of treatment with all tested compounds Ayyangar and Rao (1990) reported that injection of azadirachtin into 6th instar larvae of *S. littoralis* resulted in decreasing activities of alkaline phosphatase.

REFERENCES

- Abbott, W.S. (1925): A method of computing the effectiveness of an insecticide. *J. Econ. Entomol.* 18: 265 - 267.
- Abdel-Hafez, M. M.; Shaaban, M. M.; El-Malla, M. A.; Farag, M. and Abdel-Kawy, A. M. (1988). Effect of insect growth regulators on the activity of transaminases with reference to protein and amino acids in Egyptian cotton leafworm, *Spodoptera littoralis* (Boisd.). *Minia J. Agric. Res. Dev.*, 10 : 1357-13.

- Ayyangar, G. S. G. and Rao, P. J. (1990). Azadirachtin effects on consumption and utilization of food and midgut enzymes of *Spodoptera littora* (Fabr.). Ind. J. Entomol., 51: 373-376
- El-Defrawi, M.E.; Topozada, A.; Mansour, N. and Zeid, M. (1964): Toxicological studies on the Egyptian cotton leafworm, *Prodenia littora* (L.). I. Susceptibility of different larval instars of *Prodenia* to insecticides. J. Econ. Entomol., 57: 591-593.
- El-Ghar, G.E.S.A.; Radwan, H.S.A.; El-Bernawy, Z.A. and Zidan, L.T.M. (1995): Inhibitory effect of thuringiensis and abamectin on digestive enzymes and non-specific esterases of *Spodoptera littoralis* (Boisd.) (Lepidoptera: Noctuidae) larvae. J. App. Entomol., 119(5): 355-359.
- El-Mageed, -A-E-M-A; Elgohary, -L-R-A. (2006): Impact of spinosad on some enzymatic activities of the cotton leafworm. Pakistan-Journal-of-Biological-Sciences. 2006; 9(4):NO 713-716.
- Finney, D.J. (1971): Probit analysis. A Statistical Treatment of the Sigmoid Response Curve. 7th Ed., Cambridge Univ. Press, England.
- Florkin, M. and Jeanuiaux, C. H. (1964). Haemolymph composition. In "Physiology of Insecta": (Edited by Rockstein, M.), Vol. 3, pp. 109-152. Academic Press, New York & London.
- Hill, D.S. (1975) . Agriculture onect prts of tropic and their control. Cambridge univ., 516 pp.
- Ishaaya, I. and Swirski, E. (1976): Trehalase, invertase and amylase activities in the black scale, *Saissetia oleae* and their relation to host adability. J. Ins. Physiol., 16 : 1025-1025.
- Laufer, H. and Schin, K. S. (1971).
- Quantitative studies of hydrolytic enzymes activity in the salivary gland of *Chironomus tentans* (Diptera: Chironomidae) during metamorphosis. Can. Entomol., 103: 454-457
- Mordue, A. J. and Blackwell, A. L. (1993). Azadirachtin : an update. J. Insect Physiol., 39: 903-924.
- Tsumuki, H. and Kanehisa, K. (1984). Phosphatases in the rice stem borer, *Chilo suppressalis* Walker (Lepidoptera, Pyralidae) : Some properties and changes of the activities during hibernation. Cryobiology, 21: 177-182.
- Wyatt, G. R. (1967): In Insect Biochemistry and function (Candy and Kilby) 89-154, Chapman and Hall, London.

دراسات بيولوجية و كيميائية حيوية و هستولوجية للباسيليس ثيورجينسيس و سبينوساد على دودة ورق القطن سيدوبتيراليتوراليس و المقارنه مع بعض المواد البيروثرويدية.

سلامة زيدان احمد سلامة* - هدى عبد الحسيب احمد* - احمد عبد اللطيف عبيد* و طارق عفيفي عبد الحميد الشيخ**
*كلية العلوم - جامعة المنصورة
**مدرس- معهد بحوث وقاية النبات-مركز البحوث الزراعية وزارة الزراعة

التأثيرات البيوكيميائية إنزيمات الفوسفاتيز : أدت المعاملة بالباسيليس ثيورجينسيس Diple2x إلى زيادة غير معنوية في نشاط إنزيم الفوسفاتيز الحامضي بينما أدت المعاملة بسبينوساد (Spinosad) إلى نقص معنوي في نشاط إنزيم الفوسفاتيز القلوي. ومن ناحية أخرى أدت المعاملة بالثيرميثرين (Cypermethrin) إلى نقص معنوي في نشاط هذين الإنزيمين. الإنزيمات المحللة للكاربوهيدرات: وجدت زيادة معنوية لإنزيم الأميليز في حالة سبينوساد (Spinosad) بينما وجدت نقص غير معنوية لنشاط هذا الإنزيم في حالة المعاملة بكل من بالباسيليس ثيورجينسيس Diple2x والثيرميثرين (Cypermethrin) . أدت المعاملة بالباسيليس ثيورجينسيس Diple2x والثيرميثرين (Cypermethrin) إلى نقص غير معنوية في نشاط إنزيم الترباليز وعلني العكس أدى استخدام سبينوساد (Spinosad) إلى نقص معنوي في نشاط ذلك الإنزيم. والواضح أيضا من هذه الدراسة أن جميع المواد الثلاثة أدت إلى حدوث نقص معنوي في نشاط إنزيم الإنفرتيز.