

FACTORS AFFECTING MICROTUBER FORMATION IN SOME POTATO GENOTYPES

El-Sayed, S.F.¹; A.A. Gharib¹; O.A Elshihy^{2,3} and Rasha R. Eid³

- 1- Vegetable Crops Department, Faculty of Agriculture, Cairo University, Giza, Egypt.
- 2- Botany Department, Plant Physiology Section, Faculty of Agriculture, Cairo University, Giza, Egypt.
- 3- Plant Biotechnology Research Lab, Faculty of Agriculture, Cairo University, Giza, Egypt.

ABSTRACT

The effect of some factors, namely BA concentration (5 and 10 mg/l), sucrose concentration (8, 10, 12, 13, 14 and 15%) and presence of active charcoal in medium (0 and 5 g/l) on microtuber formation of six potato genotypes, four tetraploid, viz. Diamant, Picasso, Santa and Forella and two dihaploid, namely 805 and 808, was studied in two experiments. Potato genotypes responded differently to such factors, where Forella cv. showed the best performance regarding microtuber number per jar, average weight of microtuber, and microtuber yield per jar. Increasing sucrose concentration up to 12% and adding active charcoal at concentration of 5 g/l in medium having 10 mg/l BA caused a remarkable increase in microtuberization parameters in all cultivars. On the other hand, increasing sucrose concentration or adding 5 g/l active charcoal to MS medium having 5 mg BA /l promoted microtuber formation in some cultivars, and inhibited this process in others.

INTRODUCTION

The production of microtubers has got significant importance in many countries of the world for the purpose of the production of disease-free seed tubers, germplasm exchange, and conservation (Roca *et al.* 1979; Ranalli *et al.* 1994). Mass production system of potato microtuber can revolutionize the world potato agriculture in the near future. Moreover, many laboratories are producing microtubers commercially for their own national programs as well as for export. In South Korea, they are producing millions of microtubers annually (Hyoun *et al.* 1994). In the Philippines, the national breeding project on conservation and multiplication has a strong microtuber production programme in order to produce millions of minitubers every year (Paet and Zamora 1994). Baoqing *et al.* (1991) harvested at least ten usable size microtubers per flask after 30 days of culture. They tried to give a picture of a large scale and economic production of microtubers for the national programme with a comprehensive research outline. In USA, some commercial laboratories are producing millions of microtubers, which are sold with 15 cent each, while in Indonesia and China it costs only 5 cent due to cheap labor cost.

In vitro microtuberization is a complex process regulated by different factors. However, work on micro tuberization in potato has mainly focused on

the use of growth regulators, such as benzyl amino purine (Wang and Hu 1982; Hussey and Stacey 1984; Abbott and Belchel 1986), 2-chloroethyl trimethyl ammonium chloride (CCC) (Tovar *et al.* 1985), coumarin (Dodds *et al.* 1988), and jasmonoid (Pelacho *et al.* 1993), etc. Cytokinins are commonly included in media for in vitro tuberization of potatoes (Wang and Hu 1982), but their role is still obscure (Ewing 1985). Doubts have been raised whether cytokinins are directly responsible for tuber initiation (Ewing 1987). Addition of cytokinin to the agar medium promoted tuberization of stolons subcultured in darkness (Palmer and Smith 1970) and cytokinin like activity was nearly twice as high in induced as in non-induced tissues (Forsline and Langille 1976; Mauk and Langille 1978). Generally, the response to growth regulators depended upon a range of factors including sucrose concentration and cultivars (Hussey and Stacey 1984; Ortiz- Montiel and Lozoya-Saldana 1987; Garner and Blake 1989). Concerning the effect of sucrose concentration, Yan-Zheng and Guo-Dezhang (2004) found that the best medium for inducing microtuber production consisted of MS, 8% sucrose, 0.1% Active carbon, 30 mg l⁻¹ coumarin and 3 mg l⁻¹ 6-BA. Also Gopal *et al.* (2004) reported that higher concentration of sucrose (60-80 g/liter) promoted biomass production, microtuber production and microtuber dry matter content. Tariq *et al.* (2004) reported that MS medium supplemented with higher concentration of sucrose (6%) and 1 µ M of BA recorded the maximum number of micro tubers. Sidikou *et al.* (2003) studied the effect of sucrose level at 2-12% and BA at 0 - 5 mg/ liter on tuberization and found that tuberization increased with increasing concentration of BA and sucrose. Islam *et al.* (1999) investigated the effect of sucrose level (6, 8, 10 and 12%) and BA (4, 5 and 6 mg /liter) on the tuberization, and they observed that the highest number of tubers per flask and average tuber weight were upon treatment with 6% sucrose, 5 mg/liter BA. Alix *et al.* (2001) studied the effect of sucrose level (1, 2, 4 and 8%) on potato microtuber initiation, and found that microtuber initiation was 100% at 8% sucrose falling to 40-50% at 4% sucrose and was absent at 1 or 2% sucrose. Regarding cultivar response, Hossain and Sultana (1994) reported that the preliminary studies on microtuberisation showed that the potato cultivars differed widely in this respect, i.e. the cultivar Patrones produced about 11 tubers per plant as against one - two for Heera, Dheera or Chamak. Medium containing activated charcoal gave the highest rate of tuberization and the largest microtubers in potato. It thus played a role in optimizing conditions for rapid, mass tuberization of these cultivars, and produced large microtubers for field planting (Bizarrri. *et al.* 1995; Wang *et al.* 2006). Bulblet growth was also improved in liliium by supplement medium of AC. The medium with 2 g/L AC was most effective for bulblet growth (Bong *et al.* 2005). As it was reported by Hossain (2005), most of the studies were based on a few genotypes only. Thus, the present experiment was aimed at investigating the effect of different factors, namely, BA concentration, sucrose concentration, and presence of active charcoal on microtuberization over a w

MATERIAL AND METHODS

These experiments were carried out during the period from 2005-2006 in Plant Biotechnology Research Lab, Faculty of Agriculture, Cairo University, Giza, Egypt.

Six potato genotypes, namely Santa, Picasso, Diamant, Forella, 808, and 805 were used in the present experiments. The cultivars Santa, Picasso, and Diamant were received from Agriculture Research Center, Giza, Egypt, while Forella cv. and the dihaploid genotypes 805 and 808 were received from IPK Gene bank, Aussenstelle Nord, Gross Lusewitz, Germany.

Experiment treatments:

This work included two experiments as follows:

1- Effect of sucrose and active charcoal concentrations on microtubers production in six different genotypes.

This experiment consisted of 96 treatments, i.e., 8 sucrose concentrations (8, 9, 10, 11, 12, 13, 14 and 15%) X 2 active charcoal levels (0 and 5 g/l) X 6 potato genotypes (4 tetraploid, namely Diamant, Picasso, Santa, Forella, and 2 dihaploid, namely 808 and 805).

2- Effect of BA, sucrose concentration and active charcoal level on microtuber production in three different potato cultivars.

This experiment included 12 treatments which was a combination of 3 sucrose concentrations (8, 10 and 12%), 2 active charcoal levels (0 and 5 g/l) X 2 BA (benzyl adenine) concentrations (5 and 10 mg/l). This experiment was conducted separately on 3 different potato cultivars, namely Diamant, Picasso and Santa.

Plant material

Potato tubers (*Solanum tuberosum* L. cvs Diamant, Picasso, and Santa) were established from apical meristems of sprouted tubers on MS medium (Murashige and Skoog, 1962) with 3% sucrose, 100mg⁻¹ myo-inositol, 0.1mg⁻¹ BA and solidified with 0.7% agar. pH was adjusted to 5.8 before autoclaving at 121°C for 25 min. Sprouts were grown under 16h photoperiod at 2500 lux of white cool light and incubated at 24°C for 4 weeks in growth chambers. Forella, 805 and 808 genotypes were produced from Gen bank, Germany as plantlets in tubes. Thereafter, Shoot of all genotypes, i.e., Santa, Picasso, Diamant, Forella, 808 and 805 were subcultured every 28 days in baby gars (250 ml capacity) containing 40 ml of the same previously described culture medium via nodal cuttings as described by Espinoza *et al.* (1985). Subcultures were repeated for 4 times before cultured on tuberization medium.

Tuberization medium and incubation condition

Tuberization medium consisted of MS macro and micro elements salts, 0.7% agar and the additives mentioned in the experiments, i.e., 6 sucrose concentrations (8, 9, 10, 11, 12, 13, 14 and 15%) as well as two active charcoal levels (0 and 5 g/l) in the first experiment and two BA concentrations (5 and 10 mg/l), 3 sucrose concentration (8, 10 and 12%)

and two active charcoal levels (0 and 5 g/l) in the second experiment. pH was adjusted to 5.8 before autoclaving at 121°C for 25 min.

Three shoots, each shoot containing 6 nodal, were cultured on different media cultures which were kept at 17°C in continuous dark.

Recording data

Data were recorded after 3 months from culture in the two experiments as follows:

- 1- Number of microtubers per jar.
- 2- Average weight of microtuber.
- 3- Yield of microtubers per jar.

The experimental design

The experiments were set in a three factors completely randomized design (CRD) in four replicates, each replicate consisted of 3 jars, each jar contained 3 plants. The three factors of the first experiment were 6 cultivars X 8 sucrose concentrations X 2 active charcoal levels, while in the second experiment, they were 2 BA concentrations, 3 sucrose concentration X 2 active charcoal levels. The data on different parameters were collected and statically analyzed. Thereafter, the significant differences between means of treatments were recorded at 5% level using L.S.D as described by Steel and Torrie (1960).

RESULTS

Experiment 1: Effect of sucrose concentration and active charcoal on microtubers production in six potato genotypes.

There were significant differences in microtuber number, microtuber average weight and yield per jar among the different genotypes. In this respect, Forella cultivar showed the best performance in all tuber characters and followed by genotype 808 in respect of microtuber number and followed by cv. Picasso regarding average weight of microtubers, and jar yield of microtubers. On contrast, genotype 805 showed the lowest value of microtuber number per jar, average microtuber weight and jar yield of microtubers, were recorded by genotypes Santa, 808 and 805, respectively. Increasing sucrose concentration from 80 to 120 g/l significantly increased number of microtubers, microtuber average weight and jar yield per jar. This was true in all genotypes (Table 1, 2 and 3). Further increase in sucrose concentration up to 13% led to death of all genotypes, except Forella cultivar (Data are not shown). Data presented in the same table reveal that adding active charcoal to the tuberization medium led to production of higher microtuber number per jar, and markedly increased both of average microtuber weight and microtuber yield per jar, as compared with medium without active charcoal. These results were true in all genotypes and at all sucrose concentrations of 80, 90, 100, 110 and 120 g/l.

The highest values of average microtuber weight (1.933 g) and microtuber yield per jar (18.027 g) were noticed in cv. Forella by using 12% sucrose in presence of active charcoal. Meanwhile, the highest number of microtubers per jar (11.3) was produced in genotype 808 in medium containing 12% sucrose + 5 g/l active charcoal.

Table. 1: Effect of concentrations of active charcoal (A.C), and sucrose on number of tuber / jar of some potato genotypes.

Genotypes	A.C (g /l)	Sucrose level (g/l)					Mean
		80	90	100	110	120	
Diamont	0	1.7	3	2	3.6	2.3	2.5
	5	3.3	4	5.3	5.6	7	5.1
Average		2.5	3.5	3.6	4.6	4.6	3.8
Picasso	0	2	2.7	2	2.6	3	2.5
	5	3.3	4	4	4	4.6	4
Average		2.7	3.3	3	3.3	3.8	3.2
Santa	0	1	1.7	1.3	2.3	2.3	1.7
	5	2.7	3	2.6	2.6	3.6	2.9
Average		1.8	2.3	2	2.5	3	2.3
Forella	0	3.3	3.7	5	6.3	7.3	5.1
	5	3.7	4	5.6	6.6	9.3	5.8
Average		3.5	3.4	5.3	6.5	8.3	5.5
805	0	1	2.3	2.3	2.6	1.6	2.2
	5	2	2	2.6	3.3	5.6	3.1
Average		2	2.2	2.5	3	3.7	2.6
808	0	2.3	2	2.6	4	6	3.4
	5	2.3	2.3	4	4	11.3	4.8
Average		2.3	2.6	3.3	4	8.7	4.1
0	0	2.1	2.5	2.5	3.6	3.8	2.9
	5	2.9	3.2	4	4.4	1.1	4.4
Mean		2.5	2.9	3.31	4	5.4	

L.S.D. at 0.05 for Factor A (genotypes) 0.7
 Factor B (A.C) = 0.4 Factor C (sucrose) = 0.7
 A X B = 1.1 B X C = 0.9 A X C = 1.7 A X B X C = 2.4

Table. 2: Effect of concentrations of active charcoal (A.C) and sucrose on tuber average weight (g) of some potato genotypes.

Genotypes	A.C (g /l)	Sucrose level (g/l)					Mean
		80	90	100	110	120	
Diamont	0	0.06	0.09	0.10	0.11	0.12	0.10
	5	0.11	0.12	0.13	0.14	0.15	0.13
Average		0.09	0.11	0.12	0.13	0.14	0.12
Picasso	0	0.10	0.11	0.12	0.13	0.14	0.12
	5	0.11	0.12	0.13	0.14	0.15	0.13
Average		0.11	0.12	0.13	0.14	0.15	0.13
Santa	0	0.11	0.12	0.13	0.14	0.15	0.13
	5	0.12	0.13	0.14	0.15	0.16	0.14
Average		0.12	0.13	0.14	0.15	0.16	0.14
Forella	0	0.12	0.13	0.14	0.15	0.16	0.14
	5	0.13	0.14	0.15	0.16	0.17	0.15
Average		0.13	0.14	0.15	0.16	0.17	0.15
805	0	0.13	0.14	0.15	0.16	0.17	0.15
	5	0.14	0.15	0.16	0.17	0.18	0.16
X		0.14	0.15	0.16	0.17	0.18	0.16
808	0	0.14	0.15	0.16	0.17	0.18	0.16
	5	0.15	0.16	0.17	0.18	0.19	0.17
Average		0.15	0.16	0.17	0.18	0.19	0.17
0	0	0.15	0.16	0.17	0.18	0.19	0.17
	5	0.16	0.17	0.18	0.19	0.20	0.18
Mean		0.15	0.16	0.17	0.18	0.19	0.17

L.S.D. at 0.05 for Factor A (genotypes) = 0.07
 Factor B (A.C) = 0.224 Factor C (sucrose) = 0.035
 A X B = 0.055 B X C = 0.050 A X C = 0.087 A X B X C = 0.123

Table. 3: Effect of concentrations of active charcoal (A.C), and sucrose on jar yield (g) of some potato genotypes.

Genotypes	A.C	Sucrose level (g/l)					Mean
		80	90	100	110	120	
Diamont	0	.118	.211	.143	1.074	.297	.278
	5	.807	.787	.700	2	2.094	1.089
Average		.487	.097	.424	1.032	1.947	.983
Picasso	0	.703	.309	.270	4	1.000	.020
	5	2.719	2.770	1.820	1.777	2.842	2.402
Average		1.437	1.489	1.022	1.032	2.427	1.489
Santa	0	.109	.218	.327	1.020	1.722	.780
	5	.422	.004	.080	1.790	2.771	1.192
Average		.266	.231	.203	1.308	2.242	.937
Forella	0	.322	.848	2.048	2.220	0.729	2.018
	5	1.077	1.777	2.002	0.477	18.027	0.904
Average		.849	1.300	2.770	4.270	11.882	4.276
805	0	.212	.327	.377	.470	.403	.370
	5	.309	.328	.490	.822	2.277	.800
Average		.261	.322	.431	.648	1.377	.788
808	0	.170	.148	.322	.094	.987	.042
	5	.327	.372	.777	.992	7.171	1.720
Average		.251	.270	.049	.792	2.079	1.087
0	0	.307	.302	.079	1.140	1.709	.818
	5	.877	.877	1.072	1.214	7.112	2.290
Mean		.092	.712	.947	1.722	2.910	

L.S.D. at 0.05 for Factor A (genotypes) = 0.580

Factor B (A.C) = 0.335 Factor C (sucrose) = 0.530

A X B = 0.821 B X C = 0.749

A X C = 1.298 A X B X C = 1.836

Experiment 2: Effect of BA concentration, sucrose concentration and active charcoal level on microtubers production in three different potato cultivars.

BA at 10 mg/l significantly increased number of microtubers per jar and microtubers yield per jar in cv. Picasso and Diamant as well as average microtubers weight in cv. Diamant as compared with the concentration of BA at 5 mg/l. On the contrary, in cv. Santa the reverse was true, where number of microtubers /jar, average microtubers weight and jar yield of microtubers were greater when BA was used at 5 mg/l, as compared with the concentration of 10 mg/l. Adding active charcoal to medium in presence of BA at 10 mg/l significantly increased number of microtubers /jar, average microtubers weight and jar yield of microtubers. This was true in all cultivars. In contrast, adding active charcoal to MS medium having 5 mg BA/l led to significant reduction in number of microtubers /jar in cv. Santa and Picasso and jar yield of microtubers in cv. Santa. Meanwhile, cv. Diamante responded differently to active charcoal in presence of BA at the concentration of 5 mg/l, where active charcoal promoted greater number of microtubers /jar, average microtubers weight and jar yield of microtubers. Increasing sucrose concentration significantly increased number of microtubers per jar in cv. Picasso at both concentrations of BA; i.e., 5 and 10 mg/l. Meanwhile, increasing sucrose concentration in the medium of the cultivars Diamante and Santa had a different effect, depending on BA concentration. Whereas

the increase in sucrose concentration led to an increase in the number of microtuber in presence of 10 mg BA/l, it caused a decrease in microtuber numbers in presence of 5 mg BA/l. Concerning effect of sucrose concentration on average microtubers weight and microtubers yield per jar, data in the same Tables reveal that both of average microtubers weight and microtubers yield per jar were promoted by raising sucrose concentration in presence of BA at 10 mg/l in medium. These results were true in all cultivars. On the other hand, in presence of BA at 5 mg/l medium increasing sucrose concentration from 80 to 120 g/l increased average tuber weight in the three cultivars. Meanwhile, such increase in sucrose concentration at the concentration of 5 mg/l BA led to increment, reduction and no effect in microtubers yield per jar in cv. Picasso, Santa and Diamant, respectively. The highest values of number of microtubers per jar, average microtubers weight and microtubers per jar were recorded by using 12% sucrose + 10 mg BA/ l in presence of active charcoal in cv. Diamant and Picasso. On the other hand, in cv. Santa the highest values of microtubers number per jar, average microtubers weight and microtubers per jar were produced by the utilization of 5 mg BA/l +8% sucrose without active charcoal, 5 mg BA + 12% sucrose + 5 g active charcoal and 10 mg BA+ 12% sucrose + 5 g active charcoal, respectively.

Table. 4: Effect of concentrations of benzyl adenine (BA), active charcoal (A.C) and sucrose on tuber number per jar, average tuber weight (g) and jar yield (g) of Santa cultivar.

BA (mg / l)		Tuber number/ jar	Average tuber weight (g)	Jar yield (g)
10		2.4	.,2.2	.,032
5		2.0	.,271	.,229
L.S.D		.,4	.,.26	.,.79
A.C (g / l)				
.		2.7	.,182	.,012
0		2.7	.,292	.,248
L.S.D at 0.05		.,4	.,.47	.,.79
Sucrose (g / l)				
8.		2.1	.,174	.,06.
10.		2.6	.,2.6	.,001
12.		2.2	.,23.	.,279
L.S.D at 0.05		.,0	.,.22	.,978
BA X AC				
10	0	1.8	.,12.	.,192
	5	2.0	.,286	.,871
5	0	2.7	.,204	.,822
	5	2.2	.,298	.,650
L.S.D at 0.05		.,7	.,267	.,112
BA X Sucrose				
1.	8.	2.1	.,126	.,271
	10.	2.2	.,2.7	.,482
	12.	2.8	.,238	.,841
8.	8.	4.1	.,214	.,849
	10.	2.0	.,2.6	.,62.

Table 5: Effect of concentration of benzyl adenine (BA), active charcoal (A.C) and sucrose on tuber number per jar, average tuber weight (g) and jar yield (g) of Diamant cultivar.

BA (mg / l)	Tuber number/ jar	Average tuber weight (g)	Jar yield (g)
10	4.8	0.262	1.914
5	3.4	0.202	0.731
L.S.D	0.0	0.061	0.128
A.C (g / l)			
.	3.0	0.147	0.260
0	0.2	0.410	2.280
L.S.D at 0.05	0.0	0.061	0.128
Sucrose (g / l)			
8.	4.2	0.262	1.110
10.	3.9	0.266	1.209
12.	4.1	0.219	1.700
L.S.D at 0.05	0.6	0.061	0.129
BA X AC			
10	0	3.6	0.179
	5	6.1	0.047
	0	3.0	0.112
	5	0.2	0.287
L.S.D at 0.05	0.7	0.094	0.190
BA X Sucrose			
10.	8.	4.0	0.272
	10.	4.4	0.242
	12.	0.6	0.270
	8.	4.1	0.102
	10.	3.0	0.189
	12.	2.0	0.212
L.S.D at 0.05	0.9	0.066	0.228
A.C X Sucrose			
.	8.	3.4	0.160
	10.	2.6	0.129
	12.	3.0	0.124
	8.	0.2	0.26
	10.	0.2	0.292
	12.	0.1	0.044
L.S.D at 0.05	0.9	0.066	0.228
BAX AC X Sucrose			
10	0	80	3.0
		100	3.2
		120	4.0
	5	80	0.0
		100	0.0
		120	7.2
5	0	80	3.2
			0.089
			0.202
			0.066
			0.044
			2.731
			2.809
			4.484
			0.208

Table 6: Effect of concentration of benzyl adenine (BA), active charcoal (A.C) and sucrose on tuber number per jar, average tuber weight (g) and jar yield (g) of Picasso cultivar

BA (mg / l)		Tuber number/ jar	Average tuber weight (g)	Jar yield (g)
10		3,3	0.303	1,328
5		1,8	0.367	0,682
L.S.D		0,4	0,030	0,123
A.C (g / l)				
.		2,4	0,292	0,687
o		2,6	0,437	1,223
L.S.D at 0.05		0,4	0,030	0,123
Sucrose (g / l)				
8.		2,3	0,229	0,042
10.		2,3	0,294	0,700
12.		3,0	0,070	1,773
L.S.D at 0.05		0,4	0,042	0,213
BA X AC				
10	0	2,7	0,187	0,487
	5	3,9	0,020	2,170
5	0	2,2	0,397	0,687
	5	1,3	0,304	1,323
L.S.D at 0.05		0,0	0,049	0,174
BA X Sucrose				
10.	8.	2,9	0,269	0,762
	10.	2,3	0,271	0,883
	12.	3,0	0,021	2,340
5.	8.	3,0	0,190	0,323
	10.	3,8	0,318	0,017
	12.	0,0	0,619	1,207
L.S.D at 0.05		0,7	0,070	0,213
A.C X Sucrose				
.	8.	2,4	0,189	0,432
	10.	2,1	0,267	0,049
	12.	2,8	0,421	1,079
o	8.	2,3	0,269	0,602
	10.	2,4	0,321	0,801
	12.	3,3	0,719	2,467
L.S.D at 0.05		0,7	0,070	0,213
BAX AC X Sucrose				
10	0	80	0,181	0,471
		100	0,147	0,324
		120	0,233	0,660
5	80	3	0,306	1,003
		100	0,390	1,441
		120	0	4,010
5	0	80	0,197	0,394

DISCUSSION

In the present investigation, there were significant differences among the different genotypes in the number of microtubers per jar, average microtubers weight and microtubers yield per jar (Table 1, 2 and 3). The average microtubers number per jar ranged from 1 in cv. Santa to 11.3 in the genotype 808. Meanwhile, the average microtubers weight ranged from 0.056g in cv. Diamant to 1.933 g in cv. Forella. On the other hand, microtubers yield per jar ranged from 0.109 g in cv. Santa to 18.027 g in cv. Forella. Such levels were produced from culturing 3 shoots per jar. The previous studies revealed that Zakaria (2003) harvested 8.12 microtubers per Erlenmeyer flask of cv. Diamant. Hossain (2005) Obtained from 0.5 to 4 microtubers per test tube having three micro plants and found that the average microtubers weight ranged from 0.066 g to 0.105 g and the yield of each tube ranged from 0.103 g to 0.407 g, depending on genotype. The differences in genotype response are due to the varietal differences and their interaction to the different environmental condition. The highest values of the number of microtubers per jar, average microtubers weight and microtubers yield per jar in the present work was obtained in all cultivars, except in cv. Santa, in medium contented 10 mg BA / l + 12% sucrose and 5g/l active charcoal. Hossain (2005) used in his study MS medium containing 5 mg BA/l + 500 mg CCC/l + 8% sucrose in absence of active charcoal. The result in respective of average microtubers weight indicated obtaining high value up to 1.933 g (in cv. Forella). Simko (1990), Hossain and Sultan (1998), Zakaria (2003) and Yassmin (2005) obtained a mean microtubers weight of about 100 mg from culturing their cultivars in the dark. In contrast, Randhawa and Chamder (1990) obtained much higher microtubers weight ranging from 0.850 to 2.060 g in MS +10 mg BA/l + 8% sucrose in six Indian potato cultivars, which is contradictory to the findings of the present investigation. Probably, varietal differences and their response are main reasons for such a good performance for some genotypes (Tovar *et al.* 1985).

The present study proved that increasing sucrose concentration up to 12% in medium containing BA at concentration of 10 mg/l led to improving of microtuberization. This concentration of sucrose caused a significant increase in number of microtubers per jar, microtubers average weight and microtubers yield per jar. These results were achieved in all tested genotypes in the two conducted experiments (Table1-6). On the other hand, in medium containing 5 mg BA/l, increasing sucrose concentration increased number of microtubers in cv. Picasso, whereas this concentration of sucrose decreased number of microtubers in cv. Diamant and Santa. The previous study indicated that the highest microtubers per flask were obtained by using 6% sucrose in presence of 5 mg BA/l as compared with sucrose level at 8, 10 and 12 % (Islam *et al.* 1999). In other studies, the best medium for inducing microtubers production contained 8% sucrose + 3 mg BA/l (Yan-Zhong and Guo –Dezhang, 2004; Gopal *et al.* 2004), 6% sucrose + 1 μ M BA (Tariq *et al.* 2004). Also, Sidikuo *et al.* (2003) studied the effect of sucrose level at 2 -12% and BA at 0-5 mg /l on tuberization and found that tuberization increased with

increasing concentration of BA and sucrose. The results indicated the importance of the interaction between sucrose level and BA concentration. The present investigation also indicated that BA at 10 mg/l increased microtuberization in cv. Picasso and Diamant, as compared with BA at 5 mg/l, whereas the reverse was achieved in cv. Santa. As it was mentioned before addition of cytokinin to agar medium promoted tuberization in the darkness (Palmer and Swith 1970). The optimum concentration of BA for microtuberization was 10 mg for some cultivars (Randhawa and Chamder 1990) and 5 mg/l for other (Randhawa and Chamder 1990; Sidikuo *et al.* 2003). This clearly indicates that different genotypes are mainly responsible for such a wide variation. Adding active charcoal to medium in presence of BA at 10 mg/l significantly increased number of microtubers /jar, average tuber weight and jar yield of microtubers per jar. This was true in all cultivars. In contrast, active charcoal adding to MS medium having 5 mg BA/l led to significant reduction of number of microtubers /jar and jar yield of microtubers in cv. Santa and Picasso. Meanwhile, cv. Diamante responded differently to active charcoal in presence of BA at the concentration of 5 mg/l, where active charcoal promoted greater number of microtubers /jar, average microtubers weight and jar yield of microtubers. The previous investigations indicated that medium containing activated charcoal gave the highest rate of tuberisation and the largest microtubers in potato. It thus played a role in optimizing conditions for rapid, mass tuberization of potato cultivars, and produced large microtubers for field planting (Bizarri. *et al.* 1995; Wang *et al.* 2006). Generally, it was previously proved that media with charcoal had substantially lower levels of phenylacetic and O-OH benzoic acids, compounds that may inhibit tuberization (Fridborg *et al.* 1978). Activated charcoal also has been shown to absorb 5- hydroxymethyl furfural, an inhibitor formed by sucrose degradation during autoclaving, as well as substantial amounts of auxin and cytokinins. Thus it may absorb inhibitors that would prevent growth as well as reduce the level of growth promoters that would cause continued proliferation.

REFERENCES

- Abbott, A.J. and Belcher, A.R. (1986). Potato tuber formation *in vitro*. In: Plant tissue culture and its agricultural applications. LA Withers and PG Alderson (eds.). Butterworths, London, pp. 113-122.
- Alix, M.J; Savvides, S; Blake, J; Herrmann, R. and Hornung, R. (2001). Effects of illumination source, culture ventilation and sucrose on potato (*Solanum tuberosum* L.) microtuber production under short days. *Ann. appl. Biol.* 139: 175-187.
- Baoqing, L.; Miaochang, Z.; Jin L.X.; Jun W. and Bofu, S. (1991). Large scale production of *in vitro* potato tubers: An example. *Asian Potato J.*, 2: 13-15.
- Bizarri, M.; Borghi, L. and Ranalli, P.(1995). Effects of activated charcoal effects on induction and development of microtubers in potato (*Solanum tuberosum* L.). *Ann. appl. Biol.* 127: 175-181
- Bong Hee Han ; Byeoung Woo Yae ; Hee Ju Yu and Kee Yeoup Peak. (2005). Improvement of *in vitro* micropropagation of *Lilium oriental*

- hybrid 'Casablanca' by the formation of shoots with abnormally swollen basal plates. *Sci. hortic.* 103: 351-359.
- Dodds, J.H.; Tovar, P.; Chandra, R.; Estrella, D. E. and Cabello, R. (1988). Improved methods for in vitro tuber induction and use of in vitro tubers in seed programs. Proc. of the symp.on improved potato planting material. Kunming : China. June 21-24, 1988, pp. 49-64.
- Espinoza N, Lizarraga R, Siguenas, C Buitron, F Bryan J and Dodds JH (1992). Tissue culture, micropropagation, conservation and export of potato germplasm. CIP Research Guide 1, 16 p.
- Ewing, E.E. (1985). Cuttings as a simplified models of the potato plant. In: Li PH, ed. *Potato Physiology*. Orlando, Florida: Academic Press, 154-207.
- Ewing, E.E. (1987). The role of hormone in potato(*Solanum tuberosum* L.) tuberization. In: Davies PJ, ed. *Plant hormones and their role in plant growth and development*. Dordrecht, The Netherlands: Martinus Nijhoff Publishers, 515-538.
- Fridborg, G.; Pedersen, M.; Landstron, L.E. and Eriksson, T. (1978). The effect of activated charcoal on tissue cultures. A adsorption of metabolites inhibitors morphogenesis. *Plant.* 43: 104-106.
- Forsline, P. and Langille, A.R. (1976). An assessment of the modifying effect of kinetin on in vitro tuberization of induced and non-induced tissue of *Solanum tuberosum*. *Canadian J. of Bot.* 54: 2514-2516.
- Garner, N. and Blake, J. (1989). The induction and development of potato microtubers in vitro on media free of growth regulating substances. *Ann. Bot.* (London), 63: 663-674.
- Gopal, J.; Chamail Anjali and Sarkar Debabrata. (2004). In vitro production of microtubers for conservation of potato germplasm: Effect of genotype, Abscisic acid and sucrose. *In vitro Cell & Develop. Biol. Plant.*, 40: 485 – 490.
- Hossain, M. J., (2005). In vitro microtuberisation in potato obtained from diverse sources. *Plant Tissue Cult. & Biotech.* 15: 157-166.
- Hossain, M. J. and Sultana, N. (1994). Research report on tissue culture research tuber crops. Tuber Crops Research Centre, BARI, 23-25 July, pp. 1-21.
- Hossain, M.J. and Sultana, N. (1998). Effect of benzylaminopurine (BAP) and chlorocholinechloride (CCC) on in vitro tuberisation of potato. *Bangladesh J. Agr. Res.*, 23: 685-690.
- Hussey, G.; and Stacey, N.J. (1984). Factors affecting the formation of *in vitro* tubers of potato (*Solanum tuberosum* L.). *Ann. Bot.*, 53: 565-578.
- Hyoun, J.; Jeong, K.S.; Jeon, H.J. and Jang, R.L. (1994). Mass production of potato microtubers by novel tissue culture techniques and its agricultural applications. Proc. of the 4th Triennial Conf. and Symp. of the Asian Pot. Assoc., 4-9 July, South Korea. p. 21.
- Islam, M.S; Chowdhury, A.R. and Hossain, M.M. (1999). Microtuber production of six potato varieties affected by temperature, sucrose and BAP. *Ann. Bangladesh Agricul.*, 9: 91-97.
- Mauk, C.S. and Langille, A.R. (1978). Physiology of tuberization in *Solanum tuberosum* L.: cis zeatin ribosid in the potato plant – its identification

- and changes in endogenous levels as influenced by temperature and photoperiod. *Plant Physiol.* 62: 438-442.
- Ortiz-Montiel, G.; and H Lozoya-Saldana. (1987). Potato minitubers: technology validation in Mexico. *Am. Potato J.*, 64:535-544.
- Paet, C.N.; and Zamora, A.B. (1994). Production of *in vitro* tuberlets and greenhouse planting. Proc. of the 4th Triennial Conf. and Symp. of the Asian Pot. Assoc., 4-9 July, South Korea, pp. 17-18.
- Pelacho, A.M.; Viscasillas, T. and Mingo-Castel, A.M. (1993). Photoperiod dependence of *in vitro* jasmonic acid induced tuberisation in (*Solanum tuberosum*) ssp. *andigena*. Proc. Of the 1th Triennial Conf. of the European Assoc. for Pot. Res. Paris, France, July 18-23, 1993, pp. 249-250.
- Palmer, C.E. and Snlittl, O.E. (1970). Effect of kinetin on tuber formation on isolated stolons of *Solanum tubelusum* L. cultured *in vitro*. *Plant Cell Physiol.*, 11: 303-314.
- Randhawa, G.J. and Chandra, R. (1990). *In vitro* tuberisation in six potato cultivars. *J. Indian Potato Assoc.*, 17: 199-201.
- Ranalli P., Bassi, F; Ruaro, MR.; Del Re P.; Dicandilo, M. and Mandolino, G. (1994). Microtuber and minituber production and field performance compared with normal tuber. *Potato Res.*, 37:383-391.
- Roca, W.M.; Bryan, J.E. and Roca, M.R.(1979). Tissue culture for international transfer of potato genetic resource. *Am. Pot. J.*56: 1-10.
- Sidikou, R.D.S; Siha chakr, D; Lavergne, D; Nato, A; Ellisseche.D; Jouan, B. and Ducreux, D. (2003). Contribution of microtubersation to the adaptation of potato culture in the sahel. *Cahiers Agri*, 12: 7- 14.
- Simko, I. (1990). Tuberizacia {emiakov v podmilenkach *in vitro* poosetreni Paklobutrazolon. *Biologia (Bratislava)*, 46: 251-256.
- Steel, R. G. D. and Torrie, J. H. (1960). Principles and Procedures of Statistics with Special Reference to the Biological Sciences. McGraw Hill Book Co., Inc., New York 481p.
- Tariq, R.; Jaskani, M. J.; Hasnain R. and Abbas, M. (2004). Intro studies on microtuber induction in potato. *Inter. J. Agri. & Biolo.*, 6: 375-377.
- Tovar, P.; Estrada, R.; Schilde-Rentschler, L. and Dodds., J.H. (1985). Induction and use of *in vitro* potato tubers. CIP Circular, International Potato Centre., 13: 1-4.
- Wang, P.J. and Hu, C.Y. (1982). *In vitro* mass tuberisation and virus-free seed potato prodcution in Taiwan. *Amer. Potato. J.*, 59: 33-37.
- Wang Rui-bin, Wang Di Si and Huai-jun (2006). Study on induction of potato microtuber by inte raction of edible sugar with activated charcoal. *J. Gansu Agri. Univ.*, 4:35-40
- Yasmin, L. (2005). Production of microtubers in potato (*Solanum tuberosum* L.) MS thesis, Depar of Biotech, Bangladesh Agri Univ, Mymensingh. pp.1-81.
- Zakaria, M. (2003). Induction and performance of potato microtuber. Ph.D. thesis, Depar of Hort, BSMRKB, Gazipur, Bangladesh.
- Yan-Zhen and Guo-Dezhang (2004). Effect of plant hormones on microtuber production of potato. *Chinese potato J.*, 23: 84-86.

بعض العوامل المؤثرة علي عملية تكوين الدرينات في البطاطس

سيد فتحى السيد^١، احمد علي غريب^١، اسامة محمد الشحي^٢ و رشا رمضان عيد^٣

١- قسم الخضر كلية الزراعة - جامعة القاهرة

٢- قسم النبات الزراعي فرع فسيولوجيا النبات كلية الزراعة - جامعة القاهرة

٣- مشروع زيادة تحمل الارز و القمح للملوحة باستخدام تقنيات الهندسة الوراثية - كلية الزراعة جامعة القاهرة

اجريت تجربتين لدراسة تأثير بعض العوامل المؤثرة علي عملية تكوين الدرينات في البطاطس مثل استخدام البنزايلى اندين بتركيز (10,5 mg /l)، للسكروز بتركيز (٨، ١٠، ١٢، ١٣، ١٤ و ١٥%) و اضافة الفحم للنشط في بيئة الزراعة بتركيز (صفر، ٥ g/l) و ذلك باستخدام ٦ من انواع و اصناف البطاطس، اربعة tetraploid هم : Santa، Picasso، Diamant، و اثنان dihaploid: ٨٠٥ و ٨٠٨. انواع و اصناف البطاطس كان لها رد فعل مختلف لكلا من هذه العوامل، الصنف Forella . اظهر افضل اداء من ناحية عدد الدرينات لكل برطمان، و متوسط وزن الدرينات لكل برطمان، و محصول البرطمان من الدرينات. زيادة تركيز السكروز إلى ١٢% و اضافة الفحم للنشط بتركيز ٥ g /l الي بيئة الزراعة المحتوية علي mg /l ١٠ بنزايلى اندين ادي الي حدوث زيادة ملحوظة في لوزان و اعداد الدرينات في كل الاصناف. من ناحية اخرى، زيادة تركيز السكروز أو اضافة ٥ g /l من الفحم نشط إلى بيئة MS المحتوية علي mg /l 5 بنزايلى اندين حفز عملية تكوين الدرينات في بعض الاصناف و ثبطها في البعض الاخر