

EFFECT OF SOME NITROGENOUS AND PHOSPHATIC FERTILIZERS SOURCES AND VA-MYCORRHIZA INOCULUM ON GROWTH, PRODUCTIVITY AND STORABILITY OF GIRLIC (*Allium sativum* L.) ;

1-Vegetative growth and chemical constituents.

Dawa, K. K.*; E. A. A. Tartoura*; A. M. Abdel-Hamed**and A. E. A. I. Gouda**

* Hort. Dept., Fac. of Agric., Mansoura Univ.

** Hort. Res. Inst., Agric. Res. Center

ABSTRACT

Two field experiments were carried out at a private farm at El-Manial village, Talkha province, Dakahlia governrate during winter seasons of 2004/2005 and 2005/2006. The study aimed to investigate the effect of N-sources (calcium nitrate, ammonium sulphate, sulphur coated urea and bentonite coated urea), P-sources (super phosphate either powder and granules and rock phosphate) as well as vesicular arbuscular mycorrhiza (VAM) fungi and their interactions on the vegetative growth (plant height, number of leaves/plant, dry weight, leaf area/plant, chlorophyll content/plant, neck diameter, bulb diameter and bulbing ratio) and chemical constituents (N, P, K, NO₃ and NO₂) of garlic plants (Sids-40).

The obtained results showed that plant height, dry weight, leaf area/plant, total chlorophyll content, neck diameter and bulb diameter were significantly increased by application of ammonium sulphate, in both seasons, while, number of leaves/plant was not significantly affected by N-sources, in both seasons. Plant height, dry weight, leaf area, total chlorophyll and bulb diameter were significantly increased by application of powder super phosphate, in both seasons, while, number of leaves/plant, neck diameter and bulbing ratio were not significantly affected by P-sources, in both seasons. All measured parameters were significantly increased by inoculation of VAM fungi except of number of leaves/plant and bulbing ratio were not significantly affected by VAM fungi, in both seasons.

Application of sulphur coated urea significantly increased the percentage content of N, P and K and significantly reduced NO₃ and NO₂, in both seasons and the lowest values of NO₃ and NO₂ obtained by application of bentonite coated urea, in both seasons. Application of granules super phosphate significantly increased N % and K % and the concentration of NO₃ and NO₂ reduced by using rock phosphate followed by using powder super phosphate, in both seasons, while, P % was not significantly affected by P-sources. Inoculation of VAM fungi significantly increased the concentration of N %, P % and K % and this treatment significantly reduced the concentration of NO₃ and NO₂, in both seasons.

INTRODUCTION

Garlic (*Allium sativum* L.) as a member of the *Alliaceae* Family is one of the most important vegetable bulb crop. It is commonly used as a spice or for many medicinal purposes. In Egypt, it has been generally cultivated for both local consumption and export. Increasing garlic productivity and improving bulb quality are important aims. Growing desired genotypes and using suitable form of nitrogen could be good tools to

accomplish these aims. Use of slow release nitrogenous fertilizers (SRNF) is one of the most important alternatives to rationalize the use of soluble nitrogenous fertilizers, besides to protect the environment from nitrogenous residues pollution, whether for soil or water or atmosphere. Where nitrogen element is released at a slow rate throughout one season or more, the plants are able to take up most of it without waste. SRNF also promotes steady and uniform growth for guarantees high N-efficient use and reduces N-losses either by volatilization or leaching (*Abbadly et al.*, 1997 and *El-Mallah et al.*, 1998).

Phosphorus is one of the most important nutrients for garlic growth, productivity and storability in heavy soil. In Egypt the most important problem of phosphorus fertilization is that the amount available for plant is usually low since the phosphorus of the applied fertilizers could be converted to unavailable form for plant absorption by its reaction with the soil constituents (*Kumar and Sharma*, 2004).

Vesicular-Arbuscular Mycorrhiza (VAM) is the symbiotic association of a fungus and the roots of plant. This association alters the root structure and helps plants to obtain water and minerals, particularly phosphorus. In addition, mycorrhizal fungi (VAM) are extremely widespread in the most of plants grown in Egypt (*El-Shaikh*, 2005).

Application of ammonium sulfate as a source of nitrogen gave the maximum growth rate, photosynthetic pigments, fresh and dry weight, leaf area, chlorophyll content/plant and bulb diameter, followed by application of sulphur coated urea, *El-Beheldi et al.* (1985), *Abdel-Hamed et al.* (1996) and *Gad El-Hak and Abd El-Mageed* (2000). *Sang et al.* (2001) found that the slow release fertilizers, urea formaldehyde slightly increased the plant height, leaf sheath diameter and leaf number compared with conventional fertilizer treatment. *Tartoura et al.* (2003) showed that rock phosphate coated urea gave the significant increases in most of vegetative characters followed by bentonite coated urea.

Wani and Konde(1998) found that application of different P-sources with VAM fungi to garlic plants significantly increased plant height, number of leaves/plant and plant weight over the non-mycorrhizal treatment and *Nagaraju et al.*(2001) on onion.

Inoculation of VAM fungi to garlic plants significantly increased all measured parameters (vegetative growth and chemical constituents) and significantly reduced NO₃ and NO₂, in both seasons (*Koch et al.* 1997, *Wani and Konde* 1998, *Lubraco et al.* 2000 on garlic, *Hammad and Abdel-Ati*, 1998 on potato *Nagaraju et al.* 1999 on onion, *Awad* 2002 on potato and *El-Morsy et al.* 2002 on sweet potato.

MATERIALS AND METHODS

Two field experiments were carried out during the two successive winter seasons of 2004-2005 and 2005-2006 at EL-Manial village, Talkha province, Dakahlia governorate to study the effect of some nitrogenous and phosphatic fertilizers sources and Vesicular Arbuscular Mycorrhiza (VAM)

fungi inoculum on growth and chemical constituents of garlic. Physical properties for experiment soil

(texture class) was clay loam and pH = 8.1 and 7.9 in the first and second year, respectively.

The experimental design and treatments:-

The experiment was carried out in a split split- plots design with three replicates. The four nitrogen fertilizers sources occupied the main plots which were subdivided to three sub plots each contained one of the phosphatic fertilizers sources while the VA- Mycorrhiza fungi treatments were assigned to the sub-sub plots. The sub-sub plots area was 10.5 m² which contained five rows, 3.5 m long and 0.6 m wide. The full dose of recommended chemical fertilizers rates (120 kg N + 75 kg P₂O₅ + 72 kg K₂O \ fed.) the experiment includes 24 treatments, which were the combination of 4 nitrogen sources x 3 phosphorus sources x 2 VAM fungi. Four sources of chemical nitrogenous fertilizers were used, soluble as: Calcium nitrate (15.5 % N), ammonium sulfate (20.5 % N) and slow release N fertilizers as: Sulphur coated urea (37.8 % N) and bentonite coated urea (37.8 % N) which were taken from Egyptian Fertilizers Development Center (EFDC). Also three sources from phosphatic chemical fertilizers were used, soluble as:- calcium super phosphate powder (15.5 % P₂O₅), calcium super phosphate granules (15.5 % P₂O₅) and insoluble rock phosphate (28 % P₂O₅) which was taken from Abo-Tartor, Suez governorate. The soluble N fertilizers were applied as 50 % at 30 days after planting and 50 % one month later while the slow release N fertilizers were applied as one dose at 30 days after planting. The soluble and insoluble phosphatic fertilizers were applied as one dose at 30 days after planting. VAM fungi are the most widespread associations between fungi and plant. Three species of indomycorrhizal fungi (*Glomus fasciculatum*, *Glomus mosseae* and *Glomus monosporum*) were supplied by Botany Department, Faculty of Science, Mansoura University, Egypt. A fifty grams of VAM inoculum were added to the root absorption zone of each plant at 21days after planting before irrigation. The planting was carried out during the first week of October for both seasons of study. Uniformed cloves were hand planted on both sides of the ridges at 10 cm apart. All field plots were fertilized with 72 kg k₂o/fed. as potassium sulfate(48 % k₂O) equally divided and added after 30 and 60 days after planting. The other cultural practices for garlic commercial production were used according to the instruction laid down by the Ministry of Agriculture, Egypt. The harvesting was done 180 days after planting for both seasons.

Data recorded and statistical analysis

Vegetative growth characteristics:-

At 120 days after planting, a random of 5 guarded plants were taken from each plot to estimate traits of plant vegetative growth as follow:-

a-Plant height:It was measured in cm starting from tip of the longest leaf blade to the base of bulb.

b-Number of leaves per plant:All visible leaves of chosen plants were counted (dry and undifferentiated leaves were excluded).

c- Leaf area per plant:The leaves of chosen plants were cleaned from dust and given 10 disks (1.54 cm diameter) from the leaves of each plant were

taken, dried and weighted up to 0.1 mg. Plant leaf area was calculated according to the method described by Kollar (1972) using following formula:

$$\text{Leaf area in m}^2 = \frac{\text{Leaves dry weight}}{\text{Disks dry weight}} \quad (\text{No. of disks} \times \text{disk area})$$

d-Dry weight of whole plant:The chosen plants of each treatment were cleaned from the dust, and dried at 70 °C till constant weight.

e-Total chlorophyll:Chlorophyll contents were determined by A Minolta SPAD chlorophyll Meter (Yadava, 1986). Chlorophyll meter reading were taken on 6th leaf of plant. A Minolta SPAD chlorophyll meter uses light sources and detectors to

measure the light transmitted by a plant leaf at two different wavelengths (one in the red and one in the infrared region of the spectrum). The ratio of the light transmittance at these wavelengths, in addition to the ratio determined with no sample, is processed by the instrument to produce a reading shown on a digital display. This reading is in SPAD units, which are values, defined by Minolta to indicate the relative amount of chlorophyll contained in plant leaves (Piekielek and Fox, 1992).

f- Bulbing ratio: It is measured as reported by Mann(1952).

Neck diameter (cm)

$$\text{Bulbing ratio} = \frac{\text{Neck diameter (cm)}}{\text{Bulb diameter (cm)}}$$

Diameters of both plant neck and bulbs were determined by using caliper.

Chemical constituents:-

Sample (100 gm) of dried cloves of each treatment was ground and wet digested as described by Hesse (1971) to determined:-

a- Total nitrogen % determined according to the method described by Chapman and Partt(1961).

b- Phosphorus (%) determined as reported by John (1970).

c- Potassium (%) determined by the method of Brown and Lilleland (1946)

d- Concentrations of NO_3^- and NO_2^- (ppm) were determined by the modified method of Singh (1988).

Data statistical analysis:-All obtained data were subjected to statistical analysis of variance and the least significant difference (L.S.D.)was calculated as mentioned by Gomez and Gomez (1984).

RESULTS AND DISCUSSION

Vegetative growth characteristics :

1- Effect of N – sources :

Data presented in Table(1) show that plant height, dry weight, total chlorophyll content/plant, neck diameter and bulb diameter were generally significantly increased, in both seasons with application of ammonium sulphate fertilizer, followed by sulphur coated urea, in both seasons, while number of leaves significantly not affected by N – sources, in both seasons, and bulbing ratio was significantly reduced by application of calcium nitrate followed by ammonium sulphate and sulphur coated urea, in the first season.

Leaf area was significantly increased by application of sulphur coated urea, in both seasons. These increases occurred on plant vegetative growth characteristics as a result of application ammonium sulphate and sulphur coated urea fertilizers could be attributed to that two fertilizers contain N – element plus S – element, whereas calcium nitrate and bentonite coated urea are not have. N and S elements are presented in the molecule of most amino acids, whereas, Ca present very less in the minority of the amino acids, this in turn, stimulating division and elongation of cells and hence plant growth. Further more, nitrogen is an essential component of many organic compounds in plant, such as proteins, enzymes, chlorophyll, vitamins, carotenoids, hormones and nucleic acids (Russell, 1950). These compounds may be play an important role in cell formation and plant development. The obtained data are in harmony with those of EL- Beheidi *et al.* (1985), Abdel-Hamed *et al.* (1996), Gad El-Hak and Abdel-Mageed(2000) and Sang *et al.* (2001).

2 – Effect of P – sources :-

Data presented in Table (1) show that there were significantly increases on plant height, dry weight, total chlorophyll content/plant and bulb diameter by application of super phosphate powder followed by super phosphate granules and rock phosphate at the last. Leaf area/plant was significantly increased by application of super phosphate granules, in both seasons. While, number of leaves / plant, neck diameter and bulbing ratio were not significantly affected by the P – sources . These increases due to that super phosphate powder fertilizer has the best solubility, it makes better uptake by the plant roots and the beneficial effect of P – element on the activation of photosynthesis and metabolic processes of organic compounds in plant and hence increasing plant growth whereas, super phosphate granules has the best guard from contact with the soil particles, so, it was away from fixation. These results are in agreement with those of Nagaraju *et al.* (2001) on onion , Kumar and Sharma (2004) on cabbage and Shabana (2004) on tomato.

3 – Effect of VAM fungi : -

Data presented in Table (1) reveal that inoculation of garlic plants by VAM fungi led to significant increases in plant height, dry weight, leaf area/plant, total chlorophyll content/plant, neck diameter and bulb diameter in both seasons, while, number of leaves / plant and bulbing ratio were significantly not affected by inoculation of VAM fungi, in both seasons.

The superiority effect of VAM fungi could explained based on their role in supplying the growing plants with available phosphorus needs, some micronutrients and phytohormones, such as gibberellins, auxins and cytokinins which promoted plant growth, In addition to root development and thereby enhanced nutrient uptake (Marschner 1995), Obtained results go well with those of Koch *et al.* (1997), Wanl and Konde (1998) on garlic and El – Morsy *et al.* 2002) on sweet potato.

Table 1: Plant height, number of leaves/plant, dry weight /plant, leaf area/plant, total chlorophyll content/plant SPAD unit, neck diameter/plant, bulb diameter and bulbing ratio of garlic at 120 days after planting as affected by some nitrogen and phosphorus fertilizer sources as well as VAM fungi during 2004/2005 and 2005/2006 seasons.

Characters Treatments	Plant height (cm)		Number of leaves /plant		Dry weight /plant (g)		Leaf area/ plant (m ²)		Total Chlorophyll content/plant SPAD unit		Neck diameter/plant (cm)		bulb diameter (cm)		bulbing ratio	
	2004/ 2005	2005/ 2006	2004/ 2005	2005/ 2006	2004/ 2005	2005/ 2006	2004/ 2005	2005/ 2006	2004/ 2005	2005/ 2006	2004/ 2005	2005/ 2006	2004/ 2005	2005/ 2006	2004/ 2005	2005/ 2006
A- Nitrogen fertilizer sources																
Calcium nitrate	78.41	79.34	10.04	9.64	12.48	12.95	0.149	0.157	100.42	99.39	1.08	1.11	3.50	3.42	0.309	0.326
Ammonium sulphate	82.07	82.69	10.28	10.22	13.91	13.95	0.170	0.176	101.86	101.22	1.37	1.35	4.13	4.11	0.334	0.332
Sulphur coated urea	80.44	81.27	10.32	10.35	13.06	13.95	0.173	0.176	100.35	99.54	1.30	1.35	3.85	3.94	0.337	0.343
Bentonite coated urea	79.85	80.64	10.46	10.46	12.70	12.67	0.164	0.167	99.08	98.32	1.32	1.29	3.59	3.73	0.368	0.346
L.S.D. at 5 %	0.38	0.39	N.S.	N.S.	0.07	0.15	0.005	0.005	0.17	0.52	0.06	0.06	0.13	0.05	0.017	N.S.
B- Phosphorus fertilizer sources																
Super phosphate powder	80.71	81.44	10.21	9.89	13.56	13.73	0.166	0.170	101.82	100.68	1.30	1.26	3.86	3.85	0.337	0.328
Rock phosphate	79.34	80.13	10.41	10.24	12.58	13.06	0.159	0.164	99.24	98.60	1.24	1.27	3.62	3.71	0.343	0.345
Super phosphate granules	80.53	81.38	10.20	10.37	12.98	13.35	0.167	0.174	100.22	99.58	1.27	1.29	3.83	3.84	0.331	0.337
L.S.D. at 5 %	0.18	0.22	N.S.	N.S.	0.11	0.10	0.004	0.004	0.22	0.19	N.S.	N.S.	0.09	0.08	N.S.	N.S.
C- VAM fungi																
Without	79.14	80.06	10.32	10.15	11.27	11.53	0.138	0.141	99.17	98.53	1.104	1.11	3.31	3.27	0.333	0.340
With	81.25	81.91	10.23	10.18	14.81	15.23	0.190	0.197	101.69	100.70	1.441	1.44	4.23	4.33	0.341	0.334
F. test	*	*	N.S.	N.S.	*	*	*	*	*	*	*	*	*	*	N.S.	N.S.

Table 2: Plant height, number of leaves/plant, dry weight /plant, leaf area/plant, total chlorophyll content/plant SPAD unit, neck diameter/plant, bulb diameter and bulbing ratio of garlic at 120 days after planting as affected by the interaction between nitrogen and phosphorus fertilizer sources during 2004/2005 and 2005/2006 seasons.

Characters		Plant height (cm)		Number of leaves/plant		Dry weight /plant (g)		Leaf area/ plant (m ²)		Total Chlorophyll content/plant SPAD unit		Neck diameter/plant (cm)		bulb diameter (cm)		bulbing ratio	
		2004/2005	2005/2006	2004/2005	2005/2006	2004/2005	2005/2006	2004/2005	2005/2006	2004/2005	2005/2006	2004/2005	2005/2006	2004/2005	2005/2006	2004/2005	2005/2006
Calcium nitrate	Super phosphate powder	77.93	78.82	10.00	8.73	13.09	13.43	0.141	0.144	101.82	99.98	1.05	1.11	3.58	3.35	0.293	0.335
	Rock phosphate	77.67	78.99	10.40	10.10	12.49	12.95	0.149	0.159	99.54	98.99	1.04	1.06	3.26	3.27	0.318	0.326
	Super phosphate granules	79.63	80.21	9.73	10.10	11.87	12.47	0.158	0.169	99.90	99.21	1.15	1.15	3.67	3.65	0.315	0.318
Ammonium sulphate	Super phosphate powder	83.86	84.34	10.16	10.20	14.20	14.64	0.176	0.180	103.29	102.70	1.40	1.38	4.30	4.30	0.326	0.322
	Rock phosphate	80.85	81.11	10.33	10.06	13.33	13.26	0.161	0.169	100.64	100.15	1.35	1.34	4.02	4.06	0.339	0.338
	Super phosphate granules	81.52	82.60	10.36	10.40	14.20	13.97	0.171	0.177	101.64	100.82	1.37	1.33	4.09	3.99	0.337	0.336
Sulphur coated urea	Super phosphate powder	80.89	81.82	10.33	10.16	13.57	14.00	0.179	0.185	101.70	100.74	1.41	1.33	4.00	4.05	0.353	0.329
	Rock phosphate	79.83	80.81	10.30	10.30	12.39	13.73	0.162	0.165	99.11	98.25	1.25	1.36	3.70	3.82	0.339	0.356
	Super phosphate granules	80.60	81.38	10.33	10.60	13.23	14.13	0.178	0.178	100.26	99.64	1.24	1.35	3.87	3.95	0.319	0.345
Bentonite coated urea	Super phosphate powder	80.16	80.79	10.36	10.46	13.37	12.85	0.166	0.169	100.49	99.30	1.33	1.22	3.56	3.72	0.376	0.327
	Rock phosphate	79.02	79.82	10.63	10.50	12.12	12.31	0.164	0.162	97.66	97.02	1.32	1.33	3.51	3.70	0.375	0.361
	Super phosphate granules	80.37	81.32	10.40	10.41	12.63	12.85	0.162	0.170	99.09	98.64	1.31	1.32	3.70	3.77	0.354	0.349
L.S.D. at 5 %		0.37	0.43	N.S.	N.S.	0.22	0.20	0.007	0.007	N.S.	0.38	N.S.	N.S.	0.17	0.15	N.S.	N.S.

4-Interaction effect between N-sources and P-sources :

The effect of interaction of N – sources with P – sources on plant vegetative growth parameters of garlic plants at 120 DAP, data in Table (2) show that there were significant differences among the studied treatments. Plant height, dry weight, bulb diameter were significantly increased by application of ammonium sulphate with powder super phosphate, followed by application of ammonium sulphate with granules super phosphate, in both seasons. Leaf area/plant significantly increased by application of sulphur coated urea with powder super phosphate, in both seasons. Total chlorophyll content/plant was significantly increased by application of ammonium sulphate with powder super phosphate, in the second season only, while, number of leaves/plant, neck diameter and bulbing ratio were not significantly affected by the interaction, in both seasons and total chlorophyll significantly not affected by the interaction, in the first season. The interaction had a positive effect on the most studied parameters. These results are in agreement with those of Maksoud *et al.* (1983 & 1984), Oh *et al.* (1991), Ashok *et al.* (1996), El – Gamal and Salim (2005) and Sardi and Timar (2005) on garlic, Rizk (1997) on onion and Awad (2005) on potato.

5-Interaction effect between N-sources and VAM fungi :

Data in Table (3) show that there were significant increases at most of measured parameters, i.e., plant height and number of leaves/plant in the first season only were significantly increased by application of sulphur coated urea with inoculation of VAM fungi, dry weight and bulb diameter were significantly increased by application of ammonium sulphate with VAM fungi, in both seasons and neck diameter and bulbing ratio, in the second season were significantly reduced by using calcium nitrate without VAM fungi. While, plant height in the second season, number of leaves/plant in the second season, leaf area/plant and total chlorophyll content/plant, in both seasons, neck diameter and bulbing ratio, in the first season were not significantly affected by the interaction. These increases might be ascribed to the beneficial effect of VAM fungi and N – element on absorption and efficiency of plant nutrition. These results are in agreement with that of Baath and Spokes (1989) who suggested that supplying chives plants by combined N and P fertilization in the presence of VAM fungi stimulated plant growth.

6-Interaction effect between P-sources and VAM fungi :

Data in Table (3) show that there were significant increases at most of measured parameters, i.e., dry weight, in both seasons by application of powder super phosphate with VAM fungi, leaf area/plant, in both seasons by application of granules super phosphate with VAM fungi and neck diameter and bulb diameter, in both seasons were significantly increased by application of granules super phosphate with VAM fungi. While, plant height, number of leaves/plant and bulbing ratio, in both seasons and total chlorophyll content/plant in the second season were not significantly affected by the interaction. These results are in agreement with those of Sari *et al.* (2002) on garlic, Awad (2002) on potato and El – Morsy *et al.* (2002) on sweet potato.

Table 3: Plant height, number of leaves/plant, dry weight /plant, leaf area/plant, total chlorophyll content/plant SPAD unit, neck diameter/plant, bulb diameter and bulbing ratio of garlic at 120 days after planting as affected by the interaction between nitrogen fertilizer sources and VAM fungi as well as phosphorus fertilizer sources and VAM fungi during 2004/2005 and 2005/2006 seasons.

Characters	Treatments	Plant height (cm)		Number of leaves/plant		Dry weight /plant (g)		Leaf area/ plant (m ²)		Total Chlorophyll content/plant SPAD unit		Neck diameter/plant (cm)		bulb diameter (cm)		bulbing ratio	
		2004/2005	2005/2006	2004/2005	2005/2006	2004/2005	2005/2006	2004/2005	2005/2006	2004/2005	2005/2006	2004/2005	2005/2006	2004/2005	2005/2006	2004/2005	2005/2006
<i>Interaction between nitrogen fertilizer sources and VAM fungi</i>																	
Calcium nitrate	Without	77.29	78.51	10.20	10.13	10.73	11.28	0.124	0.128	99.27	98.51	0.96	1.01	3.14	2.99	0.308	0.338
	With	79.54	80.17	9.88	9.15	14.23	14.81	0.174	0.186	101.57	100.27	1.20	1.21	3.87	3.86	0.310	0.314
Ammonium sulphate	Without	80.80	81.60	10.08	10.00	11.88	11.99	0.141	0.147	100.51	100.00	1.19	1.18	3.55	3.43	0.337	0.348
	With	83.34	83.77	10.48	10.44	15.94	15.92	0.198	0.204	103.20	102.44	1.56	1.52	4.71	4.80	0.331	0.317
Sulphur coated urea	Without	79.38	80.30	10.42	10.24	11.28	11.88	0.147	0.150	99.05	98.47	1.11	1.16	3.39	3.43	0.330	0.338
	With	81.50	82.25	10.22	10.46	14.85	16.03	0.199	0.202	101.66	100.62	1.48	1.54	4.32	4.44	0.344	0.348
Bentonite coated urea	Without	79.08	79.84	10.57	10.24	11.17	10.99	0.140	0.139	97.83	97.15	1.13	1.08	3.17	3.24	0.359	0.335
	With	80.82	81.45	10.35	10.87	14.24	14.35	0.188	0.195	100.33	99.49	1.51	1.50	4.01	4.21	0.377	0.356
L.S.D. at 5 %		0.36	N.S.	0.36	N.S.	0.13	0.14	N.S.	N.S.	N.S.	N.S.	N.S.	0.05	0.13	0.14	N.S.	0.020
<i>Interaction between phosphorus fertilizer sources and VAM fungi</i>																	
Super phosphate powder	Without	79.70	80.61	10.21	10.10	11.81	11.71	0.142	0.144	100.64	99.55	1.14	1.09	3.48	3.41	0.328	0.323
	With	81.72	82.28	10.21	9.68	15.30	15.75	0.189	0.195	103.01	101.79	1.46	1.43	4.24	4.30	0.346	0.333
Rock phosphate	Without	78.31	79.17	10.56	10.18	10.34	10.99	0.129	0.131	97.76	97.44	1.03	1.08	3.05	3.07	0.338	0.352
	With	80.37	81.09	10.26	10.31	14.83	15.13	0.188	0.196	100.71	99.77	1.45	1.47	4.20	4.36	0.347	0.338
Super phosphate granules	Without	79.40	80.40	10.18	10.20	11.65	11.90	0.143	0.148	99.10	98.59	1.13	1.15	3.41	3.35	0.334	0.344
	With	81.65	82.36	10.23	10.55	14.31	14.81	0.192	0.200	101.34	100.56	1.40	1.43	4.26	4.33	0.329	0.330
L.S.D. at 5 %		N.S.	N.S.	N.S.	N.S.	0.11	0.12	0.006	0.004	0.29	N.S.	0.08	0.05	0.11	0.12	N.S.	N.S.

7-Interaction effect between N-sources, P-sources and VAM fungi

As for the interaction effect of the three studied factors, data in Table (4) reveal that dry weight significantly influenced by the interaction, in both seasons. However, plant-height and number of leaves / plant were not significantly affected by the interaction (N – sources x P – sources x VAM fungi) in both seasons . It meaning that the parameters of dry weight / plant have a positive response to the interaction (N – sources x P – sources x VAM fungi). Generally, plants received combination of ammonium sulphate and super phosphate powder in the presence of VAM fungi resulted in the highest vegetative growth. The leaf area was significantly increased, in both seasons by application of sulphur coated urea with powder super phosphate with VAM fungi and total chlorophyll content/plant, neck diameter, bulb diameter and bulbing ratio significantly increased in the first season only while, these parameters were not significantly affected by the interaction, in the second season. The obtained results are in harmony with that obtained by Baath and Spokes (1989) on chives and Hammad and Abdel – Ati (1998) on potato.

Chemical constituents

1-Effect of N-sources :

Data presented in Table (5) show that there were significantly increases in N %, P % and K %, in both seasons. The application of sulphur coated urea as a source of nitrogen significantly increased the concentration percentage of N %, P % and K %, and this treatment exerted the highest values, in both seasons, followed by bentonite coated urea, followed by ammonium sulphate and calcium nitrate came in the last. This treatment significantly reduced the concentration of NO₃ and NO₂ in the garlic bulb, in both seasons. The lowest values of NO₃ and NO₂ obtained by applying bentonite coated urea, in both seasons. These increment may be due to that sulphur coated urea has big value of sulphur, this converts by bacteria and other organisms to sulphuric acid in the soil, in turn, decrease the soil pH to the optimum level for the majority of micronutrients to be absorbed by plant root. These results were similar to that reported by Govind *et al.* (1976) who showed that the effect of sulphur coated urea on nitrate accumulation was minimal on cabbage and tomato. Similar results were reported by Abdel - Fattah *et al.* (2001) who found that Ca(NO₃)₂ fertilizer application exerted significantly increases on NO₃ and NO₂ concentration. These results were similar with those of El-Saei and Tartoura(2006) on cabbage, and Hussien *et al.* (2007) on Jerusalem artichoke.

2-Effect of P-sources :

Data in Table (5) clarified that the concentration percentage of macronutrients, i.e., N % and K % were significantly increased, in both seasons, while the concentration percentage of P % was not significantly affected by application of any of P – sources. The highest values obtained by application super phosphate powder or granules on N % (no significant between the two values) and super phosphate granules on K %, compared with the other sources, in both seasons. The concentration of NO₃ and NO₂ were significantly reduced by application of powder super phosphate and the

lowest values of NO_3 and NO_2 by applying rock phosphate. Several investigators reported that P is the main constituent of many organic compounds in plants. Moreover, it plays an important role in certain essential steps, such as photophosphorylation and release of energy during cellular metabolism (Russel 1950, Mengel and Kirkby 1987 and Marschner 1995). These obtained data are in harmony with those of Sanderson *et al.* (2002) on potato and Shabana (2004) on tomato.

3-Effect of VAM fungi :

As for the effect of VAM fungi inoculation on chemical constituents of garlic bulb, data in Table (5) show that macronutrients uptake percentage, i.e., N %, P % and K % were significantly increased by inoculation with VAM fungi and significantly reduced the concentration of NO_3 and NO_2 of garlic bulb, as compared with the uninoculated ones. The VAM inoculation being significantly the most effective, in both seasons of the study, the increased concentration percentage with VAM inoculation better than without by 11.29, 11.82, 47.80, 40.37 and 9.21, 9.21 % on N, P and K, in both seasons, respectively. The superiority effect of VAM fungi may be due to the small diameter and large size of their hyphae and the accumulation of polyphosphate in their vacuoles. Polyphosphate was involved in transport of P in hyphae to infect root where it is hydrolyzed in the arbuscules and most likely transported as inorganic phosphate across the plasma membrane of the host root cell. Moreover, K ions and basic amino acids may be taken up internally by absorption to polyphosphate granules (Smith and Person, 1988). These results were similar with those of Awad (2002) on potato and El-Morsy *et al.* (2002) on sweet potato, found that N, P and K concentrations markedly increased in foliage and tubers of plants inoculated with VAM fungi.

4-Interaction effect between N-sources and P-sources :

As for the effect of interaction between N-sources with P-sources on the chemical constituents of garlic bulb, data in Table (6) show that the interaction had a significant effect on the concentration of N %, in the first season and K %, in both seasons. However, P % concentration of garlic bulbs did not reflect any significant increases, in both seasons and N % concentration, in the second season of the study. The highest values obtained by application of sulphur coated urea with super phosphate powder on N %, in the first season and super phosphate granules on K %, in both seasons. The lowest values of nitrate and nitrite accumulation of garlic bulb cloves obtained by application of bentonite coated urea with super phosphate powder, in both seasons. These results are in harmony with those of Shobahalan and Arumugam (1991) and El-Gamal and Selim (2005) on garlic and Fayed (1998) on onion.

Table 4: Plant height, number of leaves/plant, dry weight /plant, leaf area/plant, total chlorophyll content/plant SPAD unit, neck diameter/plant, bulb diameter and bulbing ratio of garlic at 120 days after planting as affected by the interaction among nitrogen and phosphorus fertilizer sources as well as VAM fungi during 2004/2005 and 2005/2006 seasons.

Characters			Plant height (cm)		Number of leaves/plant		Dry weight /plant (g)		Leaf area/ plant (m ²)		Total Chlorophyll content/plant SPAD unit		Neck diameter/plant (cm)		bulb diameter (cm)		bulbing ratio	
			2004/ 2005	2005/ 2006	2004/ 2005	2005/ 2006	2004/ 2005	2005/ 2006	2004/ 2005	2005/ 2006	2004/ 2005	2005/ 2006	2004/ 2005	2005/ 2006	2004/ 2005	2005/ 2006	2004/ 2005	2005/ 2006
Calcium nitrate	Super phosphate powder	Without	76.82	78.00	10.06	10.33	11.14	11.41	0.127	0.129	100.89	98.45	0.93	1.01	3.36	2.98	0.277	0.342
		With	79.04	79.86	9.93	7.13	15.04	15.45	0.186	0.160	102.75	100.52	1.18	1.22	3.80	3.72	0.310	0.328
	Rock phosphate	Without	76.63	78.18	10.60	10.26	10.25	11.09	0.114	0.122	98.19	97.90	0.84	0.93	2.74	2.77	0.308	0.337
		With	78.72	79.80	10.20	9.93	14.74	14.81	0.183	0.196	100.89	100.07	1.24	1.19	3.79	3.77	0.327	0.315
	Super phosphate granules	Without	78.41	79.35	9.93	9.90	10.82	11.34	0.131	0.135	98.73	98.19	1.12	1.08	3.32	3.22	0.336	0.337
		With	80.86	81.06	9.53	10.40	12.92	13.69	0.185	0.203	101.07	100.23	1.17	1.22	4.03	4.09	0.292	0.300
Ammonium sulphate	Super phosphate powder	Without	82.60	83.63	10.06	10.00	11.95	12.43	0.149	0.154	101.70	101.18	1.20	1.18	3.86	3.71	0.311	0.320
		With	84.92	85.15	10.26	10.40	16.46	16.85	0.203	0.207	104.88	104.22	1.61	1.58	4.74	4.66	0.342	0.324
	Rock phosphate	Without	79.67	80.01	10.33	9.93	10.61	11.29	0.126	0.132	99.27	98.87	1.10	1.15	3.19	3.14	0.345	0.368
		With	82.12	82.22	10.33	10.20	16.15	15.23	0.193	0.205	102.02	101.42	1.60	1.53	4.64	4.66	0.332	0.306
	Super phosphate granules	Without	80.05	81.27	9.86	10.08	13.20	12.28	0.147	0.155	100.67	99.98	1.26	1.22	3.62	3.44	0.364	0.355
		With	82.99	83.94	10.66	10.73	15.20	15.68	0.196	0.199	102.71	101.88	1.46	1.44	4.56	4.54	0.320	0.316
Sulphur coated urea	Super phosphate powder	Without	79.71	80.75	10.33	9.93	11.97	11.90	0.180	0.155	100.41	99.61	1.22	1.15	3.61	3.61	0.336	0.319
		With	82.06	82.90	10.33	10.40	15.18	16.20	0.209	0.215	102.99	101.87	1.61	1.51	4.39	4.46	0.368	0.336
	Rock phosphate	Without	78.94	79.67	10.46	10.13	10.55	11.16	0.136	0.139	97.45	97.14	1.10	1.11	3.16	3.21	0.347	0.348
		With	80.72	81.54	10.13	10.46	14.24	16.32	0.186	0.191	100.78	99.37	1.40	1.61	4.24	4.43	0.330	0.365
	Super phosphate granules	Without	78.48	80.47	10.46	10.66	11.33	12.68	0.153	0.156	99.30	98.66	1.03	1.21	3.39	3.49	0.304	0.348
		With	81.71	82.30	10.20	10.63	15.13	15.68	0.203	0.201	101.23	100.82	1.44	1.50	4.34	4.41	0.334	0.341
Bentonite coated urea	Super phosphate powder	Without	79.47	80.16	10.40	10.13	12.19	11.21	0.143	0.139	99.56	98.02	1.20	1.04	3.11	3.33	0.366	0.312
		With	80.88	81.42	10.33	10.80	14.65	14.50	0.190	0.199	101.42	100.56	1.46	1.41	4.01	4.11	0.364	0.343
	Rock phosphate	Without	78.11	78.81	10.66	10.33	10.05	10.45	0.137	0.132	98.13	95.65	1.09	1.12	3.10	3.15	0.352	0.358
		With	79.94	80.83	10.40	10.66	14.19	14.17	0.191	0.192	99.19	98.20	1.56	1.55	3.92	4.25	0.396	0.364
	Super phosphate granules	Without	79.67	80.54	10.46	10.26	11.27	11.32	0.141	0.145	97.81	97.58	1.11	1.09	3.30	3.26	0.339	0.336
		With	81.06	82.11	10.33	10.56	13.99	14.38	0.183	0.195	100.37	99.70	1.52	1.55	4.11	4.26	0.370	0.362
L.S.D. at 5 %			N.S.	N.S.	N.S.	N.S.	0.22	0.24	0.011	0.006	0.58	N.S.	0.15	N.S.	0.22	N.S.	0.045	N.S.

Table 5: N %, P %, K %, NO₃ and NO₂ accumulation (ppm) of garlic as affected by some nitrogen and phosphorus fertilizer sources as well as VAM fungi during 2004/2005 and 2005/2006 seasons.

Characters Treatments	N %		P %		K %		NO ₃ (ppm)		NO ₂ (ppm)	
	2004/ 2005	2005/ 2006	2004/ 2005	2005/ 2006	2004/ 2005	2005/ 2006	2004/ 2005	2005/ 2006	2004/ 2005	2005/ 2006
A- Nitrogen fertilizer sources										
Calcium nitrate	2.83	2.85	0.501	0.492	1.42	1.42	278.3	280.2	3.61	3.64
Ammonium sulphate	3.25	3.31	0.636	0.616	1.51	1.52	240.6	242.7	3.28	3.31
Sulphur coated urea	3.62	3.64	0.648	0.644	1.83	1.81	190.9	193.6	2.79	2.80
Bentonite coated urea	3.39	3.45	0.589	0.571	1.60	1.60	188.1	188.5	2.75	2.76
L.S.D. at 5 %	0.09	0.06	0.024	0.027	0.01	0.01	1.5	1.8	0.01	0.01
B- Phosphorus fertilizer sources										
Super phosphate powder	3.31	3.34	0.595	0.583	1.58	1.59	226.4	228.2	3.13	3.16
Rock phosphate	3.22	3.25	0.592	0.574	1.55	1.55	218.2	219.2	3.01	3.04
Super phosphate granules	3.30	3.35	0.594	0.586	1.64	1.62	228.8	228.2	3.18	3.18
L.S.D. at 5 %	0.04	0.04	N.S.	N.S.	0.01	0.01	1.5	1.8	0.01	0.01
C- VAM fungi										
Without	3.10	3.13	0.479	0.483	1.52	1.52	234.6	237.7	3.25	3.26
With	3.45	3.50	0.708	0.678	1.66	1.66	214.4	214.7	2.97	2.99
F. test	*	*	*	*	*	*	*	*	*	*

Table 6: N %, P %, K %, NO₃ and NO₂ accumulation (ppm) of of garlic at 120 days after planting as affected by the interaction between nitrogen and phosphorus fertilizer sources during 2004/2005 and 2005/2006 seasons.

Characters		N %		P %		K %		NO ₃ (ppm)		NO ₂ (ppm)	
		2004/2005	2005/2006	2004/2005	2005/2006	2004/2005	2005/2006	2004/2005	2005/2006	2004/2005	2005/2006
Treatments											
Calcium nitrate	Super phosphate powder	2.89	2.92	0.490	0.487	1.41	1.41	280.3	282.1	3.65	3.67
	Rock phosphate	2.70	2.73	0.482	0.468	1.37	1.38	271.1	275.8	3.46	3.52
	Super phosphate granules	2.90	2.91	0.532	0.522	1.47	1.46	283.6	282.6	3.71	3.73
Ammonium sulphate	Super phosphate powder	3.29	3.32	0.623	0.598	1.50	1.53	241.6	244.3	3.31	3.36
	Rock phosphate	3.21	3.28	0.653	0.622	1.48	1.49	233.0	233.1	3.17	3.22
	Super phosphate granules	3.26	3.33	0.632	0.627	1.55	1.54	247.1	250.6	3.35	3.36
Sulphur coated urea	Super phosphate powder	3.68	3.68	0.672	0.667	1.83	1.81	192.5	195.1	2.80	2.83
	Rock phosphate	3.59	3.56	0.638	0.640	1.78	1.77	187.1	188.5	2.73	2.73
	Super phosphate granules	3.60	3.67	0.635	0.627	1.69	1.85	193.1	197.1	2.84	2.83
Bentonite coated urea	Super phosphate powder	3.37	3.45	0.597	0.580	1.60	1.60	191.3	191.3	2.78	2.79
	Rock phosphate	3.37	3.42	0.593	0.565	1.57	1.57	181.8	179.6	2.68	2.70
	Super phosphate granules	3.44	3.49	0.577	0.568	1.65	1.63	191.3	194.5	2.81	2.79
L.S.D. at 5 %		0.08	N.S.	N.S.	N.S.	0.01	0.01	3.0	3.6	0.01	0.03

6-Interaction effect between P-sources and VAM fungi :

Concerning the interaction effect of P-sources with VAM fungi on mineral composition of garlic bulbs, data in Table (7) indicate that there were significantly increases on P % , in the first season and K % , in both seasons. While, N % concentration was not significantly affected by the interaction, in both seasons and P % concentration, in the second season of the study. The highest values obtained by application of rock phosphate with VAM fungi on P % , in the first season, and by application super phosphate granules and inoculated with VAM fungi in case of K % concentration of garlic bulb, in both seasons. Nitrate accumulation was significantly reduced, in the first season only by rock phosphate with VAM fungi, while, in the second season was not significantly affected by the interaction. the lowest records of NO₂ accumulation, in both seasons obtained by rock phosphate with VAM fungi. The obtained results are nearly with the results obtained by Hammad and Abdel-Ati(1998) on potato and Mengistu and Singh (1999) on onion.

Table 7: N %, P %, K %, NO₃ and NO₂ accumulation (ppm) of of garlic at 120 days after planting as affected by the interaction between nitrogen fertilizer sources and VAM fungi as well as phosphorus fertilizer sources and VAM fungi during 2004/2005 and 2005/2006 seasons.

Characters		N %		P %		K %		NO ₃ (ppm)		NO ₂ (ppm)	
		2004/ 2005	2005/ 2006	2004/ 2005	2005/ 2006	2004/ 2005	2005/ 2006	2004/ 2005	2005/ 2006	2004/ 2005	2005/ 2006
<i>Interaction between nitrogen fertilizer sources and VAM fungi</i>											
Calcium nitrate	Without	2.71	2.73	0.424	0.417	1.36	1.36	288.2	291.7	3.78	3.81
	With	2.95	2.97	0.578	0.568	1.47	1.48	268.5	268.6	3.43	3.47
Ammonium sulphate	Without	3.04	3.09	0.529	0.536	1.44	1.44	252.2	256.2	3.43	3.46
	With	3.46	3.53	0.743	0.696	1.58	1.60	229.0	229.2	3.12	3.16
Sulphur coated urea	Without	3.37	3.38	0.504	0.524	1.75	1.73	199.6	204.1	2.90	2.91
	With	3.87	3.90	0.792	0.764	1.91	1.89	182.2	183.1	2.68	2.68
Bentonite coated urea	Without	3.26	3.30	0.459	0.457	1.55	1.54	198.3	198.8	2.87	2.86
	With	3.52	3.61	0.719	0.686	1.66	1.67	178.0	178.1	2.64	2.65
L.S.D. at 5 %		0.07	0.09	0.039	0.032	0.01	0.01	1.3	2.2	0.02	0.02
<i>Interaction between phosphorus fertilizer sources and VAM fungi</i>											
Super phosphate powder	Without	3.15	3.17	0.500	0.496	1.52	1.51	236.9	240.1	3.29	3.31
	With	3.47	3.52	0.691	0.670	1.64	1.66	216.0	216.3	2.98	3.02
Rock phosphate	Without	3.02	3.04	0.449	0.466	1.49	1.48	229.3	231.0	3.17	3.19
	With	3.42	3.46	0.734	0.682	1.62	1.62	207.2	207.5	2.85	2.89
Super phosphate granules	Without	3.13	3.17	0.488	0.488	1.56	1.56	237.5	242.0	3.28	3.29
	With	3.47	3.53	0.699	0.683	1.71	1.69	220.0	220.5	3.07	3.07
L.S.D. at 5 %		N.S.	N.S.	0.034	N.S.	0.01	0.01	1.1	N.S.	0.01	0.02

7-Interaction effect between N-sources, P-sources and VAM fungi :

Data presented in Table (8) show the interaction effect among N-sources, P-sources and VAM fungi on macronutrients uptake, i.e., N %, P %, K %, NO₃ and NO₂ concentration of garlic bulbs. Obtained results revealed significant increases in K % content, in both seasons. While N % and P % were not significantly affected by the above mentioned interaction, in both seasons. The best records of K % content of garlic bulbs obtained by application of sulphur coated urea with super phosphate granules in the presence of VAM fungi. The lowest values of NO₃ and NO₂ concentration obtained by application of bentonite coated urea with rock phosphate with VAM fungi, in both seasons followed by results obtained by applying of sulphur coated urea with rock phosphate with VAM fungi, in both seasons. Similar results were obtained by Hammad and Abdel-Ati (1998) on potato and Mingistu and Singh (1999) on onion and Awad(2002) on potato.

Table 8: N %, P %, K %, NO₃ and NO₂ accumulation (ppm) of of garlic at 120 days after planting as affected by the interaction among nitrogen and phosphorus fertilizer sources as well as VAM fungi during 2004/2005 and 2005/2006 seasons.

Characters			N %		P %		K %		NO ₃ (ppm)		NO ₂ (ppm)	
			2004/ 2005	2005/ 2006	2004/ 2005	2005/ 2006	2004/ 2005	2005/ 2006	2004/ 2005	2005/ 2006	2004/ 2005	2005/ 2006
Calcium nitrate	Super phosphate powder	Without	2.83	2.85	0.427	0.417	1.37	1.36	290.6	297.0	3.85	3.86
		With	2.96	2.99	0.553	0.557	1.46	1.47	270.0	267.3	3.46	3.49
	Rock phosphate	Without	2.52	2.54	0.383	0.380	1.32	1.33	281.3	286.0	3.71	3.75
		With	2.89	2.92	0.580	0.557	1.43	1.43	261.0	265.6	3.22	3.29
	Super phosphate granules	Without	2.79	2.81	0.463	0.453	1.40	1.40	292.6	292.3	3.80	3.83
		With	3.02	3.01	0.600	0.590	1.54	1.53	274.6	273.0	3.62	3.64
Ammonium sulphate	Super phosphate powder	Without	3.05	3.11	0.557	0.547	1.44	1.44	253.3	257.3	3.47	3.51
		With	3.53	3.53	0.690	0.650	1.57	1.62	230.0	231.3	3.16	3.21
	Rock phosphate	Without	3.03	3.07	0.507	0.517	1.41	1.41	245.0	246.3	3.32	3.36
		With	3.39	3.50	0.800	0.727	1.56	1.57	221.0	220.0	3.01	3.07
	Super phosphate granules	Without	3.06	3.10	0.523	0.543	1.48	1.48	258.3	265.0	3.50	3.51
		With	3.47	3.57	0.740	0.710	1.62	1.61	236.0	236.3	3.20	3.21
Sulphur coated urea	Super phosphate powder	Without	3.44	3.44	0.543	0.543	1.75	1.72	201.0	204.0	2.93	2.96
		With	3.92	3.93	0.800	0.790	1.91	1.90	184.0	186.3	2.67	2.71
	Rock phosphate	Without	3.31	3.29	0.467	0.523	1.71	1.68	197.6	200.0	2.85	2.85
		With	3.86	3.83	0.810	0.757	1.86	1.86	176.6	177.0	2.62	2.61
	Super phosphate granules	Without	3.36	3.41	0.503	0.507	1.81	1.79	200.3	208.3	2.94	2.93
		With	3.83	3.93	0.767	0.747	1.97	1.92	186.0	186.0	2.75	2.74
Bentonite coated urea	Super phosphate powder	Without	3.27	3.27	0.473	0.477	1.55	1.54	202.6	202.3	2.91	2.93
		With	3.47	3.62	0.720	0.683	1.85	1.67	180.0	180.3	2.65	2.66
	Rock phosphate	Without	3.21	3.25	0.440	0.443	1.52	1.51	193.3	192.0	2.81	2.79
		With	3.53	3.59	0.747	0.687	1.63	1.64	170.3	167.3	2.56	2.61
	Super phosphate granules	Without	3.31	3.38	0.463	0.450	1.58	1.57	199.0	202.3	2.90	2.88
		With	3.56	3.60	0.690	0.687	1.71	1.70	183.6	186.6	2.72	2.70
L.S.D. at 5 %			N.S.	N.S.	N.S.	N.S.	0.02	0.02	2.2	3.9	0.03	0.03

REFERENCES

- Abbadly, K. A. ; A. B. Barakat ; M. I. El- Mallah and A. A. Khatab (1997) .The Effect of some slow -release fertilizers on onion yield and successive Sweet corn growth . 1- The effect on onion bulb yield ; bulb quality and chemical constituents . Egypt. J. Appl. Sci., 12 (3) : 245 – 261.
- AbdEl - Fattah , A. E. ; A. H. A. El-Morsy and A. A. Salim (2001). Application effect of some nitrogen sources and Mepiquat Chloride (Pix) on plant growth , yield and accumulation of nitrate and nitrite in tuber roots of sweet potato . The Fifth Arabian Horticulture Conference , Ismailia , Egypt , March 24 – 28 , Vol . II .
- Abd El – Hamed , A. M. ; A. Z. Osman ; S. A. Ismail and F. M. Ahmed (1996b). Effect of nitrogen sources with different levels on garlic plants (*Allium sativum* L.). J. Agric. Sci. Mansoura Univ., 21 (1) :423-429.
- Ashok , K. ; G. N. Singh and A. Kumar (1998). Effect of NPK on growth , yield And quality of garlic (*Allium sativum* L.). Recent Horticulture, 3 (1) : 118 – 121
- Awad , E. M. (2002). Growth , yield and quality of potato crop as affected by the Inoculation with Vesicular Arbuscular Mycorrhizal (VAM) Fungi under different levels of phosphorus . J. Agric. Sci. Mansoura Univ., 27 (8) : 5593 – 5605 .
- Awad ,E. M.(2005).The influence of organic and mineral fertilization on growth, yield and quality of potato crop . J. Agric. Sci. Mansoura Univ., 30 (12) : 7965 – 7975 .
- Baath ,F. and J. Spokes (1989). The effect of added nitrogen and phosphorus on mycorrhizal growth response and interaction in *Allium schoenoprasum*. Can. J. Bot., 67 : 3227 – 3232 .
- Brown , J. D. ; O. Lilleland (1946). Rapid determination of potassium and sodium in plant material and soil extracts by flame photometry. Proc. Amer. Soci. Hort. Sci., 48 : 341 – 346.
- Chapman, H. D. and P. F. Partt (1961). Methods of analysis for soil, plant and water. Department of soil and plant nutrition, Univ. of California, Citrus Exp. Sta. Riverside , California , USA .
- El-Beheidi, M. A.; M. A. I. Khalil ; M. H. El-Sawah and A. A.Gad (1985).Response of some garlic cultivars to different levels of nitrogen fertilizer. Zagazig J. Agric. RES. , 12 (2) : 65 – 84 .
- El-Gamal , S. and A. H. Selim (2005).Response of garlic plants grown in sandy soils to organic and inorganic fertilizers. J. Agric. Sci. Mansoura Univ., 30 (11) 6687 – 6700 .
- El-Mallah, M. I. ; I. A. Ibrahim ; A. A. Darwish and K. A. Abbady (1998). Efficiency of slow – release and fast release N – fertilizers on spinach (growth and maturity quality) and the residual effect on corn. Egypt. J. Appl. Sci., 31 (4) : 292 – 306 .
- El-Morsy, A. H. A.; A. E. A. Abdel-Fattah and Z. S. A. El-Shal (2002).Effect of phosphate fertilizer and VA-Mycorrhizal inoculation on growth, tuber yield and quality of sweet potato. Proc.Minia Frist Conf. for Agric. & Environ. Sci. Minia Egypt. March 25 – 28 pp : 1815 – 1827 .

- El-Saei, M. A. and E. A. A. Tartoura (2006). How far bentonite coated urea and N- bio fertilizer to suppress NO₃ accumulation in cabbage ?. J. Agric. Sci. Mansoura Univ., 31 (7) : 4813 – 4823 .
- El-Shaikh, K. A. A. (2005). Growth and yield of onion as affected by biofertilization, application of nitrogen and phosphorus fertilizers under South Vally conditions. Assiut, J. of Agric. Sci., 36 (1) :37-50.
- Fayed, R. M. (1998). Effect of chicken manure and sulphur fertilizer mixture with NPK fertilizers on growth, yield and NPK contents of garlic and pea plants. J. Agric. Sci. Mansoura Univ., 23 (5) :2305 – 2313 .
- Gad El-Hak, S. H. and Y.T. Abd El-Mageed (2000). Effect of nitrogen source on growth, yield, nitrate content and storage ability of two garlic cultivars. Minia J. of Agric. Res. & Devel., Vol. 20 No. (1) : 115 – 139 .
- Gomez, K. A.; and R. Gomez (1984). " Statistical procedure for Agric. Res. " 2nd Ed. John Wiley and Sons. Inc. New Yourk, 680 pages.
- Govind, C. S. ; A. J. Patel and D. A. Mays (1976). Effect of sulfur – coated urea on yield, N uptake, and nitrate content in Turnip Greens, Cabbage and Tomato. J. Amer.Soc. Hort. Sci., 101 (2) : 142 – 145 .
- Hammad, A. M. M. and Y. Y. Abdel-Ati (1998). Reducing of nitrite and nitrite contents of potato tubers via Bio fertilization with Azospirillum and VA-Mycorrhizal fungi. J. Agric. Sci. Mansoura Univ., 23 (6) : 2597 – 2610 .
- Hesse, P. R. (1971). A text book of soil chemical analysis. John. Murray (Publish), London, Great Britain.
- Hussein, M. A. ; W. H. Mohamed ; M. G. Nasseem and O. A. A. Hediya (2007). Effect of nitrogen fertilization and sulphur application of Jerusalem artichoke (*Helianthus tuberosus*) grown in lacustrine soil. Arab Conf. of Soil and Water Management For Sustainable Agric. Development 10 – 11 April, Fac. Agric. Mansoura Univ., Egypt : 273 – 288.
- John, M. K.(1970). Colorimetric determination of phosphorus in soil and plant material with ascorbic acid. Soil Sci., 109 : 214 – 220 .
- Koch, M. ; Z. Tanami ; H. Bodani ; S. Wininger and Y.Kapulnik (1997). Field application of Vesicular-Arbuscular Mycorrhizal fungi improved garlic yield in disinfected soil. Mycorrhiza (7) : 47 – 50 .
- Koller, H.R. (1972): Leaf area – leaf weight relationship in the soybean canopy. Crop Sci., 12: 180-183.
- Kumar, P. and S. K. Sharma (2004). Effect of phosphorus sources on cabbage – tomato cropping sequence at Solan. Haryana J. of Horticultural Sciences, 33 (3 / 4) : 272 – 273.
- Lubraco, G. ; A. Schubert and A. Previati (2000). Micropropagation and mycorrhization of *Allium sativum* L. Acta Horticulture, 530 : 339 – 343
- Maksoud, M. A. ; Foda, Sherifa ; A. El-Gizawi and E. M. Taha (1983). Response of garlic plants to fertilization treatments. Egypt. J. Hort., 10 No. (2) pp. 159 – 165 .
- MAksoud, M. A. ; Foda, Sherifa ; E. M. Taha and H. M. Ibrahim (1984). Effect of different fertilizers on quality and yield of garlic. Egypt. J. Hort. , 11 , No.(1) pp. 51 – 58 .

- Mann, L. K. (1952). Anatomy of the garlic bulb and factors affecting bulb development. *Hilgardia*, 21 : 195 – 228 .
- Marschner, H. (1995). Mineral nutrition of higher plants. 2nd Ed. Acad. Press. Limited, Text Book, pp. 252 .
- Mengl, K. and E. A. Kirkby (1987). Principles plant nutrition, publisher. International potash institute publisher long Druck AG. Liebefeld / Bern, 203 – 210 .
- Mengistu, F. and N. Singh (1999). Effects of bio fertilizers on growth, yield and nutrient uptake of onion (*Allium cepa* L.). *Vegetable – Science* , 26 (2) : 193 – 195 .
- Nagaraju, R. ; K. Haripriya and K. Purushotham (2001). Influence of VAM mycorrhizal inoculation under varying phosphorus levels and sources on onion. *South – Indian Hort.*, 49 (special) : 204 – 207 .
- Nagaraju, S.; G. V. Rajalingam ; R. S. Kumar and K. Haripriya (1999). Screening of efficient VAM Fungi for aggregatum onion . *South Indian Hort.*, 47 (1/6) : 255 – 258 .
- Navale, A. M. ; P. V. Wanl and A. S. Patel (2004). Effect of VAM and Azospirillum inoculation to onion (*Allium cepa* L.) cv. B – 780 with respect to N , P and micronutrient uptake. *Orissa J. of Hort.*, 32 (1) : 82 – 88 .
- Oh, D. H. ; N. Y. Hwang ; J. S. Na ; B. J. Choi and J. D. Soh (1991). The responses and optimum rates of NPK fertilizer application to garlic. *Research Reports of the Rural Development Administration Soil and Fertilizer*, 33 (1) : 46 – 51 .
- Piekielek, W. P. and R. H. Fox (1992). Use of a chlorophyll meter to predict sidedress nitrogen requirements for plants. *Agron. J.* , 84 (1/2) : 59 – 65 .
- Rizk, F. A. (1997). Productivity of onion plant (*Allium cepa* L.) as affected by method of planting and NPK application. *Egypt. J. Hort.* , 24, No. (2) pp. 219 – 238 .
- Russell, E. J. (1950). Soil conditions and plant growth. Longmans, Green and Co. London; pp. 30 – 36 and 38 – 39 .
- Sari, N. ; I. Ortas and H. Yetisir (2002). Effects of mycorrhizae inoculation on plant growth, yield and phosphorus uptake in garlic under field conditions. *Communications in Soil Science and Plant Analysis*, 33 (13/14) : 2189 – 2201
- Shabana, A. E. A. (2004). Effect of some biological treatments on tomatoes under saline condition. Ph. D. Thesis, Fac. Agric. Mansoura Univ., Egypt.
- Shobahalan, U. and R. Arumugam (1991). Chemical changes in the quality parameters of garlic (*Allium sativum* L.) during growth and development. *South Indian Hort.*, 39 (2): 93 – 95 ..
- Singh, I. P. (1988): A rapid method for determination of nitrate in soil and plant extracts. *Plant and Soil.*, 110: 137-139.
- Sanderson, J. B. ; T. W. Bruulsema ; R. Coffin ; B. Douglas and J. A. Macleod (2002). Phosphorus sources for potato production. *Better Crops with Plant Food*, 86 (4) : 10 – 12.

- Sang S. Y. ; Hu, C. I. ; Choon, C. B. and Yul, C. W. (2001). Effects of applying slow - release fertilizer on southern type garlic (*Allium sativum* L.) cultivation . Korean J. of Hort. Sci. and Tech., 19 (4) : 471 - 475 .
- Smith, J. H. and U. C. Pearson (1988). Physiological interaction between symbiotics in vesicular arbuscular mycorrhiza plants. Ann. Rev. Plant Physiology Plant Biol., 39 : 221 - 244 ..
- Tartoura, E. A. A. ; M. A. El-Saei ; E. M. El-Said and G. A. Baddour (2003). Suppressing the nitrate pollution in plant and soil using a clean performance in N - fertilization on potato. J. of Envir. Sci. , Vol. 25 : pp. 313 - 331 .
- Wani, P. V. and B. K. Konde (1998). Effects of *Glomus mosseae* inoculation using different P - sources on garlic. J. of Maharashtra Agric. Univ., 23 (1) : 39 - 42
- Yadava, U.L. (1986). A rapid and non-destructive method to determine chlorophyll in intact leaves. HortScience, 21: 1449-1450.

تأثير بعض مصادر الأسمدة الأزوتية والفوسفاتية والتلقيح بالميكوريزا على نمو وانتاجية وتخزين الثوم

١- النمو الخضري و المحتوى الكيميائي.

- كوثر كامل ضوه* ، السيد أحمد طرطورة* ، عبد المنعم محمد عبد الحميد** و
أنور الدسوقي على اسماعيل جوده.
* قسم البساتين - كلية الزراعة - جامعة المنصورة.
** معهد بحوث البساتين - مركز البحوث الزراعية.

لجريت هذه الدراسة على محصول الثوم (سمن - ٤٠) خلال موسمي للدراسة ٢٠٠٤ - ٢٠٠٥ ، بهدف دراسة تأثير بعض مصادر الأسمدة الأزوتية مثل نترات الكالسيوم سلفات الأمونيوم ، اليوريا المغلفة بالكبريت واليوريا المغلفة بالبنتونات وبعض مصادر الأسمدة الفوسفاتية مثل سوبر فوسفات الكالسيوم كل من الناعم والمحب وصخر الفوسفات وكذلك التلقيح بفطر الميكوريزا والتفاضل بين هذه العوامل على النمو الخضري والمكونات الكيميائية لأبصال الثوم. وتشير النتائج المتحصل عليها الى أن طول النبات والوزن الجاف والمساحة الورقية للنبات ومحتوى الكلوروفيل الكلي وقطر الحنق وقطر البصلة زادت معنوياً بإضافة سماد سلفات الأمونيوم كمصدر للنترجين في كلا موسمي للدراسة. كما أدت إضافة سماد السوبر فوسفات الناعم الى حدوث زيادة معنوية في كل من طول النبات والوزن الجاف والمساحة الورقية للنبات ومحتوى الكلوروفيل الكلي وقطر البصلة في كلا موسمي للدراسة، بينما عدد الأوراق وقطر الحنق ونسبة التوصيل لم تتأثر معنوياً بأي من مصادر الفوسفور في موسمي للدراسة. أدى التلقيح بفطر الميكوريزا الى نباتات الثوم الى حدوث زيادة معنوية في معظم الصفات المدروسة ما عدا عدد الأوراق وقطر الحنق ونسبة التوصيل لم تتأثر معنوياً بأي من مصادر الفوسفور في كلا موسمي للدراسة. زادت النسبة المئوية لمحتوى كل من النترجين والفوسفور والبوتاسيوم بإضافة سماد اليوريا المغلفة بالكبريت كما أن هذه المعاملة قللت تركيز النترات والنترات في أبصال الثوم ولكن التركيز الأقل حدث بإضافة اليوريا المغلفة بالبنتونات، كما لم تتأثر النسبة المئوية لمحتوى الفوسفور بأي من مصادر الفوسفور. وكان التلقيح بفطر الميكوريزا أعطى زيادات معنوية في النسبة المئوية لمحتوى كل من النترجين والفوسفور و البوتاسيوم وقللت جدا من تركيز النترات والنترات في أبصال الثوم في كلا موسمي للدراسة.