



RESPONSE OF FABA BEAN (*VICIA FABA* L.) TO COMBINED INOCULATION WITH RHIZOBIA, VA-MYCORRHIZAE AND PHOSPHATE SOLUBILIZING BACTERIA UNDER SANDY SOIL CONDITIONS

[24]

Saleh¹, S.A.; M.A. Nassef²; F.T. Mikhaeel² and A.A. Abou El-Soud²

1- National Gene Bank and Genetic Resources, ARC, Giza, Egypt

2- Soils, Water and Environ. Res. Instit., ARC, Giza, Egypt

Keywords: Faba bean, (*Vicia faba* L.), *Rhizobium leguminosarum* bv. *viciae*, *Bacillus megaterium* var. *phosphaticum*, VA- mycorrhizal fungus, Phosphorus-deficient, sandy soils

megaterium did not record significant influence on the abovementioned parameters as compared to inoculation with *Rhizobium* alone.

ABSTRACT

A pot experiment was conducted at controlled greenhouse of Bio-fertilizers production Unit; Soils, Water and Environment Research Institute, ARC, Giza through the winter growing season of 2006/2007 to evaluate the effects of symbiotic nitrogen fixing bacteria (*Rhizobium leguminosarum* bv. *viciae*), phosphate solubilizing bacteria (*Bacillus megaterium* var. *phosphaticum*) and VAM fungus (*Glomus* sp.) on nodulation, N₂-fixation, growth, N and P uptake of faba bean plants grown in phosphorus-deficient sandy soils. The obtained data resulted in significant increases in VA- Mycorrhizal colonization ratios when soil inoculated with rhizobia and VA-Mycorrhizae as compared with the other treatments. In addition, rock-P fertilized plants exhibited significant increases in VAM colonization ratios when compared with those supplemented with super-P. Concerning of dual inoculation with *Rhizobium* and *Bacillus*, the results indicated higher increases in VA- mycorrhizae colonization ratios as compared with the uninoculated or inoculated treatments with rhizobia alone, being most pronounced in the rock-P amended soil. The obtained results showed clearly that combined inoculation of faba bean with rhizobia and VA-Mycorrhizae in a soil amended with rock-P or super-P, gave the highest significant increases in nodulation, nitrogen fixation, plant biomass, N and P uptake when compared with the plants inoculated with *Rhizobium* alone. On the other hand, combined inoculation with *Rhizobium* and *Bacillus*

INTRODUCTION

Significant areas of cultivated sandy soils in Egypt are deficient of available N, P and K. In modern agriculture, the replenishment of soil nitrogen involves most commonly extensive application of chemical fertilizers (Peoples *et al* 1995), an approach which has several serious drawbacks, high cost and severe negative environmental impacts. These considerations stimulated research on other alternatives, such as biological nitrogen fixation which contributes more than 170 million tons of fixed nitrogen to the biosphere. Eighty percent of the stable biologically fixed nitrogen is a direct result of the rhizobiaceae and some actinomycetes symbioses with leguminous and certain non-leguminous plants, respectively (Earl and Ausubel, 1993).

The dual inoculation with both rhizobia and VA mycorrhizae induced more significant increases in plant dry weight, N and P content of lentil and faba bean as well as seed yield of soybean than inoculation with either VA mycorrhizae or *Rhizobium* alone (Badr El-Din and Moawad, 1988).

The use of plant growth promoting rhizobacteria (PGPR), including phosphate solubilizing bacteria as biofertilizers was suggested as a sustainable solution to improve plant nutrient and production (Vessey, 2003). Direct application of rock phosphate may be agronomically more useful and environmentally more feasible than soluble P (Rajan *et al* 1996).

Phosphorus is one of the major nutrients limiting plant growth. Most soils throughout the world

(Received April 16, 2008)

(Accepted June 7, 2008)

are P deficient (Batjes, 1997) and therefore require P to replenish the P demand by crop plants. To circumvent the P deficiency in soils, P fertilizers could be applied. However, after application, a considerable amount of P is rapidly transformed into less available forms by forming a complex with Al or Fe in acid soils (Norrish and Rosser, 1983) or Ca in calcareous soils (Lindsay *et al* 1989) before plant roots have a chance to absorb it. Further, the use of rock phosphate as a phosphate fertilizer and its solubilization by microbes (Kang *et al* 2002), through the production of organic acids (Maliha *et al* 2004), have become a valid alternative to chemical fertilizers. Rock phosphate is widely distributed throughout the world, both geographically and geologically (Zapata and Roy, 2004). In conjugation with phosphate solubilizing microorganisms (PSM), rock phosphate provides a cheaper source of P fertilizer for crop production, however most of them are not readily available to a plant. In this regard, several studies showed that PSM solubilizes the fixed soil P and applied phosphates, resulting in higher crop yields (Zaidi, 1999 and Gull *et al* 2004). The alternative approach is to use these PSM along with other beneficial rhizospheric microflora to enhance crop productivity. In this context, the simultaneous application of *Rhizobium* and PS microorganisms (Perveen *et al* 2002) and arbuscular mycorrhizal VAM fungi (Zaidi *et al* 2003) has been shown to stimulate plant growth more than inoculation of each microorganism alone in certain situations when the soil is P deficient. On the other hand, VAM fungi encourage the plant roots to absorb rapidly the solubilized P. Accordingly, the increase in plant growth may be due to the release of certain plant growth promoting substances (Kucey *et al* 1989) by the PS organisms or VAM development and mycorrhizal formation (Azcon-Aguilar and Barea, 1985). However, the inoculation effect of the tripartite interaction between N₂-fixing bacteria, PSM and VAM fungus on legume crops are relatively scarce (Zaidi *et al* 2004).

Because the minerals are released slowly and their use as fertilizers often causes insignificant yield increases of current crop (Zapata and Roy, 2004). Phosphate solubilizing bacteria (PSB) have been used to improve rock P value because they convert insoluble rock P into soluble forms available for plant growth (Bojinova *et al* 1997). This conversion is through acidification, chelation and exchange reactions (Gerke, 1992) and produces strong organic acids, which have become indicators for routine isolation and selection procedures

of PSB (Illmer *et al* 1995). *Bacillus megaterium* var. *phosphaticum* is known for its ability to solubilize rock P material (Schilling *et al* 1998).

The aim of the present study was to evaluate the response of faba bean to *Rhizobium* inoculation combined with VA-mycorrhizae and phosphate solubilizing bacteria under sandy soil conditions.

MATERIALS AND METHODS

Microbial cultures

Rhizobium leguminosarum bv. *viciae*, strain ICARDA 441; *Bacillus megaterium* var. *phosphaticum* (local isolate) and VA- mycorrhizal fungus (local isolate of *Glomus* spp.) were supplied by Biofertilizer Production Unit; Soils, Water and Environment Research Institute; ARC; Giza; Egypt.

Preparation of microbial inoculants

Rhizobium strain was grown on yeast extract mannitol broth medium (Vincent, 1970) at 28 °C under shaking at 250 rpm for 3 days until early log phase (5×10^9 cfu/ml culture). Vermiculite mixed with 10% peat was packed in polyethylene bags (180 g carrier per bag), then sealed and sterilized by gamma irradiation (5.0×10^6 rads). *Rhizobium* culture was injected into the carrier to satisfy 60% of the maximal water holding capacity.

Bacillus megaterium was grown on nutrient broth medium under shaking at 250 rpm for 48 hrs at 28 °C. Viable bacterial cells (4×10^8 cfu/ml culture) was injected in the abovementioned carrier materials used in *Rhizobium* inoculants.

Vascular Arbuscular mycorrhizal (VAM) fungi spores were propagated for 16 weeks on *Sorghum bicolor* in sterilized pot cultures prior to the start of the experiment. Soil from 16 week-old pot culture was used as a crude inoculum consisted of soil, mycorrhizal spores and infected roots.

Pot experiment

Pot experiment was conducted at the controlled greenhouse of Biological Nitrogen Fixation Unit; Soils, Water and Environment Research Institute, Agricultural Research Center, Giza through the winter growing season of 2006/2007 to study the Effect of *Rhizobium* inoculation combined with VM-mycorrhizae and phosphate dissolving bacteria on growth, nodulation, nitrogen fixation and phosphorus uptake in faba bean plants under sandy soils.

The soil used in pot experiment was collected from Ismailia Research Station. The main physico-chemical characteristics of the soil were determined according to Jackson (1973) and Page *et al* (1982), (Table 1).

Table 1. Some physical and chemical properties of the used soil

Physical properties		Chemical properties	
Sand	90.20%	pH	7.30
Silt	5.68%	E.C. (dS m ⁻¹)	0.35
Clay	4.12%	<u>Soluble cations (meq l⁻¹)</u>	
Textural class	Sandy	Ca ⁺⁺	0.57
CaCO ₃	0.57%	Mg ⁺⁺	0.48
SP	19.65%	Na ⁺	1.27
		K ⁺	1.10
		<u>Soluble anions (meq l⁻¹)</u>	
		CO ₃ ⁻	0.00
		HCO ₃ ⁻	1.09
		Cl ⁻	0.76
		SO ₄ ⁻	1.58
		Total Solble N ppm	25.3
		Available P ppm	8.2

Plastic pots (30 cm diameter) were filled with 10 kg of the soil. All pots received the recommended dose of potassium sulphate (48.5% K₂O) at the rate of 100 kg fed⁻¹ applied at two equal doses (10 and 20 days of planting). The pots were divided into two main plots. The first main group was fertilized with super-phosphate (15.5% P₂O₅) at a rate of 200 kg fertilizer/fed. The second group was fertilized with rock phosphate (27% P₂O₅) at a rate of 115 kg fertilizer/fed. All treatments received ammonium sulphate (20.5%N) at a rate of 20 kg N fed⁻¹ as a starter dose of nitrogen. The nitrogen control treatment fertilized with 40 kg N fed⁻¹. The N-fertilizer applied at two equal doses (10 and 20 days of planting). Pots were arranged in a complete randomized design with four replicates.

Faba bean seeds variety Giza 40 were supplied from Food Legume Research Department, Field Crops Research Institute, ARC and inoculated with vermiculite-based inoculants containing *Rhizobium* or *Bacillus megaterium*. Each inoculant was applied at the rate of two bags per seeds of feddan using 16% Arabic gum solution as a sticking agent.

Mycorrhizal inoculation of experimental pots were achieved by placing 75g of the crude inoculum, 2-4 cm below the soil surface before planting. The used inoculum contained approximately 25

viable propagules per gram of mycorrhizal inoculum, which were estimated using the Most Probable Number (MPN) technique (Daniels and Skipper, 1982).

The following treatments were applied:

- 1- Uninoculated + superphosphate
- 2- Uninoculated (N-control) + superphosphate
- 3- *Rhizobium* + superphosphate
- 4- *Rhizobium* + Mycorrhizae + superphosphate
- 5- *Rhizobium* + *Bacillus megaterium* + superphosphate
- 6- Uninoculated + rock phosphate
- 7- Uninoculated (N-control) + rock phosphate
- 8- *Rhizobium* + rock phosphate
- 9- *Rhizobium* + Mycorrhizae + rock phosphate
- 10- *Rhizobium* + *Bacillus megaterium* + rock phosphate

After 60 days of planting, plants were uprooted and assayed for number and dry weight of nodules, nitrogenase activity, shoots and roots dry weight. Nitrogen and phosphorus uptake by faba bean plants were determined according to Page *et al* (1982).

Mycorrhizal colonization ratios

Representative samples of fresh roots were stained after washing several times according to the procedures of (Phillips and Hayman, 1970). Mycorrhizal colonization ratios of plant roots were determined using the gridline intersect method after staining the root samples (Giovannetti and Mosse, 1980). Data were subjected to analysis of variance according to Snedecor and Cochran (1980).

RESULTS AND DISCUSSION

Data in Table (2) clearly showed that the used soil in this study contained already indigenous Vascular Arbuscular mycorrhizal (VAM) fungi populations. However, inoculation of soil with rhizobia and mycorrhizae (*Glomus* sp.) supported higher levels of VAM colonization ratios as compared with the other treatments. Both super-P and rock-P amended soil were more pronounced with super-P amended soil. In this respect, Aryal *et al* (2003) and Jalaluddin (2005) reported that, dual inoculation of legumes with both rhizobia and VAM fungi maintained high levels of mycorrhizal colonization over than those achieved with rhizobia alone.

In the uninoculated treatments, rock-P fertilized plants exhibited significant increases in VAM colonization ratios when compared with those supplemented with Super-P, irrespective of the nitrogen

Table 2. Arbuscular-mycorrhizal colonization ratios in faba bean plant roots after 60 days of planting

Treatments	Phosphate-Source		Mean
	Super-P	Rock-P	
Uninoculated	16.24	30.30	23.27
Uninoculated (N-control)	20.50	30.45	25.48
<i>Rhizobium</i>	26.41	32.61	29.51
<i>Rhizobium</i> + AMF*	60.63	46.00	53.32
<i>Rhizobium</i> + BM**	37.52	48.33	42.92
Mean	32.26	37.54	
LSD 0.05 Inoc.			5.89
P-source	3.73		
Inoc. X P-source	8.34		

* Arbuscular Mycorrhizal Fungus

** *Bacillus megaterium* var. *phosphaticum*

dose added. These results are in harmony with those obtained by Nikolaou *et al* (2002) and Co-vacevlich *et al* (2006), who concluded that, the percent of mycorrhizal root colonization was higher in insoluble P-form treatments compared to control or to soluble P-form ones.

Irrespective of superiority of rock-P to support VAM colonization ratios in all treatments except in the case of (rhizobia + VAM) treatment which exhibited opposite trend, being 60.63 and 46.00 % for super-P and rock-P, respectively. This could be due to the high stimulation effect of rhizobia and mycorrhizal inoculation which may lead to more activation for N₂-fixation that requires more phosphorus to support plant growth. An enhancement of the N₂ fixation rates in *Rhizobium*-inoculated VAM-plants, over that achieved by *Rhizobium* in nonmycorrhizal plants was previously reported.

Concerning of dual inoculation with *Rhizobium* and *Bacillus*, the obtained data showed higher increases in mycorrhizal colonization ratios as compared with noninoculated or only rhizobial inoculated treatments, being most pronounced in the rock-P amended soil.

Data presented in Table (3) presented that N₂-ase activity of root nodules of faba bean inoculated with *Rhizobium* alone or combined with A-mycorrhizae or phosphate solubilizing bacteria gave the highest significant increases when compared to the uninoculated plants. Under rock phosphate treatment, inoculation with *Rhizobium*, *Rhizobium* plus VA-mycorrhizae and *Rhizobium* plus *Bacillus megaterium* var. *phosphaticum* stimulated N₂-ase activity of root nodules to be 32.00, 33.55 and 10.93 μmole C₂H₄ /plant/hr compared

to 11.91, 22.45 and 6.91 μmole C₂H₄/plant/hr under superphosphate fertilizer, respectively. On the contrary, the uninoculated plants gave the lowest nitrogenase activity of 0.26 and 1.06 μmole C₂H₄/plant/hr under both super phosphate and rock phosphate, respectively. On the other hand, combined inoculation of faba bean with *Rhizobium* and phosphate solubilizing bacteria did not give any significant effect on N₂-activity of faba bean plants compared to those obtained either by *Rhizobium* or inoculation alone or in combination with VA-mycorrhizae. The increased nodulation and N₂-fixation of legume crops following inoculation with N₂-fixing bacteria and phosphate solubilizing microorganisms have been reported by many workers (Algawadi and Guar, 1988; Gupta, 2004).

The obtained results in Table (4) showed clearly that combined inoculation of faba bean with *Rhizobium* and VA-mycorrhizae gave the highest significant increases in number and dry weight of nodules when compared with the plants inoculated with *Rhizobium* alone. These increases in number and dry weight of nodules reached to 32.6 and 37.1%, respectively under superphosphate fertilizer, compared to 30% and 27% under rock phosphate treatment. These results are in harmony with those obtained by Aryal *et al* (2003) they found that *Rhizobium* inoculation influenced positively nodulation of bean plants. On the other hand, combined inoculation of faba bean with *Rhizobium* and phosphate solubilizing bacteria did not record any significant effect on nodulation of faba bean plants compared to those obtained by *Rhizobium* inoculation alone.

Table 3. Effect of inoculation with rhizobia, VA-mycorrhizae and phosphate solubilizing bacteria on N₂-ase activity of faba bean plants (□ mole C₂H₄ /plant/hr)

Treatments	Phosphate-Source		Mean
	Super-P	Rock-P	
Uninoculated	1.06 ±0.15	0.26 ±0.13	0.66
Uninoculated (N-control)	0.88 ±0.10	1.91 ±0.87	1.40
<i>Rhizobium</i>	11.91 ±0.74	32.00 ±4.30	21.96
<i>Rhizobium</i> + AMF	22.45 ±3.07	33.55 ±2.90	28.00
<i>Rhizobium</i> + BM	6.91 ± 1.38	10.93 ±1.73	8.92
Mean	8.64	15.77	

Table 4. Effect of inoculation with rhizobia, VA-mycorrhizae and phosphate solubilizing bacteria on nodulation of faba bean plants

Treatments	Nodules per plant					
	Number			Dry weight (mg)		
	Super-P	Rock-P	Mean	Super-P	Rock-P	Mean
Uninoculated	5.0	2.7	3.8	14.67	4.67	9.67
Uninoculated (N-control)	3.7	3.0	3.3	13.33	6.00	9.67
<i>Rhizobium</i>	38.7	33.3	36.0	35.00	31.67	33.33
<i>Rhizobium</i> + AMF	51.3	43.3	47.3	48.33	38.67	43.50
<i>Rhizobium</i> + BM	29.3	35.0	32.2	34.67	30.00	32.33
Mean	25.6	23.5		29.20	22.20	
LSD 0.05 Inoc.	4.56			9.05		
P-source	2.89			5.72		
Inoc. X P-source	6.45			12.79		

Data in Table (5) revealed that faba bean plants inoculated with *Rhizobium* alone or combined with VA-mycorrhizae recorded the highest shoot dry weight as compared to the uninoculated plants. Combined inoculation of faba bean with *Rhizobium* and VA-mycorrhizae gave increases of 24.2% and 15% in shoots dry weight over the uninoculated ones under the application of super-phosphate and rock phosphate fertilizers, respectively. With respect to dry weight of roots, *Rhizobium* inoculation alone or combined with VA-mycorrhizae increased significantly dry weight of roots when compared to the uninoculated plants. Combined inoculation with *Rhizobium* and VA-mycorrhizae under rock phosphate treatment increased significantly upto 14.9% over those obtained in plants fertilized with superphosphate. These results are in agreement with those ob-

tained by Badr El-Din and Moawad (1988) in which the dual inoculation with both rhizobia and Mycorrhizae induced more significant increases in plant dry weight of lentil and faba bean. On the other hand, there were no significant differences in shoot or root dry weight between combined inoculation with *Rhizobium* and *Bacillus megaterium* var. *phosphaticum* compared to *Rhizobium* inoculation alone.

Combined inoculation (Table 6) with rhizobia and mycorrhizae significantly increased N-uptake in shoots up to 9.0 and 17.2% over rhizobial inoculation alone under super-P and rock-P, respectively. In addition, rhizobial inoculation increased significantly N-uptake in roots up to 62.7 and 77.5% over N-control treatments under super-P and rock-P, respectively. On the other hand, dual inoculation with rhizobia and phosphate solubilizing

Table 5. Effect of inoculation with rhizobia, VA-mycorrhizae and phosphate solubilizing bacteria on biomass of faba bean plants

Treatments	Dry wt. (g/plant)					
	Shoot			Root		
	Super-P	Rock-P	Mean	Super-P	Rock-P	Mean
Uninoculated	6.28de	5.93e	6.10	2.85	3.29	3.07
Uninoculated (N-control)	7.32ab	6.83bc	7.07	4.02	4.40	4.21
<i>Rhizobium</i>	7.80a	6.77cd	7.28	4.41	4.89	4.65
<i>Rhizobium</i> + AMF	7.62a	6.82bc	7.22	4.62	5.31	4.97
<i>Rhizobium</i> + BM	6.89bc	6.22e	6.55	3.39	4.19	3.79
Mean	7.18	6.52		3.86	4.42	
LSD 0.05 Inoc.	0.46			0.39		
P-source	0.29			0.25		
Inoc. X P-source	0.64			0.55		

Table 6. Effect of inoculation with rhizobia, VA-mycorrhizae and phosphate solubilizing bacteria on nitrogen content of faba bean plants

Treatments	Shoot			Root		
	Super-P	Rock-P	Mean	Super-P	Rock-P	Mean
	Nitrogen content (mg/plant)					
Uninoculated	187.4	162.3	174.8	29.4	30.9	30.2
Uninoculated (N-control)	240.7	236.1	238.4	50.9	55.9	53.4
<i>Rhizobium</i>	268.2	245.8	257.0	82.8	99.2	91.0
<i>Rhizobium</i> + AMF	292.4	288.0	275.2	71.8	91.5	81.6
<i>Rhizobium</i> + BM	208.2	199.5	203.9	45.9	88.8	67.3
Mean	239.4	220.3		56.15	73.26	
LSD 0.05 Inoc.	22.9			9.93		
P-source	14.50			6.27		
Inoc. X P-source	32.44			14.03		

bacteria had significant effect on N-uptake in both shoots and roots when compared to uninoculated treatments, received the same dose of N-fertilizer. With respect to P-uptake in faba bean plants (Table 7), combined inoculation with rhizobia and mycorrhizae increased significantly P-uptake in both shoots and roots of faba bean plants under the two sources of phosphorus, when compared to rhizobial inoculation alone. These increases reached to 37.5 and 31.3% in shoots and roots, under super-P, respectively. Under rock-P these increases reached to 38.0 and 47.8% in the same order. These results are in agreement with those ob-

tained by Aryal *et al* (2003) who indicated that VAM infection showed a significant positive correlation to the shoot phosphate or nitrogen in bean plants.

Also, Badr El-Din and Moawad (1988) found that the dual inoculation with both rhizobia and mycorrhizae induced more significant increases in N and P contents of lentil and faba bean. On contrary, combined inoculation with rhizobia and phosphate solubilizing bacteria did not record any significant effect on P-uptake in faba bean plants when compared to the other inoculated treatments.

Table 7. Effect of inoculation with rhizobia, VA-mycorrhizae and phosphate solubilizing bacteria on phosphorus content of faba bean plants

Treatments	Shoot			Root		
	Super-P	Rock-P	Mean	Super-P	Rock-P	Mean
	Phosphorus content (mg/plant)					
Uninoculated	9.64	9.05	9.3	1.30	1.21	1.3
Uninoculated (N-control)	37.96	33.40	35.7	2.75	2.75	2.8
Rhizobium	48.80	50.20	49.5	3.20	2.91	3.1
Rhizobium + AMF	67.10	69.3	68.3	4.20	4.30	4.2
Rhizobium + BM	35.51	30.14	32.8	2.11	2.00	2.0
Mean	39.8	38.4		2.7	2.6	
LSD 0.05 Inoc.	4.34			0.45		
P-source	2.74			0.29		
Inoc. X P-source	6.14			0.64		

REFERENCES

- Algawadi, A.R. and A.C. Guar (1988). Associative effect of *Rhizobium* and phosphate solubilizing bacteria on the yield and nutrient uptake of chickpea. *Plant and Soil*, 105: 241-246.
- Aryal, U.K.; H.L. Xu and M. Fujita (2003). Rhizobia and VAM fungal inoculation improve growth and nutrient uptake of bean plants under organic fertilization. *Journal of Sustainable Agric.*, 21(3): 215-220.
- Azcon-Aguilar, C. and J.M. Barea (1985). Effect of soil microorganisms on formation of vesicular arbuscular mycorrhizae. *Trans. Brit. Mycol. Soc.*, 83: 222-226.
- Badr El-Din, S.M.S. and H. Moawad (1988). Enhancement of nitrogen fixation in lentil, faba bean and soybean by dual inoculation with rhizobia and mycorrhizae. *Plant and Soil*, 108: 117-123.
- Barea, J.M.; M. Toro; M.O. Orozco; E. Campos and R. Azcon (2002). The application of isotopic (P-32 and N-15) dilution techniques to evaluate the interactive effect of phosphate-solubilizing rhizobacteria, mycorrhizal fungi and *Rhizobium* to improve the agronomic efficiency of rock phosphate for legume crops. *Nutrient Cycling in Agro-Eco-Systems*. 63: 35-42.
- Batjes, N.H. (1997). A World data set of derived soil properties by FAO-UNESCO soil unit for global modelling. *Soil Use Manage.* 13: 9-16.
- Bojinova, D.; R. Velkova; I. Grancharov and S. Zhelev (1997). The bioconversion of Tunisian phosphorite using *Aspergillus niger*. *Nutr. Cyc. Agro-Eco-Syst.* 47: 227-232.
- Covacevich, F.; M.A. Marino and H.E. Echeverria (2006). The phosphorus source determines the arbuscular mycorrhizal potential and the native mycorrhizal colonization of tall fescue and wheatgrass. *Eur. J. of Soil Biology.* 42: 127-138.
- Daniels, B.A. and H.D. Skipper (1982). Methods for recovery and quantitative estimation of propagules from soil. In: Schenck, N.C. (ed), *Methods and Principles of Mycorrhizal Research* (pp. 29-35). AM. Phytopathological Soc., St. Paul, MN, USA.
- Earl, C.D. and F.M. Ausubel (1993). The genetic engineering of nitrogen fixation. *Nutr. Rev.*, 41: 1-6.
- Gerke, L. (1992). Phosphate, aluminum and iron in the soil solution of three different soils in relation to varying concentrations of citric acid. *Z. Pfl.-Ernahr Bodenkde.* ISS: 17-22.
- Giovannetti, M. and B. Mosse (1980). An evaluation of techniques for measuring Vesicular-arbuscular mycorrhizal infection in roots. *New Phytol.*, 84: 489-500.
- Gull, F.Y.; I. Hafeez; M. Saleem and K.A. Malik (2004). Phosphorus uptake and growth promotion of chickpea by co-inoculation of mineral phosphate solubilizing bacteria and a mixed rhizobial culture. *Aust. J. Exp. Agric.*, 44: 623-628.

- Gupta, S.C. (2004).** Response of gram (*Cicer arietinum*) to types and methods of microbial inoculation. *India. J. Agric. Sci.*, **74**: 73-75.
- Illmer, P.; A. Barbato and E. Schinner (1995).** Solubilization of hardly soluble $AlPO_4$ with P-solubilizing microorganisms. *Soil Biol. Biochem.*, **27**: 265-270.
- Jackson, M.L. (1973).** *Soil Chemical Analysis*, p. 245. Constable Co. London, Prontice Hall Inc., Engwood Cbiffs, New Jersey.
- Jalaluddin, M. (2005).** Effect of inoculation with VAM-fungi and *Bradyrhizobium* on growth and yield of soybean in Sindh. *Pakistan J. of Botany*. **37**: 169-173.
- Kang, S.C.; C.G. Ha; T.G.; Lee and D.K. Maheshwari (2002).** Solubilization of insoluble inorganic phosphates by a soil fungus *fomitopsis* sp. PS. 102. *Curr. Sci.*, **82**: 439-442.
- Kucey, R.M.N.; H.H. Jenzen and M.E. Leggett (1989).** Microbially mediated increase in plant available phosphorus. *Adv. Agron.*, **42**: 199-228.
- Lindsay, W.L.; P.L.G. Viek and S.H. Chien (1989).** Phosphate minerals. In: *Minerals in Soil Environment*. 2nd Ed. pp. 1089-1130. (Eds. Dixon, J.B. and S.B. Weed). Soil Sci. Soc. of Amer., Madison, USA.
- Maliha, R.; K. Samina; A. Najma; A. Sadia and L. Farooq (2004).** Organic acids production and phosphate solubilization by phosphate solubilizing microorganisms under *in vitro* conditions. *Pak. J. Biol. Sci.*, **7**: 187-196.
- Nikolaou, N.; N. Karagiannidis; S. Koundouras and I. Fysarakis (2002).** Effects of different P sources in soil on increasing growth and mineral uptake of mycorrhizal *Vitis vinifera* L. (cv Victoria) vines. *J. International Des Sciences De La Vigne Et Du Vin*. **36(4)**:195-204.
- Norrish, K. and H. Rosser (1983).** *Mineral Phosphates in Soil an Australian View Point*. pp. 335-361, Academic Press. Metbourne. Csiro/London, U.K./Australia.
- Page, A.I.; R.H. Miller and D.R. Keeney (1982).** *Methods of Analysis*. Part 2, Chemical and Microbiological Properties, Second Edition, Am. Soc. Agron. Inc., Soil Sci. Soc. Am. pp. 595-624.
- Peoples, M.B.; D.F. Herridge and J.K. Ladha (1995).** Biological Nitrogen Fixation an efficient source of nitrogen for sustainable agriculture. *Plant and Soil*, **174**: 220-225.
- Perveen, S.; M.S. Khan and A. Zaidi (2002).** Effect of rhizospheric microorganisms on growth and yield of greengram (*Phaseolus radiatus*). *Ind. J. Agric Sci.*, **72**: 421-423.
- Phillips, J.M. and D.S. Hayman (1970).** Improved procedures for clearing roots and staining parasitic and vesicular-arbuscular mycorrhizal fungi for rapid assessment of infection. *Trans. Br. Mycol. Soc.*, **55**: 158-161.
- Rajan, S.S.S.; J.H. Watkinson and A.G. Sinclair (1996).** Rock Phosphate for direct application to soils. *Adv. Agron.*, **57**: 77-159.
- Schilling, G.; A. Aranse; A. Deubel; G. Lezovic and S. Ruppel (1998).** Phosphorus availability, root exudates and microbial activity in the rhizosphere. *Z. Pfl. Ernahr. Bodenkde*, **161**: 465-478.
- Snedecore, G.W. and W.G. Cochran (1980).** *Statistical Methods*, pp. 255-269. Seventh ed., Iowa State Univ. Press, Ames, Iowa, USA.
- Vessey, K.J. (2003).** Plant growth promoting rhizobacteria as biofertilizers. *Plant and Soil*, **255**: 571-586.
- Vincent, J.M. (1970).** *A Manual for Practical Study of Root-Nodule Bacteria*. IBP Handbook, No. 15, Blackwell Sci., Pub. Oxford, UK.
- Zaidi, A. (1999).** *Synergistic Interactions Of Nitrogen Fixing Microorganisms With Phosphate Mobilizing Microorganisms*. p. 150. Ph.D. Thesis, Alqath Muslim Univ. Alqath.
- Zaidi, A.; M.S. Khan and M. Amil (2003).** Interactive effect of rhizotrophic microorganisms on yield and nutrient uptake of chickpea (*Cicer arietinum* L.). *Eur. J. Agron.*, **19**: 15-21.
- Zaidi, A.; M.S. Khan and A. Amil (2004).** Bioasocitive effect of rhizospheric microorganisms on growth yield and nutrient uptake of greengram. *J. Plant Nutr.*, **27**: 599-610.
- Zapata F. and R.N. Roy (2004).** Use of phosphate rocks for sustainable agriculture. In: *Fertilizer and Nutrition Bulletin*. No. 13. A joint production of the FAO and water development division agency, Food Agriculture Organization of the United Nations, Roma.



إستجابة نباتات الفول البلدى للتلقيح المشترك بالريزوبيا والميكورهيذا والبكتريا المذيبة للفوسفات تحت ظروف الاراضى الرملية

[٢٤]

صالح احمد صالح^١ - محمود عبد المقصود ناصف^٢ - فتحى توفيق ميخائيل^٢ -

علاء الدين عبد الحميد أبو السعود^٢

١- البنك القومى للجينات والاصول الورثية - مركز البحوث الزراعية - الجيزة - مصر
٢- معهد بحوث الاراضى والمياة والبيئة - مركز البحوث الزراعية - الجيزة - مصر

بالمعاملات الاخرى. وكذلك أدى التسميد بالصخر الفوسفاتى إلى زيادة نسبة تواجد تجمعات الميكورهيذا عند مقارنتها بالتسميد بالسوبر فوسفات. كما ان التلقيح بالريزوبيا مشتركة مع (*Bacillus megaterium* var. *phosphaticum*) أدى إلى زيادة نسبة تجمعات الميكورهيذا بالمقارنة بالمعاملة الغير ملقحة او الملقحة بالريزوبيا منفردة.

كما أوضحت النتائج أيضاً ان تلقيح الفول البلدى بالريزوبيا مشتركة مع الميكورهيذا فى وجود الصخر الفوسفاتى او السوبر فوسفات أدى الى زيادة معنوية لتكوين العقد البكتيرية والتثبيت النيتروجينى والنمو الخضرى ومحتوى النبات من النيتروجين والفوسفور عند مقارنتها بالريزوبيا منفردة. بينما لم يظهر التلقيح بالريزوبيا المشتركة مع (*Bacillus megaterium* var. *phosphaticum*) تأثير معنوي للقياسات السابقة بالمقارنة بالريزوبيا منفردة.

تم إجراء تجربة أصص فى صوبة وحدة إنتاج الاسمدة الحيوية - معهد بحوث الاراضى والمياة والبيئة - مركز البحوث الزراعية - بالجيزة. أستخدم فيها التلقيح بالبكتريا المثبتة لنيتروجين الهواء الجوى (*Rhizobium leguminosarum* bv. *viceae*) مشتركة مع البكتريا المذيبة للفوسفات (*Bacillus megaterium* var. *phosphaticum*) وكذلك فطر الميكورهيذا (*Glomus* sp.) وذلك بهدف تقييم كفاءة تكوين العقد الجذرية وتثبيت النيتروجين الجوى وتيسير إمتصاص الفوسفات لنباتات الفول البلدى النامية فى اراضى رملية ذات محتوى منخفض من الفوسفور لزيادة محتوى المجموع الخضرى للنبات من النيتروجين والفوسفور وتأثيره على النمو.

وقد أظهرت النتائج زيادة معنوية لنسبة تواجد تجمعات فطر الميكورهيذا حول منطقة الجذور نتيجة التلقيح بالريزوبيا مشتركة مع الميكورهيذا بالمقارنة