

Dept. of Food Hygiene,
Fac. of Vet. Medicine, Beni-Sueif University.

**MONITORING OF PSYCHROTROPHIC BACTERIA
IN MILK AND SOME DAIRY PRODUCTS WITH
SPECIAL REFERENCE TO PSEUDOMONAS SPECIES**
(With 6 Tables)

By

**A.M. EL-KHOLY; SAADIA H.E.; G.M. HASSAN
and KHADIGA A.H.**

(Received at 9/6/2008)

**رصد البكتيريا المحبة للبرودة وبخاصة أنواع السيدوموناس في اللبن
وبعض منتجاته**

**عادل مصطفى الخولى ، سعادى حلمى الشناوى ، جمال محمد حسن ،
خديجة عبد الخالق حسن**

اجريت هذه الدراسة على ٧٥ عينة من اللبن ومنتجاته (٢٥ عينة من كل من اللبن والأيس كريم "من صغار المنتجين" والجبن الثلجة) جمعت كلها بطريقة عشوائية من أسواق ومحلات مدينة بنى سويف لمعرفة مدى تواجد ميكروبات السيدوموناس والميكروبات المحبة للبرودة فيها والتي استحوذت في السنوات الماضية على اهتمام الكثير من المشتغلين فى مجال علوم التغذية نظراً لما تتمتع به هذه الميكروبات من صفات بالإضافة إلى انتشارها الواسع فى الطبيعة وقدرتها على إفراز العديد من السموم المسببة للإسهال والنزلات المعوية فهى تستطيع أن تنمو وتتكاثر فى درجات الحرارة المنخفضة. هذا وقد تبين بالفحص البكتريولوجي أن ٢١ (٨٤%) ، ١٥ (٦٠%) و ١٨ (٧٢%) من عينات اللبن ، الأيس كريم والجبن الثلجة كانت ملوثة بالسيدوموناس وذلك باستخدام مستنبت Cetrimide agar وكان متوسط العدد الكلى لهذا الميكروب هو $10 \times 9 \pm 10 \times 3,3$ ، $10 \times 1,2$ ، $10 \times 6,4$ و $10 \times 7,6 \pm 10 \times 5,2$ لكل ملي أو جرام من العينات المفحوصة على التوالي. باستخدام مستنبت Glutamate starch phenol red agar (GSP) كان معدل وجود ميكروب السيدوموناس فى العينات اللبن ٢١ (٨٤%) ، الأيس كريم ١٦ (٦٤%) والجبن الثلجة ٢٠ (٨٠%) وكان متوسط العدد الكلى لهذا الميكروب هو $10 \times 1,7$ ، $10 \times 4,6$ ، $10 \times 2,7 \pm 10 \times 1,2$ و $10 \times 1,1$ ، $10 \times 3,7 \pm 10 \times 1,1$ لكل ملي أو جرام من العينات المفحوصة على التوالي. وقد تم عمل تصنيف لعترات السيدوموناس المعزولة على مستنبت Cetrimide agar وعلى مستنبت GSP. هذا وقد تم فحص العترات المعزولة من ميكروب السيدوموناس لمعرفة إمكانية إنتاجها لانزيمات الليباز والبروتياز ووجد أن ٧١ عترة أنتجت انزيم الليباز بينما ٧٠ عترة أنتجت انزيم البروتياز هذا وقد تم عمل عد للميكروبات المحبة للبرودة الموجودة فى العينات ووجد أن متوسط العدد الكلى لهذه

الميكروبات كان $10 \times 7,9 \pm 10 \times 1,95$ ، $10 \times 1,9 \pm 10 \times 6,7$ و $10 \times 3,5 \pm 10 \times 1,7$ لعينات اللبن ، الأيس كريم والجبن الثلجة على التوالي وكان معدل وجود هذه الميكروبات في عينات اللبن ، الأيس كريم والجبن الثلجة هو ٢١ (٨٤%) ، ١٨ (٧٢%) و ٢١ (٨٤%) . هذا وقد تم مناقشة الأهمية الصحية والإقتصادية لميكروب السيدوموناس .

SUMMARY

Seventy five random samples of milk and some dairy products including raw milk, small scale ice cream and Talaga cheese (each of 25) were collected from different localities in Bani-Suef city. Pseudomonas species could be counted, isolated and identified on cetrimide agar and GSP agar at different counts and percentages, also the incidence and count of psychrotrophic organisms was determined. The characterization of some isolated pseudomonas spp. from the examined samples for the production of extracellular virulence factors as lipolytic and proteolytic enzymes, were determined and from that 71 pseudomonas spp. showed lipolytic activity while 70 isolates of them showed proteolytic activity. The public health significance of the organism and the precautions, which should be taken to control this organism in the dairy industry as well as the recommended sanitary measures, were also discussed.

Key words: Psychrotrophos, Pseudomonas, Milk, dairy products.

INTRODUCTION

Milk and dairy products are generally very rich in nutrients which provide an ideal growth environment for many micro-organisms. The microbial quality of raw milk is crucial for the production of good quality dairy products.

Members of the Pseudomonas spp. (Yamamoto *et al.*, 2000) are aerobic gram-negative straight or slightly curved rods, non-spore forming, not-acid fast, non-fermentative bacteria that are widely distributed in nature (Migula, 1894). They are 1 to 5 μm long and 0.5 to 1.0 μm wide.

It belongs to family Pseudomonadaceae (Pitt, 1998). Some isolates grow under anaerobic conditions by using nitrate as a terminal electron acceptor. They are able to grow over a wide range of temperature (4°C to 44°C), and at neutral or alkaline PH (7-8.5) but most are not able to grow at PH 6 or below (Holt *et al.*, 1994).

Many species produce characteristic water soluble pigments. The yellow green pigment pyoverdin (Fluorescein) is produced by most *Pseudomonas* strains, giving the characteristic blue-green appearance (Meyer *et al.*, 2002). *Ps. aeruginosa* can produce the phenazines pyocyanin (blue pigment) and pyorubin (red pigment) (Lau *et al.*, 2004). Rare isolates of *Ps. aeruginosa* and *Ps. cepacia* produce the dark brown pigment pyomelanin (Gerald Colle, 1996).

Psychrotrophic bacteria typically enter processed dairy products through post-pasteurization contaminants in the milk processing plants (Moseley, 1980 and Ralyea *et al.*, 1998).

As a result of their metabolic diversity, ability to grow at low temperature and ubiquitous nature, many *Pseudomonas* spp. can reduce the shelf life of processed milk and cause food spoilage leading to significant economic loss for the food industry. Types of spoilage differ according to the species of *Pseudomonas* and fat content of milk.

Many strains of *Pseudomonas* produces heat-stable extracellular lipases, proteases and lecithinases which cause casein digestion leading to a bitter flavor and clotting and gelation of milk or rancid bitter taste and unclean and soapy appearance and many of these enzymes remain active even following thermal processing steps that can destroy the organisms which produce these enzymes (Sorhaug and Stepanik, 1997 and Dogan and Boor, 2003).

Under stressful conditions, these organisms produce an exopolysaccharides that are known as slime layers (biofilm formation). These exopolysaccharides make it difficult to be phagocytosed by mammalian white blood cells (Ryan and Ray, 2004).

Furthermore, *Ps. aeruginosa* could grow and multiply to numbers sufficient to induce food poisoning (Cheung and Westhoff, 1983).

Several food-borne diarrheal outbreaks linking these organisms were recorded in Yugoslavia (Kenderski, 1974), India (Perena *et al.*, 1977), Canada (Todd, 1981), Bangladesh (Mitra *et al.*, 1993), Taiwan (Wuby *et al.*, 1999) and Nigeria (Nzeako and Okafor, 2002).

In recognition of the economic and public health significance of *pseudomonas* spp., the present study was planed to fulfill the following:

- 1 - Enumeration, isolation and identification of *Pseudomonas* spp. from milk, ice-cream and talaga cheese.
- 2 - Enumeration of Psychrotrophic bacteria.
- 3 - Detection of Lipolytic and proteolytic activities of the isolated *Pseudomonas* species from the different examined samples.

MATERIALS and METHODS

A- Collection of samples:

75 samples were randomly collected from different localities including groceries, supermarkets, dairy shops in Bani-Suef city as raw milk, ice-cream and talaga cheese (each of 25). The samples were directly transferred to the laboratory with a minimum time of delay.

B- Preparation of samples and serial dilutions:

It was performed according to A.P.H.A. (1992).

C- Bacteriological examination:

1- Enumeration, Isolation and Identification of Pseudomonas organisms (Uroz and Citak, 1998): By using GSP and cetrimide agar plates.

2- Psychrotrophic count (SPC):

It was performed according to A.P.H.A. (1992).

3- Detection of the Lipolytic and proteolytic activities of isolated Pseudomonas spp. from the examined milk and selected dairy products were determined according to Harrigan and McCance. (1976).

RESULTS

Table 1: Statistical analytical results of the examined Milk, Talaga Cheese and Ice-Cream samples based on pseudomonas count/ ml or gm on Cetrimide and Gsp agar.

samples	No. of Examined Samples	Positive Samples					
		Cetrimide agar			GSP agar		
		No.	%	Mean ± SEM	No.	%	Mean ± SEM
Milk	25	21	84	$9 \times 10^4 \pm 3.3 \times 10^4$	21	84	$1.7 \times 10^5 \pm 4.6 \times 10^4$
Ice - Cream	25	15	60	$1.2 \times 10^5 \pm 6.4 \times 10^4$	16	64	$2.7 \times 10^5 \pm 1.2 \times 10^5$
Talaga cheese	25	18	72	$7.6 \times 10^4 \pm 5.2 \times 10^4$	20	80	$1.1 \times 10^5 \pm 3.7 \times 10^4$

Table 2: Incidence and frequency distribution of the isolated pseudomonas spp. from the examined milk and Ice-cream samples using Cetrimide and Gsp agar.

Pseudomonas spp	Milk						Ice - Cream					
	Cetrimide agar			GSP agar			Cetrimide agar			GSP agar		
	No.	Incidence %	frequency	No.	Incidence %	frequency	No.	Incidence %	frequency	No.	Incidence %	frequency
Ps.aeruginosa	3	12	14.29	7	28	33.33	4	16	26.67	6	24	37.5
Ps.fluorescens	8	32	38.09	4	16	19.05	3	12	20	4	16	25
Ps. Putida	4	16	19.05	2	8	9.52	1	4	6.67	2	8	12.5
Ps. stutzeri	2	8	9.52	4	16	19.05	4	16	26.67	1	4	6.25
Ps. mendocina	3	12	14.28	1	4	4.76	1	4	6.67	2	8	12.5
Ps. alcaligenes	1	4	4.76	3	12	14.28	2	8	13.33	1	4	6.25
Total	21	84	100	21	84	100	15	60	100	16	64	100

Table 3: Incidence and frequency distribution of the isolated pseudomonas spp. from the examined Talaga cheese samples using Cetrimide and Gsp agar.

Pseudomonas spp.	Talaga cheese					
	Cetrimide agar			GSP agar		
	No.	Incidence%	frequency	No.	Incidence%	frequency
Ps.aeurginosa	2	8	11.11	3	12	15
Ps.fluorescens	6	24	33.33	5	20	25
Ps. putida	4	16	22.22	2	8	10
Ps. stutzeri	3	12	16.67	2	8	10
Ps.mendocina	1	4	5.56	4	16	20
Ps.alcaligenes	2	8	11.11	4	16	20
Ps.vesicularis	0	0	0	0	0	0
Total	18	72	100	20	80	100

Table 4: Lipolytic and proteolytic activities of the isolated pseudomonas spp. from the examined milk and Ice- cream samples.

Pseudomonas spp.	milk					ice- cream				
	No. of tested isolates	Lipolytic		proteolytic		No. of tested isolates	Lipolytic		proteolytic	
		+ ve no.	%	+ ve no.	%		+ ve no.	%	+ ve no.	%
Ps.aeurginosa	10	10	100	10	100	10	8	80	10	100
Ps.fluorescens	12	10	83.33	9	75	7	7	100	5	71.43
Ps. Putida	6	4	66.67	0	0	3	1	33.33	0	0
Ps. Stutzeri	6	3	50	6	100	5	3	60	3	60
Ps.mendocina	4	1	25	3	75	3	1	33.33	1	33.33
Ps.alcaligenes	4	0	0	2	50	3	1	33.33	0	0
Total	42	28	66.67	30	71.43	31	21	67.74	19	61.29

Table 5: Lipolytic and proteolytic activities of the isolated pseudomonas spp. from the examined Talaga cheese.

Pseudomonas spp.	Talaga cheese				
	No. of tested isolates	Lipolytic		proteolytic	
		+ ve no.	%	+ ve no.	%
Ps.aeurginosa	5	4	80	5	100
Ps.fluorescens	11	9	81.81	8	72.73
Ps. Putida	6	5	83.33	-	-
Ps. Stutzeri	5	-	-	2	40
Ps.mendocina	5	1	20	1	20
Ps.alcaligenes	6	3	50	5	83.33
Ps. Vesicularis	-	-	-	-	-
Total	38	22	57.89	21	55.26

Table 6: Statistical analytical results of the examined milk, Talaga cheese and ice-cream samples based on their psychrotrophic count/ml or gm by using standard plate count agar (SPC).

Examined Samples	No. of examined Samples	Positive Samples		Min	Max	Mean	± SEM
		No.	%				
Milk	25	21	84	< 10	3×10^7	6.7×10^6	1.9×10^6
Ice – Cream	25	18	72	< 10	1.58×10^7	1.95×10^6	7.9×10^5
Talaga Cheese	25	21	84	< 10	3×10^7	3.5×10^6	1.7×10^6

DISCUSSION

I. Microbiological examination:

A- Incidence and count of *Pseudomonas* spp. In the examined samples:

1- Raw milk:

The results reported in Table 1 revealed that the pseudomonas species could be detected in 21 (84 %) from the examined milk samples with a mean value of $9 \times 10^4 \pm 3.3 \times 10^4$ and $1.7 \times 10^5 \pm 4.6 \times 10^4$, cfu/ml on each cetrinide and GSP agar, respectively.

The obtained incidence of pseudomonas species are in agreement to those recorded by Gennari and Dragotto (1992) while lower incidence of pseudomonas species in milk was recorded by Abdel – Khalek and El-Gamal (1998) Aly and Zaki (2001) and Ewida (2005), however, higher incidence were obtained by Ahmed (1995) and Al-Ashmawy *et al.* (1997).

As shown in Table 2, the isolated of pseudomonas spp. on cetrinide agar could be differentiated into *Ps. aeruginosa* 12%, *Ps. fluorescens* 32%, *Ps. putida* 16%, *Ps. stutzeri* 8%, *Ps. mendocina* 12% and *Ps. alcaligenes* 4%, while, on GSP agar, the incidence were 28%, 16%, 8%, 16%, 4% and 12%, respectively. The frequency distribution of different isolates of pseudomonas spp. on cetrinide agar were 14.29, 38.09, 19.05, 9.52, 14.28 and 4.76%, respectively, while on GSP agar these isolates were detected in frequency percentages of 33.33, 19.05, 9.52, 19.05, 4.76 and 14.28%, respectively.

The incidence of *Ps. aeruginosa* nearly was in harmony with the results recorded by Al-Ashmawy *et al.* (1992), Khalil (1992) and Aly and Zaki (2001), but was higher than that obtained by Ahmed *et al.*

(2002), El-Said (2002), Ahmed and Sotohy (2003) and Ewida (2005). Higher levels were recorded by Zaki *et al.* (1996), Al-Ashmawy *et al.* (1997) and Ezzeldeen *et al.* (2004).

The obtained results of the incidence of *Ps. fluorescens* was nearly similar to those recorded by Zaki *et al.* (1996) and El-Said (2002), while higher percentages were recorded by Al-Ashmawy *et al.* (1997), Sami (1999) and Dogan and Boor (2003), while the lower incidence were reported by Uraz and citak (1998), Ahmed *et al.* (2002) and Ewida (2005).

In this study it was found that *Ps. fluorescens* was the predominant bacteria which cause spoilage due to secretion of hydrolytic enzymes such as lipase and protease. Nearly similar findings were reported by Abdel-Hakiem (1996), Dieckelmann *et al.* (1998), Sobeih (2000) and Dogan and Boor (2003).

The *Ps. putida* constituted (16% and 8%) on cetrimide and GSP agar, respectively of the examined milk samples. Higher percentage was recorded by Al-Ashmawy *et al.* (1997) and Aly and Zaki (2001), but lower incidence was obtained by Craven and Macauley (1992), Uraz and Citak (1998) and Ewida (2005).

Ps. aeruginosa is the most important human pathogen in the genus *pseudomonas* with respect to both the numbers and types of caused infections and their associated morbidity and mortality (Pollack, 1990). Moreover, the presence of *Ps. aeruginosa* in the intestinal tract of both man and animal and food could be taken as an index of faecal contamination (Hoadley and McCay 1968). The contamination of milk with *Ps. aeruginosa* may be from the environment of dairy farms such as contaminated water, soil, utensils and faecal matter (Eneroth *et al.*, 1998) and this should be minimized by good manufacturing practices (GMP) at the farm and during production and handling of milk.

Milk are liable to contaminate with *pseudomonas* organisms during production, handling and processing as well as from animals suffering from mastitis. This organism could grow and multiply to numbers sufficient to induce spoilage of contaminated milk.

Pseudomonas can cause gastroenteritis if ingested in large number ($> 10^6$) but the food could be clearly spoiled by *pseudomonas* before reaching the limited number (Johnson 1990).

2- Ice-Cream:

As shown in Table 1 *pseudomonas* species could be detected in 60% and 64% of the examined ice cream samples on cetrimide and GSP

agar, respectively, with an average count of $1.2 \times 10^5 \pm 6.4 \times 10^4$ and $2.7 \times 10^5 \pm 1.2 \times 10^5$ cfu / ml, respectively .

Lower incidence were obtained by El-Bassiony *et al.* (1985), Korashy (1992) and Gomaa (1999).

The incidence of pseudomonas spp. recovered from ice cream samples using cetrimide agar was *Ps. aeruginosa* 16%, *Ps. fluorescens* 12%, *Ps. putida* 4%, *Ps. stutzeri* 16%, *Ps. mendocina* 4%, and *Ps. alcaligenes* 8%, while, on GSP agar these organisms were recovered from 24%, 16%, 8%, 4%, 8% and 4%, respectively (Table2). Lower incidence of these isolates was detected by Gomaa (1999).

In the present study Table 3 viewed that the frequency distribution of the isolates were *Ps. aeruginosa* (26.67 and 37.5), *Ps. fluorescens* (20 and 25), *Ps. putida* (6.67 and 12.5), *Ps. stutzeri* (26.67 and 6.25), *Ps. mendocina* (6.67 and 12.5) and *Ps. alcaligenes* (13.3 and 6.25) on cetrimide and GSP agars, respectively.

Most of these isolates could be isolated from ice cream samples at different percentages by Ahmed (1980), Saad (1983), Grover *et al.* (1993), Shanker *et al.* (1994) and Kasana *et al.* (2002)

Higher pseudomonas count in small vendors ice cream samples may be an index of insufficient heat treatment, improper freezing, contaminated water and using of unclean utensils as well as milker's hands (Otte *et al.*, 1978).

3- Talaga Cheese:

The obtained results in Table 1 revealed that, 72 and 80% of examined Talage cheese samples were contaminated with pseudomonas spp. on cetrimide and GSP agars with a mean value of $7.6 \times 10^4 \pm 5.2 \times 10^4$ and $1.1 \times 10^5 \pm 3.7 \times 10^4$ cfu / gm on both media respectively.

The lower incidence of pseudomonas spp. in cheese was recorded by Desmeasure *et al.* (1995).

The summarized results in Table 3 showed that *Ps. aeruginosa* could be isolated from 8% and 12% of cheese samples, *Ps. fluorescens* could be isolated from 24% and 20% samples; *Ps. putida* from 16% and 8% samples *Ps. stutzeri* from 12% and 8% samples; *Ps. mendocina* from 4% and 16% samples; while *Ps. alcaligenes* could be detected in 8% and 16% samples, respectively by using cetrimide and GSP agar.

Frequency distribution of isolated *Ps. spp.* from Talage cheese on cetrimide agar Table 3 revealed that 11.11%, 33.33%, 22.22%, 16.67%, 5.56% and 11.11% of the isolates were identified as *Ps. aeruginosa*, *Ps. fluorescens*, *Ps. putida*, *Ps. stutzeri*, *Ps. mendocina* and

Ps. alcaligenes, while, on GSP these isolates were detected at percentages of 15, 25, 10, 10, 20 and 20 respectively.

These organisms could be detected at different percentages by El-Bassiony *et al.* (1985), Kasaria *et al.* (2002), Salmeron *et al.* (2002) and Leriche *et al.* (2004).

pseudomonas induce important defects in cheese such as yellow to brown coloration (Leriche *et al.*, 2004) and lead to development of bad ripening flora, flavour and texture defects such as bitterness and running paste (Champagne *et al.*, 1994).

Presence of *Ps.* organisms in Talage cheese could be attributed to the method of manufacturing, unheat treated milk and bad hygienic measures as well as water used in cheese production (Cantoni *et al.*, 2003). On the other hand, use of pasteurized milk and applying good manufacturing practices are extremely important for production of safe white soft cheese to consumers.

B- Lipolytic and Proteolytic activities of the isolated *pseudomonas* spp. from milk and dairy products

1- Raw milk:

Out of 42 *pseudomonas* spp. isolates from raw milk samples, 28 (66.67%) showed lipolytic activity on tributyrin agar (Table 4), they differentiated into 10 (100%) *Ps. aeurginosa*, 10 (83.33%) *Ps. fluorescens*, 4 (66.67%) *Ps. putida*, 3 (50%) *Ps. stutzeri*, 1 (25%) *Ps. mendocina*, while *Ps. alcaligenes* couldn't show lipolytic activity.

These results are in agreement with those reported by El-Said (2002) and Ewida (2005). Lower results were obtained by Garg (1990), Jaspe *et al.* (1995) and Wiedmann *et al.* (2000), while higher findings were reported by Ahmed (1995) and Al- Ashmawy *et al.* (1997).

As shown in (Table 4), 30 (71.43%) isolates from 42 *pseudomonas* spp. isolates had proteolytic activity and differentiated as follow *Ps. aeurginosa* 10 (100%), *Ps. fluorescens* 9 (75%), *Ps. stutzeri* 6(100%), *Ps. mendocina* 3(75%) and *Ps. Alcaligenes* 2 (50%).

These results are nearly similar to Al- Ashmawy *et al.* (1997), El-Said (2002) and Ewida (2005).

2- Ice-Cream:

Inspection the results in (Table 4) declared that 21 (67.74%) strains out of 31 isolates from examined ice cream samples showed lipolytic activity by using tributyrin agar. The positive strains of examined *pseudomonas* for lipolytic activity were, *Ps. aeurginosa* 8(80%), *Ps. fluorescens* 7(100%), *Ps. putida* 1(33.33%), *Ps. stutzeri* 3(60%), *Ps. mendocina* 1(33.33%) and *Ps. alcaligenes* 1(33.33%). On

the other hand, out of 31 pseudomonas strains from ice cream samples, 19(61.29%) had proteolytic activity on skim milk agar and a high positive percentages as 100%, 71.43% and 60% were *Ps. aeruginosa*, *Ps. fluorescens* and *Ps. stutzeri*, respectively followed by *Ps. mendocina* (33.33%). While, *Ps. putida* and *Ps. alcaligenes* didn't show proteolytic activity.

3- Talaga Cheese:

Out of 38 pseudomonas spp. isolates from Talaga cheese samples, 22 (57.89%) showed lipolytic activity on tributyrin agar (Table 5), as 4(80%), 9(81.81%), 5(83.33), 1(20%) and 3(50%) were *Ps. aeruginosa*, *Ps. fluorescens*, *Ps. putida*, *Ps. mendocina* and *Ps. alcaligenes*, respectively.

Samples showed proteolytic activity were 21 (55.26%) from 38 pseudomonas spp. isolates and differentiated as follow 5(100%), 8(72.73), 2 (40%), 1(20%) and 5 (83.33) were *Ps. aeruginosa*, *Ps. fluorescens*, *Ps. stutzeri*, *Ps. mendocina* and *Ps. alcaligenes*, respectively.

Many studies have been reported that pseudomonas species are psychrotrophic organisms that can cause spoilage of milk and dairy products as they can produce Lipolytic and proteolytic enzymes which secreted in raw milk during preprocessing stages and survive pasteurization causing reduction in the sensory quality and shelf life of processed fluid milk products (Lopez-Fandino *et al.*, 1993) for example, digestion of casein by proteases can lead to bitter flavor, clotting and gelatin of milk. Lipases hydrolyze tributyrin and milk fat to yield free fatty defects acids, which can produce a range of flavor, described as cheesy, fishy, malty, putrid, soapy and unclean (Cox,1993 and Shah, 1994).

C- Incidence of psychrotrophic bacteria in the examined samples:

1- Milk samples:

The findings reported in (Table 6) indicated that the incidence of psychrotrophic organisms in the examined milk samples was 84% with a count ranged from < 10 to 3×10^7 cfu / ml, and a mean count of $6.7 \times 10^6 \pm 1.9 \times 10^6$

Similar results for psychrotrophic count in milk samples were reported by Prabha *et al.* (1996) and Kasana *et al.* (2002), while, lower results were reported by Malik and Mathur (1983), Garg (1990) and So *et al.* (1992).

2- Ice-Cream:

The results given in (Table 6) showed that the incidence of psychrotrophic organisms was 72% with a count ranged from < 10 to

1.58×10^7 cfu / ml with a mean value of $1.95 \times 10^6 \pm 7.9 \times 10^5$ cfu / ml. The obtained results of psychrotrophic organisms are in agreement to those recorded by El-Bassiony *et al.* (1985), while higher results were recorded by El-Bagoury (1996) and Kasana *et al.* (2002).

3- Talaga cheese:

From the obtained results in (Table 6) 84% of Talaga cheese samples were contaminated with psychrotrophic organisms with a count ranged from <10 to 3×10^7 cfu / gm with a mean value of $3.5 \times 10^6 \pm 1.7 \times 10^6$ cfu / gm. Lower results were reported by Santos *et al.* (1996) and Kasana *et al.* (2002).

Although the control of bacterial growth during storage of milk and milk products depends primarily on refrigeration, yet the psychrotrophic bacteria can grow at this low temperature. Longer the refrigerated storage before processing of raw milk, larger chances of spoilage could be caused in milk and dairy products by psychrotrophs (Kasana *et al.*, 2002).

II-Public health significance

Ps. aeruginosa is an opportunistic human pathogen, most commonly affecting immunocompromised patients such as those with cystic fibrosis, Infection can affect many different parts of the body, but typically targets the respiratory system causing chronic debilitating pulmonary infection due to mucoid variants that are now the major cause of death in patients with cystic fibrosis (Shanson, 1990 and Elkin and Geddes, 2003). Involvement of gastrointestinal tract most commonly occurs in infants and patients with hematologic malignancies and neutropenia that has resulted from chemotherapy and this mostly lead to pseudomonal bacteremia. The spectrum of disease can range from very mild symptoms to severe necrotizing enterocolitis with significant mortality. The infection can cause enteritis, with patients presenting with prostration, headache, fever and diarrhea (Shanghai fever). Young infants may present with irritability, vomiting, diarrhea and dehydration. *Ps. aeruginosa* is also a common cause of bacterial keratitis, scleral abscess, and endophthalmitis in adults and ophthalmia neonatorum in children. *Pseudomonas* when introduced produces extracellular enzymes that cause a rapidly progressive and destructive lesion. Also it can cause meningitis and brain abscess. These infections can involve the urinary tract through an ascending infection or through bacteremia spread (Pollack, 1990).

Pseudomonas typhitis typically present in patients with neutropenia resulting from acute leukemia with a sudden onset of fever, abdominal pain.

Pseudomonas is a common cause of chronic otitis media (Swimmer's ear), patients present with pain, pruritis and ear discharge (Patrick, 1995). *Pseudomonas* also has emerged as an important source of burn wound sepsis (Holder, 1993). It can also cause endocarditis and infections of bones and joints. Moreover, the organism is responsible for a number of mastitis cases and remains in the udder for a number of years (Howell, 1972).

REFERENCES

- A.P.H.A. (1992): "Standard methods for the examination of dairy products", 16th Ed., American Public Health Association, Washington, D.C.*
- Abdel-Hakiem, E.H. (1996): "The role of lipolytic psychrotrophic bacteria and their activities in cold stored raw milk", 7th Sci. Cong., Fac. Vet. Med., Assiut Univ., Egypt.*
- Abd-El-Khalek, A. and El-Gamal, A.M. (1998): "Quality control of pasteurized milk sold at Mansoura city in Egypt", Benha Vet. Med. J., 9 (1), 70-79.*
- Ahmed, A.K.S. (1995): "Assessment of Pseudomonas in farm bulk milk", PhD Thesis, Fac. Vet. Med., Zagazig Univ., Egypt.*
- Ahmed, H.F.; Ismail, A.M. and Tonios, A.I. (2002): "Occurance of Pseudomonas in raw cow's milk, and trials to control Ps. aeruginosa in butter and Tallaga cheese", Vet. Med. J., 50 (4), 589-599.*
- Ahmed, M.M. and Sotohy, S.A. (2003): "Sanitary conditions of milking environment in Assiut dairy farms and the quality of their produced milk", Assiut Vet. Med. J., 49 (97), 88-112.*
- Ahmed, S.H. (1980): Studies on acute gastrointestinal infection in Assiut Governorate. Ph. D. thesis, Assiut University.*
- Al-Ashmawy, A.M.; El-Gamal, A.M. and Abd-El-Khalek, A. (1997): "Prevalence of Pseudomonas in farm bulk milk", Assiut Vet. Med. J., 36 (2), 215-221.*
- Al-Ashmawy, A.M.; El-Sherbini, M. and El-Sayed, M.S. (1992): "Prevalence of Ps. aeruginosa and Ps. fluorescens in farm milk", Zagazig Vet. J., 20 (3), 383-388.*

- Aly, Salwa and Zaki, Hoda (2001): "Studies on Pseudomonas species in milk and feces of dairy cattle", Assiut Vet. Med. J., 46 (2), 172-186.*
- Cantoni, C.; Stella, S.; Cozzi, M.; Iacumin, L. and Comi, G. (2003): Blue colouring of Mozzarella cheese. Industrie Alimentari., 42 (428): 840 – 843.*
- Champagne, C.; Laing, R.; Roy, D.; Mafu, A. and Griffiths, M. (1994): "Psychrotrophs in dairy products; effect and control", Crit. Rev. Food Sci. Nutr., 34 (1), 1-30.*
- Cheung, B.A. and Westhoff, D.C. (1983): "Isolation and identification of ropy bacteria in raw milk", J. Dairy Sci., 66 (9), 1825-1834.*
- Cox, J.M. (1993): "The significance of psychrotrophic pseudomonas in dairy products", Aust. J. Dairy Technol., 48, 108-113.*
- Craven, H.M. and Macauley, B.J. (1992): "Psychrotrophic microorganisms in refrigerated pasteurized milk", Brief Communication of the XXVIII International Dairy Congress, Montreal, Vol. 1, 8-12.*
- Desmasures, N.; Radiguet, S.; Lejeune, J.; and Gueguen, M. (1995): "Effect of ripening on the microbiological profile of high quality raw milk for cheese making", Milchwissenschaft, 50 (4), 193-195.*
- Dieckelmann, M.; Johnson, L.A. and Beacham, I.R. (1998): "The diversity of lipases from psychrotrophic strains of Pseudomonas; a novel lipase from a highly lipolytic strains of Ps. fluorescens", J. Appl. Microbiol., 85, 527-536.*
- Dogan, B. and Boor, K.J. (2003): "Genetic diversity and spoilage potentials among Pseudomonas spp. isolated from fluid milk products and dairy processing plants", Appl. Environ. Microbiol., 69 (1), 130-138.*
- El-Bagoury, A.M. (1996): Bacteriological quality of market ice cream. Vet. Med. J., Giza. Vol 44 (2): 203 – 207.*
- El-Bassiony, T.A.; Aboul-Khier, F. and Saad, N.M. (1985): Psychrotrophic bacteria in dairy products. Assiut Vet. Med. J., Vol 15 (29): 101–105.*
- Elkin, S. and Geddes, D. (2003): "Pseudomonas infection in cystic fibrosis", Expert. Review of Anti-infection Therapy, 1 (4): 611-615.*
- El-Said, E.I. (2002): "Studies on food poisoning bacteria with special reference to Campylobacter jejuni in milk", PhD Thesis, Fac. Vet. Med., Zagazig Univ., Egypt.*

- Eneroth, A.; Christiansson, A.; Brendehaug, J. and Molin, G. (1998):* "Critical contamination sites in the production line of pasteurized milk with reference to the psychrotrophic spoilage flora", *Int. Dairy J.*, 8 (9): 829-834.
- Ewida-Rania, M.M.A. (2005):* Survey of the pseudomonas bacteria in milk sold in Assiut city. M.V.Sc. These, Fac. Vet. Med., Assiut Univ. Egypt.
- Ezzeldeen, Nashwa, A.; Abdel-Fattah, S.H. and Soliman, Y.A. (2004):* "Phenotypic and molecular characterization of *Ps. aeruginosa* of animals and human origins", *J. Egypt. Vet. Med. Assoc.*, 64 (1): 93-104.
- Garg, S.K. (1990):* "Psychrotrophs in milk", *A review, Ind. J. Dairy Sci.*, 43 (3), 278-284.
- Gennari, M. and Dragotto, F. (1992):* "A study of the incidence of different fluorescent *Pseudomonas* species and biovars in the microflora of fresh and spoiled meat, fish, raw milk, cheese, soil and water", *J. Appl. Bacteriol.*, 72 (4): 281-288.
- Gerald Collee (1996):* *Practical Medical Microbiology*. 4th Ed. Pp: 413.
- Gomaa-Maha, M. (1999):* Studies on the proteolytic psychrotrophic bacteria in ice cream. B.V.S. Thesis, Fac. Vet. Med., Zagazig Univ., Egypt.
- Grover, S.; Batish, V.K. and Srinivasan, R.A. (1993):* Growth proteolytic and enterotoxic activities of *pseudomonas aeruginosa* CTM-3 in milk, ice cream and butter. *Archive. Fur. Lebensmittel Hygiene* 44 (5): 115-117.
- Harrigan, W.E. and McCance, M.F. (1976):* "Laboratory methods in food and dairy microbiology", Academic press, London.
- Hoadley, A.W. and McCay, E. (1968):* Some observation on the ecology of *pseudomonas aeruginosa* and its occurrence in the intestinal tracts of animals. *Corne II vet.*, 58: 345-363.
- Holder, I.A. (1993):* "*Ps. aeruginosa* virulence associated factors and their role in burn wound infection", R.B. Ficks, Jr (Ed.); *Ps. aeruginosa; the opportunistic*, CRC press, Boca Roton, 235-245.
- Holt, J.G.; Krieg, N.R.; Sneath, P.H.A.; Staley, J.T. and Williams, S.T. (1994):* "*Berges's Manual of Determinative Bacteriology*", 9th Ed., Washington, D.C.
- Howell, D. (1972):* Survey on mastitis caused by environmental bacteria. *Vet. Rec.*, 90: 554-557.

- Jaspe, A.; Oviedo, P.; Fernaudez, L.; Palacios, P. and Sanjose, C. (1995): "Cooling raw milk: changes in the spoilage potential of contaminating Pseudomonas", J. Food Prot., 58 (8): 915-921.*
- Johnson, E.A. (1990): In food borne diseases. Ed. Cliver D.O., Academic press Limited, London.*
- Kasana, R.C.; Punj, V. and Matta, H. (2002): "Isolation and identification of gram-negative lipase producing psychrotrophic bacteria from some dairy products", Ind. J. Dairy Sci., 55 (1): 13-16.*
- Kenderski, S. (1974): "Cases of food poisoning in Yugoslavia", Flieschwirtschaft, 54 (5): 931-933.*
- Khalil, Nawal (1992): "Occurance, detection and significance of Ps. aeruginosa in raw milk", Assiut Vet, Med. J., 28 (2): 152-157.*
- Korashy, Eman, A. (1992): "Incidence of Ps. aeruginosa in milk and some milk products", M.V.Sc. Thesis, Fac.of Vet. Med., Assuit Univ., Egypt.*
- Lau, G.W.; Hassett, D.J.; Ran, H. and Kong, F. (2004): "The role of pyocyanin in Ps. aeruginosa infection", Trends in Molecular Medicine, 10 (2): 599-606.*
- Leriche, F.; Bordessules, A.; Fayolle, K.; Karoui, R.; Laval, K.; Leblanc, L. and Dufour, E. (2004): "Alteration of raw milk cheese by Pseudomonas spp.", J. Microbiol. Methods, 59 (1): 33-41.*
- Lopez-Fandino, R.; Olano, A.; Corzo, N. and Romos, M. (1993): "Proteolysis during storage of UHT milk; differences between whole and skim milk", J. Dairy Res., 60, 339-347.*
- Malik, R.K. and Mathur, D.K. (1983): Isolation and identification of protease producing psychrotrophic bacteria from dairy products in India. J. Soc. Dairy Technol. 36: 76-78.*
- Meyer, J.M.; Geoffrey, V.A. and Baida, N. (2002): "Siderophore typing, a powerful tool for the identification of fluorescent and non fluorescent Pseudomonas", Appl. Environ. Microbiol., 68 (6): 2745-2753.*
- Migula, W. (1894): Ube rein neues system der bakterien, Arbeiten aus dem bakteriologischen Institut der Technischen, 1, 235-238.*
- Mitra, A.K.; Albert, A.J. and Alam, A.N. (1993): Bacteraemia and meningitis among hospital patients with diarrhea. Trans. R. Soc. Trop. Med. Hyg., 87 (5): 560-563.*
- Moseley, W.K. (1980): "Pin pointing post pasteurization contamination", J. Food Prot., 43, 414-415.*

- Nzeako, B. and Okafor, N. (2002)*: Bacterial enteropathogens and factors associated with seasonal episodes of gastroenteritis in Nsukka, Nigeria. *Br. J. Biomed. Sci.*, 59 (2): 76–79.
- Otte, I.; Hahn, G. and Tollin, A. (1978)*: Detection, incidence and significance of pseudomonas aeruginosa in raw milk and the environment of dairy plants. *Dairy Sci. Abst.* 40 (9).
- Patrick, R.M. (1995)*: "Manual of clinical microbiology", 6th Ed., ASU press, Washington, D.C., chapter 3.
- Perena, S.; Mathan, M.; Bhat, P.; Albert, M.J. and Baket, J. (1977)*: "Etiology of acute gastroenteritis in fancy and early childhood in southern India", *Arch. Dis. Child.*, 52 (6): 482–485.
- Pitt, T.L. (1998)*: "Pseudomonas, Burkholderia and related genera", *Topley and Wilson's Microbiology and Microbial Infections*, 9th Ed., Oxford Univ. Press, New York, chapter 47.
- Pollack, M. (1990)*: "Principles and practice of infectious diseases", 3rd Ed., Churchill Livingstone, New York.
- Prabha, R.; Krishna, R. and Shanker, P.A. (1996)*: "Identification of gram-negative rod shaped psychrotrophic bacteria of dairy origin", *Ind. J. Dairy Sci.*, 49 (8): 517–518.
- Ralyea, R.D.; Wiedmann, M. and Boor, K.J. (1998)*: "Practical tracking in a dairy production system using phenotypic and ribotypic methods", *J. Food Prot.*, 61, 1336–1340.
- Ryan, K.J. and Ray, C.G. (2004)*: "Sherris Medical Microbiology", 4th Ed., MC Graw Hill, ISBN.
- Saad, Nagah, M. (1983)*: Psychrotrophic bacteria in milk and milk product. M.V.Sc. Thesis, faculty of vet. Med., Assiut university Egypt.
- Salmeron, J.; Devega, C.; Perez–Elortondo, F.J.; Albisu, M. and Barron, L.J.P. (2002)*: Effect of pasteurization and seasonal variations in the microflora of ewe's milk for cheese making (19): 167–174.
- Sami, Enas, M. (1999)*: "Prevalence of Aeromonas spp. in milk and some milk products in sharkia Governorate", M.V.Sc. Thesis, Fac. Vet. Med., Zagazig Univ., Egypt.
- Santos, J.A.; Lopez, Diaz, T.M.; Garcia–Fernandez, M.C.; Garcia–Lopez, M.L. and Otero, A. (1996)*: Characterization and extracellular activity of Psychrotrophic bacteria isolated from villalon cheese (Fresh variety of Spanish sheep's milk cheese). *International journal of food Microbiology.*, 33 (2): 301–306.

- Shah, N.P. (1994): "Psychrotrophs in milk", A review, Milchwissenschaft, 49 (8): 432-437.*
- Shanker, K.; Narasimhan, R. and Khan, M.M.H. (1994): Incidence, isolation and characterization of psychrotrophs in ice cream. Indian Vet. J., 71 (12): 1202-1206.*
- Shanson, D.C. (1990): "Septicaemia in patients with AIDS", Trans. R. Soc. Trop. Med. Hyg., 84, 14-16.*
- So, M.H.; Yoon, S.S. and Kim, Y.B. (1992): "Psychrotrophic microflora in raw milk and their proteolytic and lipolytic ability", Korean J. Dairy Sci., 14 (1): 43-51.*
- Sobeih, Azza, M. (2000): "Studies of raw milk quality in Kafr El-Sheikh", PhD Thesis, Fac. Vet. Med., Tanta Univ. (Kafr El-Sheikh Branch), Egypt.*
- Sorhaug, T. and Stepanik, L. (1997): "Psychrotrophs and their enzymes in milk and dairy products; quality aspects", Trends Food Sci. Technol., 8, 35 - 40.*
- Todd, E.C.D. (1981): "Food borne and water borne diseases in Canada", J. Food Prot., 44, 787-795.*
- Uraz, G. and Citak, S. (1998): The isolated of pseudomonas and other Gram (-) psychrotrophic bacteria in raw milk. J. Basic Microbiol., 38 (2): 129-134.*
- Wiedmann, M.; Weilmier, D.; Dneen, S.S.; Ralyea, R. and Boor, K.J. (2000): "Molecular and phenotypic characterization of Pseudomonas spp. isolated from milk", Appl. Environ. Microbiol., 66 (5): 2085-2095.*
- Wuby, P.C.T.; Tsai, C.H. and Chiu, H.H. (1999): Community-acquired Ps. aeruginosa bacteremia and sepsis in previously healthy infants. J. Acta paediat., Taiwan, 40 (4): 233-236.*
- Yamamoto, S.; Kasai, H.; Arndt, D.L.; Jaskson, R.W. and Harayama, S. (2000): "Phylogeny of the genus Pseudomonas; intragenetic structure reconstructed from the nucleotide sequences of gyrB and proD genes", Microbiol., 146, 2385-2394.*
- Zaki, M.S.A.; Abd-El-Khalek, A. and Mowafy, L.E. (1996): "Hygienic importance of Pseudomonas in dairy farms", 7th Sci. Cong., Fac. Vet. Med., Assiut Univ., Egypt.*