

Dept. of Biochemistry
Animal Health Research Institute, El. Mansoura Lab.

**BIOCHEMICAL AND HAEMATOLOGICAL STUDIES
ON POST PARTURIENT HAEMOGLOBINUREA
IN CATTLE REARED IN NEWLY
RECLAIMED AREA**
(With 5 Tables)

By

**H.A. SHALABY; S.A. TAWFIK; R.A. EL-RAMADY
and S.S.S. YONIS**

(Received at 18/10/2008)

**دراسات بيوكيميائية ودموية علي مرض البول المدمم بعد الولاده بالماشيه
المرباه في مناطق حديثه الاستصلاح**

**حامد عبد المجيد الامام شلبي ، سيد احمد توفيق عيد ، رأفت احمد الرمادي
السيد السيد السيد يونس**

أجريت هذه الدراسة بهدف معرفة تأثير مرض البول المدمم بعد الولاده في الماشيه في المناطق المستصلحه حديثا علي صوره الدم وبعض القياسات البيوكيميائية. وقد أجريت هذه الدراسة علي عدد ٤٠ رأسا من الماشيه تراوحت أعمارها بين ٦ الى ٩ سنوات ظهرت عليها اعراض فقدان في الشهية ورفض أكل البرسيم وتغير لون البول الي الأحمر وأحيانا يميل الي لون القهوه وايضا نقص سريع في انتاج اللبن. وكذلك اشتملت هذه الدراسة على ١٥ رأسا من الماشيه السليمه ظاهريا كانت تتغذي علي برسيم وعليه متوازنه في نفس مكان التجريه واستخدمت كضوابط. وقد تم دراسة الأعراض الأكلينيكية وأخذ عينات من دم ومصل وبول تلك الحيوانات قبل وبعد العلاج. وأسفرت نتائج التحليل انخفاض معنوي للعد الكلي لكرات الدم الحمراء وتركيز الهيموجلوبين وحجم الخلايا المضغوط وضغط الاوكسجين الوريدي وارتفاع معنوي للعد الكلي لكرات الدم البيضاء وضغط ثاني اكسيد الكربون وأيضا زياده غير معنوية في تركيز الاس الهيدروجيني والبيوكربونات في الدم في الحالات المريضة عند مقارنتها بالمجموعه الضابطه. وبالنسبه للتحليل البيوكيميائي لمصل الدم وجد انخفاض معنوي في مستوى البروتين الكلي والألبومين والجلوبيولين والجلوكوز والكوليسترول والدهون الثلاثية والفوسفور والمغنسيوم والصوديوم والبوتاسيوم والزنك والحديد والنحاس. وكذلك وجد ارتفاع معنوي في مستوى نشاط انزيمات الكبد وانزيم الفوسفاتيز القاعدي ونسبة البولينا والمولبيدنيوم في المجموعه المريضة بمقارنتها بالمجموعه الضابطه. وعند تحليل البول للحيوانات المريضة وجد ارتفاع معنوي للفوسفور والكالسيوم والمغنسيوم والبروتين واليوريا والكرياتينين بمقارنتها بالمجموعه الضابطه. وأظهرت النتائج بالنسبة لتحليل البرسيم ان تركيز النحاس والكالسيوم كان في حدود معدله الطبيعي مع

وجود نقص في تركيز عنصر الفوسفور وزيادة نسبة عنصر الموليبدينوم بمقارنته بالبرسيم المنزرع في القرى القديمة وقد ارجع ظهور الاعراض الى النقص الاولى في عنصر الفوسفور والنقص الثانوي في عنصر النحاس. وتم علاج حالات الماشية المصابة بمرض البول المدمم بحقن مركب فوسفات الصوديوم الحامضي (H_2PO_4) (٦٠ جرام في ٣٠٠ مل ماء معقم) في الوريد وتحت الجلد كل ١٢ ساعة لمدة ٣ الى ٤ ايام. بالاضافة الى الحقن العضلي للتونوفوسفان ٢٠% ٢٠ مل يوميا بالاضافة الى جلاسينات النحاس. وفي الحالات المريضة المتقدمة قمنا بنقل الدم لها. وقد عادت كل هذه القياسات في الحيوانات التي تم علاجها الى مايقرب من مستواها الطبيعي خلال شهر كما تم التحسن من الناحية الصحية حيث اعراض البول المدمم. ويجب ان ننتبه الى الحرص في التشخيص الاكلينيكي والمعمل للحيوانات المصابة وعلاجها مع توعية مالكي ومستصلي هذه الاراضي الجديدة بضرورة اتباع الاسلوب السليم في تغذية الحيوانات وعدم الاعتماد كلية على البرسيم والاعتماد على عليقة متزنة في موسم البرسيم وازضافة مخلوط الأملاح المعدنية والعناصر النادرة والفيتامينات الى هذه العلائق اذا لزم الامر حتى نتلافى الخسائر الاقتصادية الناجمة عن ذلك.

SUMMARY

The aim of the present study is to investigate the effect of post parturient haemoglobinuria (PPHU) on haematological and biochemical parameters of cattle excessively fed on *Trifolium Alexandrium* (Barseem) during the green season in newly reclaimed area. Blood and urine samples were collected from 40 diseased cattle aged from 6-9 years showing signs of loss of appetite, refuse barseem eating, red to coffee coloured urine and milk production dropped rapidly. A number of 15 clinically healthy cattle were used as control group. The haematological studies showed significant decrease in TRBCs, Hb concentration, PCV and significant increase in TWBCs. The blood hydrogen ion concentration (pH) and bicarbonate were found non significant increased. Blood gas analysis showed significant increase in PCO_2 and significant decrease in PO_2 . Serum biochemical analysis revealed that cattle with PPHU was accompanied by a marked decrease in total protein, albumin, globulins, glucose, total cholesterol, total triglycerides, inorganic phosphorus, magnesium, sodium, potassium, zinc, iron and copper levels. Moreover significant increase in AST, ALT, ALP activity levels, blood urea and molybdenum levels was noticed when compared with apparent clinically healthy ones. Moreover a marked increase in the excretion of phosphorus, calcium, molybdenum, protein, urea and creatinine through urine was observed only in PPHU affected cattle. Concerning the diet, analysis of barseem grows in newly reclaimed area, the obtained values of copper and calcium was adequate while phosphorus showed significant decrease

levels. Molybdenum level has an opposite direction to copper level and showed highly significant increases in its values when compared with barseem grow in old village. The affected PPHU cattle were treated according to the severity of the case by sodium acid phosphate and tonophosphan injection. Copper glycinate was also tried to halt haemolysis. Complicated cases were completed cured after blood transfusion. The symptoms disappeared, the appetite returned to normal and haematological as well as biochemical measured parameters in serum and urine returned nearly to normal. So attention must be paid for careful early clinical and laboratory diagnosis of diseased animals followed by therapeutic plan with good management. Further more excessive feeding on barseem must be avoided and feeding balanced ration mixed with minerals, trace elements and vitamins during the green season.

Key words: *Haemoglobinurea, blood analysis, urine analysis.*

INTRODUCTION

Post parturient haemoglobin urea (PPHU) is a major disease of dairy animals with detrimental economic consequences, Chugh *et al.* (1996). The post parturient haemoglobinurea is a type of haemolytic anaemia and excessive RBCs fragility associated with inadequate ATP formation developed in high producing multiparous cows reared on ration deficient in phosphates in early lactation, (Omran *et al.*, 1987). The exact aetiology and pathogenesis of PPHU are not known, as a variety of aetiological factors have been reported to be associated with the disease in different parts of the world, (Hussain *et al.*, 1991 and Chugh *et al.*, 1996). Dietary phosphorus deficiency and / or ration containing cruciferous plants are suspected causes of sever hypophosphataemia, (Pirzada and Hussain, 1998). Hypophosphataemia primarily affecting highly producing dairy cattle and buffaloes associated with prolonged feeding on barseem. Its prevalence during the period from March to May, (Nassif, 1995). It is hypothesised that hypophosphataemia results in decreased red blood cells glycolysis and ATP synthesis by inhibiting the glucose-6-phosphate dehydrogenase. Subnormal concentration of ATP predisposes red blood cells to altered structure and function, a loss of normal deformability and an increase in fragility and haemolysis, with resultant haemoglobinaemia and haemoglobinuria, (Ogawa *et al.*, 1989 and Radostits *et al.*, 2000). Copper deficiency is also an aetiological factor of Post PHU, as its

deficiency reduces the activity of the copper-containing enzyme superoxide dismutase, which is part of the erythrocyte protection mechanism against oxidative stress (Smith *et al.*, 1975). Phosphate deficiency also compromises oxygen release to the tissue due to the decreases in erythrocyte, (Jeffrey, *et al.*, 2005). Abdul-Samad (1997) recorded that at least 29%-42% of the cases of phosphorus deficiency, haemoglobinurea in buffaloes could be due to pregnancy and late stage of gestation. The author reported also that summer season is one of the putative causal factors in phosphorus deficiency haemoglobin urea. Bhikane *et al.* (1995) recorded that post parturient haemoglobinurea is characterized by coffee coloured urine, anaemia, loss of appetite and reduced milk yield with significant hypophosphataemia.

In view of the above speculations, the present investigation was undertaken to study the haematological and serum biochemical changes in cattle suffering from post parturient haemoglobinurea with history of excessive prolonged feeding on trifolium Alexandrium (barseem) in newly reclaimed areas of Egypt. This may help in the solution of this problem, as well as to reach ideal way for treatment and will contribute to the understanding of aetiopathology of post parturient haemoglobinurea. Suggestive measures for the prevention and control of disease in cattle was also aimed.

MATERIALS and METHODS

1- Animals:

The present investigation was carried out on 40 female multiparous dairy cattle suffering from post parturient haemoglobinurea (PPHU), aged 6-9 years, that were randomly selected from field cases that raised in newly reclaimed area in Salhia, Hosynia and the villages surrounded it districts in Sharkia Governorate, during the period between March till May 2008. All animals were heavily fed on Trifolium alexandrinum (Barseem) for at least 5 months. The disease was clinically diagnosed on the basis of specific signs during early lactation (2-4weeks) such as loss of appetite, refuse barseem eating, preferring dry feeds, red to coffee coloured urine, milk production dropped rapidly and straining while defecating. The control group included 15 clinically healthy cattle of similar description from the same location and received balanced ration (ration concentrate mixture consisted cotton seed cake 15%, soya bean meal 3%, wheat bran 55%, yellow corn 9%, rice polish 6%, molasses 7%, lime stone 3.5%, common salt 1.5%) during the green

season. according to NRC (1985). History of affected examination concerning animals' data, nutrition and management were recorded before the animals being clinically examined. A routine faecal examination was made for the detection of internal parasites. Blood film examination and standard laboratory methods were used to rule out other diseases that cause a reddish discoloration of urine such as babesiosis, leptospirosis or bacillary haemoglobinuria.

2-Sampling:

Two blood samples were collected from diseased group as well as from control animals before and after one month post treatment.

A- The first blood samples were collected in heparinized tube for the determination of total erythrocytes (TRBCs) and total leukocytes (TWBCs) counts, packed cell volume (PCV), haemoglobin concentration following the techniques described by Coles (1986) Red blood cells indices were also calculated. While blood gases and acid-base parameter (pH, HCO₃, PO₂, PCO₂) estimated by using radiometer of Copenhagen ABL810 Blood gases analyzer.

B- Another blood samples collected without anticoagulant for serum separation and preserved at -20°C. Serum samples were used for determination of Aspartate aminotransferase (AST), Alanine aminotransferase (ALT) activity level according to Reitman and Frankel (1957), Alkaline phosphatase activity level (ALP) Bowers and Mc Comb (1975), total proteins level Doumos (1974), albumin level Drupt (1974) and serum globulins levels were calculated as difference between total protein and albumin. Serum glucose level was determined after Trinder (1969), while serum calcium level after Gindler and King (1972). Serum inorganic phosphorus level after Morinal and Prox (1973), magnesium level was determined according to Gindler and Heath (1971). Serum sodium and potassium levels were determined according to the method described by Hawk (1965) by using a flame-photometer (corning model AVL 988-3, made in U.S.A). Serum iron, zinc, copper and molybdenum levels were estimated by atomic absorption spectrophotometer according to Cowell (1973). Also serum samples were used for determination of blood urea level after Patton and Crouch (1977); creatinine level after Thomas (1992); total cholesterol level was estimated according to Stein (1986) and total triglyceride was analyzed after Mgtowan *et al.* (1983).

C- Urine samples:

Fresh urine samples were obtained from both diseased and apparently healthy cattle and used for biochemical analysis of

phosphorus, calcium, magnesium, proteins, urea and creatinine levels according to the methods mentioned before in serum analysis.

D-Faecal samples:

Faecal samples were collected from each examined animal for detection of any internal parasite according to Soulsby (1986).

F-Diet samples:

Five samples of barseem from recently reclaimed area and from old villages were collected for phosphorus, calcium, copper and molybdenum analysis according to A.O.A.C. (1975).

3-Treatment trails:

The principal line of treatment based on correction of the feeding system of the affected animals and compensation of the occurred deficiency status

- 1- Correction of feeding system: avoiding the excessive feeding on barseem and prepared balanced ration during the green season.
- 2- Transfusion of large quantities of whole blood is the treatment for severely affected cows; blood transfusion was conducted between animals of the same family. Blood was collected into an anti coagulant solution in an open-mouthed vessel.
- 3- Treatment with sodium acid phosphate (60gm in 300ml of sterile water injected I.V. followed by subcutaneously injection every 12hours 3-4 days and daily I/M injection of 20 ml tonophosphan.^R 20% (Hookiest company)
- 4- Copper glycinate (120 mg available copper) one dose of intravenous injection will maintain adequate copper levels for about 60-90 days.

4- Statistical analysis:

The obtained data was statistically analyzed according to Snedecor and Cochran (1982) using the "SAS" computer program.

RESULTS

A-Clinical signs:

The main clinical signs observed included inappetence and depraved appetite with refuse barseem eating, preferring dry feeds. Pale to icteric visible mucous membranes, weakness, accelerated heart rate, rapid and shallow respiration was also detected. Red to coffee coloured urine, straining while defecating or during early lactation appeared. Body temperature ranged from 38-39.1 C and milk production dropped rapidly.

B- Haematological results

Post parturient haemoglobinurea (PPHU) induced very highly significant $P < 0.001$ decrease in total erythrocytic counts, haemoglobin concentration and packed cell volume reading Table (1). Mean corpuscular volume (MCV) and mean corpuscular haemoglobin (MCH) showed non significant increase. But mean corpuscular haemoglobin concentration (MCHC) showed non significant decrease. Concerning the total leucocytic count in the present study showed very highly significant increase. Mean while, cattle affected by (PPHU) showed highly significant increase in P_{CO_2} while P_{O_2} showed very highly significant decrease. The results referred to a type of anaemic anoxia. The blood hydrogen ion concentration (pH) and bicarbonate (Acid-base parameters) were found non significant increased. These parameters were returned to the normal levels nearly 30 days post treatment.

C- Biochemical results:

The present data (Table2) revealed significant $P < 0.001$ decreases in serum total proteins, albumin, globulins, glucose, total cholesterol, and total triglycerides levels. On contrary significant increases $P < 0.01$ in blood serum activities of AST, ALT, ALP and blood urea levels were noticed. Meanwhile creatinine level showed non significant increase. The obtained results in blood serum macro and micro elements in cattle affected with PPHU (Table 3) revealed a very highly significant $P < 0.001$ decrease in phosphorus, magnesium, sodium, potassium, iron and copper levels. But the mean values of calcium and zinc levels showed non significant decrease. While the molybdenum level recorded very highly significant $P < 0.001$ increase.

Concerning urine analysis, post parturient haemoglobinurea (PPHU) induced very highly significant increase $P < 0.001$ in phosphorus, calcium, magnesium, protein, urea and creatinine in urine of the diseased cattle when compared with healthy one Table (4). This biochemical parameters returned to the nearly normal levels 30 days after treatment. Analysis of samples of barseem from recently reclaimed area and from old agricultural area (old villages) revealed that the concentrations of phosphorus, calcium, copper and molybdenum content were suggestive of primary phosphorus and secondary copper deficiency in recently reclaimed area, while the level of molybdenum showed very highly significant increase than the old villages Table (5).

Table 1: Haematological variables (mean $\pm$ SE) in blood of healthy cattle and those affected with post parturient haemoglobinurea (PPHU) before and one month after treatment.

Items Parameters	Control healthy cattle	PPHU affected cattle	Post treatment
TRBCs ($\times 10^6$ /cum)	7.60 $\pm$ 0.41	3.92 $\pm$ 1.60 ^{***}	6.7 $\pm$ 0.35
Hb (gm/dl)	11.79 $\pm$ 0.28	6.48 $\pm$ 0.48 ^{***}	9.97 $\pm$ 0.59
PCV (%)	34.50 $\pm$ 1.04	17.92 $\pm$ 0.66 ^{***}	30.2 $\pm$ 1.75
MCV (cuu)	46.05 $\pm$ 1.87	49.66 $\pm$ 6.60 ^N	45.46 $\pm$ 1.57
MCH (ug)	15.62 $\pm$ 0.75	19.24 $\pm$ 1.87 ^N	14.75 $\pm$ 0.52
MCHC (gm/dl)	33.96 $\pm$ 0.92	30.47 $\pm$ 1.42 ^N	32.36 $\pm$ 1.37
TWBCs ($\times 10^3$ cum)	7.90 $\pm$ 0.29	10.06 $\pm$ 0.46 ^{***}	7.29 $\pm$ 0.37
Hydrogen ion (pH)	7.41 $\pm$ 0.008	7.67 $\pm$ 0.02 ^N	7.36 $\pm$ 0.009
Bicarbonate (mmol/L)	26.53 $\pm$ 2.30	31.39 $\pm$ 2.50 ^N	27.40 $\pm$ 1.41
Pco2 (mm Hg)	46.95 $\pm$ 2.02	55.45 $\pm$ 2.70 ^{**}	48.44 $\pm$ 2.0
Po2 (mm Hg)	42.90 $\pm$ 1.14	36.95 $\pm$ 0.66 ^{***}	43.10 $\pm$ 1.53

N: non significant

*: highly significant at (p<0.01)

***: very highly significant at (p<0.001)

Table 2: Biochemical variables (mean $\pm$ SE) in blood of healthy cattle and those affected with post parturient haemoglobinurea (PPHU) before and one month after treatment.

Items Parameters	Control healthy cattle	PPHU affected cattle	Post treatment
T.Proteins (gm/dl)	7.12 $\pm$ 0.19	4.82 $\pm$ 0.17 ^{***}	6.27 $\pm$ 0.26
Albumin (gm/dl)	3.40 $\pm$ 0.16	2.22 $\pm$ 0.15 ^{**}	2.86 $\pm$ 0.35
Globulins (gm/dl)	3.70 $\pm$ 0.17	2.60 $\pm$ 0.13 ^{***}	3.08 $\pm$ 0.14
Glucose (gm/dl)	80.82 $\pm$ 1.62	46.96 $\pm$ 1.75 ^{***}	71.04 $\pm$ 2.17
AST (u/L)	59.91 $\pm$ 3.17	73.62 $\pm$ 2.96 ^{**}	58.36 $\pm$ 2.42
ALT (u/L)	17.45 $\pm$ 1.11	30.65 $\pm$ 1.88 ^{***}	18.57 $\pm$ 0.78
ALP (u/L)	15.90 $\pm$ 1.69	22.62 $\pm$ 1.61 ^{**}	15.40 $\pm$ 1.87
B. Urea (gm/dl)	30.87 $\pm$ 2.26	47.89 $\pm$ 4.69 ^{**}	33.17 $\pm$ 2.47
Creatinine (gm/dl)	1.13 $\pm$ 0.07	1.84 $\pm$ 0.10 ^N	1.06 $\pm$ 0.08
T.Cholesterol (gm/dl)	126.31 $\pm$ 3.56	83.27 $\pm$ 2.49 ^{***}	118.40 $\pm$ 4.91
T.Triglycerides (gm/dl)	96.2 $\pm$ 2.30	81.67 $\pm$ 2.20 ^{***}	95.53 $\pm$ 1.50

N: non significant

*: highly significant at (p<0.01)

***: very highly significant at (p<0.001)

Table 3: Blood serum macro and micro elements levels (mean $\pm$ SE) in healthy cattle and those affected with post parturient haemoglobinurea (PPHU) before and one month after treatment.

Items Parameters	Control healthy cattle	PPHU affected cattle	Post treatment
Calcium (gm/dl)	10.0 $\pm$ 0.29	9.40 $\pm$ 0.39 ^N	9.92 $\pm$ 0.40
Phosphorus (gm/dl)	5.49 $\pm$ 0.14	2.15 $\pm$ 0.29 ^{***}	4.34 $\pm$ 0.45
Magnesium (gm/dl)	2.76 $\pm$ 0.10	2.42 $\pm$ 0.09 ^{***}	2.56 $\pm$ 0.09
Sodium (mmol/L)	142.96 $\pm$ 1.18	134.3 $\pm$ 1.60 ^{***}	140.72 $\pm$ 1.46
Potassium (mmol/L)	5.28 $\pm$ 0.14	3.18 $\pm$ 0.21 ^{***}	4.64 $\pm$ 0.17
Zinc (ug/L)	161.13 $\pm$ 5.69	143.19 $\pm$ 7.15 ^N	165.15 $\pm$ 4.14
Iron (ug/L)	156.24 $\pm$ 1.26	142.20 $\pm$ 2.66 ^{***}	152.59 $\pm$ 1.11
Copper (ug/dl)	72.14 $\pm$ 1.35	54.54 $\pm$ 3.45 ^{***}	62.88 $\pm$ 2.27
Molybdenum (ug/dl)	55.54 $\pm$ 0.89	72.11 $\pm$ 1.47 ^{***}	57.29 $\pm$ 0.87

N: non significant

***: very highly significant at (p<0.00)

Table 4: Mean $\pm$ SE of minerals, total proteins and protein non nitrogenous compounds in the urine in healthy cattle and those affected with post parturient haemoglobinurea (PPHU) before and one month after treatment.

Items Parameters	Control healthy cattle	PPHU affected cattle	Post treatment
Phosphorus (gm/dl)	14.91 $\pm$ 0.43	18.60 $\pm$ 0.51 ^{***}	16.38 $\pm$ 0.46
Calcium (gm/dl)	11.85 $\pm$ 0.21	15.01 $\pm$ 0.37 ^{***}	10.12 $\pm$ 0.38
Magnesium (gm/dl)	28.56 $\pm$ 0.43	31.42 $\pm$ 0.32 ^{***}	30.32 $\pm$ 0.19
Protein (gm/dl)	31.54 $\pm$ 0.16	295.53 $\pm$ 0.39 ^{***}	37.48 $\pm$ 0.50
Urea (u/L)	0.398 $\pm$ 0.052	1.11 $\pm$ 0.107 ^{***}	0.424 $\pm$ 0.008
Creatinine (u/L)	29.15 $\pm$ 0.26	37.32 $\pm$ 0.52 ^{***}	27.42 $\pm$ 0.30

***: very highly significant at (p<0.001)

Table 5: Mean $\pm$ SE of phosphorus, calcium, copper and molybdenum of barseem from newly reclaimed area and old agricultural area (old villages) on dry matter basis.

Items Parameter	DM basis			
	Copper mg/kg	Molybdenum mg/kg	Phosphorus %	Calcium %
Barseem from newly reclaimed area	5.78 $\pm$ 0.14 ^N	4.9 $\pm$ 0.27 ^{***}	0.17 $\pm$ 0.01 ^{**}	3.43 $\pm$ 0.15
Barseem from old villages	7.1 $\pm$ 0.02	0.86 $\pm$ 0.01	0.31 $\pm$ 0.02	3.51 $\pm$ 0.02

N: non significant m

**: highly significant at (p<0.01)

***: very highly significant at (p<0.001)

DISCUSSION

The clinical observation of cattle suffering from post parturient haemoglobinurea (PPHU) during early lactation (2-4 weeks) showed decreased and depraved appetite, refuse barseem eating, preferring dry feeds, pale to icteric visible mucous membranes, weakness, accelerated heart rate, increase in pulse rate, rapid and shallow respiration, red to coffee coloured urine, straining while defecating. Body temperature ranged from 38 – 39.1 C and milk production drop rapidly. The clinical signs were similar to those reported by Abdel – Maksoud and Abdle – Raoef (1998), Radostits *et al.* (2000), Mobark and El-Bealawy (2002) and Emam *et al.* (2005). The above mentioned observations were also supported by previous studies by Wang *et al.* (1985) who mentioned that the phosphorus deficiency leading to hypophosphataemia may play a part in haemoglobinurea by decreasing erythrocyte glycolysis. The increase in pulse rate might be attributed to the decrease blood volume following the intra vascular haemolysis and anaemia, while the accelerated respiratory rate also due to the anaemic hypoxia (Radostits *et al.*, 2000). Analysis of samples of barseem from newly reclaimed area and from old agricultural area, revealed that the concentrations of phosphorus, copper, molybdenum and calcium content were suggestive of primary phosphorus and secondary copper deficiency.

There would appear significant difference at ($P < 0.01$) in phosphorus content between both samples. This low levels of phosphorus in barseem from newly reclaimed area may be attributed to the low phosphorus content in the parent rock from which the soil are derived (Radostits *et al.*, 2000). Meanwhile no significant difference was detected (Table 5) in copper content between both diet samples.

Obtained values of copper content of barseem from newly reclaimed area, where a deficiency occurred, are generally considered adequate for cattle unless complicating factors cause secondary deficiency (Under wood and Suttle, 1999). In this study it appeared that copper deficiency is caused by the relatively high concentration of molybdenum in barseem of recently reclaimed area, which effectively reduce the availability of dietary copper in cattle. In this regard, molybdenum and sulphur react to form tetrathiomolybdates that then react with copper resulting in the formation of highly stable compounds that can not be digested and absorbed. Such correlation go in parallel with the results previously recorded by Ward (1978) and Allen and Gawthoren (1987). The fodders in particular barseem grown on newly

reclaimed area have high molybdenum content, the excess of this element reduces phosphorus content of the body by interfering with its absorption from the gastro - Intestinal tract and increasing phosphorus elimination through urine (Dhillon *et al.*, 1972). Analysis of blood in this study showed very highly significant decreases ($P < 0.001$) in TRBCs count, Hb concentration and PCV% values in cattle affected by PPHU indicating severe anaemia. This could be attributed to intra vascular haemolysis (Smith, 1990; Digraskar *et al.*, 1991 and Akhtar *et al.*, 2007) due to an impaired glycolytic pathway and depletion of ATP in erythrocytes which results from phosphorus deficiency. This ATP is essential for maintenance of shape and deformability of erythrocyte (Kaneko *et al.*, 1997). Subnormal concentration of ATP predisposes red blood cells to alter function and structure, a loss of normal formability, increased osmotic fragility and shortened life span ultimately leading to haemolysis (Wang *et al.*, 1985 and Ogawa *et al.*, 1989). El-Amrousi *et al.* (1977) attributed these change to destruction of erythrocyte with release of haemoglobin content causing decrease in haemoglobin concentration accompanied by oligocythemia. The higher MCV with lowered MCHC indicate that the anaemia is of macrocytic hypochromic type which may be due to malnutrition and malabsorption (Abdel-Maksoud and Abdel-raoef, 1998). The explanation of the anaemic condition in hypocupremic cattle based on the role of copper in the production of haemoglobin through the reutilization of iron liberated from normal breakdown of haemoglobin by the activation of ferroxidase enzyme, (Radostits *et al.*, 2000 and Shalaby, 2003). Concerning, the total leucocytic count in the present study showed very highly significant increases in PPHU cattle in comparison to control group. The increased TWBCs might be due to the endogenous release of corticosteroids. Increase stress due to PPHU (a metabolic disorder) is the source of the release of corticosteroids. These observation are in agreement with those mentioned by (Singari *et al.*, 1991; Abdel-Maksoud and Abdel-Roaef, 1998 and Akhtar *et al.*, 2007).

The blood hydrogen ion concentrations (pH) value showed non significant increase in PPHU cattle. These results similar to that obtained by Barzanji and Daniel (1987) who attributed the increase level of pH to the reduced oxygen tension as a sequel to the alteration in the acid base balance. The obtained results in this study revealed highly significant increase in P_{CO_2} and non significant increase in bicarbonate. While P_{O_2} showed very highly significant decrease in PPHU cattle. There is a type of anaemic anoxia which lead to decrease in oxygen level in the blood.

These results were supported by previous studies Barzanji and Daniel (1987) who mentioned that the up take of O₂ by the peripheral tissues was reduced during metabolic disorders. It was found also that the decreases in blood flow to tissues have been shown to cause increases in blood PCO₂ levels.

The highly significant decreases in serum total proteins, albumin and globulins levels in PPHU cattle in present study were nearly in accordance to the observed results of Abd-Allah and El-Fadali (1996); Abdel-Aal (1997) and Abdel-Maksoud and Abdel-Roaef (1998), who stated that the decreased values could be attributed to the loss of protein from the destructed RBCs. The case may be also due to the tubular degeneration in the kidney during haemoglobinurea or/poor nutritional status as represented by Kurundikney *et al.* (1981), and /or related to deprivation of the diet as reported by Abdel-Maksoud and Abdel-Roaef (1998) and Tawfik *et al.* (2004). Meanwhile Scott (1988) stated that the decreased levels of total proteins in PPHU might be attributed to the diet which was poor in proteins or due to the inhibition of the liver to synthesis more proteins. Mohga (2000) reported that the reduction of albumin concentration might be attributed to either a decreased food intake or albumin loss as a result of increased capillary permeability in copper deficient animals due to abnormalities in the blood vessel wall elastin and collagen structures. The highly significant decrease in the mean values of serum globulin level is similar to the opinion of Abdel-Aal (1997) and Tawfik *et al.* (2004) as they found no significant decrease in the serum globulins fractions in cattle in the green season.

The present investigation revealed that AST, ALT and ALP enzymes activity levels were highly significantly increased in cattle suffering from PPHU. The increases, in these enzymes, were previously reported by Wikes *et al.* (1992) and Tawfik *et al.* (2004), who recorded that the changes in the activities of the liver specific enzymes in combined phosphorus and copper deficiency in cattle may be due to alteration in the metabolic rate which accompanied that deficiency. Another explanation, for such increase in enzymes activities, was reported previously by Mohga (2000) and Shalaby (2003), who attributed to the hepatic dysfunction and skin lesions. Hypoglycaemia was noticed in this study in PPHU may be due to loss appetite or loss of blood through the urine and / or depraved metabolic processes which consequently reflected upon glucose metabolism and on its levels. Same results were previously reported by Mobark and El-Bealawy (2002); Tawfik *et al.* (2004) and Emam *et al.* (2005).

A high significant increase in blood urea was noticed in PPHU cattle when compared with control healthy ones. Such increases can be explained by the state of negative protein balance in the affected animals which may leads to catabolism of body protein, producing excess of urea (Coles, 1986). Or could be attributed to the endogenous release of corticosteroids, starvation and tubular epithelial necrosis (Kurundikney *et al.*, 1981 and Digraskar *et al.*, 1991). Additionally, dehydration usually occurs with PPHU which is a source of decreased renal perfusion, resulting in a reduced glomerular filtration rate and increased blood urea level Stogdale, (1981 and Latimar *et al.*, 2003). Alternatively, increased blood urea could be due to the failure of the urea recycling process through salivary glands and its non-utilisation by microbes in the rumen during digestive disorders. Most of the urea formed by the liver circulation in the circulatory system and remain unutilised (Singh *et al.*, 1989). In the present study, creatinine was increased in PPHU cattle when compared with control healthy once. This increased level may be attributed to the deficiency of inorganic phosphate which plays an important role in the intermediary metabolism of creatinine in the chemical reaction occurring in the muscle contraction (Radostits *et al.*, 2000). Both urea and creatinine levels were elevated and positively correlated to each other in PPHU affected cattle. Urea and creatinine are waste products that the kidneys normally filter from the blood and these are interrelated if the kidneys are not working properly (Kurundikney *et al.*, 1981; Digraskar *et al.*, 1991 and Mousa, 1998).

A high significant decrease in serum total cholesterol and total triglycerides levels was observed in PPHU when compared with control ones. The obtained results were similar to those previously reported by Tawfik *et al.* (2004) who found reduction in serum cholesterol level in copper deficient animals. These reductions of cholesterol and triglycerides may reject either reduced food intake or early hepatic damage (Uivund, 1990).

The inorganic phosphorus level in the serum showed very highly significant decrease in the PPHU cattle in this study. This result is in agreement with that reported by (Abdel-Maksoud and Abdel-Roaeef, 1998; Mobark and El-Bealawy, 2002; Emam *et al.*, 2005 and Akhtar *et al.*, 2007), who mentioned that the decreased phosphorus level may be due to long feeding on barseem with low level in phosphorus content. Also due to heavy drainage of phosphorus through milk particularly in high milk yielding animals (Bhikane *et al.*, 1995 and Ivanov *et al.*, 1995). Moreover, soils in the newly reclaimed area have high

molybdenum content. The fodders in particular barseem grown on such soils have high molybdenum content. The excess of this element reduces phosphorus content of the body by interfering with its absorption from the gastro-intestinal tract and increasing phosphorus elimination through urine (Dhillon *et al.*, 1972).

The non-significant changes in serum calcium levels in PPHU cattle in this study were in agreement with the opinion of Brain and Eric (1996) who reported that, the serum calcium levels were at the reference limits in hypophosphataemia. The obtained results revealed that the magnesium level was highly significantly decreased in cattle affected by PPHU. Present results were in agreement with those obtained by Mobark and El-Bealawy (2002) and Emam *et al.* (2005).

Serum electrolytes levels including sodium and potassium were highly significant decrease in cattle affected by PPHU. The obtained results were similar to those previously reported by El-Amrousi *et al.* (1977), who reported that the decreases may be due to loss of these electrolytes as a result of diarrhoea which was observed in same individuals of this group.

The non-significant change in serum zinc level in PPHU cattle was in agreement with the results, noticed on phosphate deficient buffaloes-cows (Ismaeil and Hussein, 1988).

Concerning the blood serum iron and copper levels the obtained results in PPHU cattle, suffering from primary phosphorus and secondary copper deficiencies behaved very highly significant decrease when compared with control. Nearly similar data were recorded by Georgievskij (1982) and proved that there are direct interactions between copper and iron in the formation of haemoglobin, copper is essential for iron metabolism. Therefor the serum iron level is decreased in copper deficient animals (Nabila, 1983). Also the decreased serum copper level might be due to the poor-nutrition, which accompanied by decreased appetite and gradual starvation of the diseased animals (Abdel-Maksoud and Abdel-Roaf, 1998).

The serum molybdenum levels were highly significantly increase in PPHU affected cattle suffering from combined phosphorus and copper deficiency if compared with clinically healthy ones. Similar results were previously obtained by Tawfik *et al.* (2004). Such elevation can be explained by the antagonist relationship between copper and molybdenum which was previously discussed, (Allen and Gawthoren, 1987 and Shalaby, 2003).

The result of urine analysis showed very highly significant increase in the concentrations of urine phosphorus and calcium in PPHU affected cattle. Similar observation was recorded by Singh *et al.* (1974) and Mousa (1998). The increased excretion of these substances through urine indicated renal damage (Teotia *et al.*, 1973). Depressed tubular reabsorption of phosphorus in case of renal damage was reported by Shtacher (1969).

The elevated protein concentration in urine of diseased cattle supports the involvement of kidney glomeruli. Proteinuria occurs as a result of injury of glomeruli (Kaneko, 1997). The high levels of urea and creatinine in the urine might be the result of kidney dysfunction. Estimation of creatinine is a valuable diagnostic test for haematouria (Singh *et al.*, 1974).

After treatment:

An observable advance in both clinical and laboratory results of cattle suffered from PPHU after correction of the diet system was achieved by avoiding the excessive feeding on barseem and providing the animals with balanced ration. Transfusion of large quantities of whole blood is the treatment for severely affected cows. The diseased cases were treated by sodium acid phosphate injection, plus to daily injection with 20 ml tonophosphan 20%. Also one dose of copper glycinate injection (120 mg available copper) halt haemolysis. This result agreed with the obtained by Mousa (1998); Radostits *et al.* (2000); Mobark and El-Bealawy (2002) and Emam *et al.* (2005). Who observed disappearance of clinical signs, normalization of macro and micro elements and improvement in haematological values after treatment with sodium acid phosphate and copper glycinate.

It could be concluded that PPHU noticed in cattle fed excessively on barseem from recently reclaimed area could be attributed to phosphorus deficiency which plays a key role in causing PPHU in cattle. Post parturient haemoglobinurea affected cattle showed hypocupraemia and reduce erythrocytic counts and this in turn reflects the disturbance in general healthy condition of the affected animals. So attention must be paid for careful early clinical and laboratory diagnosis of diseased animals followed by therapeutic plan with good management. Furthermore excessive feeding on barseem must be avoided and feeding balanced ration during the green season is also utmost.

REFERENCES

- Association of Official Analytical Chemists, A.O.A.C. (1975):* 12th Ed. Washington, DC.
- Abd El-Aal, A.A. (1997):* Seasonal variation electrophoretic pattern of serum protein of fertile and infertile buffaloes. M. Vet. Sc. Fac. Vet. Med. (Moshtohor) Zag. Univ.
- Abd-Allah, Amany, A.M. and El-Fadali, M. (1996):* Clinico-haematological changes in cattle fed clover and cotton seed meal. *Assuit Vet. Med. J.* 35(69): 123-131.
- Abdel-Maksoud, H. and Abdel-Raoef, Y.M. (1998):* Clinico-biochemical studies on blood of field cases of hypophosphataemic buffaloes. *Vet. Med. J. Giza* 46, 4 A: 427-441.
- Abdul-Samad, A. (1997):* Host and environmental factors associated with phosphorus deficiency haemoglobinuria in buffaloes, *Buffalo-Journal*, 13: 3, 385-395.
- Akhtar, M.Z.; Khan, A.; Khan, M.Z. and Muhammad, G. (2007):* Haemoglobin uria in buffalo. *Turk. J. Vet. Anim. Sci.*, 31 (2): 119-123.
- Allen, M.W. and Gawthoren, J.M.P. (1987):* Copper in animals and man. Vol. I, eds Howel. J. MCC.124. Boca Raton. FL. CRC Press.
- Barzanji, A.A.H. and Daniel, R.C.W. (1987):* Induced hypocalcaemia in ruminants, changes in some parameters of circulatory function. *Br. Vet. J.*, 144: 229-235.
- Bhikane, A.U.; Mirsalabat Ali; Narladkar, B.W.; Kawitkar, S.B. and Ali, M. (1995):* Post parturient haemoglobin urea in a crossbred cow and its treatment. *Indian Vet. J.* 72, 7: 734-736.
- Bowers, G.N. and Mc Comb, R.B. (1975):* Measurement of the total alkaline phosphatase activity in Human serum. *Clinical Chemistry* 21: 1988-1995.
- Brain, J.G. and Eric, P.S. (1996):* "Acute recumbancy and marginal phosphorus in dairy cattle" *J. A. V. M. A.*, 208, 5: 716-719.
- Chugh, S.K.; Mata, M.M. and Malik, K.S. (1996):* Epidemiological observations on post-parturient haemoglobinuria in buffaloes. *Indian J. Anim. Sci.*, 66: 1123-1125.
- Coles, F.H. (1986):* *Veterinary Clinical Pathology* 4th Ed. W.B. Saunders Company, Philadelphia, London and Toronto.
- Cowell, D.C. (1973):* *Medical laboratory technology* 30.133.

- Dhillon, K.S.; Sing, J. and Bajwa, R.S. (1972):* Treatment of haemoglobin uria due to molybdenum induced phosphorus deficiency in buffaloes. *J. Anim. Sci.*, 42: 996-998.
- Digraskar, S.B.; Singh, B. and Deshpande, B.B. (1991):* Epidomiology and clinico-pathology of haemoglobin urea in buffaloes (*Bubalus bubalis*). *Live Stock Advisor*, 16: 32-38.
- Doumos, B.I. (1974):* A buiret colorimetric method for determination of total proteins *Clin. Chem.*, (21) 1159-1166.
- Drupt, F. (1974):* Colorimetric method for determination of albumin. *Phar. Bio. Vol. (9). Tone VIII.* 777.
- El-Amrousi, S.M.; El-Allawy, T. and Moteilb, T. (1977):* Some studies on the causes of haematuria and haemoglobinurea in cattle and buffaloes in Assiut. *Assiut Vet. Med. J.* 4 (8): 164-176.
- Emam, E.E.; Shadia, A.R. and Alam, T.H. (2005):* Some trials for treatment of hypophosphatemia in buffaloes in Sharkia Governorate. *J. Egypt Vet. Med. Asso.* 65, 6: 105-118.
- Georgievskij, V.L. (1982):* Mineral composition of bodies and tissues of animals and physiological role micro elements. In "Mineral Nutrition of Animals". Freund Publishing House, London, Boston, Sydney, Toronto.
- Gindler, E. and Heath, D. (1971):* Determination of serum magnesium *Clin. Chem.* (17) 662.
- Gindler, E.M. and King, J.D. (1972):* Rappid colorimetric determination of calcium in biological fluids with methyl thymol blue. *Am. J. Cli. Path.* 58, 376-382.
- Hawk, P.B. (1965):* Hawk's Physiological Chemistry 4th Ed. Osier, B.L., MC. Grow-Hill, Inc. New York.
- Hussain, A.; Maqbool, C.A. and Khokhur, M.A. (1991):* Incidence and aetiology of parturient haemoglobin urea in buffaloes. *J. Anim. Hlth. Prod.*, 11:39-42.
- Ismaeil, M. and Hussein, F.M. (1988):* Studies on the relation ship between trace elements and some blood components and fertility Egyptian buffalo ". *Alex. J. Vet. Sci.* 4(1): 555-559.
- Ivanov, I.; Rajie, I. and Jovanovie, M.J. (1995):* Phosphorus content of diet for intensively kept cows during late pregnancy and lactation. *Vetrinarski Glasnik*, 49(5) 319-322.
- Jeffrey, L.; Arnold, M.D. and Facep, M. (2005):* Hypophosphatemia *J. Emerg. Med.* 37(4): 37: 39-45.
- Kaneko, J.J.; Harrey, B.D. and Bruss, M.L. (1997):* Clinical Biochemistry of Domestic Animals. 5thEd., London.

- Kurundikney, V.B.D.; Singh, B. and Ananiwar, L.G. (1981):* Biochemical and pathological changes in clinical cases of hypophosphatemia in buffaloes. *Indian J. Anim. Sc.* (51):35-38.
- Latimar, K.S.; Mahaffey, E.A. and Prasse, K.W. (2003):* Duncan and Praise's Veterinary Laboratory Medicine: Clinical Pathology. 4th Ed. Iowa State Press, Ames., Iowa, USA.
- Mgtowan, M.W.; Artiss, I.D.; Stand Bergh, D.R. and Zark, B. (1983):* "A peroxide-coupled method for colorimetric determination of serum triglycerides." *Clin. Chem.*, 29: 538.
- Mobark, M.G. and El-Bealawy, M.A. (2002):* Hypophosphatemic haemoglobin uria and gastro- intestinal nematode correlation in buffaloes and approach to treatment. *J. Egypt Vet. Med. Ass.* 62, 1:179-186.
- Mohga, S.Z. Abdel-Razik (2000):* An investigation on clinical and some biochemical alteration in unthriftiness sheep suffering from copper deficiency and therapy. *Zag. Vet. J.* 28, 3: 191-201.
- Morinal, L. and Prox, J. (1973):* New rapid procedure for serum phosphorus using O-Phenylene diamine as reductant. *Chin. Chem. Acta.*, 46: 113-117.
- Mousa, Sh.M. (1998):* Diagnosis of hypophosphatemia in buffaloes with special reference to the biochemical constituents of urine. *Zag. Vet. J.* 26, I: 63-68.
- Nabila, G. (1983):* Copper status and its blood level in sheep and cattle with special references to Egyptian water buffaloes. *Assiut Vet. Med. J.*, 10(20): 33-36.
- Nassif, N.M. (1995):* Nutritional problems affecting calcium, phosphorus and magnesium metabolism in Egyptian cattle and buffaloes. Ph.D. Thesis Fac. Vet. Med. Zag. Univ.
- NRC (1985):* Nutrient requirement of domestic animals. Nutrient requirements of cattle National Research Council, Washington.
- Ogawa, E.; Kobayashi, K. and Yoshiura, N. (1989):* Haemolytic anaemia and red blood cell metabolic disorder attributable to low phosphorus intake in cows. *Am. J. Vet. Res.* (5) 3: 388-392.
- Omran, H.; Abdel-Razik Waffa and El-Nagar, E. (1987):* Some clinical and biochemical studies on blood and urine of buffaloes in some diseased condition accompanied with haemoglobin uria. *Alex. Vet. Sci. J.* 3: 345-456.
- Patton, C.J. and Crouch, S.R. (1977):* Determination of serum urea. *Anal. Chem.*, 49: 464-469.

- Pirzada, W.H. and Hussain, S.Z. (1998):* Parturient haemoglobinuria in buffaloes. *Tropical Animal Health and Production*, 30, 4: 209-215.
- Radostits, O.M.; Gay, C.C.; Blood, D.C. and Hinchliff K.W. (2000):* Veterinary Medicine, 9th Ed. W.B. Saunders Company Ltd., London, New York, Philadelphia, San Francisco St Louis, Sydney, PP. 1366-1453.
- Reitman, S. and Frankel, S. (1957):* Determination of AST and ALT. *Amer. J. Clin.Path.*, 28: 56.
- Scott, D.W. (1988):* Large Animal Dermatology, W.B. Saunders Company, Harcour Brace, Jovanovich, Inc. Philadelphia, London, Toronto, Montreal, Sydney, Tokyo. Nutritional diseases, 358-373.
- Shalaby, H.A. (2003):* Studies on some haematological and biochemical changes associated with alopecia in calves. Ph. D. Vet. Thesis. Fac. Vet. Med. Zagazig Univ.
- Shtacher, G. (1969):* Selective renal involvement in the early development of hyper calcaemia and hypo-phosphataemia in x-z carcinoma bearing rabbits. *Cancer Res.*, 29(1): 512.
- Singari, N.A.; Bhardwaj, R.M.; Chugh, S.K. and Bhandwaj, S. (1991):* Status of erythrocytic glucose -6-phosphate dehydrogenase (G6PD) in phosphorus deficiency haemoglobin uria of buffaloes. *Indian Vet. J.* 68: 226-230.
- Singh, B.P.; Joshi, H.C. and Prasad, B. (1974):* Haematological and biochemical investigation in chronic bovine haematuria. *Indian Vet. J.*, 51:365-370.
- Singh, N.; Kumari, R. and Akbar, M.A. (1989):* Biochemical changes in blood metabolites in buffaloes with indigestion. *Indian Vet. J.* 66: 923-926.
- Smith, B.; Wood house, D.A. and Fraser, A.J. (1975):* The effect of parenteral copper on incidence of disease. Haematological changes and blood copper levels in a dairy herd with hypocuprosis. *N.Z Vet. J.*, 23: 109-112.
- Smith, B.P. (1990):* Internal Large Animals Medicine. 1st Ed. C.V. Mosby Co. Philadelphia, USA.
- Snedecor, G.W. and Cochran, W.G. (1982):* Statistical methods 7th ED., Iowa state. Ames, Loea, USA.
- Soulsby, E.J.L. (1986):* Helminths Arthropodies and Protozoa of Domesticated Animals. 1th Ed. Bailliere, Tindell, London.

- Stein, E.A. (1986):* In Text Book of Clinical Chemistry, Saunders, Philadelphia. P. 879-886.
- Stogdale, L. (1981):* Correlation of changes in blood chemistry with pathological changes in the animal's body. "Electrolytes, kidney function tests, serum enzyme and liver function tests". J. S. Afr. Vet. Assoc., 52:155-164.
- Tawfik, S.A.; El-Ramady, R.A. and Rawia, Kh.E. (2004):* Some Haematological and biochemical changes associated with deviated appetite in growing cattle. 7th Vet. Med. Zag. Conference. 770-784.
- Teotia, D.S.; Joshi, H.C. and Ray, S.N. (1973):* Studies on bovine haematuria. 2- Biochemical analysis of blood and urine of cattle suffering from haematuria. Indian J. Anim. Sci., 43(7):620-624.
- Thomas, L. (1992):* Lobar und Diagnose, 4th Ed, Medizinische Verlagsgesellschaft, Marburg.
- Trinder, P. (1969):* Determination of glucose in serum. Ann. Clin. Bio. (6)24-26.
- Uivund, M.J. (1990):* "Ovine white liver disease (OWLD): Serum copper and effects of copper and selenium supplementation. Acta Vet. Scand, 31:287-295.
- Under Wood, E.J. and Suttle, N.F. (1999):* The Mineral Nutrition of Livestock, 3rdEd. A.B. Publishing, New York, USA, PP. 520-887.
- Wang, X.L.; Gallagher, G.H.; Mc-Chre, T.J.; Reeve, V.E. and Canfield, P.J. (1985):* Bovine post parturient haemoglobin effect of inorganic phosphorus on red cell metabolism. Res. Vet. Sci. 39:333-339.
- Ward, G.M. (1978):* Molybdenum toxicity and hypocuprosis in ruminations. A review J. Anim. Sci., 46: 1078-1085.
- Wikse, S.E.; Herd, D.; Field, R. and Holland, P. (1992):* Diagnosis of copper deficiency in cattle. J. Am. Vet. Med. Assoc., 200: 1625-1629.