

STUDIES ON SOME PHENOLIC AND FLAVONOID COMPOUNDS OF RED SORGHUM BRAN AND THEIR ANTIBACTERIAL ACTIVITY

(Received:1.7.2007)

By

S. M. Morsy , M. S. Halaby* and R.N. Sandak

*Department of Crops Technology, Food Technology Research Institute, Agricultural Research Center, Giza ,Egypt. *Department of Nutrition & Food Science, Faculty of Home Economics, Helwan University,Cairo, Egypt.*

ABSTRACT

Flavonoid and phenolic compounds of red sorghum bran were investigated through ethanolic 70% extraction and subsequent extraction by diethyl ether and ethyl acetate. The diethyl ether extract contained three aglycons, *i.e* identified kaempferol, quercetin and apigenin. The ethyl acetate extract proved to contain luteolin, astragtein (Kaempferol-3-glucoside), rhoifolin (Apigenin-7-rutinoside), gallic acid and ellagic acid. The residual material after solvent fractionation contained vicenin II (Apigenin-6-8-di-C-glucoside). All isolated and purified compounds were identified using both physical and chemical methods and were further confirmed by the spectral measurements (U.V. Spectrum and H-NMR spectra). The antibacterial effects of the compounds and extracts were studied by using the agar diffusion technique on four gram positive bacteria namely *Staphylococcus aureus*, *Sarcina lutea*, *Bacillus pumilus* and *Bacillus subtilis* and two gram negative bacteria *Escherichia coli* and *Broodetella brochiseptica*.

Key words: antioxidants, antibacterial effects, bran ,flavonoid, phenolic compounds ,red sorghum.

1. INTRODUCTION

Sorghum bicolor is the fifth most important cereal crop after wheat, rice, maize and barley in terms of production (FAO, 2005). The total world annual sorghum production is over 60 million tons from a cultivated area of 46 millions ha. Moreover, some red sorghum varieties have higher antioxidant activities than the most important sources of natural antioxidants. Oxidation products of peroxidase and polyphenol oxidase (benzoquinones and polymeric compounds) affect food quality (Dicko *et al.*, 2006).

Phenolic compounds are commonly found in both edible and non-edible plants and they have been reported to have multiple biological effects including antioxidant activity. Crude extracts of vegetables, herbs, fruits, cereals and other plant materials rich in phenolics are increasingly of interest in the food industry because they retard oxidative degradation of lipids and thereby improve the quality and nutritional value of food. The importance of the antioxidant constituents of plant materials in the maintenance of health and protection from coronary heart disease and cancer is also raising interest among scientists and food manufactures. In fact, consumer as the trend of the future is moving towards functional food with

specific health effects (Kahkonen *et al.*, 1999 and Sandak and El-Hadidy 2004).

Phenolic sorghum phytochemicals are important for human nutrition (Awika *et al.*, 2004). Phenolic compounds are the most widely distributed secondary metabolites, ubiquitously present in the plant kingdom. Among cereals sorghum had the highest content of phenolic compounds reaching up to 6% (W/W) in some varieties (Awika and Rooney, 2004 and Dicko *et al.*, 2005). While, the amount of phenolic compounds which present in all sorghums influence, its genotypes and environmental conditions. The main classes of phenolic compounds are simple phenols hydroxybenzoic acids, hydroxycinnamic acids, flavonoids (flavanols, flavones, flavanones, isoflavones and anthocyanins), chalcones, lignain, hydroxycoumarins and polyflavans (Chung *et al.*, 1998 and Krueger *et al.*, 2003).

These compounds are soluble in water or organic solvents. Sorghum does not contain tannic acid and hydrolysable tannins (Waniska, 2000 and Awika *et al.*, 2004). Phenolics play an important role in plant metabolism, but also, protect the plant against stresses. For instance, it has been recently shown that flavonoids, such as catechin, regulate the auxin transport in plants and

therefore, play an important role in plant development (Brown *et al.*, 2001).

Flavonoids constitute the largest class of phenolic compounds with more than 3000 structures, possessing in common a flavylum unit (C₆-C₃-C₆) (Lacobucci and Sweeny, 1983). Sorghum contains flavonoids such as flavanols (flavan-3-ols and flavan-4-ols), flavanones, flavones and anthocyanins (Haslam, 1998 and Awika and Ronny, 2004). The flavan-4-ols apiforol (pro-apigeninidin or leuco-apigeninidin) and tuteoforol (proluteolinidin or leucoluteolinidin) are abundant in sorghum and precursors of apigeninidin and luteolinidin, respectively (Ferreira and Slade, 2002, and Hagerman, 2005).

Flavonoids are widely distributed in the plant kingdom. They are mainly produced as a pigment of many shades, and play an important role in normal growth, development, and defense of plants. At the biochemical level, flavonoids act as enzyme inhibitors, provide defense against ultraviolet radiation, act as chelating agents for metals, and as reducing agents (Harris, 1992 and Rice-Evans *et al.*, 1995). Flavonoids have been reported to exhibit a wide range of biological effects, such as antibacterial, antiviral, anti-inflammatory, anti-allergic and anticancer. They are known to inhibit lipid peroxidation and platelet aggregation and to affect capillary permeability and fragility. (Chu *et al.*, 2000 and Knekt *et al.*, 2002).

Mechanisms by which flavonoids have been proposed to provide health benefits in addition to being direct chemical protectants and involve modulator effects on a variety of metabolic and signaling enzymes. Flavonoids have been shown to block the angiotensin-converting enzyme that raises blood pressure; they inhibit cyclooxygenase, which forms prostaglandin and they block enzyme that produce estrogen. The implications of these *in vitro* inhibitory actions are that certain flavonoids could prevent platelet aggregation, reducing heart disease and thrombosis also, inhibit estrogen synthesis, which binds estrogen to receptors in several tissues, thus decreasing the risk of estrogen-related cancers (Dillard and German, 2000). Phenolic acids in sorghum are found in free or bound as esters, and are concentrated in the outer layers of the grain (Waniska, 2000; Awika *et al.*, 2004 and Dicko *et al.*, 2006).

Sorghum with black pericarp or purple/red plants has higher levels of phenolic compounds (flavan-4-ols and anthocyanins) than the other varieties (Dykes *et al.*, 2005). The biological activities of flavonoids are actions against free

radicals, free radical mediated cellular signaling, inflammation, allergies, platelet aggregation, microbes, ulcers, viruses and tumors and hepatotoxins (Kinsella *et al.*, 1993).

The retardation of autoxidation is therefore a key to product quality, because most consumers prefer natural food additives to synthetic ones. Natural antioxidants are of increasing importance (Krings and Berger, 2001; and Negi and Jayaprakasha, 2003). Although significant advances have been made in food preservation, spoilage and pathogenic bacterial growth during food preparation, storage and distribution remain a serious problem. Most approaches to food preservations employ physical and/or chemical methods that reduce pre-and post-treatment bacterial contamination of foods, and include additives, packaging or storage conditions that retard or inhibit the growth of borne bacterial pathogens, and/or reduce processing requirements for their elimination in foods are desired (Bowles and Jay 1993; and Leuschner and Ielsch 2003).

Antioxidants are major ingredients, play an important role in manufacturing, packaging and storage of lipid containing foods. Synthetic antioxidants are usually used in food industry to reduce deterioration and rancidity of oils and fats. There has been growing concern regarding the possible activity of such synthetic antioxidants to cause liver damage. Therefore development of safer, inexpensive natural antioxidants is essentially (Yen and Due, 1993).

Furthermore, most consumers prefer natural food additives to synthetic ones; since natural antioxidants are of increasing importance. Recent work is beginning to shed light on the relation of flavonoids and other dietary phenolic constituents to these protective effects.

This investigation aimed to isolate and identify flavonoid and phenolic compounds of red sorghum bran as a waste and of a low cost. In addition the effect of the extracted flavonoid and phenolic compounds on six different bacteria strains was tested.

2. MATERIALS AND METHODS

2.1 Materials

Red sorghum (*Sorghum bicolor* L.) cultivar Assiut 14 was obtained from Agricultural Research Center, Giza - Egypt. Sorghum grains were finely ground then passed through 40-mesh sieve to separate the bran. Authentic phenol and flavonoids compounds were purchased from Sigma Co. and Deisenhofen, Germany.

2.2. Methods

2.2.1. Extraction, isolation and purification




Fig. (1): $^1\text{H-NMR}$ Spectroscopy of flavonoid and phenolic compounds from red sorghum bran extracts.

Air-dried bran (500g) was finely powdered and extracted with petroleum ether (40 – 60° C) to remove fats and resinous materials. The residue was exhaustively extracted with 2 liters 70% ethanol by heating on a boiling water bath for 6 hours. Extraction was repeated until a color extract then the extracts were combined and concentrated to obtain aqueous ethanolic extract. The aqueous ethanolic extract was subjected to fractional extraction using diethyl ether followed by ethyl acetate according to Mabry *et al.* (1970).

Two-dimensional paper chromatography of diethyl ether, ethyl acetate extracts and the aqueous solution of the residual material using the solvent system butanol: acetic acid: water (4:1:5) and acetic acid 15% were performed, respectively, revealed the presence of many compounds of flavonoid and phenolic nature.

The compounds were detected under U.V. light and the major compounds were eluted with 70% ethyl alcohol. Each of the isolated compounds and authentic samples phenolic and flavonoid were subjected to one-dimensional Watman 3MM papers chromatographic technique using two different solvents (acetic acid 15% and butanol: acetic acid: water, 4:1:5) for elution. The different spots (major flavonoid and phenolic compounds and authentic samples) were located by color reaction and R_f values under U.V. lights with and without the presence of NH_3 fumes according to Markham and Mabry (1968). These compounds were further purified over Sephadex LH 20 column prior to physical and chemical analysis.

2.2.2. Physico-chemical analysis

Complete acid hydrolysis was carried out by HCl 2N (Geissman, 1962). The controlled acid hydrolysis by HCL 0.1N (Seshadry, 1962) of the flavonoid glycosides under investigation yielded the sugar residues and the aglycones. Each compound was separately dissolved in pure methanol and examined for U.V. Spectrum (Shimadzu U.V. Visible. Recording Spectrophotometer Model, U.V.240) in the presence of diagnostic reagents according to Mabry *et al.* (1970).

1H -NMR apparatus examined all flavonoid and phenolic compounds separated from red sorghum bran. The 1H -NMR spectra of the trimethylsilyl ethers of all compounds were recorded in CDCL₃ at 90 MHz and reported at δ values (ppm) relative to TMS as an internal standard on a Bruker WM90 apparatus.

2.2.3. Antimicrobial activity of the flavonoid and phenolic extracts

The antibacterial effect of the extracts and some of the isolated compounds was studied using the nutrient agar (peptone 0.5%, beef extract 0.3%, Difco agar 1.5% and pH 7.0) according to A.P.H.A. (1971) on six different bacteria strains. Two gram- negative bacteria namely *Escherichia coli* and *Broodetella brochiseptica*, and four gram- positive bacteria namely *Bacillus pumilus*, *Bacillus subtilis*, *Staphylococcus aureus* and *Sarcina lutea*. The antibacterial activity was determined by the paper disc plate method described by Jacobes and Gerstin (1960). Whatman 3MM filter paper discs of 0.6mm diameter were placed in petri dishes at the rate of one disc/ plate. Each powder extracts were dissolved in 0.7 ml absolute ethanol to give 100 ppm of the crude compound. Each filter paper disc was moistened with exactly 0.2ml of tested solvent extract in two concentrations. The plates were incubated at 37°C for 24h. and the antibacterial activity was measured as growth zone of inhibition of microorganisms. All tests were run in triplicate for each sample and the mean of inhibition zones were given to asses the activity.

3. RESULTS AND DISCUSSION

3.1. Isolation and purification of flavonoids from sorghum bran

The present study deals with the polyphenolic compounds of red sorghum bran. Whereby, three flavonoid glycosides, four flavonoid aglycons and two phenolic acids were isolated from 70% ethanolic extract. From the diethyl ether extract three flavonoid compounds were isolated using preparative paper chromatography elution in 15% acetic acid and identified as kaempferol, quercetin and apigenine. Whilst, luteolin, astraglain (kaempferol-3-glucoside), rhoifolin (apigenin-7-rutinoside), gallic acid and ellagic acid were isolated from ethyl acetate extract by preparative paper chromatography using water for elution.

The residual material after solvent fraction was dissolved in water and subjected to preparative paper chromatography using butanol: acetic acid: water (4:1:5) as the solvent system for elution which revealed the presence of one flavonoid glycosides namely vicenin II (apigenin-6-8-di-C-glucoside). The structure of the isolated compounds was identified by physico-chemical analysis. Color reaction and R_f values data are outlined in Table (1).

Table (1): R_f values and color reaction of the compounds isolated from red sorghum bran.

Compounds	R _f Values			Color reactions	
	BAW	AcoH15%	H ₂ O	U.V.light	U.V NH ₃
Diethyl ether extract					
Kaempferol	83	1	19	Yellow	Bright yellow
Quercetin	64	3	17	Yellow	Bright yellow
Apigenine	87	57	-	Purple	Yellow
Ethyl acetate extract					
Luteolin	78	66	8	Dark brown	Yellow
Astraglein	70	43	13	Brown	Yellow
Rhoifolin	41	39	9	Dark	Faint yellow
Gallic acid	78	56	58	Violet blue	Deep violet blue
Ellagic acid	38	2	-	Dark	Yellow
Residual material	31	80	57	Purple	Faint yellow
Vicenin II					

BAW: Butanol:acetic acid:water (4:1:5). AcoH: Acetic acid
U.V.NH₃: U.V.light with ammonia fumes.

Table (2): U.V. spectral data (λ max. nm) of the compounds isolated from red sorghum bran extracts.

Compounds	MeoH (a)	(a)+ NaOMe (b)	(a)+ AlCl ₃ (c)	(c)+ Hcl (d)	(a)+ NaOAL (e)	(e)+ H ₃ BO ₃ (f)
Diethyl ether extract						
Kaempferol	254, 68,322,365	275,320,412	262, 270, 352, 426	260, 271, 350, 426	275, 300, 285	269, 295, 320, 370
Quercetin	265, 268, 374	249, 424	373, 305, 335, 434	266, 303, 350, 414	264, 325, 390	362, 302, 386
Apigenin	271, 336	429, 271 304, 389	268, 277 301, 352	279, 300 344, 378	261, 271 350, 392	269, 341
Ethyl acetate extract						
Luteolin	267, 296, 336	275,324,392	276, 301, 348, 384	276, 299, 340, 381	274, 301, 376	268, 301, 338
Astraglein (Kaempferol-3- glucoside)	267, 295, 352 267, 330	273,328,405 245, 273	274, 300 348, 389 275, 298	274, 300 345, 399 276, 297	273, 300 370 255, 267	267, 295 353
Rhoifolin (Apigenin-7- rutinoside)	272 255	350, 292, 245, 277	344, 382 277 247, 271	345, 380 _____ _____	337, 390 _____ 245, 277	267, 334 _____ _____
Gallic acid						
Ellagic acid						
Residual material	274,311	383,331	281,306	280,304,	383,308	276,284
Vicenin II	333	398	352, 390	346, 386	334, 395	323, 351
Apigenin-6-8-C- glucoside						

MeoH: Methanol
NaOMe: Sodium methoxide
AlCl₃: Aluminium chloride
Hcl: Hydrochloric acid
NaOAC: Sodium acetate
H₃BO₃: Boric acid.

Table (3): ¹HNMR data of the isolated compounds from red sorghum bran extracts.

Compounds	Aglycon and sugar moiety δ (ppm)
Diethyl ether extract Kaempferol	7.88(d,j=8Hz,H-2` and H-6`), 6.77(d,j=8.5 Hz,H-3` and H-5`), 6.41(d,j= 2.5Hz, H-8), 6.12(d,j=2.5Hz, H-6).
Quercetin	7.64(9dd,j=2.5Hz and j= 8.5Hz, H-6`), 7.56(d,j=2 Hz, H-2`),6.84(d,j=8.5Hz,H-5`),6.3(d,j=2Hz,H-8),6.2(d,j=2.5Hz,H-6).
Apigenin	7.41(dd,j =2.5Hz and j=9Hz,H-6),7.3(d,j=2 Hz,Hh-2`)6.9(d,j=8Hz, H-5`),6.78(9d,j=2.5Hz,H-8)0, 6.43(S,H-3),6.42(d,j=2.5Hz,H-6).
Ethyl acetate extract Luteolin	6.3(d,j=2.5Hz,H-6),6.44(d,j=2.5Hz,H-8),6.55(S,H-3),6.93(d,j=8Hz,H-5`),7.35(d,j=2Hz, H-2`), 7.44(dd,j=2.5Hz and j =9 Hz, H-6`).
Astragalin (Kaempferol-3-glucoside)	7.98(d,j=8 Hz,H-2` and H-6`), 6.87(d,j= 8.5Hz, H-3` and H-5`), 6.23(d,j=2.5Hz,H-8), 6.23(d,j = 2.5Hz, H-6). Sugar moiety:5.8(d,j=7.5Hz,H-1``)3.0-3.98(m of the six glucose protons).
Rhoifolin Apigenin-7-rutinoside	6.42(9d,j=2.5Hz,H-6),6.81(9d,j=2.5Hz,H-8),6.88(S,3-H),6.94 (9d,j=8Hz, 3-H and 5`-H), 7.41(d,j=8Hz,2`-H and 6`H) sugar moiety:5.0(d,j=7.5Hz, H-1`` of the glucose),4.3(d,j=2Hz,H-1`` of the rhamnose)0.86(d,j= 6.5Hz, methyl group of rhamnose)3.53-3.90(m of the rutinose protons).
Gallic acid	6.98 (S,H-2 and 6-H)
Ellagic acid	7.5 (S, H-2 and H-9).
Residual material	7.86(2H,d,J=9Hz,H-2`,H-6`),(7.04(2H,d,J=9Hz,H-3`, H-5`),6.4(H-1`,5,H-3`)
Vicenin II	sugar moiety:4.7(1-H,d,J=10Hz,H-1``), 4.6(1H,d,J=10Hz,H-1``)4.2-3.0(m,sugar protons).

Table (4): Anti-bacterial effects on flavonoids and phenolic compounds.

Compounds	Gram-negative bacteria		Gram-positive bacteria			
	Es.	Br.	St.	Sa.	Ba.p.	Ba.s.
Diethyl ether extract	++	-	+	++	+	+
Ethyl acetate extract	++	++	+++	+++	+++	++
Residual material	+++	++	+	++	++	++
Kaempferol	++++	++++	+++	++++	+++	++++
Quercetin	++	++	++	+	++	++
Apigenin	++	+	+++	++	+	++
Luteolin	++	++	++	+	++	+
Astragalin	+	+	+	-	++	+
Rhoifolin	+	+	+	-	+	++
Gallic acid	++++	+++	+++	++++	++++	+++
Ellagic acid	+++	++	+	+++	+++	++
Vicenin II	++	+	++	++	+	+++

Bacteria: Zone of inhibition (mm):

Es: Escherichia Coli = (+) from 11-13 mm.

Br: Brodetella broshiseptica = (++) from 14-16 mm.

St: Staphylococcus aureus = (+++) from 17-19.mm.

Sa: Sarcina lutea = (++++) from 20-24 mm.

Bap: Bacillus pumilus.

BaS: Bacillus subtilus.

3.2. Identification of flavonoids

U.V. spectroscopy is one of the most descriptive and clearly attained forms of data and is very important in the study of flavonoids.

Flavonoids exhibit two major absorption maximum with two distinctive bands at 240 - 285 nm (Band II) and at the range 300 - 400 nm (Band I). The absorption band II may be due to the A-ring benzyl system, while band I is originated from the B-ring cinnamoyl system (Harborne *et al.*, 1975; Haslam, 1998 and Krueger *et al.*, 2003). Data of U.V. Spectral analysis in methanol and by diagnostic reagents are recorded in Table (2), for further identification co-chromatography was carried out using authentic samples.

Complete controlled acid hydrolysis and ¹H-NMR spectroscopy are shown in Table (3) and Fig. (1). Modern physical tools such as proton or nuclear magnetic resonance spectroscopy (PMR) or (NMR) proved to be valuable tools for the elucidation of the structure of flavonoidies and more complex flavonoid derivatives for measuring the magnetic moments of the hydrogen atoms which are attached to different groups in such flavonoid compounds. By these means the NMR spectra provide the number of hydrogen atoms in different situation (Harborne *et al.*, 1975; Brown *et al.*, 2001 and Hagerman, 2005).

3.3. Antibacterial effect of flavonoid and phenolic compounds

The study of antibacterial effect of diethyl ether, ethyl acetate, the aqueous solution of residual material and some of the isolated compounds are reported in Table (4).

The results show that the diethyl ether extract contained flavonoids *i.e.* kaempferol, quercetin and apigenin and had more inhibitory effect on bacteria *Escherichia coli*, *Brodetella broshiseptica*, *Staphylococcus aureus*, *Sarcina lutea*, *Bacillus pumilus* and *Bacillus subtilus* (Harborne and Baxter, 1993 and Lin *et al.*, 1997). Whereas, ethyl acetate extract contained luteolin, and had a role of antibacterial. Taj and Nagarajan (1996) reported that luteolin is anti-mutagenic and had anti-inflammatory and antibacterial activities. Gallic and ellagic acids are phenolic compounds and have antimicrobial function (Kubo *et al.*, 1993 and Chu *et al.*, 2000).

From the above results it could be concluded that the extracted compounds (diethyl ether and ethyl acetate extract) from red sorghum bran contained flavonoid (*i.e.* *Escherichia coli*, *Brodetella broshiseptica*, *Staphylococcus aureus*, *Sarcina lutea* *Bacillus pumilus* and *Bacillus subtilus*) and phenolic compounds that had antibacterial activity .

These compounds are currently gaining attention as antibacterial. More research is needed to clarify that the sorghum bran could be used as a wide source of biochemical diversity, colorants and for various industrial utilizations.

4. REFERENCES

- A.P.H.A. (1971). American Public Health Association. Recommended Methods for the Microbiological Examination of Food. Amer.Public Health Assoc. Inc. New York.
- Awika J.M. and Rooney L.W. (2004). Sorghum phytochemicals and their potential aspects on human health. *Phytochemistry*.65:1199-1221.
- Awika J.M., Rooney L.W. and Waniska R.D. (2004). Properties of 3-deoxyanthocyanins from sorghum. *J. Agric. Food. Chem.* 52:4388-4394.
- Bowles B L. and Jay J.M. (1993). The effect of phenylglyoxal on *Clostridium sporogenes* . *Food microbial.*, 10:113-121.
- Bowles B.L and Miller A.J. (1993). Antimicrobial properties of aromatic and aliphatic aldehydes. *J. Food protect.*,56:788-794
- Brown D., Rashotte A., Murphy A., dormanly J., Tague B., Peer W. and Muday G. (2001). Flavonoids act as negative regulators of auxin transport *in vivo* in *Arabidopsis*. *Plant Physiol.* 126:524-535.
- Chung T., Wei C., Huang Y. and Lin Y. (1998). Tannins and human health: A Review. *Crit.Rev.Food Sci. Nutr.*38:421-464.
- Chu Y.H., Chang C.L. and Hsu H.F. (2000). Flavonoids content of several vegetables and their antioxidant activity. *Jj. Sci. Food Agric.*,80:561-566.
- Dicko M., Gruppen H., Traore A., Van-Berkel W. and Voragen A. (2005). Evaluation of the effect of germination on control of phenolic compounds and antioxidant activities in sorghum varieties. *J.Agric.Food Chem.* 53:2581-2588.
- Dicko M., Gruppen H., Traore A., Voragen A. and Van-Barkel W. (2006). Phenolic compounds and related enzymes as determinants of sorghum for food use. *Biotechnol.Mol.Biol.Rev.*1:21-38.
- Dillard C.J. and German J.B. (2000). Phytochemicals: nutraceuticals and human health. *J.Sci. Food Agric.*,80:1744-1756
- Dykes L., Rooney L., Waniska R. and Rooney W. (2005). Phenolic compounds and antioxidant activity of sorghum grains of varying genotypes. *J.Agric.Food Chem.* 53:6813-6818.

- F.A.O. (2005). FAOSTAT Agricultural database 2005.
- Ferreira D. and Slade D. (2002). Oligomeric proanthocyanidins: naturally occurring O-heterocycles. *Nat. Prod. Rep.* 19:517-541.
- Food and Agriculture Organisation (F.A.O.) (2005). FAOSTAT Agricultural database 2005.
- Geissman T.A. (1962). *The Chemistry of Flavonoid Compounds*. Pergamon Press, London, New York.
- Hagerman A.E. (2005). *Tannin Chemistry*; Hagerman, A. E. Ed., Department of Chemistry and Biochemistry. Miami University Oxford, OH 45056 USA. Consulted on 10-09-2005.
- Harborne J. and Baxter H. (1993). *Phytochemical Dictionary: A Handbook of Bioactive Compounds from Plants*, Taylor and Francis, London, UK.
- Harborne J., Mabry T. and Mabry H. (1975). *The Flavonoids*. Chapman and Hall LTP, London.
- Harris F.D. (1992). Regulation of antioxidant enzymes. *FfASABJ.*, 6 : 2675-2683.
- Haslam E. (1998). *Practical Polyphenolics: from Structure to Molecular Recognition and Physiological Action* Haslam, E. Ed, Cambridge Univ. Press. Cambridge. UK.
- Jacobson M.B. and Gerstein M.J. (1960). "Handbook of Microbiology". D. Van Nostrand Company. Inc New York p.193-207.
- Kahkonen M., Hopia A., Vourela H., Rauha J., Pihlaga K, Kujala T. and Heinonen M. (1999). Antioxidant activity of plant extracts containing phenolic compounds. *J. Agric. Food Chem.*, 47:3954-3962.
- Kinsella J., Frankel E., German B. and Kanner J. (1993). Possible mechanisms for protective role of antioxidants in wine and plant foods. *Food Technol.* 47: 85-89.
- Knekt P., Kumpulainen J., Jarvinen R., Rissanen H., Heliovaara M., Reunanen A., Hokininen T. and Aromaa A. (2002). Flavonoid intake and risk of chronic diseases. *Am. J. Clin. Nutr.*, 76: 560-568.
- Krings U. and Berger R. (2001) Antioxidant activity of some roasted foods. *Food Chem.*, 72: 223-229.
- Krueger C.G, Vestling M.A. and Reed J.D. (2003). Matrix-assisted laser desorption/ionization time-of-flight mass spectrometry of heteropolyflavan-30ols and glucosylated heteropolyflavans in sorghum (*Sorghum bicolor* L) Moench. *J. Agric. Food. Chem.* 51: 538-543.
- Kubo I., Murio H. and Himegima M. (1993). Antibacterial activity against *Streptococcus mutans* of meta flavor components. *J. Agric. Food. Chem.*, 41: 107-111.
- Lacobucci G.A. and Sweeny J.G. (1983). The chemistry of anthocyanins, anthocyanidins and related flavylum salts. *Tetrahedron*, 39:3005-3015.
- Leuschner R. and Ielsch V. (2003). Antimicrobial effect of garlic, clove and red hot chilli on *Listeria monocytogenes* in broth model system and soft cheese. *Int. J. Food Sci. Nutr.* 54: 127-133.
- Lin J.K., Chen Y.C., Huang Y.T. and Lin-Shiau S.Y. (1997). Suppression of protein Kinase C and nuclear oncogene expression as possible molecular mechanisms of cancer chemoprevention by apigenin and cumarin. *J. Cell. Biochem. Suppl.* 28 (29): 39-48.
- Mabry T.J., Markham K.R. and Thomas M.B. (1970). *The Systematic Identification of Flavonoids*, Springer Verlag, New York, Heidelberg, Berlin.
- Markham K.R. and Mabry T.J. (1968). The identification of twenty three 5-deoxy and ten 5-hydroxy flavonoid from *Baptisa lecontii*. *Phytochem.* 7, 1197.
- Negi P.S. and Jayaprakasha G.K. (2003). Antioxidant and antimicrobial activities of *Punica granatum* peel extracts. *J. Food Sci.*, 68 (4); 1473 - 1477.
- Rice-Evans C., Miller J., Bolwell P., Bramelley P. and Pridham L. (1995). The relative antioxidant activities of plant derived polyphenolic flavonoids. *Free Rad. Res.*, 22:375-383.
- Sandak R.N. and El-Hadidy E.M. (2004). Utilization of natural antioxidants and their effects on human health. *Egypt, J. Agric. Res.*, 5 (1): 47-61.
- Seshadry T.R. (1962). *The Chemistry of Flavonoid Compounds*. Oxford, Pergamon Press, p159.
- Taj, S. and Nagarajan, B. (1996). Inhibition by quercetin and luteolin of chromosomal alteration induced by salted, deep-fried fish and mutton in rats. *Mutat. Res.* 369: 1-(2) 97-106.
- Waniska R.D. (2000). Technical and institutional Options for Sorghum Grain Mold Management: Proceeding of an International Consultation. Chandrasheker, A, Bandyopadhyay, R, and Hall, A.J. (Eds) ICRISAT, Patancheru, India. P72-106.
- Yen G.C. and Due P.D. (1993). Antioxidative properties of methanolic extracts from peanut hulls. *J. AOCS*, 70 (4) 383-386.

دراسات على بعض المركبات الفينولية والفلافونيدية لقشور السورجم الاحمر
ونشاطها المضاد للبكتيريا

سمية محمد مرسى ، منى سامى حلى* و رأفت نجيب سندق

معهد بحوث تكنولوجيا الاغذية - مركز البحوث الزراعية - جيزة - مصر
* قسم التغذية وعلوم الأطعمة-كلية الاقتصاد المنزلى-جامعة حلوان-القاهرة - مصر

ملخص

تم التعرف على المركبات الفينولية والفلافونيدية المفصولة من قشور السورجم الحمراء من خلال الاستخلاص بواسطة كحول الايثانول 70%. تمت التجزئة باستخدام الايثر ثنائى الايثانول وخلات الايثانول. وقد وجد ان مستخلص الايثر ثنائى الايثانول يحتوى على ثلاثة مركبات فلافونيدية ، وقد تم التعرف عليها وهى مركبات الكامفيرول والكيورستين والابجيينين. أما مستخلص خللات الايثانول فانه يحتوى على ليتولين والاستراجلين (كامفيرول-3-جلوكوزيد) والرفولين (ابجيين-7-ريتنوزيد) واثنين من الاحماض الفينولية وهم مركبات الجالليك والابلاجيك اما المتبقى المائى فانه يحتوى على مركب الفيسيتين II (ابجيين-6-8ثنائى الجليكوزيد). تم تنقية جميع المركبات المفصولة والتعرف على التركيب الكيميائى لها باستخدام طرق الفصل الكروماتوجرافى الورقى والتحليل الكيميائى والطبيعى والطيفى (الاشعة فوق البنفسجية U.V والرنين المغناطيسى البروتونى $^1\text{H-NMR}$). وقد درس التأثير البكتيرى لكل من المركبات والمستخلصات باستخدام ستة انواع من البكتيريا منهم اربعة انواع موجبة جرام واثنين بكتيريا سالبة جرام.

المجلة العلمية لكلية الزراعة - جامعة القاهرة - المجلد (59) العدد الأول (يناير 2008): 48-56.