

## EFFECT OF CHEESE WHEY AND POLY-ETHYLENE GLYCOL ON PERFORMANCE OF GROWING LAMBS FED PEANUT HULLS ENSILED WITH LIME AND UREA

Y.I. EL-Talty<sup>1</sup>, M.S. Farghaly<sup>1</sup> and R.S. Mikail<sup>2</sup>

*1- Animal Production Department, Faculty of Agriculture, Cairo University, Giza, Egypt, 2-Animal Production Research Institute, Agriculture Research Center, Ministry of Agriculture, , Giza, Egypt*

### SUMMARY

*This study included two experiments, the first was carried out to study the effect of treating peanut hulls chemically by ensiling with 5% lime + 5% urea on its chemical composition and nutritive value by using adult crossbred (Romanov × Rahmani) rams. The second was carried out to study the effect of adding fresh sweet liquid cheese whey (FSLCW) or/and Poly-Ethylene Glycol (PEG) on the performance and feed efficiency of crossbred (Romanov × Rahmani) growing lambs fed total mixed rations (TMR) consisting of 40% roughage and 60% concentrate mixture for 70 days. Twenty male lambs with an average body weight 28.85kg and 6 months old were divided into four equal groups and randomly assigned to fed one of the following experimental rations :*

*(T<sub>1</sub>): control ration: contains wheat bran 15%, barely 29%, soya-bean meal 8.5%, clover hay (Trifolium alexandrinum) 40%, sugar cane-molasses 4%, lime stone 2%, common salt 1% and vits. & mins. mixture 0.5% .*

*(T<sub>2</sub>): T<sub>1</sub> with replacing all of the clover hay by treated peanut hulls .*

*(T<sub>3</sub>): T<sub>2</sub> plus FSLCW (free choice) .*

*(T<sub>4</sub>): T<sub>2</sub> plus FSLCW (free choice) plus 3 g PEG / head / day.*

*The ingested daily of the cheese whey expressed as TDN was replaced with a part of T<sub>2</sub> having the same value of TDN. The experimental animals were given their requirements of energy and protein to cover their average daily gain (NRC,1985). At the end of feeding trial four digestion , trials were conducted . Some rumen liquor and some blood parameters were also studied.*

*The results of the first experiment showed that total phenols and total tannins nearly disappeared in treated peanut hulls . In addition. digestibilities of CP, CF, NFE and nutritive values as TDN and DCP decreased from 36.9%, 18.55 %, 70.36 %, 31.47 % and 2.51 % to 6.31%, 10.17· %, 44.04 %,16.86 % and 0.63%, respectively. In the second experiment, there were:*

- 1) Significant increase in digestibilities of CP from 60.53 % to 75.03% vs. 70.60 % and for CF from 14.61 % to 35.77 % vs. 49.10 % in T<sub>2</sub> and T<sub>4</sub> vs. T<sub>1</sub> respectively.*
- 2) Significant increase in the feeding values in terms of TDN and DCP from 52.19 % to 63.29 % vs. 61.18 % and from 10.07 % to 12.32 % vs. 11.93 % in T<sub>2</sub> and T<sub>4</sub> vs. T<sub>1</sub> , respectively.*
- 3) Significant increase in the average daily weight gain from 97 to 197 vs. 151 g / day in T<sub>2</sub> and T<sub>4</sub> vs. T<sub>1</sub> , respectively.*
- 4) Improving in feed conversion of DM and TDN from 12.01 to 7.77 and from 6.27 to 4.91, g feed / g gain, respectively.*

- 5) *Due to the decrease in feeding costs and increase in the average daily weight gain, relative profit expressed as net profit / feeding cost increased to 151.52 % and 190.6 % in T<sub>3</sub> containing FSLCW (free choice) and T<sub>4</sub> containing FSLCW (free choice) plus 3 g PEG / head / day vs. T<sub>1</sub> (59.62 %) containing 40% clover hay , respectively.*

*Generally, it could be concluded that, in complete ration containing 40% clover hay, it is possible to replace the hay by peanut hulls ensiled with 5% lime +5% urea with adding fresh sweet liquid cheese whey (free choice) and 3g PEG / head /day in growing lambs rations to improve their performance and economic efficiency .*

**Keywords:** *Peanut hulls, lime, urea, cheese whey, Poly-Ethylene Glycol, sheep*

## INTRODUCTION

A wide variety of agricultural residues and by-products showed varying degrees of success when fed to ruminants. These feeds are imbalance in nutrients content and most of them are deficient in protein content. In addition, the abundant energy stored in the cellulose and hemi-cellulose content of the cell walls is not readily available for digestion by rumen micro-organisms. Digestion of highly fibrous materials in ruminants is severely limited by the presence of lignin (El-Talty, 1970 and 1973; Theander and Aman, 1984 and Aguilera, 1987). Moreover, the digestibility and intake of these residues tended to be lower with the higher lignifications levels. Many of tanniniferous feeds as peanut hulls could be useful as livestock fodders. But, various secondary compounds in these fodders may be toxic to animals or cause a reduction in their productivity (Barry and Blaney, 1987). Gow-Chin Yen and Pin-Der Duh (1995) reported that peanut hulls contains total phenolic compounds reaches to 10.2 mg / g . The occurrence of tannins or phenolic substances in feeds may inhibit rumen cellulolytic activity (Theriez and Boule,1970). Cheese whey which is an environmental pollutant by-product was found to have a good nutritional value in animals feeding. This by-product represents about 85-95% of the milk volume and retains 55% of whey nutrients . These nutrients are lactose(4.5 - 5%w/v),soluble proteins (0.6 - 0.8%w/v), lipids (0.4 - 0.5% w/v) and mineral salts(8 - 10% of dried extract) (Gonzalez,1996). Therefore, many studies were carried out to maximize the utilization of certain by-products as peanut hulls and cheese whey in animal feeding. Franklin *et al.*,1974 treated peanut hulls chemically and found wide variations in IVDMD (from 11.8% to 40%). Kaneko *et al.*, 1989 and Silanikove *et al.*, 2001 found that cheese whey and Polyethylene glycol have positive effects on the rumen microflora of animals fed on tanniniferous by-products .

Therefore, this study was designed to evaluate the ensiled peanut hulls with 5% lime plus 5% urea as well as to investigate the effect of its incorporating in the rations with fresh sweet liquid cheese whey (FSLCW) and polyethylene glycol (PEG) as feed additive on the performance and economic efficiency of growing lambs.

## MATERIALS AND METHODS

The present study was conducted at Mahlet-Mousa, Station, Kafr El-Sheikh Governorate.

Bulk of ground peanut hulls (3 - 5 mm screen) was treated chemically by ensiling in anaerobic condition with a mixture of 5% lime plus 5% urea (on DM basis) for 28 days. The moisture content was adjusted to be 45%. The treated materials were ensiled in ground silos (2 x 2 x 1m). After 28 days, the ensiled materials laid out in a sunny place to lose the excess of ammonia and moisture.

The first experiment was designed to evaluate the ensiled peanut hulls. Eighteen crossbred ( Romanov x Rahmani) rams weighing 55 kg in average were used to evaluate the effect of ensiling peanut hulls on its chemical composition and digestibility coefficients, through six metabolism trails. Animals were divided into six groups. Each group was allowed randomly to fed on one of the following rations:

- R<sub>1</sub> : (200 gm starch + 15g urea) / head /day + untreated peanut hulls (was offered *ad. lib.*).  
 R<sub>2</sub> : Clover hay+ untreated peanut hulls (1:1 air dry basis) (was offered *ad. lib.*).  
 R<sub>3</sub> : Barley + untreated peanut hulls (1 : 1 air dry basis) (was offered *ad. lib.*).  
 R<sub>4</sub> : (200 gm starch +15g urea) / head /day + treated peanut hulls (was offered *ad. lib.*).  
 R<sub>5</sub> : Clover hay + treated peanut hulls (1:1 air dry basis) (was offered *ad. lib.*).  
 R<sub>6</sub> : Barley + treated peanut hulls (1:1 air dry basis) (was offered *ad. lib.*).

Mixture of vitamins and minerals was added to cover the requirements according to NRC, 1985 for sheep.

The second experiment was designed to investigate the effect of adding FSLCW as a sole liquid with or without PEG to ensiled peanut hulls in lambs rations on animals performance . Four total mixed rations (TMR) were formulated to contain concentrate and roughage ratio at 60:40 as the following table.:

**Composition of the experimental rations**

Feed ingredients	Experimental rations			
	T <sub>1</sub>	T <sub>2</sub>	T <sub>3</sub>	T <sub>4</sub>
Wheat bran	15.0	15.0	15.0	15.0
Barely	29.0	29.0	29.0	29.0
Soya-bean meal	8.5	8.5	8.5	8.5
Clover hay	40.0	-	-	-
Treated peanut hulls	-	40.0	40.0	40.0
Sugar cane-Molasses	4.0	5.0	5.0	5.0
Lime stone	2.0	-	-	-
Common salt	1.0	1.0	1.0	1.0
Vits. & mins. mix.	0.5	0.5	0.5	0.5
Urea	-	1.0	1.0	1.0
TDN(calculated)	58.6	50.6	50.6	50.6
Poly Ethylene Glycol (PEG)	-	-	-	+
Fresh sweet liquid cheese whey	-	-	<i>ad-lib</i>	<i>ad-lib</i>

Vitamins and minerals mixture per Kg contained: vit.A 2,000,000 IU, vit.D<sub>3</sub> 150000 IU, vit. E 8.33g , vit.B<sub>1</sub> 0.33g , B<sub>6</sub> 1.7mg , B<sub>2</sub> 1.0 g , B<sub>3</sub> 8.33g , vit. K 0.33mg , Pantothenic acid 3.33 g , Biotin 33mg , Folic acid 0.83mg , Choline chloride 200mg , Mg 66.7g , Cu 0.5g , Se 16.6 mg , Zn 117g and Fe 12.5g.

+ : 3 g PEG/ head/ day.

The control ration (T<sub>1</sub>) was formulated to contain 15% wheat bran, 29 % barley, 8.5% soybean meal, 4% molasses, 2% limestone, 1% common salt, 0.5 % vits.& mins. mixture and 40% clover hay. Treatments T<sub>2</sub>, T<sub>3</sub> and T<sub>4</sub> were formulated to replace all of clover hay by treated peanut hulls . Fresh sweet liquid cheese whey was offered in T<sub>3</sub> and T<sub>4</sub> free choice. Polyethylene glycol was given to the lambs in T<sub>4</sub> at 3g / head /day with the drinking water. Twenty males of growing crossbred ( Romanov x Rahmani) lambs were about 6 months old and weighing 28.85±1.39 kg in average. Animals were allocated equally to four groups using a complete randomized design and housed in separate pens for 70 days. The daily requirements of the experimental rations (DM basis) were calculated according to (NRC, 1985).The air dried whey has TDN 78 % and DM 94% (CLFF, 2001) :

$$\text{AveragedailyDM (Kg)} = \frac{0.03W^{0.75} (1 + 5.5A) - Vx \text{ liquid whey TDN \%}}{\text{Ration TDN\%}}$$

W= Empty body weight

A= Average daily gain

V= Volume of liquid whey in liter

Drinking water or FSLCW was available freely for animals all the time after feeding. At the end of feeding trials, three animals from each group were randomly chosen for a digestion trial in which the FSLCW was offered daily as the recorded average during the whole feeding period (T<sub>3</sub> and T<sub>4</sub>). At the end of digestion trials, samples of rumen liquor were taken from the same animals just before morning feeding and then at two and four hours after feeding. Rumen samples were taken through stomach tube then filtered through two layers mesh cloth. Collected samples of rumen liquor were immediately determined for pH, then NH<sub>3</sub>- N concentration according to Conway (1962) and TVFA's concentration according to Warner (1964). Blood samples were taken at 24 hrs. post morning feeding from the jugular vein. Blood serum was obtained and tested for total protein (Henery, 1964) and albumin (Doumas, 1971). Globulin was calculated by the difference between total protein and albumin. Blood levels of GOT and GPT transaminases were determined according to Reitman and Frankel (1957). Samples of feeds and feces were analyzed according to AOAC (1990), while total phenols and tannins of untreated and ensilage treated peanut hulls were determined according to Folin-Ciocalteu method (Slinkard and Singleton, 1977).

Data obtained in this study was statistically analyzed according to SAS (1990). Significant differences were tested among means according to Duncan's multiple range test (1955).

## RESULTS AND DISCUSSION

### *First experiment:*

Effect of ensiling peanut hulls with 5% lime + 5% urea on chemical composition is presented in Tables (1) and (2). Results indicated that the content of total phenols and tannins decreased from 1.61 % to 1.47 % and from 1.33% to 1.11% by storing while, they nearly disappeared . In this connection, Russel and Lolley (1989) concluded that , tannin in high tannin milos could be deactivated rapidly and completely by reconstitution with aqueous urea. Price *et al.* (1979) and Reichert *et al.*

(1980) reported similar results. On the other hand, it could be noticed that, ash and CP contents increased while CF decreased by ensiling with lime and urea being, 7.52, 6.80 and 64.08 vs. 12.47, 9.93 and 58.11%, respectively. The increasing in ash and CP might be due to adding 5% lime and 5% urea (Sirohi and Rai, 1995 and Farghaly *et al.*, 2003) while, the decrease in CF might be due to the alkali treatments which reduces the strength of intermolecular hydrogen bonds (Granzin and Dryden, 2003).

**Table 1. Total phenols and tannins of untreated and ensilage treated peanut hulls (DM basis)**

Items	Peanut hulls		
	A	B	C
Total phenols %	1.61	1.47	0.15
Total tannins %	1.33	1.11	Nile

A. Untreated (after oil extraction directly).

B. Untreated (after storing for 3 months in a sunny place).

C. Ensiled with 5% lime + 5% urea.

**Table 2. Chemical composition, digestion coefficients and nutritive value of peanut hulls**

Items	DM	(DM basis)							
		OM	CP	CF	EE	NFE	Ash	TDN	DCP
Untreated:									
Chemical Comp.	92.05	92.48	6.80	64.08	1.89	19.71	7.52		
Digestion coefficients and nutritive values:									
By direct method	29.60	31.93 <sup>b</sup>	36.90 <sup>b</sup>	18.55 <sup>b</sup>	82.34 <sup>a</sup>	70.36 <sup>a</sup>		31.47	2.51
By difference (1)	31.08	33.41 <sup>b</sup>	70.64 <sup>a</sup>	19.90 <sup>b</sup>	97.35 <sup>a</sup>	54.85 <sup>b</sup>		33.20	4.80
By difference (2)	40.00	43.09 <sup>a</sup>	64.19 <sup>a</sup>	28.85 <sup>a</sup>	87.18 <sup>a</sup>	71.95 <sup>a</sup>		41.91	4.36
Treated									
Chemical Comp.	91.30	87.53	9.93	58.11	1.14	18.35	12.47		
Digestion coefficients and nutritive values:									
By direct method	16.00	18.21 <sup>d</sup>	6.31 <sup>c</sup>	10.17 <sup>c</sup>	64.30 <sup>b</sup>	44.04 <sup>bc</sup>		16.86	0.63
By difference (1)	21.47	24.39 <sup>c</sup>	11.98 <sup>c</sup>	23.02 <sup>ab</sup>	91.94 <sup>a</sup>	29.79 <sup>cd</sup>		22.66	1.19
By difference (2)	24.68	28.49 <sup>bc</sup>	28.60 <sup>b</sup>	30.52 <sup>a</sup>	65.25 <sup>b</sup>	23.46 <sup>d</sup>		25.87	2.84

a, b, c, d means in the same column with different superscripts differ at ( $P < 0.05$ ).

(1) With clover hay.

(2) With barley.

Digestibility coefficients and nutritive value of peanut hulls were affected by ensiling treatment as shown in Table (2). Ensiling peanut hulls caused a significant decrease ( $P < 0.05$ ) in digestion coefficients of DM, OM, CP, CF, EE, and NFE, being 16, 18.21, 6.31, 10.17, 64.3 and 44.04 % vs. 29.6, 31.93, 36.9, 18.55, 82.34 and 70.36% for treated and untreated peanut hulls respectively. On the other hand TDN and DCP were decreased from 31.47 and 2.51 to 16.86 and 0.63. The severe drop in CP, CF and NFE digestibilities of peanut hulls due to the ensiling treatment may be explained by one or more reason. Strong alkali treatment (by ensiling peanut hulls with 5 % lime + 5 % urea) of the dietary protein may be depressed CP digestibility through formation of uncommon toxic amino acid which known as lysinoalanine (De Groot and Stump, 1969). Isomerization of the dietary protein (Provansal *et al.*, 1975) and cross-linking within or between polypeptide chains (Aymard *et al.*, 1979) may be another reason. Ammoniated of roughages may be produce sugar-ammonia compounds, which had toxic effects on animals (Nishie *et al.*, 1969). Formation of

these types of compounds during ensiling peanut hulls could be a reason for the noticeable severe drop in CP digestibility. The noticeable reduction in CF digestibility of peanut hulls (10.17%) as affected by the ensiling treatment might be attributed to formation of potentially hazardous compounds which produce toxic effects especially on microflora in the rumen. Probably the most of these compounds resulted from the so-called nonenzymatic browning reaction (Maillard reaction) involving nitrogenous compounds (as urea, ammonia and dietary protein), carbohydrates (reducing sugars and water soluble carbohydrates), heat (as that produced during the ensiling which it was about 60 °C) and high pH values as adding lime and urea. In this connection Lee *et al.* (1975) found that the decrease in nutritional value by browning reaction was not only due to the degradation of amino acids, but also to the appearance of toxic materials. Moreover, Eskin *et al.* (1971) cited that high pH enhances nonenzymatic browning reactions (Maillard reaction) involving sugar by catalyzing their conversion to the highly reactive *enediol* form. Lignin that released by effect of ensiling peanut hulls with 5% lime + 5% urea is another compound (phenolic compound) which might have had hazardous effects on the ruminal microflora (Hartley, 1981) and it might be associated with the other toxic compounds in reducing CF digestibility. The reduction in CF digestibility might also be explained by a decrease in ruminal cellulolytic activity as a result of alkali-ensiled peanut hulls with 5% lime + 5% urea (Klopfenstein *et al.*, 1979), or inhibition of microbial activity by phenolic acids which released by alkali. The reduction in CF digestibility also might be related to the rate of passage of digesta through the reticulo-rumen which decreased by ensiling peanut hulls with 5% lime + 5% urea (Ali *et al.*, 1977). The observed severe reduction in NFE digestibility could be explained by forming undigested sugar-ammonia compounds (Maillard reaction). These results confirmed with the findings by Nishie *et al.* (1969), Eskin *et al.* (1971) and Lee *et al.* (1975).

#### **Second experiment:**

Chemical composition of the experimental rations presented in Table (3) shows that the content of CP is nearly similar. Adding FSLCW in T<sub>3</sub> and T<sub>4</sub> caused a marked decrease in CF content by 21.8% and an increase in EE and ash contents by 142% and 24.8%, respectively.

Data concerning digestibility and feeding value of the experimental rations presented in Table (4), indicates that digestibility coefficients of DM, OM, CP and CF of T<sub>4</sub> were better ( $P < 0.05$ ) than those in T<sub>2</sub> and T<sub>3</sub>. Using FSLCW in T<sub>3</sub> improved ( $P < 0.05$ ) digestibility of DM and CP by 10.06% and 17.92% against unsupplemented ration (T<sub>2</sub>) while the differences in digestibility of OM, EE and NFE were insignificant. Adding PEG in T<sub>4</sub> improved ( $P < 0.05$ ) digestibility of DM, OM and CF by 8.44%, 10% and 227.86% against unsupplemented ration (T<sub>3</sub>) while the differences in digestibilities of CP, EE and NFE were insignificant. Comparing T<sub>4</sub> with T<sub>1</sub>, indicated that, the digestibilities of DM, OM and EE were nearly similar being 63.10, 65.76 and 91.72 vs. 62.20, 64.50 and 89.33% respectively, while digestibilities of CP was higher ( $P < 0.05$ ) and of CF was lower ( $P < 0.05$ ) being 75.03 and 35.77 vs. 70.60 and 49.10% in T<sub>4</sub> and T<sub>1</sub> respectively. There was no significant difference between the experimental rations T<sub>4</sub> and T<sub>1</sub> in the nutritive values expressed as TDN (63.29 vs. 61.18) and DCP (12.32 vs. 11.93).

Table 3. Chemical analysis of the experimental rations (DM basis)

Items	Experimental rations			
	T1	T2	T3	T4
DM	88.90	90.30	90.30	90.30
OM	91.50	92.23	90.30	90.30
CP	16.90	16.63	16.42	16.42
CF	18.10	28.78	22.50	22.50
EE	2.10	1.45	3.51	3.51
NFE	54.40	45.37	47.87	47.87
Ash	8.50	7.77	9.70	9.70

Rations T3 and T4 contain 22 % cheese whey (DM basis). T4 contain 3g PEG / head / day.

Table 4. Digestion coefficients and nutritive values of the experimental rations (DM basis)

Items	Experimental rations			
	T1	T2	T3	T4
<b>Apparent digestibility:</b>				
DM	62.20 <sup>ab</sup>	52.87 <sup>d</sup>	58.19 <sup>bc</sup>	63.10 <sup>a</sup>
OM	64.50 <sup>a</sup>	56.22 <sup>c</sup>	59.78 <sup>bc</sup>	65.76 <sup>a</sup>
CP	70.60 <sup>b</sup>	60.53 <sup>c</sup>	71.38 <sup>ab</sup>	75.03 <sup>a</sup>
CF	49.10 <sup>a</sup>	14.61 <sup>c</sup>	10.91 <sup>c</sup>	35.77 <sup>b</sup>
EE	89.33 <sup>b</sup>	90.98 <sup>ab</sup>	92.11 <sup>a</sup>	91.72 <sup>ab</sup>
NEF	66.70 <sup>c</sup>	77.04 <sup>a</sup>	76.40 <sup>a</sup>	74.77 <sup>ab</sup>
<b>Nutritive values:</b>				
TDN	61.18 <sup>ab</sup>	52.19 <sup>c</sup>	57.91 <sup>b</sup>	63.29 <sup>a</sup>
DCP	11.93 <sup>ab</sup>	10.07 <sup>c</sup>	11.80 <sup>ab</sup>	12.32 <sup>a</sup>

a, b, c, d means in the same raw with different superscripts differ at (P< 0.05).

Effects of feeding the experimental rations containing 40% ensiled peanut hulls, with FSLCW or/ and PEG on performance of growing lambs are shown in Table (5). The obtained results revealed that lambs which drank FSLCW (T<sub>3</sub>) appeared higher (P<0.05) total live weight gain (10.6 Kg) and average daily live weight gain(151g) than those fed ration without FSLCW(T<sub>2</sub>) which detected the least total live weight(6.8 Kg )and average daily live weight (97g).The differences between T<sub>3</sub> and T<sub>1</sub> were not significant (P<0.05). Lambs in T<sub>4</sub> which given 3 g PEG/head/day showed significantly (P<0.05) higher total live weight gain (13.8Kg) and average daily live weight gain(197g) than rations T<sub>3</sub> and T<sub>1</sub>. Data concerning feed intake and feed conversion as shown in Table (5) indicated that lambs in treatments T<sub>3</sub> and T<sub>4</sub> ingested relatively high quantity of FSLCW (6.1 and 6.6 L / day or 311and 337 g based on DM) detected more daily total dry matter intake. This observation was in agreement with those obtained by Anderson *et al.* (1974). Data concerning feed conversion indicated that lambs fed on T<sub>4</sub>and T<sub>1</sub> had lower values being, 7.77 and 8.60 than those fed on T<sub>3</sub> being 9.33 g DMI / g gain, while lambs fed on T<sub>2</sub> had the worst feed conversion being 12.01 g DMI /g gain. Feed conversion ratio expressed as g TDN /g gain had the same trend. These results are in harmony with those suggested by Galloway *et al.* (1992) and Titus *et al.* (2000).

Data of economical evaluation of feeding growing lambs on experimental rations are summarized in Table(5) . It was found that , the relative profit expressed as net

profit / feeding cost increased to 151.52 % and 190.6 % in T<sub>3</sub> containing FSLCW ( free choice) and T<sub>4</sub> containing FSLCW( free choice) plus 3 g PEG / head / day vs. T<sub>1</sub> (59.62 %) containing 40% clover hay , respectively.

From the previous results it could be concluded that the improving rule of FSLCW and PEG on performance of growing lambs fed rations containing peanut hulls ensiled with lime and urea is summarized as follows: Phosphopeptide complexe as a combination between whey protein and phosphorus may have an interesting effect on the intestinal absorption of minerals (Moulin and Galzy, 1984). Formation of lactosylurea may be maximizes the utilization of urea in the rumen (Sienkiewicz and Riedel, 1990). The bacterial content of cheese whey also may have a role in increasing of feed efficiency the experimental rations( Bolsen *et al.*, 1995).Supplementation of PEG has good ability to combine with the dietary tannins and phenolic compounds instead of binding with the dietary protein, digestive enzymes, minerals, cellulosic bacteria and proteolytic bacteria, and so it can alleviate the reductions in digestion associated with the presence of tannins and phenolic compounds (Scalbert, 1991; Aerts *et al.*, 1999 and Molan *et al.*, 2001).

**Table 5. Average live body weight, feed intake, feed conversion and economic efficiency of growing lambs fed the experimental rations**

Items	Experimental rations			
	T <sub>1</sub>	T <sub>2</sub>	T <sub>3</sub>	T <sub>4</sub>
No. of animals	5	5	5	5
Experimental period (days)	70	70	70	70
Initial live body weight (IBW), (kg)	28.4	28.2	29.2	29.6
Final live body weight (FBW), (kg)	39.0 <sup>ab</sup>	35.0 <sup>b</sup>	39.8 <sup>a</sup>	43.4 <sup>a</sup>
Total live weight gain, kg	10.6 <sup>b</sup>	6.8 <sup>c</sup>	10.6 <sup>b</sup>	13.8 <sup>a</sup>
Average daily live weight gain, gm	151 <sup>b</sup>	97 <sup>c</sup>	151 <sup>b</sup>	197 <sup>a</sup>
<b>Feed intake:</b>				
Feed mixture g/d (DM)	1299	1165	1098	1193
Cheese whey (liquid) L/d	0.0	0.0	6.1	6.6
Cheese whey (DM) g/d (w/v)	0.0	0.0	311	337
Total dry matter intake (g/d)	1299	1165	1409	1530
TDN (g/d)	795	608	816	968
<b>Feed conversion (g feed /g gain):</b>				
DMI	8.60	12.01	9.33	7.77
TDN	5.27	6.27	5.40	4.91
<b>Economical evaluation:</b>				
Feeding cost (LE /head/day)	2.08	1.40	1.32	1.49
Gain price (LE /head/day)	3.32	2.13	3.32	4.33
Profit (LE /head/day)	1.24	0.73	2.00	2.84
Relative profit ( profit / feeding cost % )	59.62	52.14	151.52	190.60
Economic improvement (%)	100	87	254	320

a, b, c, d means in the same raw with different superscripts differ at (P< 0.05).

RationsT3 and T4 contain 22 % cheese whey (DM basis).T4 contain 3 PEG g/head/day.

The relative profit was calculated as a percentage ratio between profit and feeding cost /head /day. Based on market prices at the beginning of experiment, the price (LE/ton) were 1600 for T<sub>1</sub> , 1200 for T<sub>2</sub> and T<sub>3</sub> , 1250 for T<sub>4</sub> and FSLCW has no market price. The price of one Kg live body weight on selling was LE 22.



Data of rumen liquor parameters presented in Table (6) shows a significant decrease ( $P<0.05$ ) in  $\text{NH}_3\text{-N}$  concentration of rations  $T_3$  and  $T_4$  compared with  $T_2$  after 2 hrs. of feeding being 31.27, 26.87 and 41.33 mg/100ml rumen liquor respectively. After 4 hrs.  $T_3$  showed significant ( $P<0.05$ ) higher concentration of  $\text{NH}_3\text{-N}$  compared with  $T_2$  and  $T_4$ , being 26.23, 16.80 and 19.13 mg/100ml respectively. These results might be due to the effects of FSLCW and PEG on the activity of rumen bacteria which consume much amount of ammonia for its growth or/ and to the formation of lactosylurea (direct reaction between lactose and urea) which acts as a reservoir of ammonia for long time. These results agreed with those obtained by Wright (1967), Moulin and Galzy (1984), Stock *et al.* (1986), Sienkiewicz and Riedel (1990) and Gaggiotti *et al.* (2002). The values of TVFA's were significantly higher ( $P<0.05$ ) in FSLCW and FSLCW + PEG groups ( $T_3$  and  $T_4$ ) compared with  $T_2$ , being 10.33 and 10.17 vs. 8.83 meq/100mL rumen liquor after 2 hrs. while after 4 hrs., concentration of TVFA's was the highest significantly ( $P<0.05$ ) in FSLCW + PEG ( $T_4$ ), being 13.0 meq /100 mL rumen liquor. Higher TVFA's concentrations as a result to FSLCW and PEG supplementation may be attributed to their ability to improve the digestibility of most nutrients in the rumen (Galloway *et al.*, 1992 and Salawu *et al.*, 1997). Supplementation of whey protein as a rapid degradable protein may be a supplier of a specific amino acids or peptides (limiting bacterial growth) which are used to form branched-chain VFA that are required by certain bacteria for optimum growth and so optimum microbial protein synthesis (Stock *et al.*, 1986).

Table 6. Mean values of rumen and blood parameters of growing lambs fed the experimental rations

Items	Time (hours)	Experimental rations			
		T1	T2	T3	T4
<b>Rumen parameters:</b>					
pH	0	6.97	7.08	6.99	7.30
	2	6.18	6.23	6.21	5.94
	4	6.87 <sup>ab</sup>	7.05 <sup>a</sup>	6.26 <sup>b</sup>	5.56 <sup>c</sup>
	mean	6.67	6.79	6.48	6.27
TVFA's (meq/100mL)	0	6.50	5.83	6.17	5.50
	2	11.17 <sup>a</sup>	8.83 <sup>b</sup>	10.33 <sup>a</sup>	10.17 <sup>a</sup>
	4	9.67 <sup>b</sup>	6.83 <sup>c</sup>	7.17 <sup>c</sup>	13.00 <sup>a</sup>
	mean	9.11	7.16	7.89	9.56
NH3- N (mg/100mL)	0	15.40	16.33	18.67	17.73
	2	16.33 <sup>d</sup>	41.33 <sup>a</sup>	31.27 <sup>b</sup>	26.87 <sup>c</sup>
	4	16.33 <sup>b</sup>	16.80 <sup>b</sup>	26.23 <sup>a</sup>	19.13 <sup>b</sup>
	mean	16.02	24.82	25.39	21.24
<b>Blood parameters:</b>					
Total protein (g / dL )	0	8.27 <sup>d</sup>	14.63 <sup>a</sup>	12.30 <sup>b</sup>	10.00 <sup>c</sup>
Albumin (g / dL )	0	3.57 <sup>b</sup>	4.68 <sup>a</sup>	3.53 <sup>b</sup>	4.80 <sup>a</sup>
Globulin (g / dL )	0	4.7 <sup>b</sup>	9.97 <sup>a</sup>	8.77 <sup>a</sup>	5.20 <sup>b</sup>
GOT ( U / L )	0	40.07 <sup>c</sup>	79.4 <sup>a</sup>	80.17 <sup>a</sup>	51.00 <sup>b</sup>
GPT ( U / L )	0	20.33 <sup>c</sup>	30.33 <sup>a</sup>	30.67 <sup>a</sup>	25.63 <sup>b</sup>

a, b, c, d means in the same raw with different superscripts differ at ( $P<0.05$ )

Data of blood serum parameters presented in Table ( 6 ) shows higher ( $P < 0.05$ ) levels in total protein , albumin , globulin , GOT and GPT by feeding on ration  $T_2$  (containing ensiled peanut hulls) than  $T_1$  (containing clover hay), being 14.63 , 4.68 , 9.97 , 79.4 and 30.33 vs. 8.27 , 3.57 , 4.7 , 40.07 and 20.33, respectively. Feeding lambs on rations  $T_3$  and  $T_4$  supplemented by FSLCW or / and FSLCW + PEG decreased significantly blood parameters to be around normal levels. From the previous results it could be concluded that the high concentrations over the normal limits  $T_2$  may refer to that the animals were under stress by feeding rations containing in anti-nutritional factors as tannins derivatives or high level of urea which may induce over limit globulin (immunoglobulin), GOT and GPT. These results are in harmony with the results of Dodak and Zboril (1976) who stated that, GOT and GPT were increased by coumarin and Arunachalam and Sivakumar (2003) also found that total protein , albumin and globulin increased significantly under stress. On the other hand, the decreasing in values of blood parameters by adding FSLCW and PEG ( $T_4$ ) might indicate a low stress in the liver to detoxicate any harmful effect due to feeding of tanniferous feeds ensiled with urea (Russel and Lolly, 1989) .

Generally, the performance and economic efficiency of growing lambs fed total mixed rations containing 40% treated peanut hulls could be improved by used fresh sweet liquid cheese whey in rations and adding polyethylene glycol with drinking water compared with the ration containing 40% clover hay.

## REFERENCES

- Aerts, R.J., W.C. McNabb; A Molan; A Brand; T.N. Batry and J.S. Peters, 1999. Condensed tannis from *Louts Corniculatus* and *Luts Pedunculatus* exert different effects on the *in vitro* rumen degradation of ribulose 1-5 biophosphate carboxylase/oxygenase protein. J. Sci. Food Agric., 79: 79-85.
- Aguilera, JF., 1987. Improvement of olive cake and grape by products for animal nutrition. Degradation of lignocellulosics in ruminants and in industrial processes. Nutrition- Abstracts- and Reviews. Series-B, 1988, 058:03320.
- Ali, C.S., V.C. Mason and J. Waagepetersen, 1977. The voluntary intake of pelted diets containing sodium hydroxide-treated wheat straw by sheep. 1. The effect of the alkali concentration in the straw. Anim. Feed Sci. and Techno., 14 : 139-149.
- Anderson, M.J., R.C. Lbmb; C.H. Mickelson and R.L. Wis-Combe, 1974. Feeding liquid whey to dairy cattle. J.Dairy Sci., 57 : 1206.
- A.O.A.C., 1990. Official Methods of Analysis.(15<sup>th</sup> Ed.). Association of Official Analytical Chemists , Arlington , VA.,USA .
- Arunachalm,s. and T. sivakumar , 2003 . Blood profile consituents associated with production diseases in Jersey cross breed cows . Indian J. Anim. Sci. , 73 : 44 – 47.
- Aymard, C., J.L. Cuq and J.C. Chefield, 1979.A textbook of Toxic constituents of plant foodstuffs.2<sup>nd</sup> Ed, Chapter 12. Liener, I.E.(ed.),New York ,N.Y.,USA (ISBN 0-12-449960-0).
- Barry, T.N. and B.J. Blaney, 1987. Secondary compounds of forages. Anim. Feed Sci. and Techno., 91 :41-57.
- Bolsen,K.K., M.A. Young, G.L. Huck, M.K. Siefes, J.E. Turner, S.A. Anderson, R.V. Pope and J.S. Pendergraft, 1995. Effect of bacterial inoculation on

- preservation efficiency and nutritive value of alfalfa silage for growing cattle. J. dairy Sci., 80( Suppl.:1).
- C.L.F.F., 2001. Feed Composition Tables for Animal and Poultry Feedstuffs used in Egypt. Edited by Central Lab for Food & Feed., Agricultural Research Center, Ministry of Agriculture.
- Conway, E.F., 1962. Microdiffusion Analysis And Volumetric Error. Rev. Edt. Lockwood, London, UK.
- De Groot, A.P. and P. Stump, 1969. Toxic constituents of plant foodstuffs J. Nutr.98, 45-56.
- Dodak , V. and F. Zboril , 1967 . Toxic action of plant phenolic groups . Chem. Coumer. , 32: 411 .
- Doumas, B., 1971. Colorimetric determination of albumin. Clin. Chem. Cat., 31: 87.
- Duncan, D.B., 1955. Multiple F-test Biometrics, 11:1-42.
- El-Talty, Y.I., 1973. Some factors affecting the feeding value of roughages with reference to lignin. Ph. D. Thesis, Fac. of Agric Cairo Univ.
- El-talty, Y.I., 1970. Some nutritional studies on the lignin fraction. M. Sc. Thesis, Fac. of Agric. Cairo Univ.
- Eskin, N.A.M., H.M. Henderson and R.J. Townsend, 1971. A textbook of plant foodstuffs. 2<sup>nd</sup> Ed, Chapter 12. Liener, I.E.(ed.), New York ,N.Y., USA (ISBN 0-12-449960-0).
- Farghaly, M.S., I.M. Awadalla and M.A. Ali, 2003. Effect of urea and lime treatments on feeding value of wheat straw used in growing lambs ration. J. Agric. Mansoura Univ., 28 : 6645 – 6654.
- Franklin E. Barton, II, Henry E. Amos, William J. Albrecht and Donald Burdick, 1974. Treating peanut hulls to improve digestibility for ruminants. J. Anim. Sci., 38:860-846.
- Gaggiotti, M.C., M.R. Gallarod, A.A. Abdela, C. Arakaki, L. Burdisso and A.R. Castillo, 2002. Effect of feeding dairy cows with whey permeate on ruminal environment under alfalfa grazing conditions. J. Anim. Sci., 79( Suppl.: 1).
- Galloway, D.L. Sr., A.L. Goetsch, W. Sun, L.A. Forester, Jr., G.E. Murphy, E.W. Grant, and Z.B. Johnson, 1992. Digestion, feed intake, and live weight gain by cattle consuming Bermuda grass hay supplemented with whey. J. Anim. Sci., 70:2533-2541.
- Gonzalez Siso, M.I., 1996. The biotechnological utilization of cheese whey. Bioresource Technology ,57 : 1-11.
- GOW-CHIN YEN and Pin-Der Duh, 1995. Antioxidant activity of methanolic extracts of peanut hulls from various cultivars. Journal of the American Oil Chemist's Society ,72: 1065-1067.
- Granzin, B.C. and G.M. Dryden, 2003. Effects of alkalis , oxidants and urea on the nutritive value of rhodes grass (*Chloris gayana cv. Acllide*). Anim. Feed Sci. and Techno. , 103: 113.
- Hartley, R.D., 1981. Chemical constitution, properties and processing of lignocellulosic wastes in relation to nutritional quality of animal. Agric. Anim. Feed Sci. and Techno., 14 :139-149.
- Henery, E.J., 1964. Colorimetric determination of total proteins and calcium. Clin.Chem.Principles and Techniques. Harper and Row, YN, USA.

- Kaneko, T., K. Ushida and Y. Kojima, 1989. Effect of starch on cellulolysis by rumen microbial populations with or without protozoa. *J. Anim. Sci.*, 70:2533-2541.
- Klopfenstein, T.J., V.E. Krause, H.J. Jones and W. Woods, 1979. Chemical treatment of low quality roughages. *J. Anim. Sci.*, 35:418.
- Lee, C.M.; T.C. Lee and C.O. Chichester, 1975. Toxic factors induced by processing. *Food Sci. and Technol.*, 1: 587-603.
- Molan, A.L.; G.T. Attwood; B.R. Min. and W.C. McNabb (2001). The effect of condensed tannins from *Lotus pedunculatus* and *Lotus corniculatus* on the growth proteolytic rumen bacteria *in vitro* and their possible mode of action. *Can.J.Microbial.*, 47:626-633.
- Moulin, G. and P. Galzy (1984). Whey, a potential substrate for biotechnology. *Bioresource Technology*, 57:1-11.
- Nishie, K.; A.C. Waies and A.C. Keyl (1969). Toxic constituents of plant foodstuffs. *Toxicol. Appl. Pharmacol.*, 14: 301-307.
- NRC, (1985). National Research Council, 6<sup>th</sup> ed. Nutrient Requirements of Sheep. National Academy Press, Washington, DC., USA.
- Price, M.L.; L.G. Butler ; J.C. Rogler and W.R. Featherston (1979). Overcoming the nutritionally harmful effects of tannins in sorghum grains by treatment with inexpensive chemicals. *J. Agric. Food Chem.*, 27: 441.
- Provansal, M., J.L. Cuq and C. Chefiel, 1975. Toxic constituents of plant foodstuffs. *Agric. Food Chem.*, 23:938-943.
- Reichert, R.D.; S.E. Fleming and D.J. Schwab, 1980. Tannin deactivation and nutritional improvement of sorghum by anaerobic storage of H<sub>2</sub>O-, HCL- or NaOH- treated. *J. Agric. Food Chem.*, 28: 824.
- Reitman, S. and S. Frankel, 1957. Colorimetric determination of AST and ALT activity. *Amer. J. Clin. Path.*, 28:56.
- Russel, R.W. and J.R. Lolley, 1989. Deactivation of tannin in high tannin milos treatment with urea. *J. dairy sci.*, 72: 2427-2730.
- Salawu, M.B., T. Acamovic, C.S. Stewart and F.D. DeB. Hovell, 1997. Quebracho tannins with or without Browse plus (a commercial of polyethylene glycol) in sheep diets: effect on digestibility of ruminants *in vivo* and degradation of grass hay in sacco and *in vitro*. *Anim. Feed Sci. and techno.*, 69: 67-78.
- SAS, 1990. User's Guide: Statistic SAS Inst. Gray, NC.
- Scalbert, A., 1991. Antimicrobial properties of tannins. *Phytochemistry*, 30: 3875-3883.
- Sienkiewicz, T. and C.L. Riedel, 1990. Whey and whey utilization. *Bioresource Technol.*, 57 :1-11.
- Silanikove, N., A. Perevolotsky and F.D. Provenza, 2001. Use of tannin-binding chemicals to assay for tannins and their negative postingestive effects in ruminants. *Anim. Feed Sci. and Technol.*, 91 :69-81.
- Sirohi, S.K. and S.N. Rai, 1995. Associative effect of lime plus urea treatment of paddy straw on chemical composition and *in-vitro* digestibility. *Indian J. Anim. Sci.*, 65:775.
- Slinkard, K. and Singleton V., 1977. Total phenol analysis: Automation and comparison with manual methods. *Am. J. Eno. Vitic.*, 28, 49-55.

- Stock R., T. Kiopfenstein, D. Brink, R. Britton and D. Harmon, 1986. Whey as a source of rumen-degradable protein.1. Effects on microbial protein production. *J. Anim. Sci.*, 63: 1561- 1573.
- Theander, O. and P. Aaman, 1984. Antimicrobial and chemical characteristics. *Anim. Feed Sci. and Techno.*,73: 243-2610.
- Theriez, M. and G. Boule, 1970. Upgrading feeding value of olive cake . *Anim. Feed Sci. and Techno.*, 14 :139-149.
- Titus,C.H., F.D. Provenza, E.A. Burritt, A. Perrvolotsky, N. Silanikove, 2000. Performance for foods varying in macronutrients and tannins by lambs supplemented with polyethylene glycol. *J. Anim. Sci.*, 78: 1443-1449.
- Warner, A.C.I., 1964. Production of volatile fatty acids in the rumen: Methods of measurements. *Nutr. Abst. and Rev.*,34:339.
- Wright, D.E., 1967. Metabolism of peptides by rumen microorganisms., *Appl. Microbial.*15:547.

## تأثير شرش الجبن و البولي ايثيلين جليكول علي أداء الحملان النامية المغذاة علي قشر الفول السوداني المعامل بالجير و اليوريا

يحيي ابراهيم التلتي<sup>1</sup> , محمد سيد فرغلي<sup>1</sup> , رأفت شحاتة ميخائيل<sup>2</sup>

1-قسم الإنتاج الحيواني، كلية الزراعة، جامعه لقااهرة، الجيزة، جمهورية مصر العربية، 2- معهد بحوث الإنتاج الحيواني، مركز البحوث الزراعية، وزارة الزراعة، الجيزة، جمهورية مصر العربية

اشتمل هذا البحث علي جزئين : الأول يهدف إلي دراسة تأثير كمر قشر الفول السوداني لا هوانيا بخلط من الجير 5% + اليوريا 5% علي كل من التركيب الكيماوي و معاملات الهضم و القيمة الغذائية باستخدام كباش خليط من الرحماني و الروماتوف . الجزء الثاني يهدف إلي دراسة تأثير إضافة شرش الجبن السائل أو الشرش مع البولي ايثيلين جليكول علي معدلات الأداء الانتاجي و الكفاءة الاقتصادية للحملان النامية ( الخليط من الرحماني و الروماتوف ) المغذاة علي علائق متكاملة ، تحتوي علي 40% مائه خشنه (قشر الفول السوداني المعامل كيماويا بالجير 5% + اليوريا 5% ) و 60% علف مركز ، حيث تم استخدام 20 حمل متوسط وزن 28.85 كجم عمر 6 اشهر ، و التي قسمت عشوائيا لأربعة مجموعات متساوية ، و غذيت علي الأربعة معاملات التجريبية لمدة 70 يوما و ذلك علي النحو التالي :

المعاملة الأولى T1 (الكنترول): تحتوي علي 15% نخاله قمح، 29% شعير، 8.5% كسب فول الصويا، 40% دريس برسيم، 4% مolas قصب السكر، 2% حجر جيرى، 1% ملح طعام، 0.5% مخلوط فيتامينات و معادن .

المعاملة الثانية T2 : هي تقريبا نفس المعاملة T1 مع إستبدال كل دريس البرسيم بقشر الفول السوداني المعامل .

المعاملة الثالثة T3 : T2 + شرش الجبن السائل بحرية بدلا من ماء الشرب .

المعاملة الرابعة T4: T2 + شرش اللبن السائل بحرية بدلا من ماء الشرب + 3 جم بولي ايثيلين جليكول / راس / يوم .

تم توفير الاحتياجات الغذائية من الطاقة و البروتين لنمو الحيوانات لتغطي متوسط الزيادة اليومية طبقا لمقررات NRC 1985 . وكان المتناول اليومي من شرش الجبن محسوبا كمركبات غذائية كلية مهضومة (TDN) يحل محل جزء من T2 بنفس القيمة من ال TDN . و في نهاية تجربة النمو أجريت أربعة تجارب هضم للأربعة معاملات التجريبية و قدرت فيها بعض قياسات الكرش و الدم، وكانت النتائج كالتالي:-

أولا : التجربة الأولى ( تأثير معاملة قشر الفول السوداني بالجير و اليوريا ) .

1- حدوث انخفاض ملحوظ في الفينولات و التانينات الكلية ( من % 1.61 السلي 0.15% و من 1.33% الي اثار علي التوالي )

- 2- انخفاض معنوي لمعاملات هضم البروتين الخام و الالياف الخام و المستخلص الخالي من الازوت والمركبات الغذائية الكلية المهضومة و البروتين الخام المهضوم ( من 36.9 % ، 18.55 % ، 70.36 % ، 31.47 % ، 2.51 % الي 6.31 % ، 10.17 % ، 44.04 % ، 16.86 % ، 0.63 % علي التوالي ) .
- ثانيا : التجربة الثانية ( تأثير اضافة شرش الجبن السائل و البولي ايثيلين جليكول )
- 1- زيادة معنوية لمعاملات هضم البروتين الخام و الالياف الخام و المركبات الغذائية الكلية المهضومة و البروتين الخام المهضوم ( من 60.53 % ، 14.61 % ، 52.19 % ، 10.07 % الي 75.03 % ، 35.77 % ، 63.29 % ، 12.32 % علي التوالي )
- 2- زياد معنوية في متوسط وزن الجسم الحي المكتسب يوميا من 97 جم/ يوم الي 197 جم/ يوم.
- 3- تحسن في معامل التحويل الغذائي للمادة الجافة والمركبات الغذائية الكلية المهضومة ( من 12.01 % ، 6.27 % الي 7.77 % ، 4.91 % علي التوالي ) .
- 4- أدت التغذية الحرة للحملان الناميه علي شرش الجبن السائل وإضافه 3 جم بولي ايثيلين جليكول/ راس/ يوم مع العلائق المتكاملة و المحتوىه علي 40% قشر الفول السوداني المعامل بالجير واليوريا الي تحسين الكفاءة الاقتصادية ( الربح / تكلفة التغذية ) الي 151.52 % و 190.60 % للمعاملة الثالثة و الرابعة مقابل 59.62 % للمعاملة الاولى .
- مما سبق يتبين أن تغذية الحملان النامية علي شرش الجبن السائل تغذية حرة و إضافة 3 جم بولي ايثيلين جليكول / راس / يوم قد حسن من معدلات الأداء الانتاجي و الكفاءة الاقتصادية للحملان النامية المغذاه علي علائق متكاملة تم فيها استبدال 40% دريس البرسيم بقشر الفول السوداني المعامل بالجير 5% + اليوريا 5% .