

Control of Potato Bacterial Soft Rot Disease Caused by *Erwinia carotovora* subsp. *carotovora* with *Streptomyces sioyaensis* and Cinnamon Oil.

F. A. Mansour, A. H. Mohamedin^{*}, A. E. Esmaeel¹ and Huda H. Badr¹

Botany Department, Faculty of Science, Mansoura, University, Mansoura and ¹Agricultural Research Center, Cairo, Egypt.

SOFT rot of potato is one of the most important diseases caused by *Erwinia carotovora* subsp. *carotovora*. Possibilities of its biological control were investigated by the use of some actinomycetes antagonists for controlling *E. carotovora* subsp. *carotovora*. A total of 40 different actinomycetes strains were isolated from the rhizosphere soil of healthy potato plants collected from different localities of Dakahlyia Province, Egypt. These were then tested, *in vitro*, for their antibacterial activity against *E. carotovora* subsp. *carotovora* using an agar disc method. Eight isolates were active against *E. carotovora* subsp. *carotovora* and the most active isolate was selected for further tests and characterized by conventional methods. Morphological, physiological and cultural characterization of the most active isolate established that it was closely related to *Streptomyces sioyaensis*.

On the other hand, 28 commercial plant essential oils were screened, *in vitro*, to evaluate their inhibitory activities against *E. carotovora* subsp. *carotovora*. Nine out of 28 oils exhibited considerable inhibitory activities and the oil of *Cinnamomum zeylanicum* (cinnamon oil) was the most active.

Controlled bacterial soft rot of potatoes were carried out with *S. sioyaensis* and cinnamon oil in greenhouse experiments. Treatment of potato tubers with *S. sioyaensis* or cinnamon oil either individually or in combination, caused significant decrease in the percentage of soft rot disease at harvest time and after storage compared with control treatment. Different formula of *S. sioyaensis* (powder, suspension and granules) showed different effects on reducing disease incidence. Powder formulation of *S. sioyaensis* showed the highest reduction in disease incidence at harvest time and after storage. Increasing dose of *S. sioyaensis* in granules formula enhanced its efficacy in reducing disease incidence of potato tubers.

Keywords: Potato, Soft rot, Biological control, *S. sioyaensis*, Cinnamon oil, *E. carotovora* subsp. *carotovora*.

Potato (*Solanum tuberosum* L.) is affected by five bacterial diseases namely: bacterial wilt or brown rot caused by *Ralstonia solanaceum*, soft rot of tubers

* Corresponding author. e-mail: amohamedin@mans.edu.eg

and stem caused by *E. carotovora* subsp. *carotovora* and *E. carotovora* subsp. *atroseptica*, ring spot caused by *Clavibacter michiganense* subsp. *sepedonicus* and common scab caused by *Streptomyces scabies*.

In Egypt, ring rot does not occur and the common scab is a minor disease. However, bacterial wilt and soft rot diseases are the most destructive and wide spread in tropics, subtropics and warm temperate regions (Bradbury, 1977).

Bacterial soft rot is considered one of the most important potato diseases causes losses in the field and one of the few diseases that can spread extensively in stores (Lund, 1979) and also cause significant losses during storage period (Helias *et al.*, 2000). Protection of potato plants from the aforementioned diseases using chemicals has been attempted with limited success.

Biological control has been found of immense potential in management of soft rot of potatoes. The basic idea of biological control is a reduced infection caused by rhizosphere colonizers antagonizing *E. carotovora* subsp. *carotovora* and *E. carotovora* subsp. *atroseptica* at the root infection site. Certain bacteria like *Pseudomonas fluorescens*, *Pseudomonas putida* and *Bacillus* sp. have been found to delay the development and reduce incidence of soft rot of potatoes (Tawfik *et al.*, 2001). In this respect, Sharga & Lyon (1998) reported that *Bacillus subtilis* (Bs 107) had complete activity *in vivo* against *E. carotovora* subsp. *carotovora* and *E. carotovora* subsp. *atroseptica*.

Kastelein *et al.* (1999) found that some strains of *Pseudomonas fluorescens* and *Pseudomonas putida* reduced the development of black leg disease. Also similar results have been recorded by Lucon & Melo (1999), Roberti & Selmi (1999) and Aysan *et al.* (2003). In addition, several isolates of non-pathogenic *Streptomyces* sp. have been shown to significantly reduce the severity of potato scab caused by *Streptomyces scabies* (Liu *et al.*, 1995 and Ryan *et al.*, 2004).

The most recent information that relates to some plant extracts or their active constituents are widely used as antimicrobial agents in controlling plant diseases. The use of plant extracts or purified known chemicals is a new approach that showed promising results in reducing certain plant diseases with less environmental pollution (Zedan, 1993 and Badr, 2006).

In this respect, Deans & Sviboda (1990) reported that volatile oils of *Origanum majorana* has inhibitory effect on the growth of *E. carotovora*. Barnejee & Nigam (1978) measured the antimicrobial activity of *Curcuma longa* essential oil against *E. carotovora* and *Pseudomonas solanacearum*. Similar results have been obtained by Ibrahim *et al.* (1987), Atilla (1989) and Singh *et al.* (1992).

This work was undertaken to evaluate the efficacy of suppressive bacteria belonging to actinomycetes as biocontrol agents and some commercial essential oils to control soft rot disease of potatoes.

Material and Methods

Bacterial culture

E. carotovora subsp. *carotovora* that has been isolated from infected potato tubers and completely identified using Bergey's Manual of Systematic Bacteriology (Lelliott & Dickey, 1984). Isolation and identification were performed according to Badr (2006). Pure cultures were maintained on nutrient agar slants and subcultured every month.

Pathogenicity test of the bacterial isolates

Pathogenicity tests of the bacterial isolates on potato tubers (Sponta cultivar) were carried out according to Tolba (1998). The bacteria were incubated in nutrient agar at 28°C for 48 hr, washed and resuspended in sterile water. The growth of the bacterial suspension was approximately at 10^8 c.f.u./ml (OD_{600} of 0.05) and further on diluted in sterile water to reach 10^6 c.f.u./ml. Healthy potato tubers were surface sterilized by washing with soap, rinsed in sterile water for 15 min, immersed in 95 % ethyl alcohol and instantly subjected to flame. In each tuber, a hole of 2 cm depth was made with a sterilized cork borer (5 mm in diameter) and the tuber was inoculated with 0.5 ml bacterial suspension (10^6 cells/ml). The hole of the tuber was closed with the removed potato cylinder, and then sealed with sterile vaspar. Inoculated tubers were kept in a moist condition at 28°C and examined for soft rot symptoms after 4 days. Control tubers were inoculated with sterilized distilled water. At least three tubers were used as replicates for each isolates.

Isolation of antagonistic actinomycetes

Actinomycetes were isolated from soil samples collected from the rhizosphere of healthy potato plants of different localities of Dakahlyia Governorate during 2004. Isolation was performed by serial dilution plate technique, according to Johnson *et al.* (1960), using starch-nitrate agar complemented with nystatin (50µg/ml) as antifungal agent. The plates were incubated at 28°C for 7 to 14 days and differently looking colonies were picked up. Selected colonies of actinomycetes were transferred from mixed culture of the plates onto respective agar plates and incubated at 28°C for 7 days. A single colony of each pure culture was transferred to starch nitrate agar slant for maintenance till further examinations.

In vitro antibacterial activities of actinomycetes

Determination of antibacterial activities of pure actinomycetes cultures against the soft rot causing bacteria, *E. carotovora* subsp. *carotovora*, were performed using the agar disc method. Starch-nitrate agar plates were prepared and inoculated with actinomycetes cultures and incubated at 28°C for 7 days. A disc of 5 mm diameter was cut from each actinomycetes culture, using a sterilized cork borer and placed in the center of nutrient agar plates seeded with *E. carotovora* subsp. *carotovora*. All plates were incubated at 28°C for 24 hr. The antibacterial activities were measured by determination of the diameters of the inhibition zones.

The antibacterial activity of the most active actinomycete isolate was further tested using the agar diffusion method according to Deans & Ritchie (1987). Actinomycetes isolate was grown on starch-nitrate liquid media on a rotatory shaker incubator at 200 rpm for 7 days at 28°C. The culture filtrate was sterilized through a bacteria filter and tested for its antibacterial activity as follow; 1 ml amount (10^6 cells/ml) of *E. carotovora* subsp. *carotovora* was pipetted into separate sterile Petri dishes to which 20 ml amounts of molten nutrient agar (45°C) were added. Once set, wells of 5 mm diameter were made in the center of each agar plate using sterilized cork borer, into which 100 µl sterilized culture filtrate was added. The plates were then left undisturbed to allow diffusion of the sample into the agar and then incubated at 28°C for 24 hr. Following this, zones of growth inhibition were measured.

Pathogenicity test of the most active actinomycete isolate

The selected actinomycete was examined for its pathogenicity to be sure that it does not cause any disease for potato. Pathogenicity test was carried out as previously mentioned with the use of spore suspension of actinomycete instead of bacterial suspension.

Taxonomic identification of the most active actinomycete isolate

The most active actinomycete isolate was identified according to Bergey's Manual of Determinative Bacteriology (Buchanan & Gibbons, 1974), Bergey's Manual of Systematic Bacteriology (Loëci, 1989) and the keys proposed by Shirling & Gottlieb (1968), Kuster (1972) and of Nonomura (1974). The cultural, morphological and physiological characteristics of the isolate were studied using media and methods described by Shirling & Gottlieb (1966). Micromorphology of the strain was examined microscopically using a light microscope (Carl Zeiss MC80- standard 20-DVEGS-Germany) and spores were observed by scanning electron microscopy (Jeol JSM LV 5500). Cell wall composition of the strain was analyzed by the method described by Becker *et al.* (1965) and modified by Stanek & Roberts (1974).

In vitro antibacterial activity of essential oils

Volatile oils used in this study were obtained from Badawya Company, Mansoura, Egypt. The antimicrobial activity of the volatile oils against *E. carotovora* subsp. *carotovora* were screened using an agar diffusion technique (Deans & Ritchie, 1987) as previously described.

The minimum inhibitory concentration (MIC) of the most active oil was determined using the agar dilution method according to Hammer *et al.* (1999).

In vivo antibacterial activity of Streptomyces sioyaensis and cinnamon oil

Greenhouse experiment was conducted in the winter season of 2005 to control the development of soft rot disease caused by *E. carotovora* subsp. *carotovora* using different treatments of the most active actinomycetes and the most active essential oil.

The experiment was performed in 30-cm diameter round plastic pots (7kg autoclaved soil per pot). Potato seeds of Sponta cultivar were treated with different treatments of the biocontrol agents and essential oil (see later) and planted in the pots. Potato seeds without any treatments were planted as control. Each pot was planted with a single potato seed and inoculated with bacterial suspension (10^6 cells/ml) of 24 hr old *E. carotovora* subsp. *carotovora*. Five pots were used as replicates for each treatment. Pots were arranged in a completely randomized design. Potato remained in the soil for about 100 days, before harvesting, and during which, the suitable cultural practices were applied. At the harvest time, the parameters of total number of tubers, total weight of tubers, number of diseased tubers, percentage of diseased tubers (disease incidence), and efficacy percentage were determined.

The effect of different greenhouse treatments on the development of potato bacterial soft rot during storage were carried out by storing healthy potato tubers after harvest at 22-28°C for 45 days and the values of disease incidence and efficacy percentage were estimated.

Preparation of the biocontrol agent (bioagent)

The most active isolate of the antagonistic actinomycetes was prepared in three different formulae: powder, suspension and granules. To prepare these forms, the actinomycete was grown on starch-nitrate liquid media for 7 days at 28°C. The powder form was prepared by mixing the actinomycete culture with talc powder at 1:1(v/w) and the mixture was left to dry out. The suspension formula was prepared by blending the actinomycete culture, using electric blender, for 3 min. Granules were prepared by adding 3gm of sodium alginate to 100ml of starch-nitrate liquid media and then autoclaved. Then, 100ml of actinomycete culture, grown on starch-nitrate liquid media, were added. The mixture was dropped into 3 M $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$ solution under aseptic conditions at 4°C to form granules. The alginate beads formed and entrapped actinomycetes propagules were allowed to stabilize for 1hr and dried under sterile air flow after washing with sterile distilled water.

The effect of different doses of the most active antagonistic actinomycetes on controlling soft rot was studied using granules formula in three different doses (0.5, 1.0 and 2.0gm of the most active actinomycete per pot).

Statistical analysis

The Tukey test was used for statistical analysis of the data (Tukey, 1953).

Results

Isolation and identification of potato soft rot causing bacteria

Twenty three isolates of *Erwinia* sp. were obtained from infected potato tubers showing typical soft rot symptoms. Pathogenicity test was carried out and all bacterial isolates proved to be pathogenic to potato tubers. These isolates were then subjected to morphological, cultural and biochemical studies and identified according to the

diagnostic keys of Fahy & Persley (1983), Bergey's Manual of Systematic Bacteriology (Lelliott & Dickey, 1984) and Bradbury (1986). Bacterial isolates No. 3, 6, 8, 13 and 18 were identified as *E. carotovora* subsp. *carotovora* (Badr, 2006). Regardless of the virulence of the pathogens, isolate No. (18) was used through out this study.

Isolation and in vitro screening of antibacterial activities of antagonistic actinomycetes

Fourty actinomycetes isolates were obtained from the rhizosphere soil of healthy potato plants. The actinomycetes isolates were screened for their antibacterial activities against *E. carotovora* subsp. *carotovora* using the agar disc method. Eight isolates, out of 40, exhibited inhibitory effect against *E. carotovora* subsp. *carotovora* with variable activities and actinomycete isolate No.15 showed the most pronounced antibacterial activities (Table 1).

TABLE 1. The antagonistic capacity of the active isolates of the tested actinomycetes against *E. carotovora* subsp. *carotovora* as demonstrated by diameters of inhibition zones.

Actinomycete isolate No.	Mean diameter of inhibition zones(mm)*
5	30
7	20
12	28
15	32
22	14
26	14
33	15
38	12

*The recorded value is the mean of 3 replicates.

The antibacterial activity of the cell free culture filtrate of the most active actinomycete isolate (No.15) was examined using the cup plate method and the results showed that the isolate exhibited a great antibacterial activity as indicated by measuring the diameter of inhibition zone developed (33mm).

Pathogenicity test of the most active actinomycete isolate (Streptomyces sioyaensis)

Pathogenicity test of *Streptomyces sioyaensis* showed no infection of potato tubers.

Taxonomic identification of the most active actinomycete isolate (No.15)

The actinomycete isolate No.15 formed mycelia filaments and chains of several spores. The aerial spores were not motile and were not borne on verticillate sporophores. Sugar analysis of cells showed glucose to be the main sugar, while galactose, xylose and arabinose could not be detected. The major amino acids in the cell wall hydrolyzates were glycine and L-DAP acid. Chemical analysis indicated that the isolate has cell wall type I. Chemical analysis of the cells and cell walls and morphological characteristics indicate that the isolate belongs to the genus

Streptomyces from the criteria in Bergey's Manual of Determinative Bacteriology (Buchanan & Gibbons, 1974).

The morphological, cultural, physiological and antagonistic properties of the isolate are summarized in Table 2. In general, aerial mycelium was gray to white. Sporophores were spiral (Fig. 1). The surface of conidia was smooth (Fig. 2).

TABLE 2. Morphological, cultural, physiological, and antagonistic properties of actinomycete isolate No. 15.

I. Morphological characteristics			
Sporophore (Spore chain morphology): section spirals (Fig. 1)			
Spore surface: smooth (Fig. 2)			
II. Cultural characteristics on various media			
<u>Starch-nitrate agar</u>			
Aerial mycelium: whitish grey to light grey			
Reverse side: yellowish white (not distinctive)			
Soluble pigment: none			
<u>Starch ammonium sulphate agar</u>			
Aerial mycelium: whitish grey			
Reverse side: yellowish white (not distinctive)			
Soluble pigment: none			
<u>Glycerol asparagines agar</u>			
Aerial mycelium: whitish grey			
Reverse side: yellowish white (not distinctive)			
Soluble pigment: trace of brown			
<u>Yeast-malt extract agar</u>			
Aerial mycelium: grayish white to white			
Reverse side: yellowish white (not distinctive)			
Soluble pigment: trace of brown			
<u>Oat-meal agar</u>			
Aerial mycelium: white to grayish white			
Reverse side: yellowish white (not distinctive)			
Soluble pigment: trace of brown			
<u>Czapek's solution agar</u>			
Aerial mycelium: white			
Reverse side: yellowish white (not distinctive)			
Soluble pigment: dark brown			
III. Physiological characteristics			
Melanin formation: positive on peptone-yeast extract iron agar			
Starch hydrolysis: positive			
Gelatin liquefaction: positive			
Nitrate reduction: positive			
Coagulation and peptonization of milk: positive			
Utilization of different carbon sources			
D-Glucose: +	D-galactose: +	L-arabinose: -	D-xylose: +
Raffinose: +	D-mannitol: +	D-Fructose: +	Sucrose: +
Mannose: -	Inositol: +	L-rhamnose: -	Lactose: +
IV. Antimicrobial Activity			
<i>E. coli</i> : + <i>Pseudomonas fluorescens</i> : - <i>Bacillus subtilis</i> : + <i>Staphylococcus aureus</i> : +			
<i>Alternaria alternata</i> : - <i>Fusarium oxysporium</i> : - <i>Trichoderma harzianum</i> : -			
<i>Saccharomyces cerevisiae</i> : - <i>Candida boidinii</i> : -			

From the results presented in Fig. 1 & 2 and Table 2 and following the diagnostic keys of Bergey's Manual (Buchanan & Gibbons, 1974 and Locci, 1989) and surveying the literatures of the description of *Streptomyces* sp. in the articles of ISP (Shirling & Gottlieb, 1968), Kuster (1972) and Nonomura (1974), isolate No. 15 was identified as a strain belonging to *Streptomyces siayaensis* (Nishimura *et al.*, 1961).




Fig. 1. Morphology of spore chain of isolate No. 15 (x = 1,000).



Fig. 2. Scanning electron micrograph of spore surface of isolate No. 15 (x = 15,000).

In vitro antibacterial activities of essential oils

Twenty eight commercial plant essential oils were screened for their inhibitory activities against *E. carotovora* subsp. *carotovora*. The results obtained (Table 3) showed that only 9 oils exhibited inhibitory activities against *E. carotovora* subsp. *carotovora* with variable degrees as demonstrated by measuring the diameters of inhibition zones developed by the oils and the essential oil of *Cinnamomum zeylincum* (cinnamon oil) showed the greatest activity.

TABLE 3. The inhibitory activity of the tested essential oils against *E. carotovora* subsp. *carotovora* as demonstrated by diameters of inhibition zones.

Commercial essential oil	*Mean diameter of inhibition zones(mm)	Commercial essential oil	*Mean diameter of inhibition zones(mm)
<i>Cinnamomum zeylanicum</i> Blume.	35	<i>Raphanus sativus</i> L.	0
<i>Mentha viridis</i> L.	26	<i>Allium cepa</i> L.	0
<i>Syzygium aromaticum</i> L.	25	<i>Allium sativum</i> L.	0
<i>Majorana hortensis</i> L.	22	<i>Trigonella foenum-graecum</i> L.	0
<i>Petroselinum sativum</i> L.	20	<i>Brassica nigra</i> L.	0
<i>Cuminum cyminum</i> L.	20	<i>Jasminum grandiflorum</i> L.	0
<i>Foeniculum vulgare</i> Mill.	20	<i>Anethum graveolens</i> L.	0
<i>Citrus limon</i> L.	15	<i>Rosmarinus officinalis</i> L.	0
<i>Lavandula officinalis</i> L.	14	<i>Ocimum bacilicum</i> L.	0
<i>Thymus vulgaris</i> L.	0	<i>Capsicum frutescens</i> L.	0
<i>Nigella sativa</i> L.	0	<i>Coriandrum sativum</i> L.	0
<i>Eucalyptus rostrata</i> Schlecht.	0	<i>Pinus halepensis</i> Mill.	0
<i>Eruca sativa</i> Mill.	0	<i>Daucus carota</i> L.	0
<i>Carum carvi</i> L.	0	<i>Elettaria cardamomum</i> Maton.	0

*The recorded value is the mean of 3 replicates.

The minimum inhibitory concentration (MIC) of the cinnamon oil was determined using agar dilution method and it was found to be 0.5% (v/v).

The antibacterial activity of cinnamon oil against *Streptomyces sioyaensis* was also tested using the cup plate method and no inhibition was detected.

In vivo antibacterial activities of *Streptomyces sioyaensis* and cinnamon oil

The greenhouse experiment was designed to evaluate the efficacy of different treatments of *S. sioyaensis* and cinnamon oil in controlling bacterial soft rot of potato. Disease incidence in addition to total number and total weight of

tubers/treatment were used as parameters to detect the effect of different treatments in controlling soft rot of potato.

The results presented in Tables 4 & 5 indicate that all treatments were of marked potentiality for successful control of potato soft rot; they significantly reduced disease incidence and enhanced yield comparing with the control. Cinnamon oil, *S. siوياensis* in powder formula and combination between cinnamon oil and *S. siوياensis* (powder formula) strongly inhibited (100%) soft rot caused by *E. carotovora* subsp. *carotovora* on potato tubers. *S. siوياensis* (granule formula, 1.0g) and (granule formula, 2.0g) resulted in 92.31% and 96.30%, reduction of the disease, respectively. *S. siوياensis* (suspension formula) and *S. siوياensis* (granule formula, 0.5g) prevented symptoms development at the lowest ratio, they were 88.47%.

The results in Tables 4 & 5 also indicated that *S. siوياensis* in different formulae had significant different effects on disease incidence. From this the formula with the highest effect on disease reduction was the powder formula (100%), followed by granules (96.3-88.47%), and suspension (88.47%).

TABLE 4. Effect of different greenhouse treatments on controlling bacterial soft rot.

Treatment	Total number of tubers	Total weight of tubers (g)	Number of diseased tubers	Disease incidence (%)	Efficacy (%)
Cinnamon oil	27	1134	zero	zero	100
<i>S. siوياensis</i> (powder)	27	1150	zero	zero	100
<i>S. siوياensis</i> (suspension)	26	930	3	11.53	88.47
<i>S. siوياensis</i> (granules, 0.5 g)	26	857	3	11.53	88.47
<i>S. siوياensis</i> (granules, 0.1 g)	26	996	2	7.69	92.31
<i>S. siوياensis</i> (granules, 2.0 g)	27	1044	1	3.70	96.30
Cinnamon oil + <i>S. siوياensis</i> (powder)	28	1210	zero	zero	100
Control	27	910	9	33.33	66.67

The results in Tables 4 & 5 further indicated that different doses of *S. siوياensis* have different effects on disease incidence where increasing dose of *S. siوياensis* increased its efficacy in controlling bacterial soft rot. From this, the dose with highest effect on disease development was 2g/pot of *S. siوياensis* granules (96.30%), followed by 1g/pot (92.31%), and 0.5g/pot (88.47%).

This experiment was carried out to evaluate the effect of different treatments carried out during growing season on disease incidence during storage period.

TABLE 5. Tukey test for multiple-comparison between greenhouse treatments at harvest time (significance level = 0.05)

Treatment	Cinnamon oil	<i>S.sioyaensis</i> (powder)	<i>S.sioyaensis</i> (suspension)	<i>S.sioyaensis</i> (granules, 0.5 g)	<i>S.sioyaensis</i> (granules, 1.0 g)	<i>S.sioyaensis</i> (granules, 2.0 g)	Cinnamon oil + <i>S.sioyaensis</i> (powder)	Control
Cinnamon oil	-	NS	S	S	S	S	NS	S
<i>S.sioyaensis</i> (powder)	-	-	S	S	S	S	NS	S
<i>S.sioyaensis</i> (suspension)	-	-	-	NS	S	S	S	S
<i>S.sioyaensis</i> (granules, 0.5 g)	-	-	-	-	S	S	S	S
<i>S.sioyaensis</i> (granules, 0.1 g)	-	-	-	-	-	S	S	S
<i>S.sioyaensis</i> (granules, 2.0 g)	-	-	-	-	-	-	S	S
Cinnamon oil + <i>S.sioyaensis</i> (powder)	-	-	-	-	-	-	-	S
Control	-	-	-	-	-	-	-	-

NS = non-significant difference

S = significant difference

The results presented in Tables 6 & 7 show that all treatments significantly reduced disease development during storage comparing with the control. Combination between cinnamon oil and *S. siوياensis* (powder formula) completely inhibited (100%) soft rot disease. *S. siوياensis* in powder formula and cinnamon oil had great effect in reducing disease development during storage (95.66%). *S. siوياensis* (granules, 2.0g) prevented disease development at 91.31% ratio while *S. siوياensis*(granules, 1.0g) prevented disease development at 86.96% ratio. *S. siوياensis* (suspension) and *S. siوياensis* (granules, 0.5g) prevented symptoms development at the lowest ratio, they were 82.61%.

TABLE 6. Effect of greenhouse treatments on disease development during storage.

Treatment	Total number of tubers	Number of diseased tubers	Disease incidence (%)	Efficacy (%)
Cinnamon oil	23	1	4.34	95.66
<i>S. siوياensis</i> (powder)	23	1	4.34	95.66
<i>S. siوياensis</i> (suspension)	23	4	17.39	82.61
<i>S. siوياensis</i> (granules, 0.5 g)	23	4	17.39	82.61
<i>S. siوياensis</i> (granules, 0.1 g)	23	3	13.04	86.96
<i>S. siوياensis</i> (granules, 2.0 g)	23	2	8.69	91.31
Cinnamon oil + <i>S. siوياensis</i> (powder)	23	Zero	Zero	100
Control	18	10	55.55	44.45

Different formulae of *S. siوياensis* had significant different effects on disease development during storage (Tables 6 & 7). Powder formula showed the greatest effect in reducing disease incidence (95.66%). Granules formula occupied the second rank in this respect (91.31-82.61%). Finally, suspension formula had the least effect (82.61%).

Different doses of *S. siوياensis* had significant different effects on disease development during storage, where increasing dose of *S. siوياensis* resulted in reducing disease development during storage (Tables 6 & 7). 2g/pot of *S. siوياensis* granules resulted in 91.31% reduction of the disease while 1.0g/pot of *S. siوياensis* granules resulted in 86.96% disease reduction. 0.5g/pot of *S. siوياensis* granules prevented disease development at the lowest ratio (82.61%).

Discussion

A major limiting factor in profitable potato production is disease, which can be seed-borne, soil-borne, or both. Soft rot, caused by the bacteria *E. carotovora* subsp. *carotovora*, is the most serious of all storage diseases. This organism spreads rapidly from tuber to tuber if the conditions are appropriate.

TABLE 7. Tukey test for multiple-comparison between greenhouse treatments during storage (significance level = 0.05).

Treatment	Cinnamon oil	<i>S.sioyaensis</i> (powder)	<i>S.sioyaensis</i> (suspension)	<i>S.sioyaensis</i> (granules, 0.5 g)	<i>S.sioyaensis</i> (granules, 1.0 g)	<i>S.sioyaensis</i> (granules, 2.0 g)	Cinnamon oil + <i>S.sioyaensis</i> (powder)	Control
Cinnamon oil	-	NS	S	S	S	S	S	S
<i>S.sioyaensis</i> (powder)	-	-	S	S	S	S	S	S
<i>S.sioyaensis</i> (suspension)	-	-	-	NS	S	S	S	S
<i>S.sioyaensis</i> (granules, 0.5g)	-	-	-	-	S	S	S	S
<i>S.sioyaensis</i> (granules, 0.1 g)	-	-	-	-	-	S	S	S
<i>S.sioyaensis</i> (granules, 2.0 g)	-	-	-	-	-	-	S	S
Cinnamon oil + <i>S.sioyaensis</i> (powder)	-	-	-	-	-	-	-	S
Control	-	-	-	-	-	-	-	-

NS = non-significant difference

S = significant difference

Among soil microorganisms, bacteria and fungi and to a lesser extent actinomycetes, have received considerable attention as biocontrol agents of soil borne plant pathogens. Several researchers have successfully employed antagonistic bacteria, *Streptomyces* and yeasts to control plant bacterial diseases (Alvizatos & Pantazis, 1992; Aysan *et al.*, 2003 and Ryan *et al.*, 2004). One of the alternative control methods of *E. carotovora* subsp. *carotovora* is the use of biocontrol agents. In this study, actinomycetes were developed as potential biocontrol agent for controlling soft rot caused by *E. carotovora* subsp. *carotovora* on potato in greenhouse conditions.

In this investigation, 40 isolates of actinomycetes were screened in the laboratory for their antagonistic capacity against *E. carotovora* subsp. *carotovora*. Eight isolates exhibited antagonistic activity against *E. carotovora* subsp. *carotovora*. The most active isolate, identified as *Streptomyces sioyaensis*, was tested under greenhouse conditions to control potato soft rot where it showed great reduction in disease incidence at harvest time and after storage.

Streptomyces spp. have been employed by Hayashida *et al.* (1988), Tu (1988) and Liu *et al.* (1995) as a biocontrol measure against many diseases and it was stated that, many strains of *Streptomyces* spp. produce antibacterial metabolites or antibiotics which are active against several plant pathogens.

Our results suggest that *S. sioyaensis*, a beneficial antagonist, can significantly reduce soft rot disease caused by *E. carotovora* subsp. *carotovora* on potato in greenhouse experiment and may be a useful bioagent for the disease management in Egypt.

Suitable formulation of the bacterium for its efficiency should also be investigated, in order that this beneficial bacterium may be presented to farmers for commercial usage in the future.

In our greenhouse experiment, the antagonistic *S. sioyaensis* was used in different formulae to determine the correlation between different formulae of the antagonist and its efficacy in controlling the bacterial soft rot of potato. Data obtained revealed that using the antagonist as suspension showed poor effect comparing with using the antagonist as powder or granules. This is due to that either granules or powder contain organic matter supplemented with necessary nutrient substances which improve growth of the antagonist, also act as suitable food base of the antagonist. This food base leads to well establishment of the antagonist in the infection court, causing reduction in the pathogenic propagules and consequently reduce the contamination of the produced tubers with the pathogenic bacteria. This reduction in inoculum of pathogenic bacteria translated to reduction in disease incidence under store condition (Abd El-Moity *et al.*, 1991). These results are in agreement with those obtained by Phae *et al.* (1992), Kay & Stewart (1994) and Tolba (1998).

On the other hand, powder or granules contain some nutrient substances which stimulate inhibitors production by the antagonist. These inhibitors play an important role on stability of the formula and also improve establishment of antagonist (Bull *et al.*, 1991).

Effect of different doses of the antagonistic *S. sioyaensis* on disease incidence was also studied. Data obtained showed a positive correlation between dose and reduction in disease incidence at harvest time and after storage. This is due to that increasing inoculum leads to increasing the efficacy of the antagonist which consequently reduce the inoculum of pathogenic bacteria. These results are in agreement with those of Kloepper (1983), Abd El-Moity *et al.* (1991) and Tolba (1998).

The antimicrobial properties of plant volatile oils from a wide variety of plants have been reported in numerous investigations (Hammer *et al.*, 1999; Imai *et al.*, 2001 and Guynot *et al.*, 2003).

In this investigation, twenty eight commercial plant essential oils were screened for their inhibitory activities against *E. carotovora* subsp. *carotovora*. Nine essential oils exhibited antibacterial activities and the cinnamon oil had the greatest activity. The great antibacterial activity of cinnamon oil showed in this investigation is in agreement with many other investigations (Boyd & Gouk, 1997 and Baratta *et al.*, 1998).

Treatment of potato tubers with cinnamon oil prior to sowing showed great reduction in disease incidence at harvest time (0%) or after storage (4.34%).

Combination between *S. sioyaensis* in powder formula and cinnamon oil resulted in complete reduction (disease incidence 0%) of soft rot of potato tubers at harvest time and after storage.

In conclusion, the results of this study confirmed the possibility of using *S. sioyaensis* or cinnamon oil in controlling bacterial soft rot of potato. It is more useful to use them in combination.

References

- Abd El-Moity, T. H., El-Deeb, A. A. and Radwan, I. A. (1991) Biological control of seedling disease and pod rot of peanut, under greenhouse and field conditions. *Egypt. J. Appl. Sci.* 6 (1), 103.
- Alivizatos, A. S. and Pantazis, S. (1992) Preliminary studies on biological control of potato common scab caused by *Sreptomyces* sp. In: "Biological Control of Plant Diseases Progress and Challenges for the Future", Tjamos, E.C., Papavizas, G.C., Cook, R. J. (Ed.), Vol. 230, pp. 85-93, NATO ASI Series, Kluwer Academic Publishers, Dordrecht.

- Atila, A. (1989) Antimicrobial activity of black cumin (*Nigella sativa L.*). *Dept. Food Sci., Fac. Agric., Atatürk Univ.* 6 (1), 63.
- Aysan, Y., Karatas, A. and Cinar, O. (2003) Biological control of bacterial stem rot caused by *Erwinia chrysanthemi* on tomato. *Crop Protection*, 22, 807-811.
- Badr, H. H. E. (2006) Integrated control of potato soft rot in Dakahlyia Governorate. *M.Sc. Thesis*, Botany Department, Faculty of Science, Mansoura University, Mansoura, Egypt.
- Baratta, M.T., Dorman, H.J.D., Deans, S.G., Figueiredo, A.C., Barroso, J.G. and Ruberto, G. (1998) Antimicrobial and antioxidant properties of some commercial essential oils. *Flav. Fragr. J.* 13(4),235-244.
- Barnejee, A. and Nigam, S. S. (1978) Antimicrobial efficacy of the essential oil of *Curcuma longa*. *Indian J. Med. Res.* 68, 864-866.
- Becker, B., Lechevalier, M. P. and Lechevalier, H. A.(1965) Chemical composition of cell wall preparations from strains of various form-genera of aerobic actinomycetes. *Appl. Microbiol.* 13, 236-243.
- Boyd, R. J. and Gouk, S. C. (1997) Inhibitory activities of essential oils against plant pathogenic bacteria, *Hort. Research*, Ruakura Research Center, Private Bag 3123, Hamilton.
- Bradbury, J. F. (1986) "*Guide to Plant Pathogenic Bacteria*", pp.61-73, C.A.B. International, UK.
- Bradbury, J. F. (1977) *CMI Distribution Maps of Plant Disease* No. 138(5th ed.) CAB International, Oxford (GB).
- Buchanan, R. E. and Gibbons, N. E. (1974) "*Bergey's Manual of Determinative Bacteriology*", 8th ed., Williams and Wilkins, Co. Baltimore, USA.
- Bull, C. T., Weller, D. H. and Thomashow, L. S. (1991) Relationship between root colonization and suppression of *Gaeumannomyces graminis tritici* by *Pseudomonas fluorescens* strain 2-79. *Phytopathol.* 81,954.
- Deans, S. G. and Ritchie, G. A. (1987) Antimicrobial properties of plant essential oils. *International J. Food Microbiol.* 5, 165-180.
- Deans, S. G. and Sviboda, K. (1990) The antimicrobial properties of majoram (*Origanum majorana L.*) volatile oil. *Flavour and Fragrance J.* 5(3),187-190.
- Fahy, P.C. and Persley, G. J. (1983) "*Plant Bacterial Diseases: A Diagnostic Guide*", Academic Press. Sydney, New York, London.

- Guynot, M. E., Ramos, A. J., Seto, L., Purroy, P., Sanchis, V. and Marin, S. (2003) Antifungal activity of volatile compounds generated by essential oils against fungi commonly causing deterioration of bakery products. *J. Appl. Microbiol.* **94**(5),893.
- Hammer, K. A., Carson, C. F. and Riley, T.V. (1999) Antimicrobial activity of essential oils and other plant extracts, *J. Appl. Microbiol.* **86**, 985-990.
- Hayashida, S., Choi, M. Y., Nanri, N. and Miyaguchi, M.(1988) Production of potato common scab-antagonistic biofertilizer from swine feces with *Streptomyces albidoflavus*. *Agric. Biol. Chem.* **52**, 2397-2402.
- Hellas, V., Andrivon, D. and Jouan, B. (2000) Development of symptoms caused by *Erwinia carotovora* spp. *atroseptica* under field conditions and their effects on the yield of individual potato plants. *Plant Pathology*, **49**(1),23-32.
- Ibrahim, S. S., El-Samahy, S. K., Abd-El-Fadeal, M. G., Awad, K. and Abd-El-Hamid, A. A.(1987) The antimicrobial effect of some components of henna leaves (*Lawsonia inermis*). I. *The antibacterial effect 2nd Nat. Conf. of Pests and Dis. of Veg. and Fruits*. Ismailia, Oct. 1987, pp.768-786.
- Imai, H., Osawa, K., Yasuda, H., Hamashima, H., Arai, T. and Sasatsu, M. (2001) Inhibition by the essential oils of peppermint and spearmint of the growth of pathogenic bacteria. *Microbios.* **1**, 31-39.
- Johnson, I. F.,Curl, E. A., Bond, J. H. and Fibourg, H. A.(1960) *Methods for Studying Soil Microflora*. Burgess Publishing Co., p.15, Minneapolis, USA.
- Kastelein, P., Schepel, E. G., Mulder, A., Turkensteen, L. J., Vuurde, J. W. and Van, L. (1999) Preliminary selection of antagonists of *Erwinia carotovora* subsp. *atroseptica* (Van Hall) dye for application during green crop lifting of seed potato tubers. *Potato Research*, **42**(2), 161-171.
- Kay, S. J. and Stewart, A. (1994) Evaluation of fungal antagonists for control of onion white rot in soil box trials. *Plant Pathol.* **43** (2),371-377.
- Klopper, J. W. (1983) Effect of seed piece inoculation with plant growth-promoting rhizobacteria on populations of *Erwinia carotovora* on potato roots and in daughter tubers. *Phytopathol.* **73**,217-219.
- Kuster, E. (1972) Simple working key for classification and identification of named taxa included in the international *Streptomyces* project. *Intern. Assoc. Microbiol. Soc.* **22**, 139-148.
- Lelliott, R. A. and Dickey, R. S. (1984) Genus VII. *Erwinia*, "*Bergey's Manual of Systematic Bacteriology*", Williams & Wilkins Company, Vol 1, pp.469-476, Baltimore.
- Liu, D., Anderson, N. A. and Kinkehl, L. L.(1995) Biological control of potato scab in the field with antagonistic *Streptomyces scabies*. *Phytopathology*, **85**, 827-831.

- Locci, R. (1989) *Streptomyces* and related Genera, "Bergey's Manual of Systematic Bacteriology", Williams & Wilkins Company, Vol. 4, pp. 2451-2508, Baltimore.
- Lucon, C. M. M. and De Melo, I. S. (1999) Selection of antagonistic rhizobacteria to *Erwinia carotovora* subsp. *atroseptica* on potato tubers. *Summa Phytopathological*, 25(2), 132-136.
- Lund, B. M. (1979) Bacterial soft rot of potatoes. In: "Plant Pathogens", Lovelock, p. 19-49.
- Nishimura, H., Okamoto, S., Mayama, M., Ohtsuka, H., Nakajima, K., Tawara, K., Shinohira, M. and Shimaoka, N. (1961) Siomycin, a new thiostrepton-like antibiotic, *J. antibiotics, Tokyo, Ser. A*, 14, 255-263.
- Nonomura, H. (1974) Key for classification and identification of 458 species of *Streptomyces* included in ISP. *J. Ferment. Technol.* 52(2), 78-92.
- Phae, C. G., Shoda, M., Kita, N., Nokano, M. and Ushiyama, K. (1992) Biological control of crown and root rot and bacterial wilt of tomato by *Bacillus subtilis* NB22. *Annals Phytopathol. Soc. Jap.*, 58(3), 329-339.
- Roberti, R. and Selmi, C. (1999) Biological control of plant pathogens by *Bacillus subtilis*. *Informatore Fitopatologica*, 49(7/8), 15-21.
- Ryan, A. D., Kinckel, L. L. and Schottel, J. L. (2004) Effect of pathogen isolate, potato cultivar, and antagonist strain on potato scab severity and biological control. *Biocontrol Science and Technology*, 14(3), 301-311.
- Sharga, B.M. and Lyon, G. D. (1998) *Bacillus subtilis* BS 107 as an antagonist of potato blackleg and soft rot bacteria. *Can. J. Microbiol.* 44, 777-783.
- Shirling, E. B. and Gottlieb, D. (1966) Methods for characterization of *Streptomyces* species, *Intern. J. System. Bacteriol.* 16 (3), 313-340.
- Shirling, E. B. and Gottlieb, D. (1968) Cooperative description of type cultures of *Streptomyces*. II. Species description from first study. *Intern. J. System. Bacteriol.* 18(2), 69-189.
- Singh, S. P., Negi, S., Chand, L. and Singh, A. K. (1992) Antimicrobial properties of essential oils from caryophyllanthemum and Zingiber leaves and rhizomes. *Fitoterapia*, 63(1), 73-75.
- Staneck, J. L. and Roberts, G. D. (1974) Simplified approach to identification of aerobic actinomycetes by thin-layer chromatography. *Applied Microbiology*, 31, 702-706.
- Tawfik, A. E., Hanna, A. I., El-Ghareeb, L. A., Gomah, A. A. and Mahmoud, S. M. (2001) Applied approach for controlling brown rot and soft rot bacteria of potatoes. *J. Agric. Sci. Mansoura Univ.* 26(6), 3631-3642.
- Egypt. J. Microbiol.* 43 (2008)

- Tolba, A. F. (1998)** Biological control of some potato bacterial diseases, *Ph.D. Thesis*, Fac. Agric., Minufiya Univ., Egypt.
- Tu, J. C. (1988)** Antibiosis of *Streptomyces griseus* against *Colletotrichum lindamuthianum*. *J. Phytopathol.* **121**,97-102.
- Tukey, J. W. (1953)** Some selected quick and easy methods of statistical analysis. *Acad. Sci.* **16**(2),88-97.
- Zedan, A. M. (1993)** Antifungal properties of certain plant extracts with special reference to the possibility of controlling onion white rot disease using *Eucalyptus robusta* leaves. *Egypt. J. Appl. Sci.* **8** (12), 574.

(Received 16/5/2007;
accepted 18/2/2008)

مكافحة مرض العفن الطري في البطاطس باستخدام بكتريا استريبتومييسيس سيوينسيس وزيت القرقة

فتحي عواد منصور ، عطية حامد محمدين، عادل الصادق إسماعيل و هدى
حسين بدر
قسم النبات- كلية العلوم- جامعة المنصورة- المنصورة والمعهد بحوث أمراض
النباتات- مركز البحوث الزراعية- القاهرة- مصر.

مرض العفن الطري هو أحد أهم الأمراض البكتيرية التي تصيب محصول
البطاطس و الذي تسببه بكتريا *ايرونييا كاروتوفورا* تحت النوع *كاروتوفورا*. تم
دراسة امكانية استخدام مكافحة الحيوية لمقاومة هذا المرض باستخدام
الاكتينومييسيتات. تم اختيار أربعين عزلة من الاكتينومييسيتات، جمعت من أماكن
مختلفة من محافظة الدقهلية، لتقييم كفاءتها التضادية ضد *ايرونييا كاروتوفورا*
كاروتوفورا تحت ظروف المعمل و قد أبدى ثماني عزلات منها نشاط تضادي
و عرف أفضلها على أنه *استريبتومييسيس سيوينسيس*. على الجانب الآخر تم اختيار
ثمانية و عشرين زيت من الزيوت العطرية لتقييم تأثيرها ضد بكتيري على
ايرونييا كاروتوفورا *كاروتوفورا* تحت ظروف المعمل، أبدت تسعة زيوت منها
نشاط تضادي و قد كان زيت القرقة هو أفضل الزيوت المختبرة. اختبر كائن
استريبتومييسيس سيوينسيس و زيت القرقة لمقاومة العفن الطري في البطاطس في
الصوبة، و قد وجد أن معاملة درنات البطاطس قبل الزراعة بكائن *استريبتومييسيس*
سيوينسيس و زيت القرقة كلا على حده أدى إلى اختزال كبير في نسبة حدوث
المرض وقت الحصاد و بعد التخزين و استخدامهما معا كان له أفضل النتائج. أدى
استخدام *استريبتومييسيس سيوينسيس* بصور مختلفة (معلق – مسحوق – حبيبات)
إلى تأثيرات مختلفة على نسبة حدوث المرض وقت الحصاد و بعد التخزين ، حيث
كان استخدامه في صورة مسحوق له أفضل النتائج. كما وجد أن زيادة الجرعة
المستخدمة من *استريبتومييسيس سيوينسيس* أدت إلى زيادة تأثيره على اختزال نسبة
حدوث المرض .