

EFFECT OF DIETARY METABOLIZABLE ENERGY AND MICROBIAL PHYTASE LEVELS ON BROILER PERFORMANCE, NUTRIENTS DIGESTIBILITY AND MINERALS UTILIZATION

by

Abdallah A. Ghazalah* and Mohamed A. Alsaady**

*Anim. Prod. Dept. , Fac. Agric. , Cairo University, Egypt.

** Anim. Prod. Dept., Fac. Agric. , University of Damascus. Syria.

Received: 3/8/2008

Accepted: 18/8/2008

ABSTRACT: *An experiment designed in a 2 x 4 factorial arrangement was conducted to study the effect of using different levels of metabolizable energy (3100 and 2900 kcal/kg starter diets and 3200 and 3000 kcal/kg grower diets) , along with different levels of phytase (0, 500, 750 and 1000 FTU/kg diet) in broiler diets on their growth performance , nutrients digestibility, minerals utilization and economical efficiency . Diets were formulated to be isonitrogenous , being contain 21% and 18% CP during starter and grower periods, respectively . A total number of 240 one-day old Arbor Acres broiler chicks were randomly and equally divided into 8 groups , each in three replicates . The experimental birds were reared under similar managerial and veterinarial conditions and offered feed and water ad libitum up to 42 days of age . At this age , a digestion trial was carried out to study the effect of dietary treatments on nutrients digestibility , blood constituents and tibia analysis*

The obtained data showed that the group of chicks fed high ME diets supplemented with 750 or 1000 FTU/kg recorded heavier live body weight (LBW) and body weight gain (BWG) values than those fed the corresponding low ME diets. However, phytase addition at 750 or 1000 FTU/kg gave similar LBW and BWG values and were higher ($P \leq 0.05$) than the control group . Although, the dietary ME combined with phytase additions did not significantly affect the amount of feed intake , however, the group of chicks fed high energy level supplemented with 750 FTU/kg of phytase recorded ($P \leq 0.05$) the best feed conversion while those fed the phytase – free low energy diets had recorded the worst value . The interaction between dietary energy level and phytase supplementation significantly affected both ether extract and crude fiber digestibility . In addition, such interaction significantly affected tibia weight and its calcium

content, while tends to decrease ($P \geq 0.05$) the activity of alkaline phytase of plasma. As a result to the former findings, the group of chicks fed the high energy diets supplemented with phytase at 750 FTU/kg recorded the lowest feed cost needed to obtain one kilogram of body weight gain compared to the control. Therefore, from the economical and nutritional points of view, it is concluded that phytase addition at 750 FTU/kg to adequate energy diets is more valuable and economical.

INTRODUCTION

Most of poultry feed ingredients are from plant origin which contains some components that are hardly digested by monogasterics because of the lack of endogenous enzyme secretions. These compounds also decrease the utilization of other dietary nutrients, leading to depress performance. Phytic acid is one of these compounds that may cause various problems due to its mostly presence in cereals, oil seeds and their by-products (Cromwell *et al.*, 1993). Therefore, it is well documented that microbial phytase supplementation enhances phytate hydrolysis and increases the availability of nutrients bound to the phytic molecule (Sebastian *et al.*, 1997). As a result, phytase addition to broiler diets can improve body weight and feed utilization

(Waldroup *et al.*, 2000; Desouky, 2001; Lan *et al.*, 2002; Wu *et al.*, 2003; Abd El-Hakim and Abd Elsamee, 2004; El-Ghamry *et al.*, 2005 and Ghazalah *et al.*, 2006).

It is well known that poultry tend to eat to satisfy their energy needs, because energy is necessary for providing the body with heat needed for maintenance and doing many physiological functions (Attia *et al.*, 2001; Shaapan, 2004 and Ramadan, 2005). Studies reported that energy utilization could be improved by phytase addition into broiler diets, that may be attributed to liberation of calcium ions necessary for α -amylase activity which is involved in starch digestion (Ravindran *et al.*, 1999; Kies *et al.*, 2001; Attia, 2003a,b). It has been postulated also that calcium phytate may increase the formation of metallic soaps in the gut lumen, with a corresponding reduction in the utilization of energy from saturated fats (Ravindran *et al.*, 2000).

Due to this confliction and bearing in mind the economic aspects, this study aimed to study the interaction effect of dietary energy and phytase levels in view of broiler performance, nutrients digestibility, blood and tibia analysis.

MATERIALS AND METHODS

This experiment was conducted to study the effect of using two energy levels being high

(HME) and low (LME) , each with four levels of microbial phytase in 2 x 4 factorial arrangement on growth , feed utilization , nutrients digestibility , tibia analysis and economical efficiency of broiler chicks. Energy levels applied were 3100 and 2900 kcal ME/kg during starter (0 – 3 wks) and 3200 and 3000 kcal ME/kg during grower (4 – 6 wks) periods representing high and low energy levels , respectively. Phytase levels supplemented to each energy level were 0 , 500 , 750 and 1000 FTU/kg . Diets were formulated to contain 21 and 18 % crude protein for the starter and grower periods , respectively, being less than the strain recommendation in order to benefit from phytase addition . The composition and calculated analysis of the experimental diets (without phytase supplementation) are presented in Table (1) . Accordingly , a total number of 240 one-day old Arbor Acres broiler chicks were randomly and equally divided into 8 groups , each in three replicates . The experimental birds were reared under similar managerial and veterinarial conditions and offered feed and water *ad libitum* up to 42 days of age . At this age , a digestion trial was carried out to study the effect of dietary treatments on nutrients digestibility , blood constituents and tibia analysis . The criteria of growth performance in terms of live body weight (LBW) and feed intake (FI) were recorded per period in order to calculate body weight gain (BWG) and feed conversion (FC). Digestibility of nutrients were carried out using individual three birds from each treatment in which feed intake and excreta voided were recorded along 3 days collection period . The collected excreta were sprayed by 2% boric acid solution to prevent any loss in ammonia, then dried in an oven at 60 °C for 24 hours, thenafter weighed, finely ground and kept for chemical analysis according to AOAC (1990) methods. Fecal nitrogen was determined in dried excreta according to the method of Jakobsen *et al.* (1960). At the end of the experiment, four birds from each treatment were randomly taken , fasted for about 12 hours and slaughtered . Blood samples were taken from the slaughtered birds and plasma was separated by centrifugation at 3000 r.p.m/15 minutes and assigned for subsequent determination of calcium , phosphorus and alkaline phosphatase activity using Sigma procedures No. 586, 360-UV and 104, respectively, Sigma Diagnostics, St. Louis, Mo., USA. The left tibia was also removed, cleaned from all adhering tissues, fat extracted, dried , weighed and then ashed for determination of calcium (Lehman and Henry, 1984) and phosphorus (Tietz, 1986). Finally, the experimental treatments were economically

evaluated based upon the price of local market at the year 2008 , in terms of the feed cost needed to obtain one kilogram of live body weight gain as described by Ghazalah *et al.* (2005).

The obtained data were subjected to analysis of variance using the general model of SAS software (SAS,1990) and treatment means were compared using Duncan's new multiple range test (Duncan,1955).

RESULTS AND DISCUSSION

Growth performance :

Data presented in Table (2) indicate the main effects of either dietary energy level or phytase supplementation on growth performance of broilers. At 6 weeks of age, the results cleared that chicks fed HME diets recorded significantly ($P \leq 0.05$) higher live body weight (LBW) and body weight gain values than those fed LME diets . Although, no significant differences were found in feed intake (FI) between chicks fed HME and those given LME diets , however, the former had recorded significantly ($P \leq 0.01$) better feed conversion (FC) than the other group which fed LME diets. Regarding the phytase level, no significant differences were detected in all previous parameters within treatments (Table 2) . Data of the chicken growth performance in terms of LBW, BWG, FI and FC as affected by the interaction between dietary energy level and phytase supplementation are presented in Table (3). The initial LBW for all treatments was nearly similar (about 49 g) which may create a suitable condition to appraise the effect of dietary treatments during the subsequent period. However, at 42 days of age, significant differences ($P \leq 0.05$) had been observed with regard to LBW and BWG values among treatments. Moreover, supplementing broiler diets with microbial phytase either at 750 or 1000 FTU/kg surpassed those groups which fed diets supplemented with microbial phytase at 500 FTU/kg or without phytase supplementation . At 6 weeks of age, the addition of microbial phytase to the HME diets at 500, 750 and 1000 FTU/kg increased LBW by 2.05, 6.35 and 7.77% and BWG by 2.20, 6.51 and 8.02%, respectively, compared to their corresponding control diet without phytase supplementation. While with LME diets, the improve in LBW and BWG at 6 weeks of age amounted to 2.85, 7.83 and 7.26% for LBW and 2.98, 8.09 and 7.5 % for BWG , respectively compared to the control diet .These findings declared the adequacy of phytase addition to broiler diets up to 750 FTU/kg diet.

As was evidenced in Table (3), the interaction between dietary energy and phytase levels did not significantly affect the amount of feed intake, however, the group of chicks fed the HME

diets supplemented with 750 FTU/kg recorded ($P \leq 0.01$) the best feed conversion (1.91). In addition, no significant differences in feed conversion values were detected when phytase was supplemented at 1000 FTU/kg either to HME or LME diets , compared to the corresponding control diets, confirming the adequacy of 750 FTU/kg to be used in broiler diets. On the other hand, the worst feed conversion value was obtained by the group of chicks fed the LME diets without phytase supplementation, being 2.1 . In this connection, Johnston and Southern (2000) reported that phytase supplementation into broiler diets did not affect their feed consumption, while improved feed/ gain ratio. Attia *et al.*, (2001) found that phytase addition to high energy diets of broilers resulted in the best feed conversion value. Such improvement in feed conversion of corn-soybean meal based diets may be attributed to an increase in absorbed phosphorus (Desouky, 2001; Lan *et al.*, 2002), release of other minerals affecting feed utilization(El -Deeb *et al.*, 2000) and to the increase in nutrients digestibility(Camden,2001) .

It is worthy to note that no mortality due to any of the experimental treatments were observed during the all period of feeding . Therefore, all the experimental chicks were in a good health recording 100% viability.

Nutrients digestibility :

Data of nutrients digestibility of the experimental diets as affected by the interaction between dietary energy and phytase levels are presented in Table (4). It is worthy to note the values of dry matter ratio (dry matter excreta / dry matter fed) were nearly similar and ranged between 0.21 and 0.24 indicating the similarity in feeding value among the dietary treatments. The interaction effect appeared significantly only on ether extract (EE) and crude fiber (CF) digestibility. In this respect, Attia *et al.*, (2001) observed a significant improvement in CF digestibility with phytase addition to broiler diets, which was explained by most of phytic acid located within cell walls. Moreover, the positive effect on EE digestibility was in agreement with the findings of Kies and Van Hermert (2000); Attia *et al.*, (2001); Lan *et al.*, (2002) and Shirely and Edwards (2003) who stated that phytase may prevent the formation of insoluble metallic soaps in the gastrointestinal tract, which may improve lipid utilization of the diets. On the other hand, no significant effects were observed among dietary treatments regarding crude protein (CP), nitrogen free extract (NFE) and organic matter (OM) digestibility. This could be explained based upon the experimental diets which were isonitrogenous and their contents of all the nutrients were similar either at starter or grower periods, respectively.

Blood and Tibia analysis :

The effect of dietary energy and phytase levels on plasma content of calcium , phosphorus and alkaline phosphatase activity at 6 weeks of age are shown in Table(5). Results of plasma calcium and phosphorus content were nearly similar with no significant differences within all the treatments. However, it seems from the data obtained that whatever the dietary energy level (HME or LME) , phytase supplementation tend to decrease significantly the activity of alkaline phosphatase . This can be explained by increasing homeostatic balance between phosphorus and calcium by increasing the absorbed phosphorus in order to elevate plasma phosphorus content , which lead to decrease the activity of alkaline phosphatase in blood. These results are in agreement with the findings of Huff *et al.*, (1998) who reported that the decreasing in plasma alkaline phosphatase activity associated with the diets supplemented with microbial phytase might reflect the down regulation of this enzyme resulting from the increasing availability of phosphorus. Similarly, Lan *et al.*, (2002) showed that there was no significant effect of using phytase on the average values of plasma calcium , while such addition of phytase may lower the alkaline phosphatase activity.

The effect of interaction between dietary ME and phytase levels on tibia analysis are shown also in Table(5). The obtained data nearly indicated similar values of tibia ash and phosphorus content of tibia among all the dietary treatments. In this respect, Ahmed *et al.*, (2000); Lan *et al.*, (2002) and El-Medany and El-Afifi (2002) found that phytase supplementation to broiler diets had no effect on tibia ash percent. However, the results showed that when microbial phytase had been added at 750 FTU/kg either to HME or LME broiler diets, both tibia weight and calcium content were significantly increased . This confirmed the effective role of phytase in releasing calcium from phytate complex. In this connection, Viveros *et al.*, (2002); Shirley and Edwards (2003); Augspurger and Baker (2004) and Onyango *et al.*,(2004) reported that phytase addition to broiler diets revealed significant increase in tibia ash and minerals content compared to those unsupplemented.

Economical Efficiency :

It is worthy to note that the experimental diets were adjusted to be similar in all nutrients except the ME level. According to guide lines of economical evaluation, the production costs include chicks price, feed cost, medicine and medical care, body weight gain, labor cost, fuel, electricity, depreciation cost for equipments and other miscellaneous items. However, in this study, all the formentioned items are similar per chick within the

different treatments except for the feed cost and the final live body weight that had been affected either positively or negatively by treatments. Therefore, the economical efficiency of the present study could be calculated based mainly upon the total feeding cost and gain in live body weight. Results of the economical evaluation as affected by the dietary treatments are illustrated in Table (6). The group of chicks fed the HME diet supplemented with 750 FTU/kg (T₃) recorded the lowest feed cost needed to obtain one kg of BWG(4.497 L.E), while those fed the LME diet without phytase supplementation (T₅) had gave the worst value(4.912 L.E). Compared to the control group of each energy set, the relative economical value of T₃ (HME + 750FTU/kg) surpassed the other treatments having the HME set. While, the group of chicks of T₇ (LME +750FTU/kg) and T₈ (LME + 1000FTU/kg) surpassed the other treatments of LME set. However, assuming T₁ (HME phytase free) is a general control, the best economical value had been obtained by T₃ (HME + 750 FTU/kg) which is better by 4.0% than the control. However, the all other treatments except T₄ (HME +1000 FTU) gave economically lower values than the control. This means that feeding chicks HME diets with added microbial phytase at 750FTU/kg is more successful in view of growth , feed utilization and economical evaluation of broiler chicks.

Table (1) : Composition and calculated analysis of the experimental diets

Ingredients (%)	Starter (0 – 3 weeks)		Grower (4 – 6 weeks)	
	Yellow corn	61.66	60.34	68.15
Soybean meal 44%	24.16	23.00	18.95	24.30
Corn gluten meal 60%	7.80	7.85	5.84	2.00
Vegetable oil	2.01	0.05	2.89	0.59
Wheat bran	0.50	4.90	0.50	0.50
Dicalcium phosphate	1.75	1.75	1.65	1.61
Limestone	1.15	1.13	1.10	1.09
Vit. & Min. mixture ¹	0.30	0.30	0.30	0.30
NaCl	0.35	0.35	0.35	0.35
DL- methionine	0.12	0.12	0.09	0.12
L-lysine HCl	0.20	0.21	0.18	0.05
Total	100	100	100	100
Calculated analysis ² :				
ME kcal / kg	3100	2900	3200	3000
CP %	21.02	21.12	18.00	18.02
Ca %	0.90	0.90	0.85	0.85
Av. Phos %	0.45	0.45	0.42	0.42
Meth. %	0.49	0.49	0.41	0.42
Meth. + Cyst. %	0.86	0.86	0.74	0.74
Lys. %	1.06	1.06	0.90	0.90

⁽¹⁾ Supplied per kg of diet : Vit.A, 12000 IU; Vit.D₃, 2200IU; Vit.E, 10mg; Vit.K, 2mg; Vit.B₁, 1mg; Vit.B₂, 5mg; Vit.B₆, 1.5mg; Vit.B₁₂, 10µg; Nicotinic acid, 30mg; Folic acid, 1mg; Pantothenic acid, 10mg; Biotin, 50µg; Choline chloride, 250mg; Copper, 10mg; Iron, 30mg; Manganese, 60mg; Zinc, 50mg; Iodine, 1mg; Selenium, 0.1mg and Cobalt, 0.1mg.

⁽²⁾ According to NRC (1994).

Table (2) : Main effects of dietary energy and phytase supplementation on live body weight (LBW) , body weight gain (BWG) , feed intake (FI) and feed conversion (FC) of broilers.

Item	Age (wk)	Energy level		Sig.	Phytase level (FTU/kg)				Sig.
		HME	LME		0.0	500	750	1000	
LBW	6	2129 ^a	2016 ^b	*	1988	2037	2129	2138	NS
BWG	0 – 6	2079 ^a	1966 ^b	*	1938	1987	2079	2088	NS
FI	0 – 6	4069	4056	NS	3944	4045	4106	4157	NS
FC	0 - 6	1.96 ^b	2.07 ^a	**	2.04	2.04	1.98	1.99	NS

a...b Means within row with different superscripts are significantly different .

* = significant (P<0.05) ** = significant (P<0.01) NS = not significant

Table (3): Effect of ME x phytase interaction on live body weight (LBW), body weight gain (BWG), feed intake (FI) and feed conversion (FC, feed /gain) of broilers.

Tr. No.	ME Kcal/kg	Phytase FTU/kg	LBW , g at day old	LBW , g at 6 weeks	BWG , g 0 - 6 weeks	FI , g 0 - 6 weeks	FC 0 - 6 weeks
T ₁	HME	zero	49.4 ± 0.03	2046 ^b ± 34.3	1996 ^b ± 34.4	3942 ± 45.6	1.97 ^{ab} ± 0.01
T ₂	HME	500	48.8 ± 0.21	2088 ^{ab} ± 26.1	2040 ^{ab} ± 26.3	4030 ± 66.6	1.97 ^{ab} ± 0.01
T ₃	HME	750	49.3 ± 0.03	2176 ^a ± 80.1	2126 ^a ± 80.1	4068 ± 83.2	1.91 ^a ± 0.03
T ₄	HME	1000	49.3 ± 0.20	2205 ^a ± 61.1	2156 ^a ± 61.3	4236 ± 113.7	1.96 ^{ab} ± 0.01
T ₅	LME	zero	49.5 ± 0.05	1929 ^c ± 66.1	1879 ^c ± 66.1	3944 ± 95.8	2.10 ^c ± 0.02
T ₆	LME	500	49.2 ± 0.21	1984 ^{bc} ± 41.4	1935 ^{bc} ± 41.3	4059 ± 79.6	2.09 ^c ± 0.02
T ₇	LME	750	49.1 ± 0.20	2080 ^{ab} ± 95.3	2031 ^{ab} ± 95.3	4143 ± 115.6	2.04 ^{bc} ± 0.06
T ₈	LME	1000	49.3 ± 0.06	2069 ^{ab} ± 46.7	2020 ^{ab} ± 46.6	4078 ± 51.7	2.02 ^{abc} ± 0.05
Sign.			NS	*	*	NS	**

a ... c : Means within the same column with different superscripts are significantly different ,

* = significant (P ≤ 0.05) NS = not significant HME : High metabolizable energy

** = significant (P ≤ 0.01) LME : Low metabolizable energy .

Table (4): Effect of ME x phytase interaction on nutrients digestibility of the experimental diets.

Treat No.	ME Kcal/kg	Phytase FTU/kg	DMR	Digestibility of nutrients (%)				
				CP	EE	CF	NFE	OM
T ₁	HME	zero	0.23	73.37	89.67 ^a	18.33 ^c	89.33	83.33
T ₂	HME	500	0.21	71.15	90.33 ^a	19.33 ^{abc}	89.33	82.67
T ₃	HME	750	0.24	74.33	89.67 ^a	26.33 ^{ab}	87.67	82.00
T ₄	HME	1000	0.21	69.87	89.00 ^{ab}	18.33 ^{bc}	88.33	81.67
T ₅	LME	zero	0.23	67.49	82.67 ^c	25.00 ^{ab}	88.33	80.67
T ₆	LME	500	0.21	69.85	83.33 ^c	27.67 ^{ab}	89.00	82.00
T ₇	LME	750	0.22	72.92	81.33 ^c	29.33 ^a	89.33	83.00
T ₈	LME	1000	0.21	72.17	85.33 ^{bc}	25.00 ^{ab}	90.33	83.00
Sign.			NS	NS	*	*	NS	NS

ac : Means in the same column with different superscripts are significantly different ,

NS = not significant * = significant (P ≤ 0.05)

HME : High metabolizable energy

LME : Low metabolizable energy

DMR : Dry matter ratio CP : Crude protein EE : Ether extract

CF : Crude fiber NFE : Nitrogen free extract OM : Organic matter

Table (5): Effect of ME x phytase interaction on plasma constituents and tibia analysis of the experimental broilers .

Treat No.	ME Kcal/kg	Phytase FTU/kg	Plasma constituents			Tibia analysis		
			Ca (mg/100ml)	P (mg/100ml)	Alk. Phos. (U/L)	Weight (g)	Ash (%)	Ca (%)
T ₁	HME	zero	12.69	16.00	3.87 ^a	8.83 ^{bc}	42.19	39.97 ^{bc}
T ₂	HME	500	13.43	16.99	3.31 ^{ab}	8.89 ^{abc}	43.35	40.16 ^{bc}
T ₃	HME	750	13.10	16.52	2.72 ^c	9.12 ^a	44.01	41.97 ^a
T ₄	HME	1000	11.23	15.37	2.73 ^c	9.13 ^a	44.02	41.74 ^a
T ₅	LME	zero	13.95	14.84	4.09 ^a	8.72 ^c	41.74	39.77 ^c
T ₆	LME	500	13.66	16.05	3.01 ^b	8.84 ^{bc}	42.69	41.14 ^{ab}
T ₇	LME	750	12.44	15.58	2.84 ^{bc}	9.04 ^{ab}	43.57	42.09 ^a
T ₈	LME	1000	13.30	14.95	2.72 ^c	9.01 ^{ab}	43.60	41.77 ^a
Sign.			NS	NS	*	*	NS	*

ac : Means in the same column with different superscripts are significantly different ,

NS = not significant * = significant (P ≤ 0.05)

HME : High metabolizable energy

LME : Low metabolizable energy

Ca : Calcium P : Phosphorus

Alk. Phos. : Alkaline Phosphatase

Table (6) :Economic evaluation of the experimental treatments (ME x phytase interaction) .

Treat No.	ME Kcal/kg	Phytase FTU/kg	FI / bird Kg / 6 wks	Feed cost L.E / bird	BWG/bird Kg / 6 wks	Feed cost / Kg BWG	Relative feed cost / kg BWG	
T ₁	HME	zero	3.942	9.35	1.996	4.680	100 ^a	100 ^{aa}
T ₂	HME	500	4.030	9.62	2.040	4.720	100.8	100.8
T ₃	HME	750	4.068	9.56	2.126	4.497	96.0	96.0
T ₄	HME	1000	4.236	9.89	2.156	4.587	98.0	98.0
T ₅	LME	zero	3.944	9.23	1.879	4.912	100 ^b	104.9
T ₆	LME	500	4.059	9.40	1.935	4.858	98.9	103.8
T ₇	LME	750	4.143	9.56	2.031	4.707	95.8	100.6
T ₈	LME	1000	4.078	9.48	2.020	4.693	95.5	100.3

HME : High metabolizable energy

LME : Low metabolizable energy

FI : Feed intake

BWG: Body weight gain

a : Assuming that T₁ is the control of HME set experimental diets

b : Assuming that T₅ is the control of LME set experimental diets

aa : Considering T₁ as a general control for all the experimental diets

REFERENCES

- Abd El-Hakim, A. S. and M. O. Abd El-Samee (2004).** *Effect of feeding systems and phytase supplementation on the performance of broiler chicks during summer season. Worlds Poultry Congress, 8 – 13 June, Istanbul, Turkey.*
- Ahmed, T. S.; M. Rasool; M. Sarwar; U. Ahson and Z. U. Hasan(2000).** *Effect of microbial phytase produced from a fungus Aspergillus niger on bioavailability of phosphorus and calcium in broiler chickens. Anim. Feed Sci. Tech., 83 : 103 – 114.*
- Association of Official Analytical Chemists (1990), A. O. A. C. Official Methods of Analysis, 15th ed. Published by the A.O.A.C., Washington, D.C.**
- Attia, Y. A. (2003a).** *Responses of growth performance, carcass characteristics, meat quality and plasma constituents of male Campbell ducks to dietary levels of methionine and phytase and their interactions. Egypt. Poult. Sci. J., 23 (III) : 557 – 580.*
- Attia, Y. A. (2003b).** *Performance, carcass characteristics, meat quality and plasma constituents of meat type ducks fed diets containing different levels of lysine with or without a microbial phytase. Arch. Anim. Nutr., 57 : 39 – 48.*
- Attia, Y. A.; S. A. Abd El-Rahman and E. M. Qota (2001).** *Effects of microbial phytase without or with cell- wall splitting enzymes on the performance of broiler fed marginal levels of dietary protein and metabolizable energy. Egypt. Poult. Sci. J., 21(II): 521– 547.*
- Augspurger, N. R. and D. H. Baker (2004).** *High dietary phytase levels maximize phytate-phosphorus utilization but not affect protein utilization in chicks fed phosphorus or amino acid- deficient diets. J. Anim. Sci., 82: 1100 – 1107.c*
- Camden, B. J. (2001).** *Effectiveness of exogenous microbial phytase in improving the bioavailability of phosphorus and other nutrients in maize- soybean meal diets for broilers. J. Anim. Sci., 79: 289 – 297.*
- Cromwell, G. L.; T. S. Stahly; R. D. Coffey; H. J. Monegue and J. H. Rondolph (1993).** *Efficacy of phytase in improving the bioavailability of phosphorus in soybean meal and corn-soybean meal diets for pigs. J. Anim. Sci., 71 : 1831 – 1840.*

- Desouky, Zeinab, M. A. (2001).** *Effect of supplemental microbial phytase on productive performance of broiler chickens. Ph. D Thesis, Faculty of Agric. Ain Shams Univ., Egypt.*
- Duncan, D. B. (1955).** *Multiple Ranges and Multiple F- Tests. Biometrics, 11: 1 – 42.*
- El-Deeb, Mariam, A. ; H. Sharara and M. N. Makled (2000).** *Enhance calcium and Phosphorus utilization by enzyme phytase supplemented to broiler diet containing rice bran. Egypt. Poult. Sci. J., 20: 546 – 566.*
- El-Ghamry, A. A.; M. A. Al-Harthi and Y.A. Attia (2005).** *Possibility to improve rice polishing utilization in broiler diets by enzymes or dietary formulation based on digestible amino acids. Archiv. Fur Gelfugelkunde, 69: 49 – 56.*
- El-Medany, N. M. and S. F. El-Afifi (2002).** *The effect of microbial phytase on utilization of protein and phosphorus in broiler rations. Egypt. Poult. Sci. J., 22: 427 – 443.*
- Ghazalah, A. A.; A. H. Abd El-Gawad; M. S. Soliman and Amany W. Youssef (2005).** *Effect of enzyme preparation on performance of broilers fed corn-soybean meal based diets. Egypt. Poult. Sci. J., 25: 295 – 316.*
- Ghazalah, A. A.; M. O. Abd-Elsamee; M. A. El-Manyalawi and Eman S. Moustafa (2006).** *Response of broiler chicks to microbial phytase supplementation in diets differ in available phosphorus sources and levels. Egypt. Poult. Sci. J., 26: 1321 – 1341.*
- Huff, W. E.; P. A. Moore, Jr.; G. R. Huff; J. M. Balog; N. C. Rath; P. W. Waldroup; A. L. Waldroup; T. C. Daniel and V. Raboy (1998).** *Reducing inorganic phosphorus in broiler diets with high available-P-corn and phytase without affecting broiler performance or health. Poult. Sci., 77: 1899 – 1904.*
- Jakobsen, P. E.; S. G. Kirston and H. Nelson (1960).** *Digestibility trials with poultry. 322 bertning fraforsgs laboratoriet, udgivet of stants. Husdyrbugsud valy-Kabenhaven.*
- Johnston, S. L. and L. L. Southern (2000).** *The effect of varying mix uniformity (simulated) of phytase on growth performance, mineral retention and bone mineralization in chicks. Poult. Sci., 79: 1485 – 1490.*

Metabolizable Energy , Phytase , Broiler , Growth , Feed Utilization , Tibia .

- Kies, A. K. and K. H. F. Van Hemert (2000).** *Phytase: a remarkable enzyme, in: selected topics in animal nutrition, biochemistry and physiology. Reviews presented at the symposium on the occasion of the retirement of Dr. R. R. Marquardt, W. C. Sauer and J. He. EDS., PP. 119 – 136 Winnipeg, Edmonton, Canada.*
- Kies, A. K.; K. H. F. Van Hemert and W. C. Sauer (2001).** *Effect of phytase on protein and amino acid digestibility and energy utilization. World's Poult. Sci., 57: 109 – 125.*
- Lan, G. Q.; N. Abdallah; S. Jalaludin and Y. W. Ho (2002).** *Efficacy of supplementation of phytase producing bacterial culture on the performance and nutrient use of broiler chickens fed corn soybean meal diets. Poult. Sci., 81: 1522 – 1532.*
- Lehman, H. P. and J. P. Henry (1984).** *Calcium determination. In clinical diagnosis and management by laboratory methods. 17th ed. J. P. Henry, Edsunders, Philadelphia, PP.1431 – 1438.*
- National Research Council , NRC (1994).** *Nutrient Requirements of Poultry. 9th Rev. ed. National Academy Press, Washington, DC.*
- Onyango, E. M. ; M. R. Bedford and O. Adeola (2004).** *The yeast production system in which Echerichia coli phytase is expressed may affect growth performance, bone ash and nutrient use in broiler chicks. Poultry Sci., 83 : 421 – 427 .*
- Ramadan, Nehad A. (2005).** *Broiler performance as affected by dietary energy source and level. Ph.D Thesis, Fac. Agric., Cairo Univ., Egypt.*
- Ravindran, V.; S. Cabahug; G. Ravindran and W. L. Bryden (1999).** *Influence of microbial phytase on apparent ileal amino acid digestibility of feedstuffs for broilers. Poult. Sci., 78: 699 – 706.*
- Ravindran, V.; S. Cabahug; P. H. Selle and W. L. Bryden (2000).** *Response of broiler chickens to microbial phytase supplementation as influenced by dietary phytic acid and non- phytate phosphorus levels. II. Effects on apparent metabolizable energy, nutrient digestibility and nutrient retention. Br. Poult. Sci., 41: 266 – 274.*
- SAS, Institute (1990).** *SAS/STAT User's guide: Statistics, 8eds. SAS Institute INC., Cary NC., USA.*
- Sebastian, S,; S. P. Touchburn; E. R. Chavez and P. C. Lague (1997).** *Apparent digestibility of protein and amino acids in broiler*

chickens fed a corn-soybean meal diets supplemented with microbial phytase. Poult. Sci., 76: 1760 – 1769.

Shaapan, M. H. M. (2004). *Response of broiler performance to dietary microbial phytase, energy levels, amino acids and phosphorus sources. Ph.D Thesis, Fac. Agric., Cairo Univ., Egypt.*

Shirley, R. B. and H. M. Edwards, Jr. (2003). *Graded levels of phytase past industry standards improves broiler performance. Poult. Sci., 82: 671 – 680.*

Tietz, N. W. (1986). *Textbook of Clinical Chemistry. N. B. Saunder Co., London, Philadelphia, 18.1.*

Viveros, V.; A. Bernes; I. Arija and C. Centano (2002). *Effect of microbial phytase supplementation on mineral utilization and serum enzyme activities in broiler chicks fed different levels of phosphorus. Poult. Sci., 81: 1172 – 1183.*

Waldroup, P. W.; J. H. Kersey; E. A. Saleh; C. A. Fritts; F. Yatf; H. L. Stilborri; J. R. Crum and V. Raboy(2000). *Non-phytate phosphorus requirement and phosphorus excretion of broiler chicks fed diets composed of normal or high available phosphate-corn with or without microbial phytase. Poult. Sci., 79: 1451 – 1459.*

Wu, Y. B.; V. Ravindran and W. H. Hendriks (2003). *Effect of microbial phytase produced by solid-state fermentation, on the performance and nutrient utilization of broilers fed maize and wheat based diets. Br. Poult Sci., 44: 710 – 718.*

الملخص العربي

تأثير استخدام مستويات مختلفة من الطاقة الممتلئة وإنزيم الفيتيز الميكروبي

على الأداء الإنتاجي لحجاج اللحم و معاملات هضم المركبات الغذائية

والإستفادة من العناصر المعدنية

عبدالله على غزاله* - محمد أيمن السعدى *

*قسم الإنتاج الحيوانى - كلية الزراعة - جامعة القاهرة - مصر

* قسم الإنتاج الحيوانى - كلية الزراعة - جامعة دمشق - سوريا

أجريت تجربة بتصميم إحصائى 2×4 لدراسة تأثير استخدام مستويات مختلفة من الطاقة الممتلئة (٣١٠٠ ، ٢٩٠٠ ك.كالورى / كجم من علائق البادئ) و (٣٢٠٠ ، ٣٠٠٠ ك.كالورى / كجم من علائق النامى) كل مع أربع مستويات من إنزيم الفيتيز الميكروبي (صفر ، ٥٠٠ ، ٧٥٠ ، ١٠٠٠ وحدة إنزيم / كجم عليقة) على الأداء الإنتاجي ، معاملات هضم المركبات الغذائية ، الإستفادة من العناصر المعدنية والكفاءة الإقتصادية لكتاكيت اللحم . تكونت علائق التجربة بحيث تحتوى على ٢١% ، ١٨% من البروتين الخام فى مرحلتى البادئ (عمر يوم - ٣ أسابيع) والنامى (٤ - ٦ أسابيع) على الترتيب وهى أقل قليلا من إحتياجات السلالة وذلك للإستفادة من إضافة إنزيم الفيتيز . أستخدم فى التجربة عدد ٢٤٠ ككتوت أربور إيكروز عمر يوم قسمت عشوائيا وبالتساوى إلى ٨ معاملات كل منها فى ٣ مكررات . ربيت الكتاكيت تحت نفس ظروف الرعاية مع تقديم الغذاء والماء بصورة حرة حتى عمر ٤٢ يوم . وفى نهاية التجربة أجريت تجربة هضم لدراسة تأثير المعاملات الغذائية على معاملات هضم المرذبات الغذائية وبعض مكونات بلازما الدم وتحليل عظمة الساق . تم وزن الكتاكيت والغذاء المستهلك أسبوعيا لحساب الزيادة فى الوزن والكفاءة التحويلية للغذاء .

أظهرت النتائج المتحصل عليها أن الطيور المغذاة على العلائق العالية فى مستوى الطاقة الممتلئة مع إضافة إنزيم الفيتيز الميكروبي بمستويات ٧٥٠ أو ١٠٠٠ وحدة إنزيم / كجم قد سجلت قيمسا أعلى لكل من وزن اللحم والزيادة فى وزن الجسم مقارنة بالطيور التى غذيت على العلائق المقابلة المحتوية على مستوى الطاقة المنخفض ، ومع ذلك فقد أعطى الفيتيز بمستويات ٧٥٠ ، ١٠٠٠ وحدة إنزيم / كجم قيمة متشابهة ولكنها أعلى (غير معنوية) من الكنتترول. لم يكن

لمستوى الطاقة الممتلئة مصحوبا بإضافة إنزيم الفيتيز أى تأثير معنوى على الغذاء المستهلك ورغم ذلك عندما أضيف الفيتيز الميكروبي بمستوى ٧٥٠ وحدة إنزيم / كجم من العليقة المحتوية على المستوى العالى من الطاقة الممتلئة سجلت الطيور معنويا أفضل كفاءة تحويلية للغذاء ، بينما سجلت اقل كفاءة تحويلية للغذاء بواسطة الطيور المغذاة على العلائق ذات المستوى المنخفض من الطاقة الممتلئة بدون إضافة الفيتيز . كان للتداخل بين مستوى الطاقة وإضافة الفيتيز تأثيرا معنويا على معامل هضم كل من الألياف الخام ومستخلص الإثير ، وزن عظمة الساق ومحتواها من الكالسيوم بينما أدى هذا التداخل إلى خفض نشاط إنزيم الفوسفاتيز القلوى فى بلازما الدم . وتبعاً للنتائج السابقة فقد سجلت الكتاكيت المغذاة على العلائق المرتفعة فى مستوى الطاقة الممتلئة مضافا إليها الفيتيز الميكروبي بمستوى ٧٥٠ وحدة إنزيم / كجم عليقة أقل تكلفة للغذاء اللازم للحصول على كيلوجرام من الزيادة فى وزن الجسم الحى . و من هذه الدراسة يتضح من الناحية الإقتصادية أنه لا جدوى من إضافة الفيتيز الميكروبي لعلائق كتاكيت التسمين المنخفضة فى محتواها من الطاقة الممتلئة ومن الأفضل غذائيا وإقتصاديا إضافة الفيتيز الميكروبي للعلائق المتزنة فى محتواها من الطاقة .