

EFFECT OF EXTENDER TYPE ON FREEZABILITY OF RAM SEMEN

A.F.Nebar, H.Abdel Rahman, I.M. Seddik and M.A. El-Ghonamy

Animal Production Department, Faculty of Agriculture, Menofiya University.

(Received: May 20, 2008)

ABSTRACT: Semen was collected from 4 adult Ossimi rams (2-4 years) by artificial vagina. Ejaculates having $\geq 80\%$ motility and sperm concentration of $\geq 3 \times 10^9/ml$ were pooled together and gently mixed to study the effect of selected five extenders on freezability of ram spermatozoa by split-sample technique. Two extenders (A and B) were free of skim milk and three extenders (C, D and E) contained skim milk at various levels (35, 50 and 65%, respectively). The final glycerol level was 14.5% and the dilution rate was 1:6 (v/v). Semen was equilibrated for 3 h at 4-5 °C. The frozen semen was thawed in water bath at 42 °C for 35 seconds. Sperm motility, livability and abnormality were evaluated just before and after glycerolation, after equilibration and post-thawing of frozen semen.

Results indicated that extenders A and B have yielded significantly ($P < 0.01$) the highest percentage of post-thawing motility (35.6 and 42.5%), livability (37.0 and 42.1%), and the least sperm abnormality (23.8 and 19.7%), as compared to 31.3, 23.1 and 21.3% for motility, 31.9, 25.4 and 23.3% for livability and 21.51, 24.99 and 26.74% for abnormality of extenders C, D and E, respectively. Post-thawing motility and viability progressively decreased as the skim milk content in the extender increased, while the post-thawing abnormality showed an opposite trend. The differences were significant ($P < 0.01$). B-extender maintained superior post-thawing recovery rate of ram spermatozoa, being 58.6% for motility and 58.2% for livability relative to equilibrated values and 46.5 for motility and 46.1% for livability relative to initial values. Total abnormality increment was 17.68%. The least respective values for motility and viability and the higher value for abnormality were expressed by the E-extender, which contained the greatest skim milk. The loss in motility between equilibration and thawing ranged from 28.8 to 36.9% and the least values (5.6-8.7%) were obtained between glycerolation and equilibration. The corresponding values were 28.5-33.4% and 5.5-9.5% for livability and 12.37-15.72 % and 1.68-2.78% for abnormality, respectively. In conclusion, tris-based extenders have higher freezability for ram semen as compared to skim milk extenders.

Key words: Sheep, semen, freezing, motility.

INTRODUCTION

Successful manipulation of artificial insemination through applying frozen semen is unfortunately still being beyond optimum. Improvement of such technology would accelerate the genetic improvement in sheep. One of the important factors influencing frozen storage of semen is the composition of the extender used for dilution of semen before freezing (Salamon and Maxwell, 2000). However, with adaptation of the early tested ram extenders, which were based on utilization of citrate buffer combined with monosaccharide (glucose), the results had some limitations and most have been modified later (Salamon and Maxwell, 1995). Milk is an isotonic medium containing many components favorable to the maintenance of sperm viability. Milk was used extensively by several investigators for liquid and frozen ram semen, mostly combined with fructose, sodium citrate or tris (Nebar, 1989; Söderquist *et al.*, 1997; El-Alamy and Foote, 2001; Gil *et al.*, 2003 and Paulenz *et al.*, 2003). Moreover, Jones and Martin (1965) found that egg yolk extenders were inferior to skim milk extenders for freezing and *in vitro* incubation of ram spermatozoa after thawing. The efficiency of tris and milk extenders over egg yolk – citrate and egg yolk glucose to maintain ram spermatozoa was reported by Deka and Rao (1980). Similarly, Nebar (1989) found that tris-based extenders maintained, in general progressive motility and acrosome integrity of ram frozen spermatozoa over skim milk – based extenders. Gil *et al.* (2000) recommended the utilization of the milk-based extenders for freezing of ram spermatozoa with low concentration of egg yolk (5–10%) and addition of glycerol at 5°C on two-step dilution method and 2 h-equilibrated period. The present work aimed to study the effect of five different extenders on freezability of ram spermatozoa, based on the percentage of motility, livability and abnormality of spermatozoa at various stages of freezing and thawing processing.

MATERIALS AND METHODS

Semen was collected twice weekly from 4 adult healthy Ossimi rams (2-4 years) by artificial vagina at the Experimental Farm, Department of Animal Production, Faculty of Agriculture Minufiya University. Semen was evaluated from the rams for two weeks prior to the initiation of the experiment. Rams were fed a balanced ration meeting the NRC requirements for adult rams (NRC, 1987).

Immediately after semen collection, every ejaculate was evaluated for color, motility and sperm concentration. Ejaculates having 80% or more progressive motility and sperm concentration of not less than 3×10^9 /ml were pooled together and gently mixed. The pooled semen sample was split into 5 aliquots to study the effect of the selected five extenders on freezability of ram semen. The composition of these extenders is shown in Table 1.

Effect of extender type on freezability of ram semen

Table 1: Composition of semen extenders.

Component	Tris-based extenders		Skim milk-tris-based extenders		
	A	B	C	D	E
Tris, g	2.422	2.422	1.574	1.211	0.848
Citric acid, g	1.340	1.340	0.871	0.670	0.469
Na ₃ - Citrate, g	-	0.145	0.094	0.073	0.051
Fructose, g	1.000	0.750	0.111	0.085	0.060
Egg yolk, ml	20.0	20.0	13.0	10.0	7.0
Skim milk, ml	-	-	35	50	65

Tris: Hydroxymethyl aminomethane.

- Distilled water was added up for 100 ml to extenders A, B, and which mixed with skim milk (C, D and E).
- Antibiotics were added as 75 mg penicillin plus 50 mg streptomycin

Each semen extender was divided into two equal fractions namely; fraction I being free of glycerol and fraction II which was glycerolated. Glycerol was added to fraction II at a level of 29 %. In the meantime, the pooled semen sample was divided into five equal aliquots, and each aliquot was partially diluted by adding three volumes of fraction I of one of the selected extenders, so that the preliminary dilution rate was 1: 3 (volume/volume). The temperature of both semen and fraction I was 30-32 °C at such partial dilution. Then the partially diluted semen was gradually cooled to 4-5 °C over 1.5 h. Glycerolation was carried out at 5 °C by adding an equal volume of the appropriate glycerolated extender (fraction II) to the unglycerolated fraction I of diluted semen in 3 parts over an interval of 15 minutes. The final glycerol concentration after adding fraction II was 14.5% and the final dilution rate was 1:6 (volume/volume). Thereafter, the glycerolated semen was equilibrated at 4 - 5 °C for 3 hours. After equilibration, the semen was packed in the 0.5 ml French plastic straws, which were sealed using polyvinyl powder. Three straws for each extender were prepared, labeled and coded. The cold straws were mounted on metal rack and frozen in ethyl alcohol to -16.5 °C over 4 minutes and the freezing rate is shown in Table (2). Thereafter, the straws were dipped in the liquid nitrogen (-196°C) and stored for 24 hours. The frozen semen was thawed in water bath at 42 °C for 35 seconds.

Table 2: Preliminary freezing rate of the filled straws in ethyl alcohol and time spent to reach the subzero temperature -16.5 °C.

Temp., °C	zero	-4.0	-0.7	-0.9	-10.5	-12.0	-13.5	-15.0	-16.5
Time, min	0	0.5	1.0	1.5	2.0	2.5	3.0	3.5	4.0

The percentages of sperm motility, livability and abnormality were determined at the following stages during semen processing:

1. Fresh semen: after pooling satisfactory ram ejaculates and gently mixed (Initial motility, livability or abnormality).
2. After glycerolation (G).
3. After equilibration (E).
4. Post thawing (PT).

Reduction in the percentage of sperm motility and livability during the various stages was calculated as follow:

Stage 1: The difference between the % of initial motility or livability and % of the motility or livability after glycerolation = (I-G).

Stage 2: The difference between the % of motility or livability after glycerolation and % of the motility or livability after equilibration = (G-E).

Stage 3: The difference between the % of motility or livability after equilibration and the % of motility or livability post thawing = (E-PT).

Total reduction: the difference between the % of initial motility or livability and the % of motility or livability post thawing = (I-PT).

All the processing steps and the previous evaluations were replicated eight times. The obtained results were statistically analyzed according to Snedecor and Cochran (1980).

RESULTS AND DISCUSSION

Data listed in Table (3) indicate that the type of extender significantly ($P < 0.01$) affected the percentage of sperm motility and livability. In this respect, the tris-citric acid-fructose-egg yolk-glycerol-based extenders (B and A extenders) yielded significantly ($P < 0.05$) higher post-thawing motility (42.5 and 35.6%, respectively) and livability (42.1 and 37.0, respectively) than that resulted after extension with skim milk-tris-citric acid-based extenders (C, D and E, being 31.3, 23.1 and 21.3% for motility and 31.9, 25.4 and 23.3% for livability, respectively; Table 3). The superiority of B-extender in maintaining post thawing motility (42.5%) and livability (42.1%) of spermatozoa than A extender (35.6 and 37.0%, respectively) could be due to difference in its fructose content (1.0 g in A vs. 0.75 g in B) and sodium citrate existing only in B-extender but was not present in A extender (Table 1). It means that addition of sodium citrate beside the slight decrease in fructose level of the tris based egg yolk-glycerol extender seems to improve the post-thawing motility and livability. In agreement with the present differences among different types of extenders, Paulenz *et al.* (2002) found that ram spermatozoa diluted in tris based extender showed higher sperm motility and membrane integrity than those diluted in milk based extender. In the same time, Foote and Kaproth (2002) indicated that percentage of motile bull sperm significantly ($P < 0.01$) increased in response to the addition of 1.0% fructose to whole milk-glycerol diluent. In nearly similarity with the present results, El-Maghraby (2007)

Effect of extender type on freezability of ram semen

revealed that the post thawing sperm motility of goat semen diluted with tris-based extender was 42.3%.

Table (3): Effect of extender type on percentage of motility, livability and abnormality of ram spermatozoa at the various stages of freezing and thawing.

Stage of semen processing	Extender type				
	Tris-based extender		Tris-skim milk based extender		
	A	B	C	D	E
Sperm motility (%) :					
Unglycerolated (UG)	84.4 ^a	85.0 ^a	80.0 ^b	72.5 ^c	71.3 ^c
Glycerolated (G)	73.1 ^b	78.1 ^a	71.9 ^b	66.3 ^c	64.4 ^c
Equilibration (E)	64.4 ^b	72.5 ^a	65.0 ^b	60.0 ^c	56.0 ^d
Post thawing (PT)	35.6 ^b	42.5 ^a	31.3 ^c	23.1 ^d	21.3 ^d
Sperm livability (%) :					
Unglycerolated (UG)	85.8 ^a	84.6 ^a	78.9 ^b	71.8 ^c	69.4 ^c
Glycerolated (G)	75.0 ^{ab}	78.0 ^a	71.2 ^b	64.6 ^c	61.3 ^c
Equilibration (E)	65.5 ^b	72.3 ^a	64.9 ^{bc}	58.8 ^c	55.8 ^c
Post thawing (PT)	37.0 ^b	42.1 ^a	31.9 ^c	25.4 ^d	23.3 ^d
Sperm abnormality (%) :					
Unglycerolated (UG)	2.85 ^a	3.04 ^a	3.07 ^b	3.47 ^c	3.5 ^c
Glycerolated (G)	6.68 ^b	5.48 ^a	7.46 ^b	8.01 ^c	8.24 ^c
Equilibration (E)	8.82 ^b	7.22 ^a	9.14 ^b	9.66 ^c	11.02 ^d
Post thawing (PT)	23.83 ^b	19.68 ^a	21.51 ^c	24.99 ^d	26.74 ^d

Initial motility: 91.3 % Initial livability: 91.4 % Initial abnormality: 2.0 %
a, b, c, d: Values within the same row with different superscripts are significantly different (P<0.05).

It is worthy noting that, the post-thawing sperm motility and livability recorded for the extenders C, D and E progressively decreased as the skim milk increased and the citric acid, tris, sodium citrate, fructose and egg yolk decreased in the diluent (Table 1). Accordingly, it was not possible to isolate effect(s) of constituents of these diluents contents on such differences among these extenders. However, the differences in both egg yolk and skim milk content among such extenders are much more pronounced than that of the other constituents (Table 1). Thus the significant differences in percentage of post- thawing motility and livability detected in the present study among C, D, and E extenders may be attributed to differences in skim milk and/or egg yolk level rather than due to other constituents existed in such extenders.

Sansone *et al.* (2000) reported that majority of workers used egg yolk concentration around 20%. However, Gil *et al.* (2003) diluted ram semen with milk-based extenders containing four different levels of egg yolk (5, 10, 15, 20%), but they did not found significant positive effect of increasing the level of egg yolk above 10% on post-thawing motility. They added that the preparations containing 10% egg yolk gave significantly better post-thawing membrane integrity. In an earlier study, Sahní and Mohan (1990) found that egg yolk beyond 5% in tris-glycerol based freezing diluent did not show any significant improvement in respecting with the post-thaw sperm motility. However, their study clearly showed the scope of reduction of egg yolk level from 20% to 5% without any adverse effect on the freezability of buffalo semen. Kumar *et al.* (1992) reported that addition of egg yolk to the diluents stimulated an enzymatic system of the spermatozoa in addition to its protective action against cold shock. The confliction in respecting with the observed effects of addition of egg yolk at various levels on post-thaw sperm motility among the mentioned studies could be attributed to the interaction of the other components of the extender with the protective effect of egg yolk.

Akhtar *et al.* (1990) found that the tris-based extender was superior to milk diluent as determined by post-thawing motility, which agrees with the present study. An opposite trend was reported by Galli *et al.* (1993) who found that the milk-based extender was better than the tris-based extenders in respecting with the post-thawing sperm motility. In fact, the present study used the skim milk combined with tris-citrate buffer which may explain the confliction of the present finding and that of Galli *et al.* (1993).

Data in Tables (3 and 4) indicate that the sperm abnormality expressed an opposite trend to both motility and livability of ram spermatozoa with the advancement of processing stages. In other words, dilution, cooling, equilibration, freezing and thawing substantially increased the percentage of sperm abnormality. B-extender showed the least ($P < 0.05$) percentage of sperm abnormality (Table 3). The skim milk-tris based extender (C-extender) maintained abnormality at significantly lower percentage than D- and E-extenders (Table 3). Farag *et al.* (1983) reported that the percentage of sperm coiled tail greatly increased in frozen ram semen and it was correlated with increasing of Zn influx.. A similar relationship was early achieved by Blom and Wolstrup (1976). Their finding may suggest differences in such effects depended on the extenders components.

Data in Table (4) indicate that the percentage of motile ram spermatozoa progressively decreased with the advancement of semen processing. In this respect, reduction in sperm motility was the greatest (28.8-36.9%) during stage 3 due to freezing and thawing, followed by that occurred during stage 1 (due to dilution and cooling), which varied from 13.2 to 26.9%. The least values of motility loss were recorded (5.6 to 8.7%) during equilibration (stage 2). Aisen

Effect of extender type on freezability of ram semen

et al. (2000) found that the percentage of sperm motility of ram was 78.8 after cooling and 58.3% after thawing, when they used tris-citric acid-fructose-egg yolk glycerol extender. This finding is almost similar with that obtained in the present study at cooling and glycerolated but it was greater at post-thaw than that obtained in the present study. Such differences in the results may be due to differences in the extender components, glycerol level or the freezing methods.

Table (4): Reduction of motility and livability and Increment of abnormality of ram spermatozoa during freezing process.

Extender	Stages of reduction				Recovery rate (%)	
	1 (I-G)	2 (G-E)	3 (E-PT)	Total	Initial	Equilibrium
Reduction of motility:						
A	18.20	8.70	28.80	55.70	39.0	55.3
B	13.20	5.60	30.00	48.80	46.5	58.6
C	19.40	6.90	33.70	60.00	34.3	48.2
D	25.00	6.30	36.90	68.20	25.3	38.5
E	26.90	8.40	34.70	70.00	23.3	38.0
Reduction of livability:						
A	16.40	9.50	28.50	54.40	40.5	56.5
B	13.40	5.70	30.20	49.30	46.1	58.2
C	20.20	6.30	33.00	59.50	34.9	49.2
D	26.80	5.80	33.40	66.00	27.8	43.2
E	30.10	5.50	32.50	68.10	25.5	41.8
Increment of abnormality:						
	(G-I)	(E-G)	(PT-E)	Total		
A	4.68	2.14	15.01	21.83		
B	3.48	1.74	12.46	17.68		
C	5.46	1.68	12.37	19.51		
D	6.01	1.65	15.33	22.99		
E	6.24	2.78	15.72	24.74		

Initial motility: 91.3 %

Initial livability: 91.4 %

Initial abnormality: 2.0 %

Results further illustrate that the total depression of motility was the least for tris-citrate based extender either for B-extender (48.8%) or for A-extender (55.7%) compared with skim milk-tris-citrate based extenders, being 60%, 68.2% and 70.0% for C, D and E extenders, respectively. It is of interest to note that the superiority of extenders A and B in maintaining sperm motility was associated with the least depression of motility during all freezing stages (Total reduction in sperm motility). So, it could be concluded that, tris-citrate based extender (free of skim milk) provided the spermatozoa with more suitable protection and environment against the effect of dilution, cooling, equilibration, freezing and thawing than that provided by skim milk-tris-citrate based extenders (C, D and E).

The post-thaw recovery rate of motility was the best for B-extender being 46.5% in the proportion to the initial motility in fresh semen and 58.6% in the proportional to the motility just after equilibration. The corresponding figures for C-extender were 34.3% and 48.2%, respectively (Table 4). At the same time, the percentage of sperm livability decreased considerably with the advancement of semen processing, but at variable rates. The greatest rate of depression of sperm livability occurred during freezing and thawing (from 28.5 to 33.4%), followed by that during dilution and cooling (13.4 to 30.1%) and the least depression action was due to equilibration (from 5.5 to 9.5%). Values of recovery rate of post-thawing viable ram spermatozoa are very close to the corresponding values of sperm motility (Tables 3 and 4), which suggest close association of both traits as reported by Abdel-Khalek *et al.* (2008).

Vishwanath and Shannon (2000) reported that even with the best preservation techniques and all the development occurred in semen processing, the best sperm recovery post thawing is just over 50%, which agrees with the present recovery rate in proportional to sperm motility in equilibrated semen, but it is greater than the recovery rate in proportional to the initial motility. On the other hand, the post-thaw recovery rate of motility basis of data demonstrated by Aisen *et al.* (2000) was 74% in proportional to motility after cooling.

El-Alamy and Foote (2001) compared the effect of three different egg yolk ratios (i.e., 20, 30 and 40% v/v) in tris based ram extender on post-thaw motility, and they found it to be 41, 38 and 20% for Finn ram spermatozoa versus 47, 41 and 27% for Dorset ram spermatozoa, respectively. It means that the excess addition of egg yolk to tris-based extender at rates over 20% (v/v) adversely affected the post-thaw motility of ram spermatozoa. However, the recovery rate was 71, 65 and 36% for Finn ram semen versus 76, 66 and 43% for Dorset ram semen, when it was computed as percentage of initial motility (58 and 62%) for the two breeds, respectively.

In general, the depression in sperm motility occurred during semen processing may have been due to: 1- A decrease in concentrations of naturally existing seminal plasma components. 2- Disturbance in proportional

Effect of extender type on freezability of ram semen

the various electrolytes in the diluted semen, relative to that in the seminal plasma and/or addition of substances having toxic and detrimental action on sperm during semen dilution. 3- Possible effect of cold shock to which some sperms are proportionally subjected during both cooling and freezing in addition to thawing rate (e.g., thawing temperature and thawing duration). A similar assumption was also suggested by EL-Gaafary (1990). Salamon and Maxwell (1995), Maxwell and Watson (1996) and Watson (2000) demonstrated that cold shock starts soon after the semen is cooled to room temperature, although it is more critical for ram semen at temperature below 15°C (Fiser and Fairfull, 1986). It is clear from this investigation that tris-based extenders have higher freezability for ram semen as compared to skim milk extenders based on post thawing sperm motility, livability and abnormality .

REFERENCES

- Abdel-Khalek, A.; M.B. Abou-Ela; Soheir A. Fawzy and E. Dandoush (2008). An overview of filtration of semen of cows and buffaloes by sephadex column. Saudi J. Biolo. Sci., 15(1):212-221.
- Akhtar, T.; R.A. Chaudhry; L.H. Khan and T.M. Khan (1990). Extracellular release of hyaluronidase and acrosin from buffalo bull spermatozoa extended in different extenders. Recent Adv. Buffalo Res. 3: 75 – 79.
- Aisen, E.G.; H.L. Alvarez; A. Venturino and J.J. Garde (2000). Effect of trehalose EDTA on cryoprotective action of ram semen diluents . Theriogenology; 53: 1053 – 1061.
- Blom, E.; and C. Wolstrup (1976). Zinc as possible causal factor in the sterilizing sperm tail defect in Jersey bulls. VIII th Int. Cong. Anim. Reprod. And A.I. cracow, July 12-16.
- Deka, B.C. and A.R. Rao (1980). Effect of extenders on the biometrics of ram sperm head. Indian Vet. J. 57: 905 – 908.
- El-Alamy, M.A. and R.H. Foote (2001). Freezability of spermatozoa from Finn and Dorest rams in multiple semen extenders . Anim. Reprod. Sci, 65: 245 – 254.
- El-Gaafary, M.N. (1990). A diluent for freezing of ram semen . Indian. J. Anim. Sci, 60: 769 – 772.
- El-Maghraby, M.M.I. (2007). Physiological studies on reproduction in goat. Ph.D. Thesis Fact. Agric. Mansoura Univ. Egypt.
- Farag, R.S.; S.T. El-Aassar; H. El-Okash; A.A. Mohamed and A.A. El-Sharabassy (1983). Mineral changes in ram spermatozoa as affected by freezing and various extenders . Egypt. J. Anim. prod, 23: 151 -158.
- Fiser, P.S. and R.W. Fairfull (1986). Combined effect of glycerol concentration, cooling velocity effects of glycerol of skim milk diluents on cryopreservation of ram spermatozoa . Theriogenology., 25: 473 – 484.

A.F.Nebar, H.Abdel Rahman, I.M. Seddik and M.A. El-Ghonamy

- Foote, R.H. and M.T. Kaproth (2002). Large batch freezing of bull semen: Effect of time of freezing and fructose on fertility. *J. Dairy Sci.*, 85: 453-456.
- Galli, A.; V. Bornaghi; D. Balduzzi; L. Buttazzoni and R. Aleandri (1993). Sexual behaviour and semen quality relating to buffalo. *Proc. 3rd World buffalo Congr., Varna, Bulgaria vol. 1 pp. 562 – 570.*
- Gil, J., A. Januskauskas; Håård Mch., Håård MGM., A. Johannisson; L. Söderquist and H. Rodriguez-Martinez (2000). Functional sperm parameters and fertility of bull semen extended in Biocighos plus® and triladyl®. *Reprod. Dom. Anim.*, 35: 69 –77.
- Gil, J., Nils Lundeheim, Lennart Söderquist. and Heriberto Rodriguez (2003). Influence of extender, temperature and addition of glycerol on post – thaw sperm parameters in ram semen. *Theriogenology*; 59: 1241 – 1255.
- Jones, R.C. and T.C.A. Martin (1965). Deep-freezing ram spermatozoa: the effect of milk, yolk – citrate and synthetic diluents containing sugar. *J. Reprod. Fertil.*, 35: 311 – 320.
- Kumar, S.; K.L.N. Sahni and G. Mohan (1992). Effect of different levels of glycerol and yolk on freezing and storage of buffalo semen in milk, tris and Sodium citrate buffers, *Buffalo J.* 2, 151 – 156.
- Maxwell, W.M.C. and P.F. Watson (1996). Recent progress in the preservation of ram semen. *Anim. Reprod. Sci.* 42: 55 – 65.
- National Research Council (NRC) (1987). *The Nutrient Requirement of Sheep.* National Academy of Sciences, Washington. D.C.U.S.A.
- Nebar, A.F. (1989). Comparative studies on goat and ram semen in association with dilution and preservation techniques. Ph.D. Thesis, Faculty of Agriculture, Minufia University.
- Paulenz, H.; T. Adnøy; O.H. Fossen; L. Söderquist and K. Andersed Berg (2002). Effect of deposition site and sperm number on fertility in sheep inseminated with liquid semen. *Vet. Rec.*, 150: 299 – 302.
- Paulenz, H., L. Söderquist; T. Adnøy; O.H. Fossen and K. Berg (2003). Effect of milk – and tris– based extenders on the fertility of sheep insemination vaginal once or twice with liquid semen. *Theriogenology*, 60: 759 – 766.
- Sahni, K. L. and G. Mohan (1990). Effect of various levels of yolk on viability of buffalo semen at 37 °C, 5 °C and - 196 °C. In: Acharya, R. M, Lokeshwar, R.R., Kumar, S. (Eds.), *Recent Advances in buffalo Research vol. 3 pp. 63 – 65.*
- Salamon, S. and W.M.C. Maxwell (1995). Frozen storage of ram semen II. Causes of low fertility after cervical insemination and methods of improvement. *Anim. Reprod. Sci.* Vol. 38: 1 – 36.
- Salamon, S. and W.M.C. Maxwell (2000). Storage of ram semen. *Anim. Reprod. Sci.* Vol. 62: 77 – 111.

Effect of extender type on freezability of ram semen

- Sansone, G.; M.J.F. Natriand and A. Fabbrocini (2000). Storage of buffalo (*bubalus bubalis*) semen . *Anim. Reprod. Sci*; 62: 55 – 76.
- Snedecor, G. W. and V.S. Cochran, Raina (1980). *Statistical Methods*. 7th Ed. Iowa State University Press, Ames.
- Söderquist, L.; N. Madrid-Bury and H. Rodríguez-Martinez (1997). Assessment of membrane integrity after using different procedures to thaw ram spermatozoa frozen in mini – straws . *Theriogenology*, 48: 1115 – 1125.
- Vishwanath, R. and P. Shannon (2000). Storage of bovine semen in liquid and frozen state. *Anim. Reprod. Sci*; 62: 23 – 53.
- Watson, P.F. (2000). The causes of reduced fertility with cryopreserved semen . *Anim. Reprod. Sci*. 60: 48 – 492.

تأثير نوع المخفف على قابلية السائل المنوي للكباش للتجميد

عبد الله فتحي نيبير ، حلمى عبد الرحمن عبد الهادى ،

ابراهيم صديق ، ممدوح الغنيمى

قسم الإنتاج الحيوانى - كلية الزراعة - جامعة المنوفية.

الملخص العربى

تم جمع السائل المنوي من ٤ كباش أوسيمي ناضجة (٢-٤ سنوات) بواسطة المهبل الصناعي وتم خلط عينات السائل المنوي التى اظهرت ٨٠% فأكثر حركة تفرمية وتركيز الحيوانات المنوية بها لا يقل عن 3×10^6 /مل لدراسة تأثير خمسة مخففات على المقدرة التجميدية للسائل المنوي للكباش. مخففان منهما خاليان من اللبن الفرز (المخفف أ والمخفف ب) وثلاثة مخففات أخرى تحتوى على اللبن الفرز (ج ، د ، هـ) بنسب متباينة هي ٣٥ ، ٥٠ ، ٦٥% على الترتيب. وكانت النسبة النهائية للجلسرول ١٤,٥% وللخفيف ١ : ٦ ، وتم إجراء عملية الموازنة على درجة ٤ - ٥ م لمدة ثلاث ساعات. وتمت إسالة قصببات السائل المنوي في حمام مائي درجة حرارته ٤٢ م لمدة ٣٥ ثانية. تم تقييم النسبة المئوية لحركة الحيوانات المنوية التقدمية ، والحيوانات المنوية الحية والحيوانات المنوية الشاذة بعد مرحلة عملية التخفيف ، والموازنة ، وبعد مرحلة التجميد والإسالة.

وقد أوضحت النتائج أن تخفيف السائل المنوي بالمخففان أ ، ب (الخاليان من اللبن الفرز) نتج عنهما وبدرجة معنوية عالية (١%) أعلى نسب مئوية لحركة الحيوانات المنوية التقدمية (٣٥,٦% ، ٤٢,٥%) ، والحيوانات المنوية الحية (٣٧,٠% ، ٤٢,١%) وأقل نسب للحيوانات المنوية الشاذة (٢٣,٨% ، ١٩,٧%) بعد الإسالة ، وذلك مقارنة بالمخففات ج ، د ، هـ والتي أعطت بعد الإسالة نسب مئوية أقل لحركة الحيوانات المنوية التقدمية (٣١,٣% ، ٢٣,١% ، ٢١,٣%) ، والحيوانات المنوية الحية (٣١,٩% ، ٢٥,٤% ، ٢٣,٣%) ونسب مئوية أعلى للحيوانات المنوية الشاذة (٢١,٥١% ، ٢٤,٩٩% ، ٢٦,٧٤%) على الترتيب .

Effect of extender type on freezability of ram semen

انخفضت النسبة المئوية لحركة الحيوانات المنوية والحيوية بصورة ملحوظة مع ارتفاع نسبة اللبن الفرز في المخفف بينما ارتفعت النسبة المئوية للحيوانات الشاذة ، وكانت الفروق معنوية بدرجة عالية (١%).

حافظ المخفف (ب) على أعلى معدل لاسترداد حركة الحيوانات المنوية التقديمية (٥٨,٦%) والحيوانات المنوية الحية (٥٨,٢%) عند مقارنة تلك القيم بمثلها بعد عملية الموازنة للسائل المنوي ، وبلغت هذه القيم ٤٦,٥ و ٤٦,١% ، على التوالي ، عند مقارنة تلك القيم بمثلها للسائل المنوي الطازج قبل إجراء عمليات التجميد ، وكما حافظ على أقل زيادة كلية للحيوانات المنوية الشاذة (١٧,٦٨%) أثناء مراحل التجميد المختلفة. بينما بلغت هذه المعدلات أداها بالنسبة حركة الحيوانات المنوية التقديمية و الحيوانات المنوية الحية وأعلاها بالنسبة للزيادة الكلية للحيوانات المنوية الشاذة عند استخدام المخفف (هـ) والذي يحتوى على أعلى نسبة من اللبن الفرز.

نتج عن إجراء عمليتي الموازنة والإسالة للسائل المنوي تدهور في حركة الحيوانات المنوية بنسبة عالية تتراوح بين ٢٨,٨ و ٣٦,٩% بينما بلغت هذه النسب أداها (٥,٦ - ٨,٧%) عند إجراء عمليتي إضافة الجليسرول و الموازنة. وبلغت هذه القيم ٢٨,٥ - ٣٣,٤% ، ٥,٥ - ٩,٥% بالنسبة للحيوانات المنوية الحية ، ١٢,٣٧ - ١٥,٧٢ ، ١,٦٨ - ٢,٧٨% بالنسبة للحيوانات المنوية الشاذة ، على الترتيب . ويستخلص من هذه الدراسة أن مخففات الترس تملك القدرة على تجميد السائل المنوي للكباش بصورة أفضل من مخففات اللبن الفرز.