

EFFECT OF LONG-TERM STRESS WITH HEAVY METALS COMBINATIONS ON GROWTH AND CHEMICAL COMPOSITION OF SOME ORNAMENTAL SHRUBS

II. EFFECT ON CHEMICAL COMPOSITION

Abdalla, M. Y. A.¹ and A. M. A. Mahmoud²

1- Veget. and Flor. Dept., Fac. of Agric., Mansoura Univ.

2- Botanical Gardens Res. Dept., Hort. Res. Inst., ARC, Giza, Egypt.

ABSTRACT

Four separated pot experiments were carried out in the open field at the Experimental Station of Vegetable and Floriculture Department, Faculty of Agriculture, Mansoura University during of 2006/2007 and 2007/2008 seasons, as each season contained two periods of growth, to detect the effect of long-term stress with lead (Pb), cadmium (Cd) and nickel (Ni) added in combinations as thawing acetate salts to the soil mixture at six rates on chemical composition of six-months-old transplants of *Acalypha wilkesiana* Müll. Arg., *Asclepias curassavica* L., *Dodonaea viscosa* (L.) Jacq. and *Tabernaemontana divaricata* (L.) R. Br. ex Roem & Schult. grown in 25-cm-diameter black polyethylene bags filled with 3 Kg of a mixture composed of sand and loam at 1:1 (v/v). The chemical analysis of the studied shrubs was conducted two times in the second season only; i.e. at the terminal of October 2007 (first growth period), as well as at the end of October 2008 (second growth period).

The obtained results indicated that chlorophylls a and b, carotenoids, total carbohydrates, N, P and K contents in the leaves of all studied shrubs were progressively decreased with increasing heavy metal concentrations to reach the minimum values in transplants subjected to the highest level of toxic metals with highly significant differences compared to the means of unpolluted ones in most cases of the two periods of growth. An exception was only obtained for transplants treated with the lowest level of heavy metals, as all previous constituents were slightly increased in the leaves of both *Acalypha* and *Asclepias* transplants, while in *Tabernaemontana* and *Dodonaea* transplants many of them were increased and many others were decreased. In general, prolonging the period of subjecting to the toxic metals stress caused a gradual decrement in the content of aforementioned constituents.

On the other hand, a progressive increase was noticed in the content of Pb, Cd and Ni in the leaves and roots of the four used shrubs with elevating heavy metals concentrations. Moreover, the content of such metals was higher in the roots than in the leaves, and in the second growth period than in the first one. So, the highest content of them was observed in the roots of transplants polluted with the highest level in the second period of growth.

From the above mentioned results, it could be recommended to use *Acalypha wilkesiana* and *Asclepias curassavica* transplants to landscape areas suffering from Pb, Cd and Ni pollution, as they absorbed the highest amounts of these metals under the conditions of the present study, followed by *Tabernaemontana divaricata* then *Dodonaea viscosa*.

INTRODUCTION

Some of ornamental shrubs widely used for landscaping and gardening are sensitive to heavy metals toxicity. Among them may be the

monocious shrub with attractive foliage *Acalypha wilkesiana* Müll. Arg., Jacob's coat (Fam. Euphorbiaceae), the fragrant annual or short-lived evergreen subshrub with glowing red-flowered umbels *Asclepias curassavica* L., blood flower (Fam. Asclepiadaceae), the tropical and subtropical shrub *Dodonaea viscosa* (L.) Jacq., hop bush (Fam. Sapindaceae) and the crepe jasmine or crepe gardenia *Tabernaemontana divaricata* (L.) R. Br. ex Roem & Schult that belongs to Fam. Apocynaceae (Bailey, 1976).

The pollution with heavy metals has many deleterious effects; the most important of them is the effect on chemical composition of plant tissues. In this regard, Shahin *et al.* (2002) found that chlorophylls a and b content in the leaves of *Salvia* and *Vinca* transplants were reduced, while carotenoids content was increased in response to heavy metals pollution. A reduction was observed in total carbohydrates content in *Salvia* leaves and roots, but in *Vinca* transplants, it was increased in the leaves and decreased in the roots. Contents of both N and K in the leaves and roots were progressively decreased with increasing heavy metals level, while Pb Cd and Hg were greatly increased. Dissanayake *et al.* (2002) noticed that *Lantana camara* and *Albizzia odoratissima* nearly uptake 50% of Zn, Cd, Ni and Mn from the metal ion solutions, but the uptake of Cr, Co, Cu, Fe and Pb was less than 50%. *Wedelia trilobata*, however absorbed more than 1 mg/dm³ of Cu, Ni and Fe after 72 hours of treatment, and Mn, Zn, Pb, Co and Cr after 120 hours of treatment. In addition, Bush *et al.* (2003) revealed that Zn level in leaf tissue of *Betula nigra* and *Ulmus parvifolia* plants exceeded normal level, whereas pigments content was less due to growing in media amended with crumb rubber. Likewise, Laypheng *et al.* (2004) mentioned that the amounts of Cd, Cu, Cr, Fe, Pb and Zn in tissues of *Bougainvillea spectabilis*, *Ixora coccinea* and *Heleconia taxa* plants were significantly increased compared to the control when grown in polluted soil.

Similar observations were also recorded by Abbaas (2002) on *Casuarina glauca*, *Taxodium distichum* and *Populus nigra*; Rossini and Rautio (2004) on *Duranta repens*; Shahin and El-Malt (2006) on sant, oak and tipu; and Shahin *et al.* (2007) on stock (*Matthiola incana*) and Cape marigold (*Dimorphotheca ecklonis*).

Such trial, however aims to explore the effect of lead, cadmium and nickel combinations on chemical composition of some ornamental shrubs.

MATERIALS AND METHODS

Four independent pot experiments were carried out in the open field at the Experimental Station of Vegetable and Floriculture Department, Faculty of Agriculture, Mansoura University during 2006/2007 and 2007/2008 seasons to study the effect of long-term stress of Pb, Cd and Ni in combinations at various levels on chemical composition of Jacob's coat, blood flower, hop bush and crepe jasmine transplants.

Therefore, six-months-old transplants of Jacob's coat (55-60 cm long with 2 branches carrying about 15-17 leaves), blood flower (40-45 cm long with one branch carrying about 32-35 leaves), hop bush (25-27 cm long with

one branch carrying about 7-9 leaves) and crepe jasmine (43-45 cm long with 2 branches carrying about 16-18 leaves) were planted on April, 1st for both seasons in 25-cm-diameter black polyethylene bags (one transplant/bag) filled with 3 kg of a mixture of sand and loam (1:1 v/v), which its some physical and chemical properties are shown in Table (a).

Thawing salts (acetates) of Pb, Cd and Ni produced by Aldrich Chemical Co. Inc., USA, were thoroughly mixed in combinations with the used soil mixture before planting at the concentrations of 00.00 ppm for each metal as a control, 500 ppm Pb + 50 ppm Cd + 25 ppm Ni for treatment number one (T₁) and 2-, 3-, 4- and 5-fold of these concentrations for treatments number two (T₂), three (T₃), four (T₄) and five (T₅), respectively above the background levels of these metals in the used soil mixture. The bags (without drainage holes to prevent leaching of metals) were immediately irrigated after planting with 250 ml of fresh water/bag, but thereafter the irrigation was done once every 3 days with only 200 ml of water/bag. The transplants were not fertilized throughout the course of study, but received the usual agricultural practices recommended for such plantation (i.e. digging, weed control ... etc.). They were set out in a complete randomized design (Mead *et al.*, 1993) for each plant with 3 replicates, as each replicate contained 6 transplants.

The experiments began in the first season on April, 1st 2006 and lasted for two growing periods till the end of October 2007, whereas for the second one, it commenced on April, 1st 2007 and lasted till the end of October 2008. However, chemical analyses were measured two times in the second season only; i.e. at the terminal of October of both 2007 and 2008 years as follows:

- Chlorophyll a, b and carotenoids (mg/g fresh weight) were determined in fresh leaf samples taken from the middle parts of transplants according to the method of Moran (1982).
- In dry leaf samples, the percentages of total carbohydrates (Herbert *et al.*, 1971), N (Pregl, 1945), P (Luatanab and Olsen, 1965) and K (Jackson, 1973) were assessed.
- Another dry samples from the leaves and roots were digested in nitric and perchloric acids and analyzed for Pb, Cd and Ni (ppm) determination on a Perkin Elmer 403 atomic absorption spectrophotometer as the method described by Jackson (1973).

The data were subjected to analysis of variance and the method of L.S.D. was used to differentiate the means (Mead *et al.*, 1993).

RESULTS AND DISCUSSION

Effect of lead (Pb), cadmium (Cd) and nickel (Ni) combinations on chemical composition of the different shrubs:

1. *Acalypha wilkesiana* Müll. Arg. (Jacob's coat):

Data presented in Table (1) cleared that chlorophyll a, b and carotenoids content (mg/g fresh weight), as well as the percentages of total carbohydrates, N, P and K in the leaves of polluted transplants were

Table (a): Some physical and chemical properties of the used soil mixture during 2006/2007 and 2007/2008 seasons

Season	Properties												
	Particles size distribution (%)												
	Coarse sand		Fine sand		Clay		Silt		Texture				
2006/07	74.50		6.35		12.45		6.70		Sandy loam				
2007/08	73.85		7.00		12.25		6.90		Sandy loam				
	SR (%)	EC (mmhos/cm)	pH	Cations (meq/l)							Anions (meq/l)		
				Ca ⁺⁺	Mg ⁺⁺	Na ⁺	K ⁺	Pb ⁺⁺	Cd ⁺⁺	Ni ⁺⁺	HCO ₃ ⁻	Cl ⁻	SO ₄ ⁼⁼
2006/07	24.57	3.61	7.12	24.12	10.00	36.55	1.56	1.00	0.05	nil	3.60	28.16	41.52
2007/08	24.76	3.20	7.26	25.72	9.38	37.21	1.42	0.94	0.05	nil	3.48	27.33	43.87

gradually decreased with increasing heavy metals level to reach the minimum values in both T₄ and T₅ treatments with significant and highly significant differences compared to the unpolluted ones during the two periods of growth, except for T₁ combination, which slightly increased the content of all previous constituents with non-significant differences in the two growth periods. The only significant increment, which was achieved by T₁ treatment, was in chlorophyll b content in both growth periods. Prolonging the period of subjection to heavy metals stress caused a progressive decrement in the previously mentioned constituents, especially at the highest concentrations of toxic metals (T₄ and T₅).

Table (1): Effect of heavy metals combinations on some chemical constituents of *Acalypha wilkesiana* Müll. Arg. leaves during the two periods of growth in the second season

Treatments	Chlorophyll a (mg/g f. w.)	Chlorophyll b (mg/g f. w.)	Carotenoids (mg/g f. w.)	Total carbohydrates (%)	N (%)	P (%)	K (%)
First period of growth (terminal of October 2007)							
Control	0.475	0.197	0.407	63.75	1.84	0.063	1.36
T ₁	0.490	0.265 *	0.446	66.50	1.92	0.071	1.48
T ₂	0.468	0.183	0.369	61.63	1.80	0.067	1.39
T ₃	0.386	0.159	0.341	51.80	1.47	0.052	1.13
T ₄	0.328 **	0.130 *	0.336	43.86 *	1.28 *	0.043 *	0.95 *
T ₅	0.281 **	0.098 **	0.312 *	41.72 **	1.07 *	0.037 *	0.81 **
L.S.D 5%	0.093	0.043	0.072	12.03	0.53	0.012	0.26
1%	0.141	0.074	0.094	20.33	0.81	0.028	0.43
Second period of growth (terminal of October 2008)							
Control	0.518	0.215	0.435	64.43	1.76	0.061	1.48
T ₁	0.534	0.283 *	0.456	67.84	1.82	0.069	1.31
T ₂	0.446	0.176	0.350	58.56	1.70	0.064	1.32
T ₃	0.380	0.153	0.318 *	49.21 *	1.43	0.049	1.08 **
T ₄	0.352 *	0.128 *	0.309 *	42.75 *	1.24 *	0.042	0.91 **
T ₅	0.310 **	0.109 **	0.301 *	39.71 **	1.03 *	0.034 *	0.74 **
L.S.D 5%	0.139	0.063	0.096	14.50	0.46	0.018	0.33
1%	0.181	0.091	0.128	21.96	0.75	0.030	0.51

* Concentrations of Pb, Cd and Ni in T₁ were 500, 50 and 25 ppm, and in T₂, T₃, T₄ and T₅ were 2-, 3-, 4- and 5-fold of these concentrations, respectively

These results may be due to the indirect effects of heavy metals on photosystems related to the disturbances caused by the metals in Calvin cycle and downregulation or even feedback inhibition of electron transport by the excessive amounts of ATP and NADPH (Krupa *et al.*, 1993). Droppa *et al.* (1996), however suggested that Cd in green leaves interferes with chlorophyll biosynthesis, acts mainly by inhibiting the LHC synthesis into stable complexes required for normal functional photosynthetic activity.

A progressive increase was observed in Pb, Cd and Ni content (ppm) in the leaves and roots (Table, 2) with significant differences for T₃ combination and highly significant ones for both T₄ and T₅ combinations compared to control in the first period of growth, while in the second one, the level of significancy was elevated for T₂ treatment in the leaves and for both T₁ and T₂ treatments in the roots to become significant. In general, the

content of the three heavy metals in the roots was higher than in the leaves, and in the second period than in the first one. This may indicate that transportation of the used toxic metals to leaves was less than accumulation of them in the roots. However, absorption of metals by roots of plants grown in polluted soil may be necessary for keeping the equilibrium between metals concentration in soil medium and nutrients content in plant tissues (Foy *et al.*, 1978).

Table (2): Effect of heavy metals combinations on Pb, Cd and Ni content (ppm) in the leaves and roots of *Acalypha wilkesiana* Müll. Arg. during the two periods of growth in the second season

Treatments	First period of growth (terminal of October 2007)			Second period of growth (terminal of October 2008)		
	Pb	Cd	Ni	Pb	Cd	Ni
	In the leaves					
Control	3.76	0.50	nil	4.06	0.56	nil
T ₁	7.35	2.41	1.08	8.38	2.60	1.21
T ₂	13.50	3.28	1.67	13.78 *	3.74 *	1.88 *
T ₃	15.77 *	6.21 *	1.97	17.03 *	6.71 *	2.20 *
T ₄	23.48 **	8.92 **	2.84 **	25.21 **	9.58 **	3.19 **
T ₅	30.24 **	10.90 **	3.36 **	32.40 **	11.86 **	3.75 **
L.S.D 5%	9.76	2.86	1.98	9.05	2.31	1.66
1%	16.28	7.42	2.63	15.97	6.84	3.01
	In the roots					
Control	6.79	0.78	nil	7.35	0.93	nil
T ₁	9.68	3.56	1.19	10.50	3.98 *	1.97 *
T ₂	17.07	5.91	1.97	18.47 *	6.50 *	2.16 *
T ₃	21.86 *	8.45 *	2.63 **	23.55 *	9.20 **	2.96 *
T ₄	28.73 **	12.54 **	4.70 **	31.10 **	13.67 **	5.23 **
T ₅	34.10 **	15.82 **	5.23 **	36.78 **	17.33 **	5.78 **
L.S.D 5%	10.97	5.33	2.16	10.32	3.00	1.84
1%	17.94	10.67	3.51	18.75	6.58	3.30

* Concentrations of Pb, Cd and Ni in T₁ were 500, 50 and 25 ppm, and in T₂, T₃, T₄ and T₅ were 2-, 3-, 4- and 5-fold of these concentrations, respectively

Although the Jacob's coat transplants absorbed high amounts of Pb, Cd and Ni, the concentrations of such metals did not reach the upper critical levels, which ranged in various plant tissues between 50-60, 60-120 and 8-12 ppm for the three metals mentioned above, respectively (Macnicol and Beckett, 1985). However, these findings are in accordance with those attained by Shahin *et al.* (2002) on *Salvia* and *Vinca*; Bush *et al.* (2003) on *Betula nigra* and *Ulmus parvifolia*; Laypheng *et al.* (2004) on *Bougainvillea spectabilis*, *Ixora coccinea* and *Heleconia taxa* and Shahin *et al.* (2007) on stock and Cape marigold.

2. *Asclepias curassavica* L. (Blood flower):

Data averaged in Table (3) exhibited that the least combination of heavy metals (T₁) caused a non-significant increase in chlorophyll a, b and carotenoids contents (mg/g fresh weight), as well as in the percentages of total carbohydrates, N, P and K in the leaves during the two periods of growth, except for chlorophyll b content in the first growth period that was

significantly raised, whereas higher combinations (T₂, T₃, T₄ and T₅) induced a progressive decrement in these constituents as heavy metals concentrations increased. So, the least records were due to the highest level of toxic metals (T₅). It was noticed that chlorophylls a and b were more affected by toxicity of heavy metals than total carbohydrates, N, P and K parameters, especially under medium levels (T₂ and T₃) during the two periods of growth.

Table (3): Effect of heavy metals combinations on some chemical constituents of *Asclepias curassavica* L. leaves during the two periods of growth in the second season

Treatments	Chlorophyll a (mg/g f. w.)	Chlorophyll b (mg/g f. w.)	Carotenoids (mg/g f. w.)	Total carbohydrates (%)	N (%)	P (%)	K (%)
First period of growth (terminal of October 2007)							
Control	0.739	0.566	0.337	70.10	1.38	0.078	2.76
T ₁	0.817	0.658 *	0.361	77.68	1.51	0.086	2.89
T ₂	0.698	0.505	0.337	66.23	1.28	0.076	2.58
T ₃	0.646 *	0.456 *	0.341	60.51	1.21	0.067	2.03
T ₄	0.626 *	0.433 **	0.296	58.33 *	1.03 *	0.053	1.81 *
T ₅	0.538 **	0.326 **	0.250 *	51.08 *	0.97 *	0.046 *	1.62 **
L.S.D 5%	0.081	0.076	0.045	11.33	0.32	0.026	0.75
1%	0.143	0.114	0.094	19.18	0.49	0.037	1.01
Second period of growth (terminal of October 2008)							
Control	0.723	0.546	0.326	66.50	1.31	0.081	2.63
T ₁	0.780	0.627	0.344	70.15	1.45	0.085	2.76
T ₂	0.661	0.480	0.320	62.70	1.20	0.073	2.45
T ₃	0.615 *	0.431 *	0.324	57.00	0.98 *	0.061	1.92
T ₄	0.595 *	0.409 **	0.285 *	55.43 *	0.96 *	0.050 *	1.70 *
T ₅	0.511 **	0.312 **	0.233 *	48.53 *	0.93 *	0.042 **	1.50 *
L.S.D 5%	0.097	0.083	0.039	10.67	0.30	0.018	0.33
1%	0.148	0.129	0.087	18.42	0.46	0.030	0.51

* Concentrations of Pb, Cd and Ni in T₁ were 500, 50 and 25 ppm, and in T₂, T₃, T₄ and T₅ were 2-, 3-, 4- and 5-fold of these concentrations, respectively

On the other hand, contents of Pb, Cd and Ni (ppm) in the leaves and roots (Table, 4) were markedly increased with various significant levels in the two periods of growth. Their content in the roots was higher than that in the leaves. However, accumulation of toxic metals in plant tissues usually leads to depression of vital processes, such as photosynthesis, inhibition of some enzymatic systems and blocking the formation of proteins and chlorophylls (Mengel and Kirkby, 1979). A reduction in glutathione reductase activity in relation to Cd and Ni stress was registered by Schickler and Caspi (1999) in *Alyssum argentum*. Moreover, Salgare and Palathingal (2002) found that industrial pollution delayed flowering of *Allamanda cathartica* plants due to a reduction in cytokinin and gibberellin activities, and in sucrose and glucose contents in bud meristems during the transition to flowering. Similarly were those results pointed out by Dissanayake *et al.* (2002) on *Lantana camara*, *Albizzia odoratissima* and *Wedelia trilobata*; Rossini-Oliva and Rautio (2004) on *Duranta repens* and Shahin and El-Malt (2006) on sant, oak and tipu.

Table (4): Effect of heavy metals combinations on Pb, Cd and Ni content (ppm) in the leaves and roots of *Asclepias curassavica* L. during the two periods of growth in the second season

Treatments	First period of growth (terminal of October 2007)			Second period of growth (terminal of October 2008)		
	Pb	Cd	Ni	Pb	Cd	Ni
	In the leaves					
Control	2.18	0.78	nil	2.31	0.84	nil
T ₁	4.36	3.96	1.09	4.78	4.33	1.18
T ₂	10.90	7.49	2.27 *	11.87 *	8.64	2.49 *
T ₃	15.28 *	15.26 **	3.10 **	16.76 *	17.28 **	3.46 **
T ₄	17.85 **	17.66 **	4.91 **	19.89 **	19.44 **	5.41 **
T ₅	22.89 **	21.26 **	5.50 **	25.46 **	23.76 **	6.88 **
L.S.D 5%	8.75	6.83	1.17	9.10	7.81	1.33
1%	13.62	11.47	2.36	14.93	12.40	2.50
	In the roots					
Control	3.70	1.03	nil	4.03	1.08	nil
T ₁	5.78	4.36	1.58 *	6.28	5.19	1.73 *
T ₂	14.16 ^a	9.81 *	2.96 *	15.47 **	10.83 *	3.25 **
T ₃	21.80 **	18.60 **	4.51 **	23.76 **	20.56 **	4.86 **
T ₄	25.07 **	23.98 **	6.33 **	28.09 **	26.00 **	6.81 **
T ₅	33.21 **	31.76 **	7.29 **	34.67 **	34.56 **	7.90 **
L.S.D 5%	9.64	7.88	1.34	10.64	8.35	1.70
1%	14.10	12.35	3.00	16.73	13.46	3.16

* Concentrations of Pb, Cd and Ni in T₁ were 500, 50 and 25 ppm, and in T₂, T₃, T₄ and T₅ were 2-, 3-, 4- and 5-fold of these concentrations, respectively

3. *Dodonaea viscosa* (L.) Jacq. (Hop bush):

On the contrary of what was found in *Acalypha* and *Asclepias* transplants, data in Table (5) indicated that all heavy metals combinations declined the content of chlorophyll a, b, carotenoids, total carbohydrates, N, P and K in the leaves of hop bush transplants with significant differences in both growth periods, except for combination No. 1 (T₁), which caused a slight elevation in the content of K in the second period only. The utmost low content in the previous constituents was generally referred to the highest rates of heavy metals (T₄ and T₅).

As for the content of Pb, Cd and Ni in the leaves and roots, data in Table (6) show that they were gradually raised with increasing heavy metals concentration in either periods of growth. However, the content of these metals was higher in the roots than in the leaves, and in the second growth period than in the first one indicating their continuous absorption with prolonging the growth period. Hence, the highest contents of these metals was found in the roots of transplants polluted with the highest levels of them, especially in the second period of growth.

The aforementioned results could be explained and discussed as done before in case of both *Acalypha* and *Asclepias* transplants.

Table (5): Effect of heavy metals combinations on some chemical constituents of *Dodonaea viscosa* (L.) Jacq. leaves during the two periods of growth in the second season

Treatments	Chlorophyll a	Chlorophyll b	Carotenoids	Total carbohydrates	N	P	K
	(mg/g f. w.)	(mg/g f. w.)	(mg/g f. w.)	(%)	(%)	(%)	(%)
First period of growth (terminal of October 2007)							
Control	0.465	0.471	0.430	58.73	1.18	0.047	1.26
T ₁	0.552	0.465	0.428	57.38	1.16	0.053	1.38
T ₂	0.461	0.384	0.400	47.46 *	1.16	0.042	1.32
T ₃	0.430 *	0.306 *	0.368	44.07 *	0.89	0.036	1.10 *
T ₄	0.407 **	0.276 **	0.352 *	41.78 **	0.82 *	0.033 *	0.91 **
T ₅	0.400 **	0.306 **	0.337 *	40.67 **	0.82 *	0.030 *	0.90 **
L.S.D 5%	0.106	0.096	0.076	9.97	0.30	0.012	0.15
1%	0.151	0.167	0.098	14.86	0.39	0.018	0.24
Second period of growth (terminal of October 2008)							
Control	0.558	0.483	0.451	60.63	1.24	0.063	1.33
T ₁	0.569	0.488	0.449	59.85	1.22	0.056	1.45
T ₂	0.483	0.396	0.418	48.93 *	1.17	0.043	1.39
T ₃	0.452 *	0.327 *	0.387 *	45.21 *	0.91 *	0.035 *	1.16
T ₄	0.421 *	0.291 **	0.361 *	41.50 **	0.84 *	0.031 *	0.90 **
T ₅	0.403 **	0.263 **	0.354 *	39.81 **	0.81 *	0.028 *	0.83 **
L.S.D 5%	0.100	0.099	0.081	10.33	0.33	0.022	0.18
1%	0.139	0.170	0.103	16.21	0.45	0.033	0.31

* Concentrations of Pb, Cd and Ni in T₁ were 500, 50 and 25 ppm, and in T₂, T₃, T₄ and T₅ were 2-, 3-, 4- and 5-fold of these concentrations, respectively

Table (6): Effect of heavy metals combinations on Pb, Cd and Ni content (ppm) in the leaves and roots of *Dodonaea viscosa* (L.) Jacq. during the two periods of growth in the second season

Treatments	First period of growth			Second period of growth		
	(terminal of October 2007)			(terminal of October 2008)		
	Pb	Cd	Ni	Pb	Cd	Ni
In the leaves						
Control	1.12	0.53	nil	1.21	0.63	nil
T ₁	2.28	1.20	0.60 *	2.65	1.50	0.71 *
T ₂	4.39 *	2.16 *	0.98 *	5.18 *	2.63 **	1.15 **
T ₃	5.28 *	4.15 **	2.40 **	6.93 **	4.76 **	2.88 **
T ₄	9.27 **	5.98 **	3.31 **	11.04 **	6.98 **	3.91 **
T ₅	13.85 **	6.54 **	4.06 **	16.10 **	8.07 **	5.07 **
L.S.D 5%	2.67	0.76	0.46	2.47	0.88	0.50
1%	5.10	1.68	0.69	4.16	1.93	0.78
In the roots						
Control	2.60	0.78	nil	2.88	0.87	nil
T ₁	3.99	2.19 *	0.83 *	4.67	2.53 *	1.04 **
T ₂	5.67	3.42 **	1.25 **	6.71 *	4.03 **	1.50 *
T ₃	7.63 *	4.96 **	3.57 **	8.98 **	5.87 **	4.15 **
T ₄	10.48 **	6.86 **	5.02 **	12.20 **	8.10 **	6.33 **
T ₅	15.29 **	8.18 **	6.49 **	17.83 **	9.78 **	7.71 **
L.S.D 5%	3.26	1.08	0.71	2.80	0.97	0.77
1%	6.84	2.17	0.95	5.13	2.38	1.02

* Concentrations of Pb, Cd and Ni in T₁ were 500, 50 and 25 ppm, and in T₂, T₃, T₄ and T₅ were 2-, 3-, 4- and 5-fold of these concentrations, respectively

4. *Tabernaemontana divaricata* (L.) R. Br. ex Roem & Schult. (Crepe jasmine or Crepe gardenia):

It is obvious from data in Table (7) that chlorophyll a, b and carotenoids content (mg/g fresh weight), as well as the percentages of total carbohydrates, N, P and K in the leaves were augmentatively depressed as a result of raising the rates of heavy metals in polluted soil with significant and highly-significant differences in the most cases of the two growth periods, except for T₁ combination, which slightly improved the contents of chlorophyll b, carotenoids and nitrogen compared to those of transplants grown in unpolluted soil.

With regard to heavy metals content (ppm) in the leaves and roots, data recorded in Table (8) revealed that Cd and Ni were significantly increased in response to T₁ treatment during the two growth periods, while other treatments (T₂, T₃, T₄ and T₅) greatly elevated the content of the three metals with highly significant differences in both periods of growth. Such gains, however may be interpreted and discussed as previously mentioned in case of both *Acalypha* and *Asclepias* transplants.

Table (7): Effect of heavy metals combinations on some chemical constituents of *Tabernaemontana divaricata* (L.) R. Br. ex Roem & Schult. leaves during the two periods of growth in the second season

Treatments	Chlorophyll a (mg/g f. w.)	Chlorophyll b (mg/g f. w.)	Carotenoids (mg/g f. w.)	Total carbohydrates (%)	N (%)	P (%)	K (%)
First period of growth (terminal of October 2007)							
Control	0.578	0.433	0.375	74.39	2.80	0.134	2.25
T ₁	0.529	0.456	0.382	67.98	2.96	0.121	2.03
T ₂	0.478	0.271 *	0.366	61.57 *	2.35	0.110	1.83 *
T ₃	0.359 **	0.268 *	0.357	58.18 *	1.87 *	0.083 *	1.80 **
T ₄	0.361 **	0.221 **	0.284 *	46.23 **	1.72 **	0.081 *	1.64 **
T ₅	0.231 **	0.180 **	0.269 *	39.56 **	1.17 **	0.057 **	1.43 **
L.S.D 5%	0.109	0.160	0.072	10.76	0.48	0.051	0.40
1%	0.210	0.207	0.094	16.38	0.95	0.070	0.60
Second period of growth (terminal of October 2008)							
Control	0.549	0.421	0.395	70.30	2.67	0.117	2.13
T ₁	0.512	0.433	0.406	65.61	2.81	0.114	1.93
T ₂	0.458	0.319	0.384	58.90	2.17	0.103	1.74
T ₃	0.346 **	0.276 *	0.371	55.10 *	1.88 *	0.081 *	1.70 **
T ₄	0.352 **	0.251 *	0.294 *	47.51 **	1.65 **	0.079 *	1.52 **
T ₅	0.285 **	0.209 **	0.281 *	41.90 **	1.14 **	0.062 **	1.30 **
L.S.D 5%	0.097	0.109	0.097	11.50	0.59	0.033	0.40
1%	0.174	0.185	0.141	17.79	0.88	0.052	0.67

* Concentrations of Pb, Cd and Ni in T₁ were 500, 50 and 25 ppm, and in T₂, T₃, T₄ and T₅ were 2, 3, 4- and 5-fold of these concentrations, respectively

Table (8): Effect of heavy metals combinations on Pb, Cd and Ni content (ppm) in the leaves and roots of *Tabernaemontana divaricata* (L.) R. Br. ex Roem & Schult. during the two periods of growth in the second season

Treatments	First period of growth (terminal of October 2007)			Second period of growth (terminal of October 2008)		
	Pb	Cd	Ni	Pb	Cd	Ni
	In the leaves					
Control	1.86	0.69	nil	2.03	0.78	nil
T ₁	3.81	1.97 *	0.98 *	4.20	2.20 *	1.09 *
T ₂	7.45 *	3.88 **	2.03 **	8.27 **	4.29 **	2.40 **
T ₃	9.86 **	9.54 **	2.80 **	10.88 **	10.46 **	3.28 **
T ₄	12.60 **	10.33 **	4.43 **	13.96 **	11.33 **	5.00 **
T ₅	16.10 **	13.68 **	5.12 **	17.58 **	15.20 **	5.63 **
L.S.D 5%	2.06	0.88	0.77	2.21	0.90	0.81
1%	5.70	2.19	0.99	4.94	2.76	1.10
	In the roots					
Control	3.15	1.00	nil	3.40	1.10	nil
T ₁	5.10	2.78 *	1.43 *	5.61	3.10 *	1.57 *
T ₂	9.54 **	6.84 **	2.68 **	10.50 **	7.67 **	3.00 **
T ₃	12.78 **	10.93 **	4.02 **	14.03 **	12.00 **	4.43 **
T ₄	15.62 **	16.20 **	5.76 **	16.51 **	17.98 **	6.40 **
T ₅	20.38 **	21.07 **	6.48 **	22.00 **	23.70 **	7.36 **
L.S.D 5%	2.13	1.06	0.93	2.38	0.97	0.98
1%	6.10	3.00	1.56	5.10	3.28	1.33

* Concentrations of Pb, Cd and Ni in T₁ were 500, 50 and 25 ppm, and in T₂, T₃, T₄ and T₅ were 2-, 3-, 4- and 5-fold of these concentrations, respectively

According to the previously stated results, it could be concluded that *Acalypha wilkesiana* and *Asclepias curassavica* transplants are considered hyper accumulators for Pb, Cd and Ni toxic metals, as they absorbed more amounts of them, followed by *Tabernaemontana divaricata*, that absorbed medium amounts, and then *Dodonaea viscosa*, which absorbed the least amounts.

REFERENCES

- Abbaas, M. M. (2002). Effect of some heavy metals in the irrigation water on growth and chemical constituents of some timber trees. Ph. D. Thesis, Fac. Agric., Cairo Univ.
- Bailey, L. H. (1976). Hortus Third. Macmillan publishing Co., Inc., 866 Third Avenue, New York, N. Y. 10022, 1290 pp.
- Bush, E.; A. Owings and K. Leader (2003). Foliar accumulation of zinc in tree species grown in hardwood bark media amended with crumb rubber. J. Plant Nutrition, 26 (7): 1413-25.
- Dissanayake, U. S.; K. U. Tennakoon and N. Pryantha (2002). Potential of two invasive plant species, *Lantana camara* L. and *Wedelia trilobata* L., for selective heavy metal uptake. Ceylon J. Bio. Sci., 29 (1): 1-11.
- Droppa, M.; A. Oravec; B. Boddí and G. Horvath (1996). Heavy metals inhibition of photosynthetic membrane. Proc. 1st Egypt-Hung. Hort Conf., Hort. Dept. Fac. Agric., Kafr El-Sheikh, 1: 13-18.
- Foy, C. D.; R. L. Chaney and M. C. White (1978). The physiology of metal toxicity in plants. Ann. Rev. Plant Physiology, 29: 511-566.

- Herbert, D.; P. J. Philips and R. E. Strange (1971). Determination of total carbohydrates. *Methods in Microbiology*, 5 (8): 290-344.
- Jackson, M. L. (1973). *Soil Chemical Analysis*. Prentice-Hall of India Private Ltd. M-97m New Delhi, India, 498 pp.
- Krupa, Z.; A. Siedlecka; W. Maksymiec and T. Baszynski (1993). *In vivo* response of photosynthetic apparatus of *Phaseolus vulgaris* L. to nickel toxicity. *J. Plant Physiol.*, 142 (6): 664-668.
- Laypheng, T.; H. Jie and L. Singkong (2004). Growth of some tropical ornamental plants on artificial topsoil derived from mixtures of fly ash, sludge, biochips and rengam series subsoil. *J. Plant Nutrition*, 27 (1): 75-94.
- Luatanab, F. S. and S. R. Olsen (1965). Test of an ascorbic acid method for determining phosphorus in water and NaHCO₃ extracts from soil. *Soil Sci. Soc. Amer. Proc.*, 29: 677-678
- Macnicol, R. D. and P. H. T. Beckett (1985). Critical tissue concentrations of potentially toxic elements. *Plant and Soil*, 85: 107-129.
- Mead, R.; R. N. Curnow and A. M. Harted (1993). *Statistical Methods in Agriculture and Experimental Biology*, 2nd Ed., Chapman & Hall Ltd., London, 335 pp.
- Mengel, K. and E. A. Kirkby (1979). *Principles of Plant Nutrition*. 2nd Ed., International Potash Inst., P. O. Box CH-3048 Worblaufen, Bern, Switzerland, 593 pp.
- Moran, R. (1982). Formula for determination of pigments extracted with N, N-dimethyl formamide. *Plant Physiology*, 69: 1376-81.
- Pregl, F. (1945). *Quantitative Organic Micro-Analysis*, 4th Ed., J. and A. Churchill Ltd., London, p. 203-209.
- Rossini-Oliva, S. and P. Rautio (2004). Could ornamental plants serve as passive biomonitors in urban area?. *J. Atmospheric Chemistry*, 49 (1/3): 137-148.
- Salgare, S. A. and T. Palathingal (2002). Effect of industrial pollution at Sewri-Mumbai. II. Changes in resting period of pollen of successive flowers of *Allamanda cathartica* L. *Advances in Plant Science*, 15 (2): 645-647.
- Schickler, H. and H. Caspi (1999). Response of antioxidative enzymes to nickel and cadmium stress in hyperaccumulator plants of genus *Alyssum*. *Physiologia Plantarum*, 105 (1): 39-44..
- Shahin, S. M. and A. El-Malt, Azza (2006). A study on usage of sant, oak and tipu trees for arboriculture of a polluted sandy soil. *Egypt. J. Appl. Sci.*, 21 (7): 192-214.
- Shahin, S. M.; A. M. A. Mahmoud and A. H. El-Feky (2007). How far stock and Cape marigold plants can tolerate toxicity of some heavy metals combinations?. *J. Bio. Chem. Environ. Sci.*, 2 (2): 415-433.
- Shahin, S. M.; M. H. El-Shakhs and M. H. Abdel-Salam (2002). Impact of Pb, Cd and Hg combinations on growth, flowering and chemical composition of *Salvia splendens* L. and *Vinca rosea* L. *Proc. 2nd Inter. Conf. Hort. Sci.*, 10-12 Sept., Kafr El-Sheikh, Tanta Univ.: 911-920.

تأثير الإجهاد طويل المدى بتوليفات من بعض المعادن الثقيلة على النمو والتركيب الكيميائي لبعض شجيرات الزينة

٢- التأثير على التركيب الكيميائي

محمد يونس على عبد الله^١ و أحمد محمد علي محمود^٢

١- قسم الخضار والزينة - كلية الزراعة - جامعة المنصورة

٢- قسم بحوث الحدائق النباتية- معهد بحوث البساتين- مركز البحوث الزراعية-الجيزة- مصر

لجريت أربع تجارب لاصص منفصلة تحت ظروف الحقل في محطة تجارب قسم الخضار والزينة - كلية الزراعة - جامعة المنصورة خلال الموسمين المتتاليين ٢٠٠٦/٢٠٠٧ و ٢٠٠٧/٢٠٠٨ حيث اشتمل كل موسم على فترتين للنمو، وذلك لمعرفة تأثير الإجهاد طويل المدى بتوليفات من عناصر الرصاص، الكاديوم والنيكل عند إضافتها لمخلوط التربة قبل الزراعة في صورة أملاح الخلات سريعة الذوبان بستة تركيزات مختلفة على التركيب الكيميائي لشتلات الأكاليفيا (*Acalypha wilkesiana* Müll. Arg.) ، الدفلة التركية (*Asclepias curassavica* L.) ، السودونيا (*Dodonaea viscosa* (L.) Jacq.) ، والتابريتا (*Tabernaemontana divaricata* (L.) R. Br. ex Roem & Schult.) عمر ستة أشهر المزروعة في أكياس بلاستيك سوداء غير متقبة قطرها ٢٥ سم ملئت بحوالي ٣ كجم من مخلوط متساوي من الرمل والطيني (بالحجم). ولقد اجري تحليل التركيب الكيميائي لهذه الشجيرات مرتين في موسم الزراعة الثاني فقط وذلك في نهاية أكتوبر ٢٠٠٧ (فترة النمو الأولى) ونهاية أكتوبر ٢٠٠٨ (فترة النمو الثانية).

ولقد أوضحت النتائج المتحصل عليها أن محتوى أوراق الشجيرات الأربعة موضع الدراسة من الكلوروفيل أ و ب ، الكاروتينويدات، الكربوهيدرات الكلية، النتروجين، الفوسفور والبوتاسيوم قد انخفضت بشكل متصاعد بزيادة تركيز العناصر الثقيلة لتصل إلى أدنى القيم في الشتلات التي تعرضت للمستويات العالية من تلك العناصر، ويفروق عالية المعنوية عند مقارنتها بمتوسطات شتلات المقارنة (الكنترول) في معظم الحالات لكلا فترتي النمو باستثناء واحد يمكن الحصول عليه فقط من الشتلات التي عوملت بالمستوى الأدنى من العناصر الثقيلة، حيث زاد محتوى جميع المكونات الكيميائية سالفة الذكر بدرجة طفيفة في أوراق شتلات الأكاليفيا والدفلة التركية، بينما في شتلات السودونيا والتابريتا فقد زاد محتوى بعض تلك المكونات وانخفض البعض الآخر. وبصفة عامة تحدثت إطالة مدة التعرض لإجهاد العناصر السامة بإنخفاضاً متزايداً في المكونات الكيميائية سالفة الذكر، خاصة عند التعرض للمستويات الأعلى من تلك العناصر.

وعلى الجانب الآخر، فقد حدثت زيادة متصاعدة في محتوى لورق وجذور شتلات الأنواع الشجرية الأربعة موضع الدراسة من عناصر الرصاص، الكاديوم والنيكل كلما زاد تركيز تلك العناصر في التربة الملوثة. علاوة على ذلك، فإن محتوى هذه العناصر كان أعلى في الجذور عنه في الأوراق، وفي الفترة الثانية للنمو عنه في الفترة الأولى. لذلك كان أعلى محتوى من تلك العناصر في جذور الشتلات التي لوشت تربتها بالمستوى الأعلى من هذه العناصر في الفترة الثانية للنمو.

من النتائج سالفة الذكر، يمكن للتوصية باستخدام شتلات الأكاليفيا والدفلة التركي عمر ستة أشهر لتجميل وتنسيق المناطق التي تعاني من التلوث المرتفع بعناصر الرصاص، الكاديوم والنيكل، حيث قاما بامتصاص أعلى كميات من تلك العناصر تحت ظروف هذه الدراسة وسجلتا في نفس الوقت نسبة حياة ١٠٠% خلال مرحلتى النمو لكلا الموسمين، تليهما التابريتا ثم السودونيا.