

EFFECT OF ADDING CHROMIUM PICOLONATE INTO BROILER DIETS ON THEIR PERFORMANCE AND DIGESTIVE ENZYME

EI-Bander, Luma K.; Nagla K. Soliman and Sonble J. El-Hamody
Dept. of poultry prod. Faculty of Agriculture, Ain Shams University

ABSTRACT

An experiment was conducted to investigate the effect of different graded levels of chromium picolonate on performance of broiler chicks, some digestive enzymes activity, carcass yield and some blood constituents. For this purpose three hundred day old Cubb broiler chicks were allocated into five treatment groups. For five weeks experimental period, four groups of birds were fed on corn soy-basal diets supplemented with 0.25mg, 0.5mg, 0.75mg or 1mg/kg diet chromium of chromium picolonate whereas the fifth group was the control.

Average body weight improved due to feeding supplemental chromium diets. The group of birds fed 0.5mg chromium picolonate recorded the highest body weight (1888g) followed by those fed 0.75mg Cr (1872g) compared with (1788g) for control. Feed consumption values were not affected by treatments for the entire experimental period. Feed conversion (g feed/g gain) improved as a result of feeding dietary chromium. The beneficial effects of chromium supplementation may be related to its effect on stimulating digestive enzymes activity and immune response. Amylase enzyme activity was significantly elevated in jejunum and ileum due to feeding chromium enriched-diets, while lipase enzyme was slightly increased due to treatment in jejunum only. Trypsine and chemotrypsine enzymes were not affected by treatments. Dressed carcass and breast meat yield percentages were increased due to adding chromium into diets while; thigh, drumstick or back percentages were not affected.

Supplemental organic chromium significantly increased plasma concentration of glucose, total lipids, cholesterol and triglycerides while plasma high density lipoprotein cholesterol (H.D.L.) was not affected. Plasma total protein and globulin increased significantly due to treatments while, plasma albumin was not affected. Albumin to globulin ratio decreased by inclusion of chromium into broiler diets that may indicate enhancement in immune response of birds.

INTRODUCTION

Trivalent chromium is well known as an essential trace element for nutrients metabolism in human and animals (Mertz 1992).

The primary role of chromium in nutrients metabolism is to potentiate the action of insulin through its presence in an organometallic molecule, called the glucose tolerance factor (GTF) (Anderson 1987; Sahin *et al.* 2001). Insulin has been shown to increase the glucose and amino acids uptake into muscle cells to regulate energy production, muscle tissue deposition, fat metabolism and cholesterol utilization. (Moat, 1994; Olin *et al.* 1994) chromium has been included in poultry researches as an inorganic compounds like chromium chloride (CrCl₃) or organic complex such as chromium picolonate, nicotine and high chromium yeast (Toghyani *et al.*; 2006).

However, chromium picolonate is the most common chromium in the poultry diets due to its high absorbability and low toxicity (Line *et al.* 1999).

Dietary supplementation of chromium picolinate could result in improved body gain, and feed conversion ratio of broiler chicks (Lien *et al.*, 1999; Toghyani *et al.*, 2006; Anandhi *et al.*, 2006). In addition it increases carcass weight breast meat yield and reduces carcass fat (Hossain *et al.*, 1998; Toghyani *et al.*, 2006). Furthermore it increases digestive enzymes activity (Karam *et al.* 2007). As well, reduces blood cholesterol, total lipids and glucose (Lien *et al.*, 1999; Sahin *et al.*, 2002).

It seems that, the favorable action of chromium picolinate may be related to its level in broiler diet. Ward *et al.*, (1995) and Lien *et al.* (1993) showed that, enriched broiler diets with 100, to 400mg chromium picolinate /kg did not affect their growth or carcass composition.

Therefore, the objectives of this study were to evaluate the effects of different graded levels of chromium picolinate on performance, carcass traits, hematological parameters and digestive enzyme activity in broiler chicks.

MATERIALS AND METHODS

This experiment was conducted at "Broiler Nutrition Unit" Faculty of Agric., Ain Shams Univ. Three hundred one-day Cubb broiler chicks were allocated randomly into five treatment groups of 60 birds and divided into three replicates with 20 birds each. Four experimental groups of birds were fed on a corn-soybean basal diet supplemented with 0.25 mg, 0.5 mg, 0.75mg or 1 mg/kg of chromium picolinate whereas the fifth group was control.

Two basal diets were formulated (Table1) to meet the nutrients requirements of chicks during the starter (0-3 wk) and grower (3-5 wk) periods according to NRC (1994). The chicks were housed in floor pens with wood-shaving litter. Electrical heaters were used to obtain suitable environmental temperature and artificial lighting was provided constantly. Water and mach feed was provided ad lib. Individual body weights were recorded weekly for each chick and the average body weight and body weight gain was calculated for each group based on the two experimental periods (0-3 & 0-5 week). Feed conversion ratio was calculated as gram feed /gram gain in the same manner.

Digestive enzymes and Carcass traits. At the end of experiment 6 birds were chosen to represent the three replicate of each treatment. The birds were slaughtered in a horizontal position to reduce the anti peristaltic movement of the intestinal segments. The slaughtered birds allowed to bleed, defeathered and internal organs were separated.

For digestive enzymes activity the content of jejunum and ileum were collected, weighed and kept in equal volumes of buffer saline solution. The contents were then individually centrifuged (6000 rpm for ten min.) and the supernatant fluids were decanted and used for the determination of some digestive enzymes activity. Amylase activity was determined using the method described by Pinshasov and Noy. (1994), lipase activity according to Skalan, *et al.* (1975) and both Trypsine and Chemotrypsine according to Skalan and Helevy (1985).The enzymes were assayed in four chicks of slaughtered birds.

Table (1): Composition of experimental basal diets

Ingredients	Starter (0-3 wks)	Grower (3-5 wks)
Yellow corn	55.8	59.71
Soybean meal (44%)	34.32	30.00
Corn gluten	3.33	2.80-
Vegetable oil	2.79	4.00
Dicalcium phosphate	1.94	1.67
Limestone	1.14	1.14
Common salt	0.25	0.25
Vit. & min. premix*	0.25	0.25
DL. Methionine	0.18	0.18
Total	100	100
Calculated composition		
Crude protein%	22.00	20.00
M.E. Kcal/kg	3000	3100
% Calcium	0.97	0.91
% Available phosphours	0.50	0.45
%Methionine + cystein	0.91	0.78
%Lysine	1.10	1.10

Composition of vitamin and minerals premix. Each 3 kg of vitamin and minerals mixture contain: 12000000 IU vitamin A; 2000000 IU D3; 10g E; 1g K; 1g B1; 5g B2; 1500mg B6; 10mg B12; 10g Pantothenic acid; 20g Nicotinic acid; 1g Folic acid; 50mg Biotin; 500g choline chloride; 4g copper; 300mg iodine; 30g iron; 60g Manganese; 50g Zinc; and 100mg selenium

For carcass traits Bursa, Spleen and Thymus were weighted. Head and Leg were separated and dressed carcass was weighed and calculated as percentage of live body weight. The dressed carcass was divided into breast, thigh, drumstick, two wings, back and neck which were weighed and calculated as percentage of dressed carcass weight.

Blood constituents. Blood samples were collected in heparin-zed tubes and centrifuged immediately for 15 minutes to separate plasma that was decanted and frozen up to chemical analysis.

Plasma total protein was determined according to Biuret method described by Henery (1964) and albumin according to Doumas *et al.* (1971). Plasma globulin was obtained by subtracting albumin from total protein values. Plasma total lipid was determined according to Knight *et al.* (1972) and total cholesterol according to Watson (1960). High density lipoprotein cholesterol (H.D.L.) and triglyceride was assayed by using kits of Diamond Diagnostics Company. Plasma concentration of Glucose was determined immediately before plasma frozen by using kit of Spectrum diagnostic company.

Statistical Analysis. Statistical analysis was computerized by statistical program SAS (1994), with using Duncan's multiple range tests to separate means.

RESULTS AND DISCUSSION

Performance aspects. Body weight of broiler chicks (Table 2) were positively affected by adding chromium picolinate into their diets either at 3 or 5 week of age. The level of 0.5mg/kg diet of cr pico was the most effective in increasing body weight which was followed by 0.75 mg/kg. This result is in agreement with those obtained by Kim *et al.* (1996) and Bhuvensh *et al.* (2004) who observed an increase in body weight of broiler chicks fed chromium-supplemented diets.

Table(2):Effect of chromium picolinate on performance of broiler hicks.

Items	Control	Cr Pico 0.25 mg	Cr Pico 0.50 mg	Cr Pico 0.75 mg	Cr Pico 1 mg
At: 0-3 weeks					
Body wt. (g)	710.3 ^b ±15.7	706.5 ^b ±13.9	758.9 ^a ±13.1	744.8 ^{ab} ±15.6	731.2 ^{ab} ±13.4
Feed intake (g)	1390 ^{ab} ±28.2	1282.5 ^b ±25.6	1368.5 ^a ±25.7	1351 ^{ab} ±25.9	1302 ^{ab} ±28.8
Feed conversion g feed/g gain	2.02±0.08 (100)*	1.93±0.01 (105)	1.87±0.03 (108)	1.92±0.01 (105)	1.89±0.06 (107)
At:0-5 weeks					
Body wt. (g)	1788.5 ^b ±30.2	1814 ^{ab} ±26.5	1888.3 ^a ±30.4	1872.7 ^{ab} ±30.1	1824.6 ^{ab} ±18.9
Feed intake (g)	3735.7 ^a ±76	3578.0 ^a ±28.3	3850.5 ^a ±95	3619.5 ^a ±70.2	3697.3±116
Feed conversion g feed/g gain	2.14 ^a ±0.02 (100)*	2.02 ^{ab} ±0.01 (106)	2.09 ^{ab} ±0.03 (102)	1.97 ^a ±0.04 (109)	2.07 ^{ab} ±0.08 (103)

a-b differences within rows, means with no common superscripts differ significantly (p≥0.05)

() * values as percentage to control.

Feed consumption values were not affected by feeding diets enriched with chromium picolinate at the 5 or 3 week of age. Anandhi *et al.* (2006) and Motozona *et al.* (1998) did not record any effect on feed consumption of broiler chicks fed excessive chromium diets.

Feed conversion ratios (g feed/g gain) were enhancing by enriching Cr picolinate into broiler diets, this improvements were related as percentage to control and ranged between 5% to 8% at 3 week of age and from 2% to 9% at 5 week of age. Kim *et al.* (1995) and Uyanik *et al.* (2002) reported that supplementation of organic chromium significantly improved feed conversion ratio of broiler chicks.

Generally, the beneficial effects of Cr picolinate supplementation may be obtained since most broiler diets are primarily composed of ingredients from plant origin which are usually low in chromium content (Anandhi *et al.* 2006).

In addition, many illustration have been introduced to reveal the beneficial action of organic chromium on performance of broiler chicks, Evans and Bowman (1992) and Linderman *et al.* (1995) demonstrated that, Cr picolinate increase amino acids and glucose uptake by muscle cells that may enhance protein and energy utilization, consequently improve performance characteristics. Karam *et al.* (2007) and Kornegay *et al.* (1997) related the beneficial action of Cr supplementation to increase digestive enzyme activity

and nutrients utilization. Furthermore, supplemental chromium may enhance the immune response (Uyanik *et al.* 2002) and ameliorate the negative effect of environmental stress (Sahin *et al.* 2002).

In the current experiment Cr inclusion increase the digestive enzyme activity (Table 3). Consequently increase nutrients utilization and improve performance aspects.

Table (3): Effect of chromium picolonate on Amylase, Lipase, Trypsine and chemotrypsine in jejunum and ileum

parameters	Control	Cr Pico 0.25 mg	Cr Pico 0.50 mg	Cr Pico 0.75 mg	Cr Pico 1 mg
In Jejunum					
Amylase (mg/dl)	1.78±0.285	2.07±0.398	2.55±0.081	2.14±0.098	2.36±0.165
Lipase (mg/dl)	10.36±0.727	12.20±1.80	13.31±0.707	10.73±0.999	12.10±1.65
Trypsine (mg/dl)	21.42±2.75	21.09±3.33	26.83±1.30	22.11±3.27	21.91±4.17
Chemotrypsine (mg/dl)	14.78±1.16	15.56±3.72	15.85±1.02	16.92±1.62	14.56±2.07
In Ileum					
Amylase (mg/dl)	1.86±0.146	2.00±0.182	2.08±0.112	2.10±0.140	1.85±0.113
Lipase (mg/dl)	9.40±0.512	8.66±1.26	9.25±0.689	8.57±0.574	8.71±0.833
Trypsine (mg/dl)	7.66±0.542	9.01±0.389	7.32±0.774	8.52±0.348	7.68±0.254
Chemotrypsine (mg/dl)	8.27 ^a ±1.09	7.78 ^{ab} ±0.479	6.19 ^b ±0.066	6.65 ^{ab} ±0.151	7.52 ^{ab} ±0.634

a-b within rows, means with no common superscripts differ significantly ($p \geq 0.05$)

Digestive enzymes activity Amylase activity was elevated in jejunum and ileum due to enriching broiler diets with chromium picolonate, however the increment lacked significant. This result is in accordance with those obtained by Karam *et al.* (2007) who observed an increase in amylase enzyme in digestive tract of broiler chicks fed diets enriched with chromium chloride. It could be suggested that the excessive chromium supplementation may potentate insulin activity and increase cell uptake of glucose which incorporated into glycogen (Wayne *et al.* 1989). This process may stimulate amylase enzyme activity to breakdown more starch into glucose for compensating the ratio of glucose used up by high insulin activity. Hossain *et al.* (1998) reported that, chromium is an integral component of glucose tolerance factor, which regulate energy production.

There was a non slight significant increase in lipase enzyme activity of jejunum by feeding chromium supplemented-diets. It may be assumed that, supplemental chromium could stimulate lipid metabolism throughout its effect on lipase enzyme (Karam *et al.* 2007; Steele and Rasebrough 1981). Trypsine and chemotrypsine enzymes were not affected by chromium addition, and their values have not a proper symmetric trend. This result is in harmony with those of Karam *et al.* (2007) who did not observe any effect on Trypsine or chemotrypsine enzymes in broiler chicks, due to feeding Cr chloride supplemented-diets. It is likely that, the beneficial effect of chromium supplementation is more related to carbohydrates and lipid metabolism via its interference with insulin secretion, since chromium influence on digestive-protein enzymes (Trypsine, chemotrypsine) was not obvious.

Carcass traits. Dressed carcass percentages were increased due to adding chromium picolonate into diets, beyond that, the increment was significant at levels of 0.5mg, 0.75mg and 1 mg Cr picolonate (Table 4). As well, breast yield percentages were increased significantly due to adding 0.75 mg Cr picolonate into broiler diets while the increment in other treatments lacked significance. Both of thigh and drumstick were not affected by treatments. There were a reduction in wings and neck percentage due to feeding chromium diets while; back portion percentage was not affected clearly.

Table (4): Effect of chromium picolonate on carcass yield percentage and some immunity and internal organs

Items	Control	Cr Pico 0.25 mg	Cr Pico 0.50 mg	Cr Pico 0.75 mg	Cr Pico 1 mg
Dressed carcass% *	68.7 ^b ± 1.14	70.98 ^{ab} ±0.28	72.72 ^a ±0.54	71.99 ^a ±0.57	71.88 ^a ±1.29
Thighs % **	16.85±0.57	16.15±0.44	15.33±0.126	14.78±1.11	16.16±0.436
Drumstick % **	12.56±0.42	12.35±0.15	12.74±0.36	12.31±0.37	11.82±0.09
Breast % **	26.55 ^c ±0.87	27.24 ^b ±1.08	26.73 ^b ±0.47	32.75 ^a ±2.04	28.06 ^b ±1.05
Neck % **	6.29 ^c ±0.69	5.48 ^{ab} ±0.26	5.37 ^{ab} ±0.79	4.24 ^b ±0.36	5.39 ^{ab} ±0.62
Wings % **	12.73±0.74	12.01±0.34	11.89±1.17	10.74±0.34	11.01±0.52
Back % **	24.39 ^c ±0.33	25.53 ^{bc} ±0.54	27.18 ^{ab} ±0.46	23.58 ^b ±1.53	28.32 ^a ±0.68
Bursa**	0.26±0.018	0.21±0.03	0.24±0.05	0.17±0.03	0.18±0.02
Thymus**	0.553±0.061	0.515 ^{ab} ±0.114	0.46±0.072	0.49±0.038	0.63±0.074
Spleen**	0.183 ^a ±0.024	0.128 ^b ±0.016	0.123 ^b ±0.009	0.135 ^b ±0.005	0.123 ^b ±0.003

a-b-c within rows, means with no common superscripts differ significantly (p≥0.05)

* % percentage of live body weight

** % percentage of dressed carcass weight

The present results are in a good agreement with the finding of Toghiani *et al.* (2006) and Sahin *et al.* (2002) who observed an increase in carcass yield of broilers fed supplemental Cr picolonate. As well, Hossain *et al.* (1998) reported that carcass percentage and breast meat yield was significantly influenced by Cr yeast supplementation.

The beneficial effect of feeding supplemental chromium on carcass and breast meat yield may be related to its role in increasing amino acids uptake, incorporation and utilization (Hossain *et al.* 1998), ultimately increase protein deposition, muscles synthesis and meat yield (Sahin *et al.* 2002).

Neither bursa nor thymus weight was affected by treatments whereas spleen weight significantly decreased due to feeding Cr picolonate supplemented diets. El-Hommosany (2008) did not find any effect on bursa or spleen weight due to adding Cr chloride into quail diets.

Blood constituents. Glucose concentration in plasma reduced as a result of feeding dietary chromium (Table 5). This result is well accepted since chromium stimulates glucose metabolism via potentiates insulin hormone action. Sahin *et al.* (2002) found that, supplementation of Cr picolonate into broiler diet, decrease plasma glucose and increase plasma insulin concentration.

Lipid derivatives. There were reductions in plasma total lipid, triglycerides and cholesterol due to feeding excessive chromium picolonate (Table 5). This reduction was significant in total lipid and lacked significance in cholesterol

and triglycerides values while H.D.L. cholesterol was not affected. This result confirms the finding of Abraham *et al.* (1982 a, b) that, chromium is essential for lipid metabolism Uyanik *et al.* (2002) observed a reduction in plasma total cholesterol, triglycerides of quails fed supplemental chromium and illustrated that, the decrease in lipid parameters could result from the increasing activity of insulin that depressing the fatty acid synthesis by increasing glycogen build up.

Table (5): Effect of chromium picolonate on some blood constituents

Items	Control	Cr Pico 0.25 mg	Cr Pico 0.50 mg	Cr Pico 0.75 mg	Cr Pico 1 mg
Glucose mg/dl	239.75 ^a ±13.29	197.25 ^b ±2.66	210.0 ^c ±3.83	213.33 ^b ±5.46	205.2 ^b ±10.82
Cholesterol mg/dl	149.50±3.48	124.25±12.11	137.0±13.65	128.33±13.98	120.75±9.61
HDL mg/dl	88.75±5.62	79.0±6.63	92.0±2.65	87.67±1.76	88.75±4.27
Total lipid mg/dl	4.78 ^a ±0.35	3.58 ^b ±0.29	4.67 ^a ±0.43	3.33 ^b ±0.13	3.38 ^b ±0.33
Triglycerides mg/dl	174.0±8.91	144.25±13.59	164.0±17.21	124.0±14.76	146.0±23.09
Total protein g/dl	2.43 ^c ±0.12	2.74 ^{bc} ±0.04	2.80 ^{bc} ±0.08	3.03 ^b ±0.26	3.60 ^a ±0.10
Albumin	1.39±0.14	1.49±0.07	1.48±0.04	1.26±0.09	1.48±0.16
Globulin g/dl	1.04 ^c ±0.19	1.26 ^{bc} ±0.10	1.33 ^{bc} ±0.09	1.77 ^{ab} ±0.34	2.12 ^a ±0.24
Alb. / glob. ratio	1.52 ^a ±0.35	1.22 ^{ab} ±0.16	1.14 ^{ab} ±0.11	0.83 ^b ±0.21	0.75 ^b ±0.17

a-b-c within rows, means with no common superscripts differ significantly ($p \geq 0.05$)

Blood protein and immunity. Plasma proteins (Table 5) increased significantly due to inclusion of Cr picolonate into broiler diets. Simultaneously plasma albumin values were not affected by treatments so that the elevation in plasma total protein was reflected as an increment in plasma globulin values which were affected significantly by treatments and were parallel to the total protein values. This result confirms the finding of Sahin *et al.* (2002) who observed an increase in blood total protein with feeding supplemental chromium. A lack of chromium effect on plasma albumin was also reported by Karam *et al.* (2007) and Ibrahim *et al.* (2005). The high concentration of globulin inherent with low ratios of albumin to globulin (A/g) may indicate an enhancement in immune function of birds fed dietary chromium. Uyanik *et al.* (2002), observed an increase in immunoglobulin levels and antibody titers of broiler fed diets supplemented with chromium.

REFERENCES

- Abraham, A. S., Sonnenblick, M., Eini, M. (1982a). The action of chromium on serum lipids and on atherosclerosis in cholesterol fed rabbit. *Atherosclerosis*, 42:185-195.
- Abraham, A.S., Sonnenblick. M., Eini, M.(1982b). The effect of chromium on cholesterol includes atherosclerosis in rabbit. *Atherosclerosis*, 42: 371-372.
- Anandhi, A.; R. Mathivanan; K. Viswanathan and B. Mohan (2006). Dietary inclusion of organic chromium on production and carcass characteristics of broilers. *International J. of poultry Sci.* 5 (9) 880-884.
- Anderson, R.A. (1987): Chromium trace elements in human and animal nutrition. New York Academic press, pp: 225-244.

- Bhuvensh Kumar, Kusum late pandey and Narendra Kumar (2004). Influence of chromium supplementation on broiler performance in Hills. *Ind. Vet. J.*, 81: 169-171.
- Doumase, B. T.; Watson, W. A.; and Biggs, H. S. (1971). Albumin standards and measurement of serum albumin with bromocresol green. *Clin. Chem. Acta.*, 31:87-96.
- El-Hommosany, Y.M. (2008). Study of the physiological changes in blood chemistry, Humoral immune response and performance of quail chicks fed supplemental chromium. *International J. of poultry Sci.* 7 (1) 40-44.
- Evans, G. W. and T. D. Bowman (1992). Chromium picolinate increases membrane fluidity and rate of insulin internalization. *J. Inorgan. Biochem.* 48: 234-350.
- Gursory, U. (2000). Chromium in broiler diets. *Feed Int.*, March, 24-26.
- Henery R. J. (1964). A calorimetric method for the determination of the total protein. *Clinical chemical* Harper and Row Publisher New York 1964.
- Hossain, Sazzad M.; Sergio L. Barreto; Carlos G. Silva (1998). Growth performance and carcass composition of broilers fed supplemental chromium from chromium yeast. *Animal feed science technology* 71: 217-228.
- Ibrahim, K. A. (2005). Effect of dietary chromium on growth performance, carcass characteristics and some blood parameters of broilers. *Egypt. Poult. Sci.*, 25: 137-185.
- Karam, M. A.; N. K. Soliman; N. M. El-Medany and I. El-Wardany (2007). Effect of supplemental chromium on performance, blood constituents and digestive enzymes activity of broiler chicks. *J. Agric. Sci. Mansoura Univ.*, 32(9): 7217-7227.
- Kim, Y. H.; I. K. Han; I. S. Shin; B. J. Chae and T. H. Kang (1996). Effect of dietary excessive chromium picolinate on growth performance nutrient utilization and serum traits in broiler chicks. *Asian Aust. J. Animal Sci.* 9: 349-354.
- Kim, J.D.; Han, I.K.; Shin, I.S.; and Chae, B.J. (1995). Effect of supplemental chromium picolinate on growth performance, carcass quality and serum traits in broiler chicks. *Asian J. Anim. Sci.* 8:341-347.
- Knight, J.; A. Andersonis and J. M. Rawal (1972). Chemical basis of the sulphate vanillin reaction for estimating total serum lipid. *J. Biol. Chem.* 226-497.
- Kornegay, E.T., Wang, Z., Wood ,C.M., and Linderman, M.d., (1997). Supplemental chromium picolinate influences nitrogen balance, dry matter digestibility, and carcass traits in growing finishing pigs. *J. Anim. Sci.* 75:1319-1323.
- Linderman, M.D., Wood, C.M., Harper, A.F., Kornegay, E.T.& Anderson, R.A.(1995). Dietary chromium picolinate additions improve gain: feed and carcass characteristics in growing finishing pigs and increase litter size in reproducing sows. *Journal of Animal Science*, 73:457-465.
- Mertz,W. (1969) Chromium occurrence and function in biological Systems. *Physiol. Rev.*, 49: 163 – 239.

- Motozona, Y.; K. Hatano; N. Sugawara and T. Ishibashi (1998). Effect of dietary picolonate on growth, carcass quality and serum lipids of female broilers. *Animal Sci. tec.*, 69: 659-665.
- Mowat, D. N. (1994). Organic chromium: a new nutrient for stressed animals. In: Lyons, T.p. and K. A. Jacques. (Eds), *Biotechnology in feed industry* University press, Nothingam, UK pp.: 227-282.
- National Research Council (1994): *Nutrition Requirements of poultry 9th Rev.* Ed. National academy press, Washington, D.C
- Olin, K.L. D. M. Starnes; W.H. Armstrong and C. L. Kern (1994). Comparative retention/absorption of Cr from Cr chloride, Cr nicotinate and Cr picolonate in a rate model. *Trace Ele. Electrol.* 11: 182.
- Pinshasov, Y. and Y. Noy (1994). Early postnatal amylolysis in the gastrointestinal tract of turkey poults. *Comp. Biochem. Physiol.*, 106: 221-225.
- Sahin, K.; Ozbey, O.; Onderci, M.; Cikim, G.; and Aysondu, M.H (2002). Chromium supplementation can alleviate negative effects of heat stress on egg production, egg quality and some serum metabolites of laying Japanese quail. *J. Nutr.* 132:1265-1268.
- SAS Institute, (1994): *SAS /STAT guide for personal computers. Version 6* (Ed. SAS Inst. Inc.,) Cary, NC.
- Skalan, D. and O. Helevy (1985). Protein digestion and absorption along the ovine gastrointestinal tract. *J. Dairy Sci.*, 68: 1678-1681.
- Skalan, D. S. Hurwitz, P. Budowski, and I. Ascarelli (1975). Fat digestion and absorption in chicks fed raw or heated soybean meal. *J. Nutr.*, 105: 57-63.
- Steele, N.C.; and Rosebrough, W. (1981). Effect of trivalent chromium on hepatic lipogenesis by the turkey poul. *Poul. Sci.*, 60: 617-622.
- Toghyani, M.; M. Shivazad; A.A. Gheisari and S.H. Zarkesh (2006). Performance, carcass traits and hematological performances of heat-stressed broiler chicks in response to dietary levels of chromium picolonate. *International J. of Poultry Sci.* 5 (1): 65-69.
- Uyanik, F.; Atasever, A.; Ozdamar, S.; and Aydin, F. (2002). Effect dietary chromium chloride supplementation on performance, some serum parameters, and immune response in broilers. *Biol.; Trace Elem. Res.* 90: 99-115.
- Watson, D. (1960): Determination of total cholesterol. *Clin. Chem. Acta*, 5: 657.
- Wayne W.C., M.P. Marilyn, A.B. Noella, H.S. Joseph and R.A. Anderson (1989). Exercise training and dietary chromium effect on glycogen, glycogen synthase, phosphorylase and total protein in rates. *American Ins. of Nutrition.* 653- 660.

تأثير إضافة بيكولونات الكروم إلى علائق كتاكيت اللحم على الاداء الانتاجي ونشاط الانزيمات الهاضمة لمى خالد بندر ، نجلاء كمال سليمان و سنبل جاسم الحمودى قسم إنتاج الدواجن - كلية الزراعة - جامعة عين شمس

اجريت هذه التجربة لدراسة تأثير المستويات المختلفه من بيكولونات الكروم على الاداء الانتاجي لكتاكيت التسمين وبعض صفات الذبيحه ونشاط الانزيمات الهاضمه.

استخدمت الدراره عدد ٣٠٠ كتكوت تسمين عمر يوم من سلاله (كب) تم توزيعها عشوائيا بالتساوى على خمس معاملات بواقع ٦٠ كتكوت بكل معامله، غذيت الطيور على عليقه قاعديه من الذرة وفول الصويا حيث اضيف لها اربع مستويات من بيكولونات الكروم وهى ٢٥ و ٥٠ و ٧٥ و ١ مجم/كجم علف والمعامله الخامسه كانت للكنترول.

أشارت النتائج الى تحسن فى وزن الجسم نتيجة لاضافه الكروم للعلائق وسجلت الكتاكيت المغذاه على ٥,٥ مجم كروم بيكولونات أعلى معدل لوزن الجسم (١٨٨٨,٣ جم) تلاها تلك المغذاه على ٥,٧٥ مجم كروم بيكولونات (١٨٧٢,٧ جم) مقارنة ب ١٧٨٨,٥ جم للكنترول. الاستهلاك الغذائى لكامل فترة التجربه لم يتأثر بالمعامله. معدلات التحويل الغذائى تحسنت بمعدل ٢٪ الى ٩٪ نتيجة التغذية على علائق الكروم وربما يرجع هذا التحسن فى الاداء الانتاجي لتأثير الكروم المنشط للانزيمات الهاضمه المحفز للمناعه مما يودى لتحسن الاستفادة من العلائق.

لم يتأثر كل من نشاط انزيم الاميليز و الليبيزو التربسين أو الكيموتربسين فى كل من اللفانفى أو الصائم نتيجة التغذية على علائق الكروم.

كان هناك زيادة معنويه فى النسبه المئويه للذبيحه ولحم الصدر نتيجة لاضافه الكروم للعلائق كذلك تأثرت النسبه المئويه للاجزاء الخلفيه للذبيحه بالمعامله.

اضافه الكروم للعلائق ادى لانخفاض مستوى الجلوكوز والليبيدات الكليه بيلازما الدم بينما لم يتأثر مستوى الكوليستيرول أو الكوليستيرول عالى الكثافه (H.D.L) وكذلك الجلسريدات الثلاثيه بالمعامله.

أرتفع مستوى البروتينات الكليه والجلوبيولين بالدم بينما لم يتأثر مستوى الاليومين وذلك ربما يشير الى وجود تحسن فى مناعه الطيور نتيجة المعامله.