

## PREVALENCE OF MOULD AND YEAST IN SOME DAIRY PRODUCTS SOLD IN MENOFIA GOVERNORATE

NAHLA, A. ABOU EL-ROOS\* and FLOURAGE, M. RADY\*\*

\* Dept. of Food Hygiene    \*\* Dept. of Mycology

Animal Health Research Institute, Shibin – El koom Branch

Received: 4. 2. 2008

Accepted: 18. 2. 2008

### SUMMARY

A total of one hundred samples (25 each) of milk powder, yogurt, cheddar cheese and Ras cheese were collected from dairy shops and super markets in Menofia Governorate. Each sample was divided into two parts, one part for mycological studies and the other part for aflatoxins. Our results revealed that, the mould and yeast could be detected in all examined samples of milk powder, yogurt, cheddar cheese and Ras cheese with mean count values of  $8.3 \times 10^2 \pm 0.2 \times 10^2$ ,  $6.9 \times 10^2 \pm 0.4 \times 10^2$ ,  $5.3 \times 10^3 \pm 2.3 \times 10^2$  and  $4.1 \times 10^3 \pm 1.6 \times 10^2$  for mould and  $0.5 \times 10^2 \pm 1.1 \times 10^2$ ,  $1.4 \times 10^2 \pm 0.5 \times 10^2$ ,  $6.5 \times 10^4 \pm 4 \times 10^4$  and  $5.4 \times 10^3 \pm 1 \times 10^3$  for yeasts respectively.

Mould and yeast were isolated at varying percentages from all examined samples. The mould genera were, *Aspergillus* spp., *Alternaria* spp., *Penicillium* spp., *Cladosporium* spp., *Mucor* spp. and *Rhizopus* spp. While yeast genera were *Candida* spp., *Rhodotorula* spp., *Saccharomyces* spp and

*Torulopsis*.

Aflatoxin M1 residues were detected in 16%, 16%, 32% and 44% of examined milk powder, yogurt, cheddar cheese and Ras cheese samples respectively. Aflatoxin B1 residues were detected in 12%, 52% and 48% of examined milk powder, cheddar cheese and Ras cheese samples respectively. On the other hand, Aflatoxin B2 were detected in 12%, 24% and 12% of examined milk powder, cheddar cheese and Ras cheese samples respectively.

From the present investigation it could be concluded that, feeding of dairy animals with aflatoxins free feedstuff, using high quality raw milk, strict hygienic measures during manufacture, packing and distribution, proper personal hygiene, adequate cleaning and sanitation of dairy equipments and proper refrigeration are vital elements in the production of high quality dairy products.

## INTRODUCTION

Milk and its products are adequate substrate for yeasts and moulds growth under suitable conditions of temperature and moisture contents ( Barrois et al , 1997).The contamination of these products may occur from the raw material or during manufacturing , storage and distribution( Kure et al., 2004). These microorganisms influence the biochemical characters and flavour of these substrate as well as their appearance rendering them commercially undesirable and often resulting in down grading of the product (Demarigny et al ,1997 and Muir and Banks,2000).Some moulds can also adversely affect human and animal health as they can produce mycotoxins which are fungal secondary metabolites formed by consecutive series of enzyme- catalyzed reactions from a few biochemically simple intermediates of primary metabolism , these mycotoxin can enter the human and animal food chain by direct or indirect contamination ( Bohra and Purohit, 2003 ) and they are related to a range of pathologies ranged from gastroenteritis to cancer as these mycotoxins are highly toxic, mutogenic,teratogenic and carcinogenic( Adams and Moss, 2000 and Li et al., 2000 and Hussein and Brasel,2001).

Mycotoxins are not destroyed by normal industrial processing or cooking since they are heat stable ( Park,2002).The real hazard of mycotoxins to human health might be due to the cumulative effect of repeated exposure to very small or even undetectable

doses of toxins in diet for comparatively long periods (Hendrickse, 1991).So that the objective of this study was to characterize the fungal contamination in some dairy products during preparation ,storage or handling to safe guard consumers from being affected and to obtain a finished product of good keeping quality.

## MATERIAL AND METHODS

### 1-Collection of samples.

A total of one hundred samples (25 each) of milk powder , yogurt , cheddar cheese and Ras cheese were collected from dairy shops and supermarkets in Menofia Governorate , each samples were transported in sterile air tight jars to the laboratory with a minimum of delay to be prepared and examined . Each sample was divided into two parts, one part for mycological studies and the other part for aflatoxin analysis

### 2- Enumeration, Isolation and Identification of fungi :

The yeast and mould counts of examined prepared samples were determined according to the technique recommended by A.P.H.A (1985). Mould isolates were identified according to Kone-man and Roberts( 1985),while yeast isolates were identified according to Looder and Kreger Van RiJ (1970).

### 3- Detection of mycotoxin:

Aflatoxin in the examined samples were extract-

ed, purified and determined by using thin layer chromatography as described by A.O.A.C (1984). **RESULTS**

**Table (1): Total mould count/ ml of examined samples:**

Type of samples	No. of samples	Positive sample		Count/ml		
		No.	%	Min	Max	Mean $\pm$ S.E
Milk powder	25	22	88	1.2 x 10	1.2x10 <sup>3</sup>	8.3x10 <sup>2</sup> $\pm$ 0.2x10 <sup>2</sup>
Yogurt	25	21	84	1.0 x 10	2.7x10 <sup>2</sup>	6.9x10 $\pm$ 0.4x10
Cheddar cheese	25	18	72	1.0x10	3.0x10 <sup>4</sup>	5.3x10 <sup>3</sup> $\pm$ 2.3x10 <sup>3</sup>
Ras cheese	25	20	80	8.2x10	9.6x10 <sup>3</sup>	4.1x10 <sup>3</sup> $\pm$ 1.6x10 <sup>2</sup>

**Table(2): Total yeast count/ ml of examined sample:**

Type of samples	No of samples	Positive samples		Count/ ml		
		No	%	Min	Max	Mean $\pm$ S.E
Milk powder	25	16	64	2.0 x10	3.9x10 <sup>2</sup>	0.5x10 <sup>2</sup> $\pm$ .1x10
Yogurt	25	20	80	3.0 x10	7.2x10 <sup>2</sup>	1.4x10 <sup>2</sup> $\pm$ .5x10 <sup>2</sup>
Cheddar cheese	25	25	100	9.0 x10	3.5x10 <sup>5</sup>	6.5x10 <sup>4</sup> $\pm$ 0.4x10 <sup>4</sup>
Ras cheese	25	19	76	6.0 x10	1.1x10 <sup>3</sup>	5.4x10 <sup>3</sup> $\pm$ .1x 10 <sup>3</sup>

**Table (3): Prevalence of individual members of fungi in examined samples of milk products**

Genera and species	Milk powder		Yogurt		Cheddar cheese		Ras cheese	
	No	%	No	%	No	%	No	%
<i>A. flavus</i>	16	64	1	4	4	16	3	12
<i>A. niger</i>	13	52	2	8	19	76	7	28
<i>A. Fumigatus</i>	5	20	5	20	3	12	2	8
<i>A. ochraceus</i>	6	24	3	12	8	32	2	8
<i>A. nidulans</i>	-	-	-	-	6	24	-	-
<i>A. candidus</i>	2	8	-	-	-	-	-	-
<i>A. Tserus</i>	1	4	-	-	-	-	1	4
<i>Alternaria spp.</i>	-	-	3	12	-	-	-	-
<i>Penicillium spp.</i>	8	32	7	28	3	12	8	32
<i>Fusarium spp.</i>	1	4	4	16	3	12	4	16
<i>Cladosporium spp.</i>	-	-	5	20	3	12	6	24
<i>Mucor spp.</i>	6	20	2	8	2	8	-	-
<i>Rhizopus spp.</i>	2	8	1	4	2	8	-	-
<i>Candida spp.</i>	-	-	4	16	13	52	5	20
<i>Rhodotorula spp.</i>	-	-	2	8	4	16	2	8
<i>Saccharomyces spp.</i>	-	-	-	-	6	24	-	-
<i>Trulopsis spp.</i>	-	-	1	4	-	-	-	-

**Table (4): Incidence and quantitative estimation of aflatoxins in examined samples (ug/kg)**

Types of samples	M1				B1				B2						
	+ve samples		Min	Max	Mean	+ve samples		Min	Max	Mean	+ve samples		Min	Max	Mean
	No	%				No	%				No	%			
Milk powder	4	16	0.6	1.5	0.9±0.06	3	12	0.5	1.9	0.8±0.01	3	12	0.9	2.1	1.9±0.1
Yogurt	4	16	2.0	3.0	2.3±0.5	-	-	-	-	-	-	-	-	-	-
Cheddar	8	32	3.1	4.0	3.5±0.8	13	52	3.1	7.0	5.4±0.6	6	2.4	2.1	3.2	3.8±0.2
Ras	11	44	1.9	5.3	3.5±0.02	12	48	2.1	4.3	3.9±0.4	3	12	1.6	3.3	2.9±0.04

## DISCUSSION

Results reported in Table (1) revealed that, the examined samples of milk powder, yogurt, cheddar cheese and Ras cheese were contaminated with moulds at the percentages of 88,84,72 and 80% respectively with mean values of  $8.3 \times 10^2 \pm 0.2 \times 10^2$ ,  $6.9 \times 10^0 \pm 4 \times 10$ ,  $5.3 \times 10^3 \pm 2.3 \times 10^2$  and  $4.1 \times 10^3 \pm 6 \times 10^2$  respectively. Nearly similar results were reported by Abou Zeid (1996), Hassan and Hammad (2001), Ali et al., (2004) and Amany et al., (2005).

As shown in Table (2) samples of milk powder, yogurt, cheddar and Ras cheese were contaminated with yeast at the percentages of 64,80,100 and 76% respectively with mean values of  $0.5 \times 10^2 \pm 1.1 \times 10$ ,  $1.4 \times 10^2 \pm 0.5 \times 10^2$ ,  $6.5 \times 10^4 \pm 4 \times 10^4$  and  $5.4 \times 10^3 \pm 1 \times 10^3$  respectively. These results were similar to these obtained by Bullerman (1981); Ismail and Saad (1995); Zin El-din and El-sawah (1997), and Dermarigny et al. (1997).

Although some yeasts and moulds may be used in the manufacture of many dairy products and milk beverages to attain specific flavour or aiding in ripening of certain type of cheese (Fadda et al., 2004). Yet many species of yeast and moulds are implicated in many defects in milk and its products, as off flavour, colour defects and spoilage of such food (Filtenborg et al., 1996 and Sarias et al., 1996).

our results in Table(3) show that in milk powder samples species of *Asperigillus* as *A.Flavus*, *A. niger*, *A. fumigatus*, *A. Ochracus*, *A. Candidus* and *A. Tserus* isolated in percentages of 64, 52, 20, 24, 8 and 4% respectively. While *penicillium* Spp was present in high percentage of 32%. *Fusarium* Spp, *Mucor* Spp and *Rhizopus* Spp were detected in 4%, 20% and 8% respectively. Similar results were obtained by Hassan and Hammad (2001). Yeasts and moulds are widely distributed as environmental contaminants of air, water, soil and dust, so the presence of moulds and yeasts in milk products may be attributed to poor sanitary practices during manufacturing, packing and distribution or the use of bad quality raw ingredients (Ray, 1996).

In samples of yogurt, *A.Flavus*, *A. niger*, *A. fumigatus*, and *A. Ochracus*, were present in percentages of 4, 8, 20 and 12% respectively. *Fusarium* Spp and *Candida* spp were present in percentages of 16%. Also *Rhizopus* Spp and *Torulopsis* spp were present in the same percentage of 4%. Moreover, *Alternata* Spp and *Rhodo Torula* Spp were present in the percentage of 8 and 12% respectively. Similar results were obtained by Eman and Abdel Satar (2003). While lower results were obtained by Hassan (1999). Yogurt is considered as an excellent medium for growth of many fungi because of its high water, low pH and its good quality protein (Marth and Steel, 2001). For samples of cheddar cheese, *Asperigallus* spp

were present as *A. Flavus*, *A. niger*, *A. fumigatus*, *A. ochraceus* and *A. nidulans* in percentages of 16, 76, 12, 32 and 24% respectively. *Candida* spp were present in high percentages of 52% followed by *Saccharomyces* in a percentages of 24% and *Rhodotorula* spp in a percentages of 16%

*Penicillium* spp, *Fusarium* spp and *Cladosporium* spp were present in the same percentage of 12% also *Mucor* spp and *Rhizopus* spp were present in percentage of 8%. Nearly similar results were reported by Lund et al (1995) and Fardos(2001).

In samples of Ras cheese, *Asperigallus* spp were represented as *A. flavus*, *A. niger*, *A. fumigatus*, *A. ochraceus* and *A. tsresus* in percentages of 12, 28, 8, and 4% respectively. Meanwhile *Penicillium* spp, *Fusarium*, *Cladosporium* spp were present in percentage of 32, 16 and 24% respectively. *Candida* and *Rhodotorula* spp were present in the percentage of 20% and 8%. Nearly similar results were obtained by Salwa(1999) and Corbo et al (2001)

Kure et al (2004) reported that, the major source of cheese contaminants were found in samples of air, equipment and plastic film. He added that air is considered as the major source of cheese contaminants so high quality air with low number of contaminants in production room especially the wrapping room is important in order to reduce mould contamination.

As shown in Table (3). *Asperigallus*, *penicillium*, *Cladosporium* and *candida* spp were the most common genera recovered from the four types of examined milk products. Similar results were recorded by Sampayo et al (1995), Elprince and Ismail (1998) and El-Sherif (2000) and Montagna et al., (2004). Growth of moulds as *Mucor*, *Alternaria*, *Penicillium* and *Geotrichum* on the surface of dairy products lead to off flavour while growth of *Penicillium*, *Cladosporium*, *Asperigillus* and *Mucor* may be responsible for bitterness and rancidity of cheese. *Penicillium* species may lead to softness of the surface of cheese (Minervini et al., 2001). Some species of *Asperigallus*, *Cladosporium*, *Penicillium* and *Fusarium* were responsible for kerato-conjunctivitis in man while *Asperigillus niger* cause otomycosis and allergic condition, some species of *penicillium* have been associated with pulmonary infections, urinary tract infections and yellow rice disease causing several deaths in man (Nilesen et al., 1998).

From the results reported in Table (4), regarding the milk powder samples, Aflatoxin M1, B1 and B2 can be detected in 16%, 12% and 12% of examined samples respectively with average concentrations of  $0.9 \pm 0.06$ ,  $0.8 \pm 0.01$  and  $1.90 \pm 1.1$   $\mu\text{g}/\text{kg}$  respectively. These findings agreed with that reported by Olivera et al (1997) and Chia et al (2005). *Penicillium* species mycotoxins was failed to be detected as that reported by Hamilton (1975) who stated that milk powder are substrates which have low carbohydrate and high protein

content which combine with toxins of penicillium species.

Neither storage or processing determine a reduction of aflatoxin content in milk powder ( Deveci and Emel, 2006).

For samples of yogurt, 16% of the examined samples proved to be contain aflatoxin M1 at an average concentration of  $2.3 \pm 0.5 \mu\text{g}/\text{kg}$ . A nearly similar results were recorded by Farag( 2002) and Martins and Martins ( 2004). A highly reduction of AFM1 in yoghurt is due to degradation of AFM1, caused by enzymatic, microbial and particularly acid coagulation of manufactured yoghurt. Moreover, the lactic acid produced in yoghurt processed a detoxification action as the toxic form of AFB1 into non toxic form "B2a" (Galvano et al ., 1996)

Aflatoxins M1, B1 and B2 can be detected in 32%, 52% and 24% of the examined samples of Ras cheeses with mean values of  $3.5 \pm 0.8$ ,  $5.4 \pm 1.6$  and  $3.8 \pm 0.2 \mu\text{g}/\text{kg}$ . Nearly similar results were reported by EL Gerbi et al ( 2004) and Marnal et al ( 2006).Aflatoxins in cheese resulted from the presence of it in raw milk as a consequence of carry over of aflatoxins M1 from contaminated cow food to milk synthesis of aflatoxins ( B1, B2,G1 and G2) by *A. flavus* and *A. parasiticus* growing in cheese and the presence of these toxins in the dried milk used to enriched the

milk used to make cheese or other dairy products( Kaya et al, 1998 and Aycicek et al., 2002).

For samples of Ras cheese, aflatoxins M1, B1 and B2 can be detected in 44%, 48% and 12% of the examined samples with mean values of  $3.5 \pm 0.02$ ,  $3.9 \pm 0.4$  and  $2.9 \pm 0.04 \mu\text{g}/\text{kg}$ . Nearly , similar results were reported by Abou Zeid et al( 1996) and Manetta et al (2005).

Milk contained mycotoxins considered as a vehicle for transferring such toxins into cheese where it appears to be associated with casein, the cheese is considered as a curdle from milk protein so, it may contain 3.5- 5 times more toxins than the level present in milk from which it was made ( Chapman and Sharpe, 1990).. Mycotoxin cause alteration in carbohydrate and lipid metabolism, toxic damage of the liver disturbance in blood clotting mechanism and with immune response. Aflatoxins were classified as group1 carcinogens in 1993 by the WHO international agency for research on cancer ( IARC) ( Pestka, 1995).

Naturally occurring toxicant contamination of foods with mycotoxins is unavoidable and unpredictable and posses a unique challenge to food safety, so practical decontamination procedures must include the following hygienic procedures: inactivate, destroy or remove the toxin, not produce or leave toxic residues in the food , retain the nutritive value of the food, not alter the ac-

ceptability or the technological properties of the product and if possible destroy fungal spores ( Park,2002). Also there is a must to increase the intake of diet components such as vitamins , antioxidants and substances known to prevent carcinogenesis( Creppy, 2002). From the present investigation it could be concluded that, feeding of dairy animals with aflatoxins free feedstuff, using high quality raw milk , strict hygienic measures during manufacture, packing and distribution , proper personal hygiene, adequate cleaning and sanitation of dairy equipments and proper refrigeration are vital elements in the production of high quality dairy products

## REFERENCES

- Abou Zeid, A.M. Hassan, A.A. and Ragheb, R.R (1996): Mycological studies on hard Rome and skim milk soft cheese( Kareish). *Vet. Med. J. Giza*. 44: 2A; 113- 121.
- Adams, M.S and Moss, M.C. ( 2000): *Food microbiology* 2nd Ed. Royal Society of chemistry. Thomas Graham house, Science Park., Milton Road Cambridge CB 40 WF, UK.
- Ali, M.M.; Nahed M. Wahaba and Seham, A. Farrag( 2004): Microbiological evaluation of Assiut Market yoghurt through the shelf life time in refrigerator. *Assiut Vet Med. J.* 50, 101.
- Amany,M. Shalaby, Marcel, F. Galab and Azza, A. EL Taby.( 2005): Prevalence of Mould and Yeast in some dairy products. *Egypt. J. Comp. Path. And Clinic. Path.* 18, 1: 132- 140.
- American Public Health Association" APHA" ( 1985): *Standard methods for the examination of dairy products*.15 th ed.,Washington, DC. Association Of official Analytical Chemists" A.O.A.C.( 1984): *Official Methods of Analysis*. 14 in ed. Washington, DC.
- Aycicek, H. Yarsan, E Sarimehmetogalu, B. and Cakmak, O. ( 2002):Aflatoxin M1 in white and butter consumed in Istanbul.Turkey,*Vet. Human Toxicol.*44 ( 5): 295-296.
- Barrios, M,T.; Medina, L.M., Cordoba, M.G. and Jordano, R. ( 1997): Aflatoxin producing strains of *Asperigillus Flavus* isolated from cheese. *J. Food Prot.* 60:192-194.
- Bohra,N.K. and Purohit, D.K.( 2003): Fungal toxicity with special reference to mycotoxins. *J. Environ. Biol.* 24( 3): 213- 221.
- Bullerman, L.B. ( 1981): Public Health significance of moulds and mycotoxins in fermented dairy products. *J. Dairy Sci.* 64: 24- 39.
- Chapman, H.R. and Sharpe, M.E.C.( 1990): *Microbiology of cheese*.*Dairy microbiology*,(2) 203-290.
- Chia, Y. Ch; Wen, J.L and Kai, Y.P ( 2005): Determination of Aflatoxin M1 in milk powder using high flow solid phase extraction and liquid chromatography tandem mass spectrometry. *J. Agri. Food Chem*, 53( 22), 8474- 8480.
- Corbo, M R. , Lanciotti, R., Abenzio, M. and Sinigaglia, M. ( 2001): occurrence and characterization of yeasts isolated from milk and dairy products of Apulia Region. *Int. J. Food Microbiol.* 69( 1-2): 147-52.
- Creppy, E.E.( 2002): Update of survey, regulation and toxic effects of mycotoxins in Europe. *Toxicollett.* 28: 127( 1-3), 19-28.
- Demarigny, Y.; Beuvier, E.; Buchin, S.; Pochets, S. and Grappin,R.( 1997) : Influence of raw milk microflora on the character of Swiss type cheese.*Biochemical and sensory characteristics Lait.* 77: 151- 167.
- Vet. Med. J., Giza. Vol.56, No. 1(2008)*



- Deveci, O and Sezgin, E. ( 2006): Changes in concentration of aflatoxin M1 during manufacture and storage of skim milk powder. *J. Food . Prot.* 69( 3), 682- 685
- El- Prince, Enas and Ismail, M.A. ( 1998): Microbiological quality of Mozzarella cheese. *Assiut Vet. Med. J.* 39( 77): 94- 109.
- EL- Shrief, Lamiaa, M.T.( 2000): Incidence of mycoflora and some mycotoxins in locally manufactured cheese. M.V.Sc. Thesis. Assiut Uni.
- EL-Gerbi, A.M.; Aidoo, K.E.; Candlish, A.A. and Tester, R.F. ( 2004): Occurrence of aflatoxins in randomly selected North African milk and cheese samples. *Food Ad-dit. Contam.* 21( 6): 592- 7.
- Eman, K. Ahmed and Abdel- Satar, M.A. ( 2003): Myco- logical quality of Laben Rayeb sold in Assiut City. *As- siut Vet. Med. J. Giza* 49 (99): 70- 80.
- Fadda, M.E.; Mossa, V.; Pisano, M.M.; Deplano, M. and Cosentin, O.S.( 2004): Occurrence and characterization of yeasts isolated from artisanal fiore sardocheese. *Int.J. Food Microbiol.* 15, 95( 1):51-59.
- Farag, M.( 2002): Occurrence of Aflatoxins in fermented Milk. MVSc. Thesis. Zagazig Uni.
- Fardos, M. Bokhari( 2001): Mycobiota associated with food stuffes commodities spread in Jeddah ( Saudia Arabia) with special reference to *Asperigallus Flavus*. *Assiut Vet. Med. J.* 45( 90) : 94- 108
- Filtenborg, O.; frivvad, J.C.; Thrane, U.; Huis, I.N.T.; Veld, J.H.J.( 1996): Moulds in food spoilage. *Int.J. of food Microbiol.* 33( 1): 85- 102.
- Galvano, F.; Galolofaro, V. and Galvano, J. ( 1996): Occur- rence and stability of aflatoxin MI in milk and milk products: A world wide Review. *J. Food Prot.* 59( 10). 1079- 1090.
- Hamilton, P.P. ( 1975): Proof of mycotox cosis being a field problem and a simple method for their control. *Poult. Sci.* 54: 1706- 1708.
- Hassan, S.A.( 1999): Mycological studies on milk and some milk products. Ph. D. Vet. Thesis, Zagazig. Uni.
- Hassan, A.A. and Hammad, A. M.( 2001): Fungi and myco- toxins in milk powder and its product( soft cheese). *J. Egypt Vet. Med. Assoc.* 61, 2: 303- 309.
- Hendbrickse, R.G.( 1991): Clinical implications of food con- taminated by aflatoxins. *Annals Academy of Medicine* 20(1): 84- 90.
- Hussein, H.S. and Brasel, J.M.( 2001): Toxicity, metabo- lism and impact of mycotoxins on human and animals toxicology.. *J Food Microbiology.* 19, 69 (1-2): 141- 146.
- Ismail, M.A. and Saad, N.M.( 1995): Studies on the myco- logical quality of milk powder. *Assiut Vet . Med. J.* 32: 64, 173- 185.
- Kaya, S., Pirincci, I. and Bilgili, A. ( 1998): " Veteriner He- kimliginde Toksikoloji". *Medisan Yayin Ser isi*, 35, An- kara.
- Koneman, E.W. and Roberts, G.D. ( 1985): *Practical Labor- atory Mycology*. 3rd ed. Williams and Wilkins, Balti- more, London
- Kure, C.F., Skaar, I and Brendehaug, J.( 2004): Mould con- tamination in production of semi-hard cheese. *Int. J. Food Microbiology.* 15, 93: 41- 49.
- Li, S., Murquardt, R.R and Abramson, D. ( 2000): Immuno- chemical detection of moulds: A review. *J. Food Prot.* 63: 281- 291.
- Looder, J and Kreger Van R. J., N.J. W.( 1970): *The yeasts, A taxonomic study* 2nd North Holland Publishing Co., Amsterdam.
- Lund, F., Filtenborg, O. and frivvad, J.C( 1995): Associat- ed mycoflora of cheese. *Food Microbiol.* 12( 2); 173-

- Marth, E.H and Steele, J. L. ( 2001): Applied Dairy Microbiology 2nd ed., Marcal Dekker, Inc., New York.
- Martins, M.L and Martins, H. M.( 2004):Aflatoxin M1 in yoghurt in Portugal. *Int. J. Food Microbiol.* 15,91( 3): 315-7.
- Manal,M.A., Hanaa, M.S. and EL- Sayed, S. E( 2006): Aflatoxin residues in some dairy products with special reference to Aflatoxin M1. *J. Egypt. Soc. Toxicol.* 34, 27-36.
- Manetta, A.C, Digiuessel., Giam marco,M., Fusaro,I., Simonella, A., Gramenzi,A and Formigon, A. (2005):HPLC with post column derivatisation and Fluorescence detection for sensitive determination. *J. Chromatogr.A.* 12, 1083( 1-2): 219-22.
- Minervini, F.; Montagna, M.T.; Spilotron,G.; Monaci, L., Santacroce, M. P. and Visconti, A. ( 2001): Survey on mycoflora of cow and buffalo dairy products from southern Italy.*Int.J. Food Microbiology.* 19( 69) 141-146.
- Montagna,M. T.; Santacroce, M. P.; Spilotres, G.; Napoli,C.; Minervini, F.; Papa, A. and Dragnoi, I. ( 2004): Investigation of fungal contamination in cheese in southern Italy. *Mycopathologia* . 158( 2); 245- 249.
- Muir, D.D and , Banks, J.M.( 2000): Milk and Milk products in the stability and shelf life of food. BocaRaton. FL. 197- 219
- Munimbazi, C. and Bullerman,L.B.( 1996): Moulds and mycotoxins in foods from Burundi.*J.Food Prot.* 59( 8): 864.
- Nielsen, M.S.; Frisvad, J.C. and Nielsen, P. V.( 1998): protection by fungal starter against growth and secondary metabolite production of fungal spoilers of cheese. *Int. Food Microbiology.* 42( 2): 91
- Oliveria, C.A.F; Germano, P.M.L., Bird, C. and Pinto, C.A( 1997): Immuno- ochemical assessments of aflatoxinsM1 in milk powder consumed by infantism. Saopaulo, Brasil, *Food Add. Cont.* 14: 1, 17- 10.
- Park, D.L. ( 2002):Effect of processing on aflatoxin. *Adv. Exp. Med. Biol.* 504: 173-179.
- Pestka,J.( 1995): Fungal toxins in raw and fermented meats. Campbell- Platt, and P.E. Cook( ed.), Blackie Academic and Professional London. New York. Tokyo.Madras.
- Ray, B.( 1996) : Fundamental Food Microbiology. CRC. Press, Inc., Tokyo, New york.
- Salwa, N.A( 1999): Studies on mycotoxins in milk and some dairy products . PhD Vet. Thesis Cairo Uni.
- Sampyo, F.; Belda, V.; FranCo, A.C.; Fernander, Q.; Rodrigues, O.J. L. and Cepada, S.A. ( 1995): Distribution of fungal genera in cheese and dairies.Sensitivity to potassium sorbate and natamcin. *Arch. für Lebensmittel Hygiene* 46( 3): 62.
- Saras, I.; Piussi, D. ; Aquill, V.and Techini,M.L.( 1996): The behaviour of yeast population in cheese packaged under various condition. *J. Food Prot.* 59( 5): 541- 544.
- Saudi, A.M.; EL- Essawy, H.A. and Hafez, N.M. ( 1989): Aspects on the microbiological quality of yoghurt plastic containers. *Vet.Med. J. Giza.* 37: 3, 397- 406.
- Zein El-din, M. and EL-Sawah, M.( 1997): Evaluation of yoghurt manufactured and consumed in Mansoura City.*Agric. Sci. Mansoura Uni. J.* 22, 12: 4427- 4442.

## مدي تواجد الفطريات والخمائر في بعض منتجات الالبان المباعة بمحافظه المنوفية

نهلة احمد ابو الروس<sup>١</sup> و فلوراج محمود راضي<sup>٢</sup>  
١- قسم المراقبة المراقبة الصحة علي الاغذية ٢- قسم الفطريات  
معهد بحوث صحة الحيوان- شبين الكوم

أجرى هذا البحث علي مائة عينة من اللبن البودرة و الزبادي و الجبن الشيدر و الجبن الرومي (٢٥) عينه من كل منتج تم تجميعها من محلات بيع الالبان و السوبر ماركت في محافظة المنوفية لمعرفة مدي تواجد الفطريات و الا فلا توكسينات بها. وقد اوضحت النتائج ان العفن و الخمائر قد تواجدت في كل عينات اللبن البودرة و الزبادي و الجبن الشيدر و الرومي بمتوسط قدره  $1.0 \times 8.3$  و  $1.0 \times 9.6$  و  $1.0 \times 6.9$  و  $1.0 \times 5.3$  و  $1.0 \times 4.1$  علي التوالي بالنسبة للعفن و كانت بمتوسط قدره  $1.0 \times 5.0$  و  $1.0 \times 6.9$  و  $1.0 \times 5.3$  و  $1.0 \times 4.1$  علي التوالي بالنسبة للعفن. وكانت بمتوسط قدره  $1.0 \times 5.0$  و  $1.0 \times 6.9$  و  $1.0 \times 5.3$  و  $1.0 \times 4.1$  علي التوالي بالنسبة للخمائر. ولقد أمكن عزل أنواع عديدة من العفن تمثلت في جنس السبرجلس و جنس الألترناريا و جنس البنسليوم و جنس الكلاوسبوريم و جنس الميوكر و جنس الريزوس. أما الخمائر فتمثلت في جنس الكانديدا و جنس الرودرتوريل و جنس السكر و مسيز و جنس التوريولوبوسس.

ومن ناحية اخرى أثبتت التحليلات الكروماتوجرافية بالفصل علي الألواح الرقيقة بالكشف علي الأفلاتوكسينات عن تواجد الأفلاتوكسين م ١ في عينات لبن البودرة و الزبادي و الجبن الشيدر و الجبن الرومي بنسب ١٦% و ١٦% و ٣٢% علي التوالي. اما بالنسبة الي الأفلاتوكسين ب ١ فقد تواجد في عينات لبن البودرة و الجبن الشيدر و الجبن الرومي بنسب ١٢% و ٥٢% و ٤٨% علي التوالي. اما بالنسبة الي الأفلاتوكسين ب ٢ فقد تواجد في عينات لبن البودرة و الجبن الشيدر و الجبن الرومي بنسب ١٢% و ٢٤% و ١٢% علي التوالي.

وقد تم استنتاج ان تغذية الحيوانات بعلائق خالية من السموم الفطرية وكذلك استخدام لبن عالي الجودة و اتباع الأساليب الصحية أثناء الصناعة و التعبئة و التوزيع و التبريد الجيد كلها اسباب اساسية للحد من التلوث بالسموم الفطرية و الحصول علي منتج عالي الجودة.