

EFFECT OF SOME ANTIOXIDANTS ON ACROSIN AMIDASE ACTIVITY OF CHILLED STALLION SPERMATOZOA

Waheed M.M.* and El-Bahr S. M. **

Dept. of Clinical Studies* and Physiology, Biochemistry and Pharmacology**, College of Veterinary Medicine and Animal Resources, King Faisal University, Kingdom of Saudi Arabia.

Received: 13. 11. 2008

Accepted: 17. 12. 2008

SUMMARY

The effect of some antioxidants on acrosin amidase activity of chilled stallion spermatozoa was investigated in this study. Nine ejaculates were collected from six Arabian horses. A comparison between four antioxidants namely, sodium pyruvate (0.5 mg/ml), sodium thiosulfate (STS, 1.0 mg/ml), bovine serum albumin (BSA, 5.0 mg/ml), zinc chloride (0.15 mg/ml) and a mixture of them was studied in a chemically-defined stallion semen extender (Tris-egg yolk) at 5°C. The comparison was based on sperm viability, acrosin amidase activity and changes in the levels of extracellular alanine aminotransferase (ALT). Results of the present experiments revealed that sodium pyruvate and the mixture of antioxidants were most effective for improving viability and acrosin amidase activity of stallion spermatozoa. Low

values of ALT in the extracellular medium were coincided with high values of acrosin amidase activity of equine stallion spermatozoa during storage at 5°C.

Keyword: Stallion, Semen, Acrosine amidase, Antioxidants

INTRODUCTION

Preservation of liquid semen at 5°C is an important technique in the breeding management of horses (Ball et al., 2001). Stallion spermatozoa have a dearth of endogenous antioxidants (by removing most seminal plasma) to protect them against reactive oxygen species (ROS) that may be present during hypothermic storage of semen (Ball et al., 2000). There is an evidence that oxidative damage of spermatozoa by ROS impairs stallion sperm function in terms of declined mo-

tility and fertility (Ball et al., 2001). Sodium pyruvate is well known as an antioxidant and has protective effects against hydrogen peroxide (Giandomenico et al., 1997). Furthermore, STS has been shown to be an antioxidant and calcium solubilizer, it produces its effect via increasing the production of hydrogen sulfide (Sen et al., 2008). Concerning BSA, it acts as a multifaceted antioxidant that binds copper and other metals and keeps them from participating in oxidation reactions (Gum et al., 2004; Bonoli-Carbognin et al., 2008). Also, BSA binds fatty acids and protects them from oxidation (Gum et al., 2004). The ability of zinc to retard oxidative processes has been recognized for many years (Powell, 2000). Zinc acts as antioxidant through protection of protein sulfhydryls or reduction of OH formation from H₂O₂ through the antagonism of redox-active transition metals, such as iron and copper (Powell, 2000). The supplement of extenders with antioxidants could induce beneficial effect on viability of chilled stallion spermatozoa (Wahed and Khalifa, 2004). Acrocin, a sperm-specific acrosomal proteinase, has an essential role in the fertilization process (Polankoski and Zaneveld, 1977; Kennedy et al., 1989). Low levels of acrosin in human semen appear to be associated with subfertility and infertility (Kennedy et al., 1989). The enzyme is localized in the rostral portion of the acrosome, but is absent from the equatorial segment (Garner et al., 1977). The total acrosin activity of ejaculates can not be predicted from standard semen parameters such as

sperm concentration, motility and morphology (Goodpasture et al., 1987).

The main objective of the present study was to evaluate the effect of some antioxidants on acrosin amidase activity in stallion spermatozoa during storage at 5°C.

MATERIALS AND METHODS

Reagents

Benzamidine hydrochloride, N-α-benzoyl-DL-arginine p-nitroanilide hydrochloride (BAPNA), dimethylsulfoxide (DMSO), HEPES (N-2-hydroxyethylpiperazine-N'-2-ethanesulfonic acid), and Triton X-100 were obtained from Sigma Chemical Company (St. Louis, MO).

Solutions

Solution A. Detergent Buffer: 0.01% Triton X-100 in 0.055 M HEPES, 0.055 M NaCl at 8.0. The solution is prepared by dissolving 1.31 g of HEPES, 0.32 g of NaCl and 1 ml of a 1% Triton X-100 stock solution (1 ml Triton in 99 ml water) in 95 ml of distilled, deionized water. The solution is adjusted to pH 8.0 (with 1 N NaOH as required) and then adjusted to a final volume of 100 ml with distilled, deionized water. The detergent buffer is stable for three days but can be stored for extended periods in the refrigerator after addition of sodium azide (0.1%, or 100 mg added to 100 ml) as preservative.

Solution B. Benzamidine: 500 mM in water. The Solution is prepared by dissolving 87.3 g benzamidine-HCl in 1 liter distilled, deionized water. The solution can be stored in the refrigerator without any additives for at least two weeks.

Solution C. Substrate: 23 mM BAPNA in DMSO. The solution is prepared by dissolving 25 mg N- α -benzoyl -DL-arginine para-nitroanilide-HCl (BAPNA) in 2.5 ml dimethylsulfoxide (DMSO). The solution should be prepared fresh on the day of the assay. Complete dissolution of the BAPNA in DMSO requires about 5 to 10 minutes.

Solution D. Substrate-detergent mixture: 22.5 ml of the detergent buffer (solution A) is mixed thoroughly with 2.5 ml of the BAPNA/DMSO substrate solution (Solution C) in a 50-ml Erlenmeyer flask. This solution should be prepared while the Ficoll centrifugation step is taking place (see step 4 of the Assay Procedure).

Animals and ejaculates:

In this study, six healthy Arabian stallions (9 - 22 years old) belonging to three Arabian horse farms (Al-Bushaier Mydod, Haleim Shah and Al-Hashem farms), Al-Ahsa, Kingdom of Saudi Arabia, were used. These stallions have been used as sires in the regular breeding program of their farms.

Nine ejaculates were collected from the stallions using the CSU model artificial vagina. Immedi-

ately after collection, semen samples were transferred to the laboratory and evaluated by the conventional method.

Experimental design

In this investigation, comparison between four antioxidants namely, sodium pyruvate (0.5 mg/ml), STS (1.0 mg/ml), BSA (5.0 mg/ml), zinc chloride (0.15 mg/ml) and their respective mixture were done. The doses were used according to Waheed and Khalifa (2004). The comparison was based on evaluation of sperm viability, acrosin amidase activity in spermatozoa. In addition, the study was extended to determine the changes in the extracellular levels of ALT.

Experimental procedure

Following initial evaluation, each semen sample was prepared for preservation in Tris-egg yolk extender (Samper et al., 1988) at 5°C for 72 hours after being supplemented with the above mentioned antioxidants (Waheed and Khalifa, 2004). Briefly, 12 ml of each semen sample were divided into six aliquots (each one 2 ml). These aliquots were diluted with Tris-antioxidants free extender (Samper et al., 1988) 1:1 (v/v) and left for 5 minutes at 30°C. Samples were centrifuged at 1000 Xg for 5 minutes. Most of the supernatant was decanted and sperm pellets were resuspended in 2 ml of Tris-egg yolk extender supplemented with the antioxidants and reconstituted gently. The diluted samples were cooled slowly to 5°C (0.05-0.1°C/min) in the refrigerator and stored

for 72 hours. Sperm progressive motility was assessed after dilution (0 hour) as well as after 6 and 72 hours of incubation period and viability indices were computed according to Milovanov et al. (1964).

Aliquot of one ml semen was removed after dilution and after 72 hours of incubation at 5°C. These aliquots were centrifuged (1000 Xg) and the supernatants were stored at - 20°C pending analysis. ALT (IU/L) was assayed spectrophotometrically under the conditions specified by the commercial kit systems (Reitman and Frankle, 1957).

Sperm pellets, following the removal of the supernatants, were stored at - 20°C until assay to evaluate the acrosin amidase activity (Kennedy et al., 1989). The assay procedure described in steps as follow:

1. The ejaculate (of known sperm concentration) is allowed to liquefy completely.
2. For each ejaculate, a volume is calculated that it will contain between 2 to 10 10^6 spermatozoa.
3. One control and 5 tests (contain each antioxidant) are preferably run simultaneously for each ejaculate. The control and testes are incubated in duplicates (0 and 72 hours).
4. 100 μ l of the benzamidine solution (solution B) is immediately added to the control tube and the contents of the tube are mixed thoroughly by vortexing.
5. To each tube, including the control, 1 ml of the substrate-detergent mixture (solution D) is added and the contents of the tube are mixed thoroughly. Save at least 1 ml of solution E for step 9.
6. The tubes are incubated at 22 to 24°C for exactly three hours after addition of the assay solution. It is optimal to mix the contents of the tubes once every hour during the incubation period by vortexing.
7. After three hours of incubation, 100 μ l of the benzamidine solution (solution B) is added to all tubes except the control.
8. All tubes are centrifuged at 1000 Xg for 30 minutes and the supernatant solutions are collected separately.
9. A spectrophotometer capable of holding 1-ml cuvettes is adjusted so that the substrate-detergent mixture (solution D) has an absorbance reading of 0.0 at 410 nm. Subsequently, the absorbance of each supernatant solution is recorded at 410 nm. The control solution should show no or almost no absorption.

Calculations

One IU of acrosin activity is defined as the amount of enzyme that hydrolyzes 1 μ mol BAPNA/min at 23°C. To obtain whole numbers, the acrosin activity is expressed in μ IU/ 10^6 spermatozoa. The activity is calculated simply by the following formula according to Kennedy et al. (1989).

$$\mu\text{U acrosin}/10^6 \text{ sperm} = \frac{[(\text{mean ODtest}) - \text{OD control}]}{1485 \times \text{numbers of sperm in mil-lions}} \times 10^6$$

Statistical analysis

Analysis of data was performed using a commercial software Statistica for Windows (1993).

RESULTS

The effect of antioxidants on viability and acrosin amidase activity of cold-stored stallion semen is declared in Table 1. Sodium pyruvate, BSA, zinc

chloride and their mixture exerted a beneficial effect ($P < 0.05$) on viability (57.41 (6.42, 60.22 (5.88, 63.67 (6.28 and 64.94 (4.83, respectively) of stallion semen stored at 5°C. Furthermore, best results of acrosin amidase activity in stallion spermatozoa were obtained when sodium pyruvate and the respective antioxidants mixture, were added to the Tris - egg yolk extender (2.42 (0.54 and 1.74 (0.28 uIU/10⁶ sperm, respectively (Table 1).

Table (1): Effect of different antioxidants on motility, viability and acrosin amidase activity (uIU/10⁶ sperm) of chilled stallion semen (mean ± SEM).

Treatments	Sperm motility (%) during incubation at 5 °C			Viability indices	Acrosin activity
	0 hour	6 hours	72 hours		
Control	72.48 ^a ± 3.69	67.48 ^a ± 3.31	38.39 ^a ± 5.31	51.59 ^a ± 5.01	1.43 ^{ac} ± 0.39
Sodium pyruvate	75.35 ^b ± 4.14	71.46 ^b ± 4.70	41.46 ^{bc} ± 6.90	57.41 ^b ± 6.42	2.42 ^a ± 0.54
Sodium thiosulfate	70.91 ^{ab} ± 3.14	66.46 ^a ± 3.38	39.24 ^{ab} ± 6.34	54.53 ^{ab} ± 5.87	1.35 ^{ab} ± 0.29
BSA	78.69 ^c ± 3.49	74.24 ^{bc} ± 4.43	45.35 ^{cd} ± 6.57	60.22 ^c ± 5.88	1.18 ^{bc} ± 0.32
Zinc chloride	78.69 ^{cd} ± 4.67	74.24 ^{bc} ± 5.21	50.35 ^d ± 6.57	63.67 ^{cd} ± 6.28	1.35 ^{ab} ± 0.43
Respective mixture	83.84 ^e ± 2.67	81.57 ^e ± 3.09	50.20 ^d ± 5.67	64.94 ^{de} ± 4.83	1.74 ^b ± 0.28

Means with dissimilar superscripts in the same column are significantly different at $P < 0.05$

As shown in Table 2, the rate of increase of ALT in the extra-cellular medium was significantly higher for controls (88.63%) in comparison to the use of antioxidants during storage of stallion semen at 5°C. Also, the lesser values of the rate of

ALT increase in the extra-cellular medium were coincided by higher values of acrosin amidase activity in the stallion spermatozoa after 72 hours at 5°C (Table 2).

Table (2): The rate of increase of ALT (IU/L) in the extracellular medium and acrosin amidase activity in spermatozoa (uIU/10⁶ sperm) during storage of stallion semen at 5°C (mean ± SEM).

Treatments	ALT after dilution	ALT after 72 hours at 5°C	Rate of ALT increase (%)	Acrosin activity
Control	12.23 ^a ± 0.93	23.07 ^a ± 0.40	88.63**	1.43 ^{ac} ± 0.39
Sodium pyruvate	20.25 ^b ± 1.65	29.51 ^b ± 1.13	45.73**	2.42 ^a ± 0.54
Sodium thiosulfate	18.08 ^{ab} ± 2.08	27.87 ^{ab} ± 2.52	54.14***	1.35 ^{ab} ± 0.29
BSA	15.18 ^{bc} ± 0.38	25.50 ^{ab} ± 0.93	67.98***	1.18 ^{bc} ± 0.32
Zinc chloride	14.31 ^{ac} ± 0.76	22.67 ^a ± 0.74	58.42***	1.35 ^{bc} ± 0.43
Respective mixture	18.66 ^{bc} ± 1.49	27.71 ^{ab} ± 1.98	48.50**	1.74 ^b ± 0.28

Means with dissimilar superscripts in the same column are significantly different at P<0.05

** p<0.01

*** p<0.001

Although, the correlation coefficient between the rate of ALT increase in the extra-cellular medium and acrosin amidase activity in spermatozoa was non-significant, there was a tendency of inverse correlation (r = - 0.55).

DISCUSSION

Mammalian sperm contain particularly high con-

centrations of polyunsaturated fatty acids in their plasma membranes (Jones et al, 1979), making them susceptible to damage caused by lipid peroxidation (Jones and Mann, 1977). Lipid peroxidation of spermatozoal membrane lipids has been reported to be the major cause for the loss of sperm motility and fertilizing ability in human spermatozoa incubated for long periods of time during in vitro fertilization procedures (Aitken et

al., 1989; Gomez and Aitken, 1996), by altering membrane fluidity and functional integrity (Aitken, 1995) as a result of ROS accumulation. Although cooling spermatozoa to 5°C reduces the rate of lipid peroxidation, it is likely that lipid peroxidation plays a significant role in the reduced fertility seen for stallion spermatozoa stored for long periods of time (Pickett et al., 1989). Hypotaurine, taurine, and glutathione are antioxidants found in stallion semen (Li, 1975; Bavister and Yanagimachi, 1977). Antioxidants are added to reduce ROS formation and lipid peroxidation, thereby increasing spermatozoal longevity (Aitken, 1995). Interestingly, till the time of the present study, there were no publications concerning the effect of antioxidants on acrosin amidase activity in stallion spermatozoa and for this reason the present study was performed.

Concerning the effect of antioxidants on sperm motility and viability, the present findings indicated that, antioxidants improved stallion sperm motility and viability during incubation at 5°C (Table 1). Similar results have been previously reported in human (De Lamirande and Gagnon, 1992; Alvarez and Storey, 1995; Gavella and Lipovac, 1998; Kouoh et al., 1999), rams (Ollero et al., 1998; Upreti et al., 1998), bulls (Bilodeau et al., 2002) and stallions (Bruemmert et al., 2002; Waheed and Khalifa, 2004). Pyruvate is a potent scavenger of H₂O₂ (Upreti et al., 1998) and its supplementation at a concentration of 5 mM (~0.55 mg ml⁻¹) to chilled-stored stallion se-

men resulted in a significant augmentation of sperm motility and ATP levels (Bruemmert et al., 2002).

Oxidation of thiols (SH) in sperm proteins by O₂- and H₂O₂ was found to be associated with inhibition of sperm motility and fertilizing ability (Mammoto et al., 1996). Supplementation of egg yolk extender with sodium thiosulfate (1mg m⁻¹), as a source of SH to sperm cells, led to a remarkable improvement of their motility and plasma membrane stability in buffalo semen (Khalifa, 2001). Luxuriant entry of Ca²⁺ into sperm cells was found to coincide with activation of membrane-bound phospholipases (Roldan, 1998) and peroxidative damage of their membranes (Megahed and Anwar, 1997). The stabilizing influence of zinc on sperm membranes was attributed to its ability to suppress lipid peroxidation via inhibition of phospholipases and to protect SH and poly unsaturated fatty acids in biomembranes from Fe²⁺- mediated oxidation (Andrews et al., 1994). Albumin is a crucial extracellular antioxidant (Alvarez and Storey, 1983) through its propensity to bind transition metal ions (Fe²⁺ and Cu¹⁺) in egg yolk extender and, thereby, to minimize formation of OH, the powerful initiator of lipid peroxidation cascade in sperm (Halliwell, 1988).

Regarding the effect of antioxidants on acrosin amidase activity, the current study revealed that, sodium pyruvate and the antioxidants mixture preserved the acrosin amidase activity in stallion

spermatozoa during liquid storage at 5°C in Tris-egg yolk extender (Table 1). On the same direction, Ball et al. (1997) found that addition of egg yolk to equine spermatozoa significantly increased acrosin amidase activity in the cryopreservation medium. Furthermore, if the spermatozoa are damaged by cold shock, the proacrosine undergoes activation within the inner acrosomal membrane and acrosin appears in the extract instead of proacrosin (Mann and Lutwak-Mann, 1981).

The superior effect of Pyruvate than the other antioxidants on acrosine amidase activity preservation (Table 1) perhaps attributed to that pyruvate is an effective energy substrate as it can freely move into the cytoplasm and the mitochondria of cells reducing the ROS at a rate higher than the other used antioxidants. However, with the exception of pyruvate, the antioxidants mixture performed better than the individual use of antioxidants at the level of acrosine amidase activity. Interestingly, pyruvate alone preserved the acrosine amidase activity better than that when it was mixed with the other antioxidants. This perhaps indicated some sort of antagonism between pyruvate and the rest of antioxidants used.

In view of our findings, antioxidants offered a better protection for stallion sperm during storage at 5°C than the antioxidant free extender (Table 2). This is confirmed by the increased rate of extra-cellular ALT in the antioxidant free extender (

Table 2). Pursel et al. (1968) reported that one of the consequences of acrosomal damage is the leakage of enzymes from the sperm. The leakage of ALT revealed a positive correlation between enzyme release and sperm cell integrity and acrosomal damage (Chauhan et al., 1993). Enzyme release has generally been recognized as an indicator of cellular injury whereby membranes become inactivated or destroyed resulting in the loss of cellular material (De Reuck and Knight, 1964). Positive correlation between enzyme release and acrosomal damage was reported with goat's sperm (Chauhan et al., 1993). Yousef and Zeitoun (1998) found that there were negative correlation coefficients between sperm motility on one side and ALT release on the other side. They reported that the activities of these enzymes could be used as an indicator of sperm integrity. Concerning the effect of antioxidants as decreasing the activity of ALT, similar results were reported previously (Waheed and Khalifa, 2004). Further, the results of this study provide clear evidence that antioxidants protect the stallion spermatozoa against damage during cold storage and subsequently keep the acrosin amidase activity with spermatozoa. This is elucidated by the significant ($p < 0.01$) decrease in the rate of extra-cellular ALT that coincided with high values of acrosin amidase activity in stallion spermatozoa after cold storage.

Accordingly, acrosin is an indicator of sperm activity (Zaneveld et al., 1996). The present study

concluded that acrosin activity of spermatozoa may be potentially used as a useful marker of semen quality .

REFERENCES

- Aitken, R.J. (1995): Free radicals, lipid peroxidation and sperm function. *Reprod. Fertil. Dev.* 7: 659-668.
- Aitken, R. J., Clarkson, J.S. and Fishel, S. (1989): Generation of reactive oxygen species, lipid peroxidation and human sperm function. *Biol. Reprod.* 41: 183- 187.
- Alvarez, J.G. and Storey, B.T. (1983): Taurine, hypotaurine, epinephrine and albumin inhibit lipid peroxidation in rabbit spermatozoa and protect against loss of motility. *Biol. Reprod.*, 29: 548 - 555.
- Alvarez, J.G. and Storey, B.T. (1995): Differential incorporation of fatty acids into and peroxidative loss of fatty acids from phospholipids of human spermatozoa. *Mol. Reprod. Dev.*, 42: 334 - 346.
- Andrews, J.C., Nolan, J.P., Hammerstedt, R.H. and Bavister, B.D. (1994): Role of zinc during hamster sperm capacitation. *Biol. Reprod.*, 51: 1238 - 1247.
- Ball, B.A., Fagnan, M.S., Dobrinski, I., (1997): Determination of acrosin amidase activity in equine spermatozoa. *Theriogenology*, 48 (7): 1191 - 1198.
- Ball, B.A., Gravance, C.G., Medina, V., Baumber, J. and Liu, I.K.M. (2000): Catalase activity in equine semen. *Am.J.Vet.Res.*, 61: 1026 - 1030.
- Ball, B.A., Medina, V., Gravance, C.G., and Baumber, J. (2001): Effect of antioxidants on preservation of motility, viability and acrosomal integrity of equine spermatozoa during storage at 5°C. *Theriogenology*, 56: 577 - 589.
- Bavister, B.D. and Yanagimachi, R. (1977): The effect of sperm extracts and energy sources on the motility and acrosome reaction of hamster spermatozoa. *Biol. Reprod.* 16: 228 - 237.
- Bilodeau, J.F., Blanchette, S., Cormier, N and Sirard, M.A. (2002): Reactive oxygen species - mediated loss of bovine sperm motility in egg yolk Tris extender: Protection by pyruvate, metal chelators and bovine liver or oviductal fluid catalase. *Theriogenology*, 57: 1105 - 1122.
- Bonoli-Carbognin, M., Cerretani, L., Bendini, A., Almajano, M.P. and Michael H. Gordon, M.H. (2008): Bovine Serum Albumin Produces a Synergistic Increase in the Antioxidant Activity of Virgin Olive Oil Phenolic Compounds in Oil-in-Water Emulsions. *J. Agric. Food Chem.*, 56 (16): 7076 - 7081.
- Bruemmert, J.E., Coy, R.C., Squires, E.L. and Graham, J.K. (2002): Effect of pyruvate on the function of stallion spermatozoa stored for up to 48 hours. *J. Anim. Sci.*, 80: 12 - 18.
- Chauhan, M.S., Kapila, R., Gandhi, K.K., Anand, S.R. (1993): Acrosome damage and enzyme leakage of goat spermatozoa during dilution, cooling and freezing. *Andrologia* 26: 21 - 26.
- De Lamirande, E. and Gagnon, C. (1992): Reactive oxygen species and human spermatozoa: 1. Effects on the motility of intact spermatozoa and sperm axonemes. *J. Androl.*, 13: 368 - 378.
- De Reuck, A.V.S., Knight, J. (1964): *Cellular Injury*, CIBA Foundation Symposium. Little Brown, Boston, MA, pp.68.
- Garner, D.L., Reamer, S.A., Johnson, L.A. and Lessley, B.A. (1977): Failure of immunoperoxidase staining to detect acrosin in the equatorial segment of spermatozoa. *J. Exp. Zool.*, 201: 309 - 315.

- Gavella, M. and Lipovac, V. (1998): In vitro effect of zinc on oxidative changes in human semen. *Andrologia*, 30: 317 - 323.
- Giandomenico, A.R., Cerniglia, G.E., Biaglow, J.E., Stevens, C.W. and Koch, C.J. (1997): The importance of sodium pyruvate in assessing damage produced by hydrogen peroxide. *Free Radic. Biol. Med.* 23 (3): 426 - 434.
- Gomez, E., and Aitken, J. (1996): Impact of in vitro fertilization culture media on peroxidative damage to human spermatozoa. *Fertil. Steril.* 65: 880- 882.
- Goodpasture, J.C., Zavos, P.N. and Zaneveld, L.J.D. (1987): Relationship of human sperm acrosin and proacrosin to semen parameters. II. Correlations. *J. Androl.*, 8: 267 - 271.
- Gum, E.T., Swanson, R.A., Alano, M., Liu, J., Hong, S., Weinstein, P.R. and Panter, S.S. (2004): Human Serum Albumin and its N-Terminal Tetrapeptide (DAHK) Block Oxidant-Induced Neuronal Death. *Stroke*, 35: 590 - 595.
- Halliwel, B. (1988): Albumin, an important extracellular antioxidant? *Biochem. Pharmacol.*, 37: 569 - 571.
- Jones, R., and Mann, T. (1977): Damage to ram spermatozoa by peroxidation of endogenous phospholipids. *J. Reprod. Fertil.* 50: 261 - 168.
- Jones, R., Mann, T. and Sherins, R. (1979): Peroxidative breakdown of phospholipids in human spermatozoa, spermicidal properties of fatty acid peroxides and protective action of seminal plasma. *Fertil. Steril.* 31: 531 - 537.
- Kennedy, W.P., Kaminski, J.M., Van der Van, H.H., Jeyendran, R.S., Reid, D.S., Blackwell, J., Bielfeld, P. and Zaneveld, L.J.D. (1989): A simple, clinical assay to evaluate the acrosin activity of human spermatozoa. *J. Androl.*, 10: 221 - 231.
- Khalifa, T.A.A. (2001): Effect of some antioxidants on viability of preserved buffalo and ram semen. Ph.D. Vet. Thesis, Cairo Uni., Egypt.
- Kouoh, F., Gressier, B., Luyckx, M., Brunet, C., Dine, T., Cazin, M. and Cazin, J.C. (1999): Antioxidant properties of albumin: effect on oxidative metabolism of human neutrophil granulocytes. *Farmacol.*, 54: 695 - 699.
- Li, T. (1975): The glutathione and thiol content of mammalian spermatozoa and seminal plasma. *Biol. Reprod.* 12: 641 - 646.
- Mammoto, A, Masumoto, N., Tahara, M., Ikebuchi, Y., Ohmichi, M., Tasaka K. and Miyake, A. (1996): Reactive oxygen species block sperm-egg fusion via oxidation of sperm sulfhydryl proteins in mice. *Biol. Reprod.*, 55: 1063 - 1068.
- Mann, T. and Lutwak-Mann, C. (1981): Biochemistry of spermatozoa: Chemical and functional correlations in ejaculated semen, andrological aspects. In *Male Reproductive Function and Semen*. T. Mann and C. Lutwak-Mann (ed.). Berlin Heidelberg New York, Springer-Verlag, pp. 195 - 268.
- Megahed, G.A. and Anwar, M.M. (1997): Effect of calcium channel blocker on quality of Egyptian buffalo liquid semen. *Assuit Vet. Med. J.*, 38: 142 - 152.
- Milovanov, V.K., Trubkin, G.D., Chubenko, N.S., Tsvetkov, I.V., Erzín, Z.K. and Meschankin, A.B. (1964): Artificial insemination of livestock in the U.S.S.R. *Israel Program For Scientific Translations*, Jerusalem, pp. 102 - 104.
- Ollero, M., Perez, P.R., Blanco, M.T. and Perez, C.J.A. (1998): Improvement of ram sperm cryopreservation protocols assessed by sperm quality parameters and heterogeneity analysis. *Cryobiology*, 37: 1 - 6.

- Pickett, B.W., Squires, E.L., McKinnon, R.O., Shideler, R.K. and Voss, J.L. (1989): Management of the Mare for Maximum Repro-ductive Efficiency. Colorado State University Animal Reproduc-tion Laboratory Bulletin 06:80, Ft. Collins.
- Polankoski, K.L. and Zaneveld, L.J.D. (1977): Biochemical examination of the human ejaculate. In Techniques of Human Andrology. E.S.E. Hafez (ed.). Elsevier/North-Holland Biomedical Press, pp. 265 - 286.
- Powell, S.R. (2000): The antioxidant properties of zinc. *J.Nutr. Suppl.* 130: 1447S - 1454S.
- Pursel, V.G., Johnson, L.A., Gerrits, E.J. (1968): Glutamic oxaloacetic transaminase and lactic dehydrogenase activities of boar spermatozoa and seminal plasma. *J. Reprod. Fertil.* 18: 176 - 177.
- Reitman, S. and Frankel, S.A. (1957): Colorimetric method for the determination of serum glutamic oxaloacetic and glutamic pyruvic transaminases. *Am. J. Clinic. Pathol.* 28: 56 - 63.
- Roldan, E.R.S., (1998): Role of phospholipases during sperm acrosomal exocytosis. *Front. Biosci.*, 3: 1109 - 1119.
- Samper, J.C., Loseth, K.J. and Crabo, B.G. (1988): Evaluation of horse spermatozoa with sephadex filtration using three extenders and three diluents. *Proc. 11th Int. Congr. Anim. Reprod & A.I.*, Dublin, Ireland, Vol. 1, pp. 295.
- Sen, U., Vacek, T.P., Hughes, W.M., Kumar, M., Moshal, K.S., Tyagi, N., Metreveli, N., Hayden, M.R. and Tyagi, S.C. (2008): Cardioprotective Role of Sodium Thio-sulfate on Chronic Heart Failure by Modulating Endog-enous H₂S Generation. *Inter. J. Expt. Clin.Pharma.*, Vol. 82, No 3: 201 - 213.
- Statistica for Windows (1993): Release 4.5 A, statsoft, Inc.
- Upreti, G.C., Jensen, K., Munday, R., Duganzich, D.M., Vishwanath, R. and Smith, J.F. (1998): Studies on aromatic amino acid oxidase activity in ram spermatozoa: role of pyruvate as an antioxidant. *Anim. Reprod. Sci.*, 51: 275 - 287.
- Waheed, M.M. and Khalifa, T.A.A. (2004): Impact of anti-oxidants on viability of chilled and cryopreserved stallion spermatozoa. *Vet. Med. J., Giza.* 52 (4): 613- 625.
- Yousef, M.I. and Zeitoun, M.M. (1998): Bovine ovarian follicular fluids modulate the release of transaminases, acrosome reaction and motility of the rabbits sperm in vitro. *Alexandria J. Agric. Res.* 43: 17 - 26.
- Zaneveld, L.J.D. Jeyendran, R.S., Vermeiden, J.P.W. and Lens, J.W. (1996): Sperm enzymes for diagnostic purposes. In *Human Spermatozoa in Assisted Reproduction.* A.A. Acosta and T.F. Kruger (ed.). The Parthenon Publishing Group. New York, London, pp. 165- 176.

تأثير مضادات التأكسد على نشاط الأكروسين أميداز في حيامن

الحصان المبردة

مجدى محمد وحيد¹ و صبري محمد بهي البحر²

قسم الدراسات الإكلينيكية¹ وقسم الفسيولوجيا والكيمياء العضوية والفارماكولوجيا² - كلية الطب البيطرى والثروة الحيوانية - جامعة الملك فيصل - المملكة العربية السعودية.

تم إجراء هذا البحث لدراسة تأثير مضادات التأكسد على نشاط إنزيم الأكروسين أميداز في حيامن الحصان أثناء حفظها مبردة. تم تجميع عدد 9 قنفات من 6 خيول عربية (فحول) تتراوح أعمارهم من 9 إلى 22 سنة. وتمت مقارنة أربعة أنواع من مضادات التأكسد هي بيروفات الصوديوم (0.5 مجم/ملي) وثيوسلفات الصوديوم (1.0 مجم/ملي) وألبومين المصل البقري (5.0 مجم/ملي) وكلوريد الزنك (0.15 مجم/ملي) ومزيج منهم عند إضافتهم إلى مخفف معروف كيميائياً في الخيل هو الترس (Tris) مع صفار البيض عند درجة 5°م. وكانت مقارنة مضادات التأكسد معتمدة على تعيين دليل حيوية الحيامن ونشاط إنزيم الأكروسين أميداز فى الحيامن وقياس معدلات الزيادة فى إنزيم الألانين أمينوترانسفيريز خارج الحيامن. وقد أظهرت نتائج هذه التجارب أن بيروفات الصوديوم ومزيج مضادات التأكسد لهما أفضل تأثير فى الحفاظ على حيوية الحيامن ونشاط إنزيم الأكروسين أميداز فى حيامن الحصان. كذلك سجلت النتائج أن انخفاض معدل الزيادة فى إنزيم الألانين أمينوترانسفيريز خارج الحيامن صاحبه ارتفاع معدل نشاط إنزيم الأكروسين أميداز فى حيامن الحصان عند حفظ السائل المنوى عند درجة 5°م.